

0040-4039(94)02464-2

A New Synthesis of (-)-Furodysin

Tse-Lok Ho,* Kwang-Yuan Lee


Department of Applied Chemistry, National Chiao Tung University, Hsinchu, Taiwan, ROC Department of Chemistry, National Taiwan University, Taipei, Taiwan, ROC

Abstract. A synthesis of furodysin from trans-limonene oxide involves transformation into the phenylearbamate of 2.8-menthadien-1-ol, allylic displacement with lithium di-(B-furylmethyl)cuprate, and cyclization.

A number of furanosesquiterpenes occur in marine invertebrates.¹ Interestingly, furodysin and furodysinin have been elaborated in both enantiomeric forms, e.g., the (+)-isomers by an Australian variety of sponges Dysidea herbacea² and the (-)-isomers by those collected in Fiji³ and by the Mediterranean species D. tupha.⁴ These metabolites show significant antiparasitic activities.

The synthesis of (-)-furodysin 1 and (-)-furodysinin 2 has been accomplished starting from (+)-9bromocamphor.⁵ and also their racemates obtained from a longer route.⁶ Here we wish to report a convenient approach to furodysin from trans-limonene oxide 3.

Reaction of 3 with dimethylamine (40% aqueous solution) in a sealed vessel at 150°C gave the amino alcohol 4^7 which was converted into the N-oxide with 30% hydrogen peroxide and then pyrolyzed to afford trans-p-2,8-menthadien-1-ol 5⁸ in an overall 44% yield. We planned to effect an S_N 2' displacement of the allylic alcohol with a furylmethyl substituent on the same side as the oxygen functionality which is cis to the isopropenyl group at C-4. Previous experience⁹ indicated that allylic carbamates undergo syn-selective $S_N 2^{1}$ reactions with cuprate reagents therefore the N-phenylcarbamate 6 was prepared (77% yield) by a DMAPcatalyzed reaction of 5 with phenyl isocyanate with the intention of using it in such a reaction. Toward this









NMe₂









7

5: B = H 6; R = CONHPh

end, the reaction of **6** with lithium di-(β -furylmethyl)cuprate¹⁰ furnished compound **7** (91% yield) whose spectral data is in total agreement with those reported.⁵ The final cyclization to culminate in a furodysin synthesis was effected following the established protocol⁴ of mercuration, trapping, and reduction.

Some Physical data: **5**: $[\alpha]_D$ +63.9° (c=0.325 CHCl₃). δ_H (300 MHz, CDCl₃) 1.29 (3H, s), 1.4-1.65 (4H, m), 1.73 (3H, s), 1.8-1.85 (1H, m), 2.65 (1H, m), 4.73 (1H, s), 4.77 (1H, s), 5.66 (1H, d, *J* 11 Hz), 5.68 (1H, d, *J* 11 Hz). δ_C (75 MHz, CDCl₃) 20.83, 24.85, 29.37, 36.68, 43.41, 67.45, 110.57, 132.13, 133.88, 148.13.

6: mp 57-59°C. $[\alpha]_D$ -55° (c=0.369 CHCl₃). M⁺ 271.1577. v 3375, 1685 cm⁻¹. δ_H 1.5-1.8 (3H, m), 1.62 (3H, s), 1.70 (3H, s), 2.18 (1H, m), 2.20 (1H, m), 2.68 (1H, m), 4.74 (2H, s), 5.72 (1H, dd, J 9, 2 Hz), 6.16 (1H, d, J 9 Hz), 6.43 (1H, br.), 6.98 (1H, m), 7.2-7.35 (4H, m). δ_C 20.56, 24.20, 26.13, 35.22, 43.38, 77.73, 110.74, 118.43, 122.92, 128.83, 130.61, 133.55, 138.22, 147.83, 152.70.

7: $[\alpha]_D + 300^\circ$ (c=0.092 CHCl₃). δ_H 1.62 (3H, s), 1.76 (3H, s), 4.70 (1H, s), 4.87 (1H, s), 5.38 (1H, m), 6.23 (1H, d, J 1.5 Hz), 7.16 (1H, d, J 0.9 Hz), 7.33 (1H, dd, J 1.5, 0.9 Hz). δ_C 22.46, 22.81, 23.54, 25.93, 30.76, 36.79, 43.79, 110.13, 111.50, 123.83, 124.93, 133.88, 139.41, 142.47, 147.66.

2: $[\alpha]_D -40^\circ$ (c=0.116 CHCl₃). δ_H 1.25 (3H, s), 1.27 (3H, s), 1.66 (3H, s), 2.19 (1H, m), 2.56 (2H, m), 5.60 (1H, m), 6.10 (1H, d, J 2 Hz), 7.23 (1H, d, J 2 Hz). δ_C 19.51, 23.21, 23.77, 27.29, 30.73, 31.58, 31.72, 34.51, 45.59, 109.78, 112.95, 126.39, 133.11, 140.40, 156.77.

Acknowledgment: We thank the National Science Council, ROC, for financial support.

References

- Faulkner, D. J. Nat. Prod. Rep., 1984, 7, 251, 551; 1986, 3, 1; 1987, 4, 539; 1988, 5, 613; 1990, 7, 267; 1991, 8, 97; 1992, 9, 323; 1993, 10, 497.
- 2. Kazlauskas, R.; Murphy, P. T.; Wells, R. J. Tetrahedron Lett., 1978, 4949, 4951.
- 3. Horton, P.; Inman, W.; Crews, P. J. Nat. Prod., 1990, 53, 143.
- 4. Guella, G.; Mancini, I.; Guerriero, A.; Pietra, F. Helv. Chim. Acta, 1985, 68, 1276.
- 5. Vaillancourt, V.; Agharahimi, M. R.; Sundram, U. N.; Richou, O.; Faulkner, D. J.; Albizati, K. F. J. Org. Chem. 1991, 56, 378.
- 6. Hirota, H.; M. Kitano, M.; K. Komatsubara, K.; Takahashi, T. Chem. Lett., 1987, 2079.
- 7. Kuczynski, H.; Zabza, A. Roczniki Chem. 1963, 37, 773.
- 8. Schenck, G. O.; Gollnick, K.; Buchwald, G.; Schroeter, S.; Ohloff, G. Liebigs Ann.Chem. 1964, 674, 93.
- Galina , C.; Ciattoni, P. G. J. Am. Chem. Soc., 1979, 101, 1035;
 Goering, H. L.; Kantner, S. S.; Tseng, C. C. J. Org. Chem. 1983, 48, 715.
- 10. Tanis, S. P.; Herrinton, P. M. J. Org. Chem. 1983, 48, 4572.

(Received in China 7 October 1994)