

# Azido-Neonicotinoids as Candidate Photoaffinity Probes for Insect Nicotinic Acetylcholine Receptors [1]

Peter Maienfisch<sup>a\*</sup>, Joerg Haettenschwiler<sup>a</sup>, Alfred Rindlisbacher<sup>b</sup>, Arnaud Decock<sup>c</sup>, Henning Wellmann<sup>c</sup>, and Hartmut Kayser<sup>c</sup>

**Abstract:** The neonicotinoids are the most successful chemical class of insecticides reaching sales of more than 630 Mio \$ in 2001, mainly due to the excellent market performance of imidacloprid and thiamethoxam. The insect nicotinic acetylcholine receptors (nAChRs) are the targets for these compounds, which are highly effective against a variety of sucking and chewing insects. Compared with the other neonicotinoid sales products, thiamethoxam binds in a different way, possibly to a different site of nAChRs in aphids. To gain further insight into the different modes of binding, a research program applying the photoaffinity labeling technique was started. A series of novel candidate photoaffinity probes containing a 5-azido-6-chloropyridin-3-ylmethyl group were prepared from 5-azido-6-chloropyridin-3-ylmethyl chloride, which was obtained in three steps from 6-chloropyridin-3-ylmethyl chloride. These probes showed good to excellent contact/feeding and systemic activity against *Myzus persicae*, however, they were at least 4- to 16-fold less effective against *Aphis craccivora*, *Nilaparvata lugens*, *Spodoptera littoralis*, and *Diabrotica balteata* than the neonicotinoid sales products. In general, the introduction of an azide group at C(5) of the 6-chloropyridin-3-ylmethyl substituent resulted in reduced potency as well as in a narrower pest spectrum. In competition binding assays with [<sup>3</sup>H]imidacloprid, analogues of imidacloprid, clothianidin, thiacloprid and thiamethoxam containing a 5-azido-6-chloropyridin-3-ylmethyl group showed high displacing potency with nAChRs from *Aphis* and *Myzus* ( $K_i$  values: 2 to 27 nM) suggesting that these compounds are valuable candidate photoaffinity probes. Taking into account the biological screening activity as well as the receptor binding potency, 1-(5-azido-6-chloropyridin-3-ylmethyl)-2-nitroimino-imidazolidine, N-(5-azido-6-chloropyridin-3-ylmethyl)-N'-methyl-N''-nitroguanidine and 3-(5-azido-6-chloropyridin-3-ylmethyl)-2-cyanoimino-thiazolidine were identified as the preferred candidate neonicotinoid photoaffinity probes to study the imidacloprid binding site.

**Keywords:** Azido-neonicotinoids · Candidate photoaffinity probes · Imidacloprid · Insecticidal activity · Insecticides · Neonicotinoids · Nicotinic acetylcholine receptors · Thiamethoxam

## 1. Introduction

An important milestone in the history of modern insect control is marked by the discovery of the neonicotinoids [2]. These compounds act on nicotinic acetylcholine

receptors (nAChRs) and are highly effective against a variety of sucking and chewing insects. As the first representative of this chemical class, imidacloprid (**1**) was introduced to the market in 1991, and since then, a series of analogues (compounds **2** to **7**) have been launched (Fig. 1) [3]. Today the neonicotinoids are the fastest growing chemical class of insecticides. This tremendous success is based on their unique chemical and biological properties, such as broad-spectrum insecticidal activity, low application rates, excellent systemic characteristics, favourable safety profile, and a new mode of action.

Our own research in this area resulted in the discovery of thiamethoxam (**5**) [4]. This compound is a second-generation neonicotinoid and belongs to the thianicotinyl

subclass. It was first synthesized in 1991 and is now developed worldwide for use in more than 100 crops. Thiamethoxam is marketed since 1998 under the trademarks Actara<sup>®</sup> for foliar and soil treatment and Cruiser<sup>®</sup> for seed treatment. In all these usages, thiamethoxam provides excellent control of a broad range of commercially important pests, such as aphids, whiteflies, thrips, rice hoppers, Colorado potato beetle, flea beetles, wireworms, leaf miners as well as some lepidopterous species [5]. Low use rates, flexible application methods, excellent efficacy, and the favourable safety profile make this new insecticide well suited for modern integrated pest management programs in many cropping systems.

All sales products **1** to **7** bind with high affinity to nicotinic acetylcholine receptors

\*Correspondence: Dr. P. Maienfisch<sup>a</sup>  
Tel.: + 41 61 323 66 47  
Fax: + 41 61 323 85 29  
E-Mail: peter.maienfisch@syngenta.com  
<sup>a</sup>Syngenta Crop Protection AG  
Optimisation Chemistry  
Schwarzwaldallee 215  
CH-4002 Basel  
<sup>b</sup>Syngenta Crop Protection AG  
Research Biology, Insect Control  
Schaffhauserstr. 101  
CH-4332 Stein  
<sup>c</sup>Syngenta Crop Protection AG  
Biochemistry  
Schwarzwaldallee 215  
CH-4002 Basel

[6][7]. To provide further evidence for our recent findings that thiamethoxam binds, compared to the other neonicotinoid sales products, in a different way, possibly to a different site of the receptor in aphids [7], we have started a research program applying the photoaffinity labeling technique to investigate neonicotinoid-receptor interactions and to identify the amino acids involved in neonicotinoid binding. This article summarises the current status relative to neonicotinoid photoaffinity probes and reports for the first time the synthesis, binding potency and biological activity of the novel candidate photoaffinity probes **13** to **16** (Fig. 5).

## 2. Photoaffinity Labeling Technique

Three major approaches are mainly used to study molecular recognition in receptor–ligand interactions [8a]:

- 1) Spectroscopic methods: X-ray crystallography, high-resolution NMR, *etc.*
- 2) Chemical methods: photoaffinity labeling, chemical affinity labeling, *etc.*
- 3) Genetic methods: site-directed mutagenesis, molecular chimera, *etc.*

Among these methods photoaffinity labeling is a very powerful tool for the identification of ligand-binding proteins in crude protein mixtures [8]. This approach uses photolabile derivatives of ligands (photoaffinity probes; see Fig. 2) for the covalent labeling of target proteins, for example enzymes or receptors. Upon absorption of UV-light, photoaffinity probes are converted into highly reactive intermediates, which can covalently bind to amino acid residues at the binding site of target proteins. Subsequently, the radiolabeled protein can be isolated, the labeled amino acids identified and thus the binding domain characterised.

Currently, aryl/heteroaryl azides and aryldiazirines are the photoreactive groups mainly used in photoaffinity labeling generating aryl/heteroaryl nitrenes and aryl carbenes, respectively, as reactive intermediates that can insert directly into most carbon-, nitrogen-, oxygen-, and sulfur-containing covalent bonds (Fig. 2).

## 3. Design of Neonicotinoid Photoaffinity Probes

As the first neonicotinoid photoaffinity probe, the 2-azido-5-<sup>125</sup>Iodobenzoyloxyethyl derivative **8** of an imidacloprid analogue (Fig. 3) has been described [9]. However, this probe is not ideal, because it is of limited hydrolytic stability and only effec-




Fig 1. Neonicotinoid sales products






Fig 2. Photoaffinity probes and examples of photoreactive groups used in photoaffinity labeling

tive with receptor proteins from the fly *Drosophila*, but not with those from other insects. Furthermore, the photoreactive group is in quite some distance from that part of the molecule essential for binding.

This is also the major disadvantage of the two recently published candidate photoaffinity probes **9** and **10**, which behaved as typical nicotinic agonists in cockroach nerve cord preparations [10].



Scheme 1. Synthesis of the key intermediate 5-azido-6-chloropyridin-3-ylmethyl chloride (**17**)

Scheme 2. Synthesis of candidate photoaffinity probes of type A

the neonicotinoid sales products **1**, **4**, **5** and **6** were obtained from competition binding assays with [ $^3\text{H}$ ]imidacloprid as a ligand that shares its binding characteristics with most other neonicotinoids, with the exception of thiamethoxam [6a][7][17]. Incubations were performed with  $100\ \mu\text{g}$  of membrane protein and  $6\ \text{nM}$  of the radioligand for 2 h at  $22^\circ\text{C}$ .

Our results, together with some data from the literature, are compiled in Table 2 as the inhibition constants  $K_i$ , which were calculated from the measured  $\text{IC}_{50}$  values

[18]. The imidacloprid analogue **11**, the clothianidin analogue **12**, the thiacloprid analogue **13** and the thiamethoxam analogue **14** possess high displacing potency with nAChRs from *Aphis* and *Myzus* ( $K_i$  values: 2 to  $27\ \text{nM}$ ) suggesting that these compounds are valuable candidate photoaffinity probes for the imidacloprid binding site. On the other hand, compounds **15** and **16** may be more suitable to study the unusual binding of thiamethoxam **5**.

In the meantime, Zhang *et al.* [15] prepared a series of further azidopyridinyl

neonicotinoids as candidate photoaffinity probes. This study demonstrated that 5-azido-6-chloropyridin-3-ylmethyl derivatives are clearly more potent than the corresponding 4-azido analogues. Compounds **35** and **38** and especially the 2-nitroimino-thiazoline **36** and the 2-nitroimino-imidazolidine **37** were found to be very potent in displacing [ $^3\text{H}$ ]imidacloprid from *Drosophila* nAChRs (Table 2, Fig. 6).

## 7. Conclusion

Candidate photoaffinity probes **11** to **16**, all containing a 5-azido-6-chloropyridin-3-ylmethyl group, were successfully prepared from 5-azido-6-chloropyridin-3-ylmethyl chloride, which was obtained in three steps from 6-chloropyridin-3-ylmethyl chloride. Taking into account the biological screening activity (Table 1) as well as the receptor binding potency (Table 2) we conclude that compounds **11**, **12**, and **13** are the preferred candidate neonicotinoid photoaffinity probes to investigate neonicotinoid-receptor interactions and to identify the amino acids involved in imidacloprid binding. Although only limited data are available compounds **36** and **37** may also be considered as valuable candidates.

Among these azido-neonicotinoids, a radiolabel is most easily introduced into the acyclic nitroguanidine derivative **12** and consequently [ $^3\text{H}$ ]-**12** (Fig. 5) was radiosynthesized with high specific activity as first azido-neonicotinoid photoaffinity probe. [ $^3\text{H}$ ]-**12** was found to bind to a single high-affinity site in *Myzus* that is competitively inhibited by imidacloprid [16].

Received: September 26, 2003

- [1] Presented in part at the 10th IUPAC International Congress of the Chemistry of Crop Protection, Basel, Switzerland, **2002**, Poster 3a.16 (book of abstracts Vol.1, p 234).
- [2] I. Yamamoto, J.E. Casida (Eds), 'Nicotinoid Insecticides and the Nicotinic Acetylcholine Receptor', Springer-Verlag, Tokyo, **1999**, 1–300; and literature cited therein.
- [3] a) P. Maienfisch, F. Brandl, W. Kobel, A. Rindlisbacher, R. Senn in: 'Nicotinoid Insecticides and the Nicotinic Acetylcholine Receptor', Eds. I. Yamamoto, J.E. Casida, Springer-Verlag, Tokyo, **1999**, 177–209; and literature cited therein. b) T. Wakita, K. Kinoshita, E. Yamada, N. Yasui, N. Kawahara, A. Naoi, M. Nakaya, K. Ebihara, H. Matsuno, H. Kodaka, *Pest Manag. Sci.* **2003**, *50*, 1016–1022.

Table 1. Biological activity of candidate photoaffinity probes **11** to **16** in comparison with the neonicotinoid standards **1, 4, 5**, and **6**

Comp.	A.c. <sup>a</sup>	A.c.	M.p. <sup>b</sup>	M.p.	N.l. <sup>c</sup>	S.l. <sup>d</sup>	D.b. <sup>e</sup>
	cf <sup>f</sup>	sy <sup>g</sup>	cf	sy	cf	cf	cf
<b>11</b>	++	++++	+++	+++++	-	++	+++
<b>12</b>	++	+++	+++	+++++	+	-	-
<b>13</b>	+	+++	++	+++	-	+	+++
<b>14</b>	-	-	+++	+++++	-	-	-
<b>15</b>	+	+++	++	+++++	+	-	-
<b>16</b>	-	-	+	+++++	-	+	-
<b>1</b>	+++	+++++	+++++	+++++	++++	++++	+++++
<b>4</b>	+++	++++	+++	+++++	+	++	+++++
<b>5</b>	+++	+++++	+++++	+++++	+++++	++++	+++++
<b>6</b>	++	+++++	++++	+++++	+++++	++++	+++++

Test insects: <sup>a</sup> *Aphis craccivora*, <sup>b</sup> *Myzus persicae*, <sup>c</sup> *Nilaparvata lugens*, <sup>d</sup> *Spodoptera littoralis*, <sup>e</sup> *Diabrotica balteata*.

Type of activity: <sup>f</sup> Contact/feeding activity, <sup>g</sup> systemic activity.

Rating system: - denotes < 80% activity at 200 ppm; + denotes ≥ 80% activity at 200 ppm; ++ denotes ≥ 80% activity at 50 ppm; +++ denotes ≥ 80% activity at 12 ppm; ++++ denotes ≥ 80% activity at 3 ppm; +++++ denotes ≥ 80% activity at 0.8 ppm or below. See [4b] for details of the assays.


Table 2. Competition binding potency of candidate photoaffinity probes of type **A** with [<sup>3</sup>H]imidacloprid ([<sup>3</sup>H]-**1**)

Comp.	K <sub>i</sub> [nM]			Reference
	A.c. <sup>a</sup>	M.p. <sup>b</sup>	D.m. <sup>c</sup>	
<b>11</b>	4.5 <sup>d</sup>	4.5 <sup>d</sup>	- <sup>e</sup>	This study
	-	1.0	15	Kagabu <i>et al.</i> [12]
	-	-	24	Zhang <i>et al.</i> [15]
<b>12</b>	22	27	-	This study
	-	4.5	13	Kagabu <i>et al.</i> [12]
	-	5.0	161	Tomizawa <i>et al.</i> [16]
<b>13</b>	2	2.5	-	This study
<b>14</b>	8	11	-	This study
<b>15</b>	125	160	-	This study
<b>16</b>	123	270	-	This study
<b>1</b>	6	3	-	This study
<b>4</b>	3	2	-	This study
<b>5<sup>f</sup></b>	2300	3200	-	Kayser <i>et al.</i> [17]
<b>6</b>	14	9	-	This study
<b>35</b>	-	-	13	Zhang <i>et al.</i> [15]
<b>36</b>	-	-	3.9	Zhang <i>et al.</i> [15]
<b>37</b>	-	-	0.72	Zhang <i>et al.</i> [15]
<b>38</b>	-	-	28	Zhang <i>et al.</i> [15]

<sup>a</sup> *Aphis craccivora*, <sup>b</sup> *Myzus persicae*, <sup>c</sup> *Drosophila melanogaster*

<sup>d</sup> See [6a] for assay description, <sup>e</sup> not tested

<sup>f</sup> Thiamethoxam (**5**) binds in a mode different from that of the other neonicotinoid sales products, *i.e.* it interferes in a non-competitive mode with [<sup>3</sup>H]imidacloprid (**1**) binding to receptor preparations from *A.c.* and *M.p.*, which explains the high K<sub>i</sub> values. The intrinsic receptor affinity of thiamethoxam, used as [<sup>3</sup>H]-**5**, is in the nanomolar range, comparable to that of imidacloprid (**1**).

Fig. 6. Structures of candidate photoaffinity probes of type **A** recently published [15]

- [4] a) P. Maienfisch, L. Gsell, A. Rindlisbacher, *Pestic. Sci.* **1999**, *55*, 351–355; b) P. Maienfisch, H. Huerlimann, A. Rindlisbacher, L. Gsell, H. Dettwiler, J. Haettenschwiler, E. Sieger, M. Walti, *Pest Manag. Sci.* **2001**, *57*, 165–176; c) P. Maienfisch, A. Rindlisbacher, H. Huerlimann, J. Haettenschwiler, A.K. Desai, V.S. Ekkundi, V.D. Gangan, *ACS Symposium Series*, **2002**, *800* (Synthesis and Chemistry of Agrochemicals VI), 219–230.
- [5] a) R. Senn, D. Hofer, T. Hoppe, M. Angst, P. Wyss, F. Brandl, P. Maienfisch, L. Zang, S. White, *Brighton Crop Protection Conference – Pests and Diseases*, **1998**, Vol. 1, 27–36; b) P. Maienfisch, M. Angst, F. Brandl, W. Fischer, D. Hofer, H. Kayser, W. Kobel, A. Rindlisbacher, R. Senn, A. Steinemann, H. Widmer, *Pest Manag. Sci.* **2001**, *57*, 906–913.
- [6] a) P. Wiesner, H. Kayser, *J. Biochem. Mol. Toxicol.* **2000**, *14*, 221–230; b) A. Zhang, H. Kayser, P. Maienfisch, J.E. Casida, *J. Neurochem.* **2000**, *75*, 1294–1303; and literature cited therein.
- [7] H. Kayser, C. Lee, H. Wellmann, 10th IUPAC International Congress of the Chemistry of Crop Protection, Basel, Switzerland, **2002**, Poster 3c.23 (book of abstracts Vol.1, p 305).
- [8] For recent reviews on photoaffinity labelling see: a) Y. Hatanaka, Y. Kanaoka, *Heterocycles*, **1998**, *47*, 625–632; b) F. Kotzyba-Hilbert, I. Kapfer, M. Goeldner, *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 1296–1312; c) H.J. Schafer, A. Schuhen, *Biol. Res.* **1996**, *29*, 31–46; d) J.E. Casida, G.B. Quistad, *Special Publication - Royal Society of Chemistry*, **1999**, 232 (Progress in Neuropharmacology and Neurotoxicology of Pesticides and Drugs), 34–45; e) C.A. Gartner, *Curr. Med. Chem.* **2003**, *10*, 671–689.
- [9] a) B. Latli, M. Tomizawa, J.E. Casida, *Bioconjug. Chem.* **1997**, *8*, 7–14; b) M. Tomizawa, J.E. Casida, *Neurosci. Lett.* **1997**, *237*, 61–4.
- [10] K. Matsuda, M. Ihara, K. Nishimura, D.B. Sattelle, K. Komai, *Biosci. Biotechn. Biochem.* **2001**, *65*, 1534–1541.
- [11] The work started at Ciba-Geigy, which became Novartis in 1996, and then Syngenta in 2000.
- [12] S. Kagabu, P. Maienfisch, A. Zhang, J. Granda-Minones, J. Haettenschwiler, H. Kayser, T. Maetzke, J.E. Casida, *J. Med. Chem.* **2000**, *43*, 5003–5009.
- [13] P. Maienfisch, H. Huerlimann, J. Haettenschwiler, *Tetrahedron Lett.* **2000**, *41*, 7187–7191.
- [14] A. Zhang, H. Kayser, P. Maienfisch, J.E. Casida, *J. Neurochem.* **2000**, *75*, 1294–1303.
- [15] N. Zhang, M. Tomizawa, J.E. Casida, *J. Med. Chem.* **2002**, *45*, 2832–2840.
- [16] M. Tomizawa, Z. Wen, H.L. Chin, H. Morimoto, H. Kayser, J.E. Casida, *J. Neurochem.* **2001**, *78*, 1359–1366.
- [17] H. Kayser, C. Lee, A. Decock, M. Baur, J. Haettenschwiler, P. Maienfisch (manuscript in preparation).
- [18] Y.-C. Cheng, W.H. Prusoff, *Biochem. Pharmacol.* **1973**, *22*, 3099–3108.