


# Organic & Biomolecular Chemistry

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: Y. Osumi, C. Liu and M. Szostak, *Org. Biomol. Chem.*, 2017, DOI: 10.1039/C7OB02269G.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



## Organic &amp; Biomolecular Chemistry

## COMMUNICATION

## N-Acylsuccinimides: Twist-Controlled, Acyl-Transfer Reagents in Suzuki-Miyaura Cross-Coupling by N–C Amide Bond Activation

Yuki Osumi,<sup>a</sup> Chengwei Liu<sup>a</sup> and Michal Szostak<sup>a,\*</sup>Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/


The palladium-catalyzed Suzuki-Miyaura cross-coupling of N-acylsuccinimides as versatile acyl-transfer reagents via selective amide N–C bond cleavage is reported. The method is user-friendly since it employs commercially-available, air-stable reagents and catalysts. The cross-coupling is enabled by half-twist of the amide bond in N-acylsuccinimides. These highly effective, crystalline acyl-transfer reagents present major advantages over perpendicularly twisted N-acylglutarimides, including low price of the succinimide activating ring, selective metal insertion under redox neutral conditions and high stability of the amide bond towards reaction conditions. Mechanistic studies indicate that oxidative addition is the rate limiting step in this widely applicable protocol.

Over the past two years, amides have emerged as important building blocks in transition-metal-catalyzed cross-coupling reactions.<sup>1,2</sup> Two general manifolds have been established, namely acyl- and decarbonylative cross-couplings,<sup>3,4</sup> wherein the facility of metal insertion is controlled by amide geometry,<sup>5</sup> electronics<sup>6</sup> and/or catalyst system<sup>7</sup> (Fig 1A). In particular, acyl-Suzuki-Miyaura cross-coupling has proved to be a popular tool for testing new amide-based reagents for selective metal insertion into the typically inert amide N–C bond ( $n_N \rightarrow \pi^*_{C=O}$  conjugation, 15–20 kcal/mol in planar amides).<sup>8</sup>

In this context, N-acylglutarimides, introduced by our group in 2015,<sup>9</sup> have been demonstrated as the most reactive amide derivatives in metal-catalyzed N–C activation to date,<sup>10</sup> owing to the perpendicular twist of the amide bond and electronic activation. A wide range of other preactivated amide-based reagents such as N-Boc-carbamates,<sup>11</sup> N-Ts-sulfonamides,<sup>12</sup> N,N-di-Boc<sub>2</sub> amides,<sup>13</sup> N-acyl-saccharins,<sup>14</sup> N-Ms-sulfonamides,<sup>15</sup> N-acyl-pyrroles<sup>16</sup> and N-Me-pyrimidines<sup>17</sup> have been successfully employed as cross-coupling partners in the acyl Suzuki reaction by N–C activation (Fig. 1B). Recently, we introduced N-acylsuccinimides as highly effective, bench-

stable, crystalline, amide-based acyl transfer reagents in Ni-catalyzed Negishi cross-coupling.<sup>18</sup> Our mechanistic studies indicated that the amide bond in N-acylsuccinimides is approximately half-twisted (Fig. 2).<sup>5a</sup> This allows for selective fine-tuning of metal insertion under redox-neutral conditions by amide bond geometry to afford acyl-metal intermediates of broad synthetic appeal.<sup>19</sup>

A. Amide bond cross-coupling



B. Amide-based precursors in Suzuki-Miyaura cross-coupling




Fig. 1 (A) Amide bond cross-coupling, (B) Amide-based precursors in Suzuki-Miyaura cross-coupling.




Fig. 2 Amide bond activation by twist: perpendicular glutarimides vs. half-twisted succinimides.

<sup>a</sup>Department of Chemistry, Rutgers University, 73 Warren Street, Newark, NJ 07102, United States. Fax: (+1)-973-353-1264; Tel: (+1)-973-353-532. E-mail: michal.szostak@rutgers.edu.

\*Electronic Supplementary Information (ESI) available: Experimental details and characterization data. See DOI: 10.1039/x0xx00000x

## COMMUNICATION

## Organic &amp; Biomolecular Chemistry


Herein, we disclose the Pd-catalyzed Suzuki-Miyaura cross-coupling of N-acylsuccinimides as versatile acyl-transfer reagents via selective amide N–C bond cleavage. The following key features render N-acylsuccinimides synthetically appealing in N–C cross-coupling reactions: (1) low price and wide-availability of succinimide (<\$0.05/gram); (2) high-bench stability; (3) N-acylsuccinimides are crystalline solids easily purified by crystallization; (4) higher stability of N-acylsuccinimides than N-glutarimide amides as a consequence of lower twist of the amide bond; and (5) the 5-membered activating ring is more sterically-accessible than the 6-membered glutarimide, while its cyclic nature prevents side reactions by the undesired cleavage of the  $\sigma$  N–C bond.<sup>20,21</sup>

During the course of this work, Zeng and co-workers reported a related process.<sup>22</sup> The catalytic system developed by our group uses air-stable reagents and is more catalytically reactive than the Zeng's system. Furthermore, we report mechanistic studies that shed light on the catalytic activation of N-acylsuccinimides in the Suzuki-Miyaura cross-coupling.

The reaction was optimized using commercially-available benzoylsuccinimide (Table 1). Our initial efforts centered on the catalytic system using H<sub>3</sub>BO<sub>3</sub> to promote switchable O-/O-coordination of the amide bond in the twisted N-acylsuccinimide (Table 1, entries 1-2).<sup>5a</sup> Encouragingly, we found that the reaction of **1a** with 4-tolylboronic acid in the presence of H<sub>3</sub>BO<sub>3</sub> and K<sub>2</sub>CO<sub>3</sub> as a base delivered the desired acylation product in quantitative yield (entry 1). Furthermore, appreciable conversion was obtained at ambient conditions, attesting to the high catalytic activity of this catalytic system (entry 3). Nevertheless, based on our previous experience in acyl Suzuki cross-couplings, we sought to develop a more general catalyst system in the absence of acidic additives. Pleasingly, we found that a catalyst system consisting of Pd(OAc)<sub>2</sub>, PCy<sub>3</sub>HBF<sub>4</sub> and Na<sub>2</sub>CO<sub>3</sub> in dioxane provided superior efficiency (entry 4). Higher temperature led to substantially higher yields (entry 5). This observation suggests that the reaction proceeds via direct metal insertion (cf. switchable activation of the succinimide under acidic conditions).<sup>23</sup> The use of Na<sub>2</sub>CO<sub>3</sub> as a base and dioxane as a solvent is superior to other base/solvent combinations (entries 6-7). Furthermore, evaluation of other ligands, including less nucleophilic PPh<sub>3</sub> and more sterically-demanding Pt-Bu<sub>3</sub>HBF<sub>4</sub> proved significantly less effective (entries 8-10). Collectively, these results confirm that PCy<sub>3</sub> serves as a privileged ligand in Pd-PR<sub>3</sub>-catalyzed Suzuki-Miyaura acyl cross-coupling of amide-based reagents by N–C activation. Importantly, our user-friendly method uses bench-stable PCy<sub>3</sub>HBF<sub>4</sub>, which is an added bonus over the air-sensitive PCy<sub>3</sub> employed by Zeng.

The scope of the Suzuki-Miyaura cross-coupling of N-acylsuccinimides was then investigated (Table 2). Importantly, the reaction tolerates electron-neutral (**3a**), electron-rich (**3b**) and electron-deficient (**3c**) boronic acids. Thus, the method is insensitive to electronic properties on the boronic acid component, resulting in a general process (cf. catalytic system reported by Zeng). Pleasingly, steric-hindrance is well-tolerated as exemplified by the coupling of ortho-tolyl boronic acid. Importantly, highly electrophilic carbonyl groups, such as


**Table 1** Optimization of Pd-Catalyzed Suzuki-Miyaura Cross-Coupling of N-Acylsuccinimides<sup>a</sup>












Entry	Catalyst	Ligand	Base	Acid	Solvent	Yield (%) <sup>b</sup>
1	Pd(OAc) <sub>2</sub>	PCy <sub>3</sub> HBF <sub>4</sub>	K <sub>2</sub> CO <sub>3</sub>	H <sub>3</sub> BO <sub>3</sub>	THF	>95
2	Pd(OAc) <sub>2</sub>	PCy <sub>3</sub> HBF <sub>4</sub>	K <sub>2</sub> CO <sub>3</sub>	-	THF	<5
3	Pd(OAc) <sub>2</sub>	PCy <sub>3</sub> HBF <sub>4</sub>	K <sub>2</sub> CO <sub>3</sub>	H <sub>3</sub> BO <sub>3</sub>	THF	54
4	Pd(OAc) <sub>2</sub>	PCy <sub>3</sub> HBF <sub>4</sub>	Na <sub>2</sub> CO <sub>3</sub>	-	Dioxane	>95
5	Pd(OAc) <sub>2</sub>	PCy <sub>3</sub> HBF <sub>4</sub>	Na <sub>2</sub> CO <sub>3</sub>	-	Dioxane	75
6	Pd(OAc) <sub>2</sub>	PCy <sub>3</sub> HBF <sub>4</sub>	K <sub>2</sub> CO <sub>3</sub>	-	Dioxane	10
7	Pd(OAc) <sub>2</sub>	PCy <sub>3</sub> HBF <sub>4</sub>	Na <sub>2</sub> CO <sub>3</sub>	-	Toluene	19
8	Pd(OAc) <sub>2</sub>	PCy <sub>2</sub> Ph	Na <sub>2</sub> CO <sub>3</sub>	-	Dioxane	47
9	Pd(OAc) <sub>2</sub>	PPh <sub>3</sub>	Na <sub>2</sub> CO <sub>3</sub>	-	Dioxane	7
10	Pd(OAc) <sub>2</sub>	PtBu <sub>3</sub> HBF <sub>4</sub>	Na <sub>2</sub> CO <sub>3</sub>	-	Dioxane	6

<sup>a</sup>Amide (1.0 equiv), 4-Tol-B(OH)<sub>2</sub> (2.0 equiv), base (2.5 equiv), Pd(OAc)<sub>2</sub> (3 mol%), ligand (12 mol%), acid (1.5 equiv), solvent (0.25 M), T, 15 h. <sup>b</sup>Determined by <sup>1</sup>H NMR and/or GC-MS. T: Entries 1-2: 60 °C, entry 3: 23 °C, entries 4, 6-10: 120 °C, entry 5: 100 °C.


**Table 2** Pd-Catalyzed Suzuki-Miyaura Cross-Coupling of N-Acylsuccinimides: Boronic Acid Scope<sup>a</sup>



		
<b>3a: 97% yield</b>	<b>3b: 93% yield</b>	<b>3c: 84% yield</b>
		
<b>3d: 65% yield<sup>b</sup></b>	<b>3e: 96% yield</b>	<b>3f: 91% yield</b>
		
<b>3g: 95% yield</b>	<b>3h: 50% yield</b>	<b>3i: 63% yield</b>

<sup>a</sup>Conditions: amide (1.0 equiv), ArB(OH)<sub>2</sub> (2.0 equiv), Pd(OAc)<sub>2</sub> (3 mol%), PCy<sub>3</sub>HBF<sub>4</sub> (12 mol%), Na<sub>2</sub>CO<sub>3</sub> (2.5 equiv), dioxane (0.25 M), 120 °C, 15 h. <sup>b</sup>ArB(OH)<sub>2</sub> (3.0 equiv), Na<sub>2</sub>CO<sub>3</sub> (4.5 equiv). See ESI for details.

aldehydes and ketones placed at various positions around the aromatic ring on the boronic acid component (**3e-3g**) are well-tolerated. It is worthwhile to note that ketones and aldehydes are incompetent substrates in the classic Weinreb ketone synthesis. Also note that the alternative method reported by Zeng does not tolerate carbonyl functional groups. Furthermore, fluorinated aromatics (**3h**) highly relevant from the medicinal chemistry standpoint and heterocycles (**3i**) are well-tolerated. In agreement with the cyclic structure of the

**Table 3** Pd-Catalyzed Suzuki-Miyaura Cross-Coupling of N-Acylsuccinimides: N-Acylsuccinimide Scope<sup>a</sup>

<sup>a</sup>Conditions: amide (1.0 equiv), ArB(OH)<sub>2</sub> (2.0 equiv), Pd(OAc)<sub>2</sub> (3 mol%), PCy<sub>3</sub>HBF<sub>4</sub> (12 mol%), Na<sub>2</sub>CO<sub>3</sub> (2.5 equiv), dioxane (0.25 M), 120 °C, 15 h. <sup>b</sup>ArB(OH)<sub>2</sub> (3.0 equiv), Na<sub>2</sub>CO<sub>3</sub> (4.5 equiv). See ESI for details.

succinimide ring, in all cases the reaction proceeded in a highly selective fashion for the activation of the N–C amide bond (cf. undesired cleavage of the σ N–C bond).




We next focused on the scope of N-acylsuccinimide coupling partner as illustrated in Table 3. Similar to the scope of the boronic acid coupling partner, the reaction delivered high yields of the coupling products irrespective of the electronic properties of the amide component, including electron-neutral, electron-donating and electron-withdrawing substituents (3a–c). Likewise, steric-hindrance on the amide is well-tolerated (3d). Perhaps most importantly, amides bearing electrophilic carbonyl groups, such as ketones and esters are also compatible (3j–k, vide supra). Finally, we were pleased to find that heterocyclic (3l) as well as aliphatic amides (3m) are competent substrates for the reaction. Overall, the scope of our method using the catalytic system based on Pd(OAc)<sub>2</sub>, PCy<sub>3</sub>HBF<sub>4</sub> and Na<sub>2</sub>CO<sub>3</sub> compares very favorably with the process reported by Zeng<sup>22</sup> in terms of reaction efficiency, generality and functional group tolerance, in particular with respect to the synthetically-valuable carbonyl groups.<sup>24</sup>

We have conducted studies to gain insight into the catalytic mechanism of this reaction:

(1) Intermolecular competition experiments with differently substituted amides revealed that electron-deficient arenes are more reactive (4-CF<sub>3</sub>:4-MeO = 97:3, cf. glutarimide, 4-CF<sub>3</sub>:4-MeO, 97:3)<sup>9a</sup> (Scheme 1A).

(2) Further competition experiments revealed preference for electron-deficient boronic acids (4-CF<sub>3</sub>:4-MeO = 73:27, cf. glutarimide, 4-CF<sub>3</sub>:4-MeO = 75:25)<sup>9a</sup> (Scheme 1B).

(3) As expected on the basis of amide twist, the reaction is slightly less selective for the cross-coupling of N-acylsuccinimides cf. N-acylglutarimides (Scheme 1C). Nevertheless, this is compensated

**A: Intermolecular competition: amides****B: Intermolecular competition: boronic acids****C: Effect of N-substitution: succinimide vs. glutarimide****Scheme 1** Mechanistic selectivity studies.

by the compelling availability and bargain price of the succinimide activating group.

(4) While the reaction proceeds more rapidly upon increasing the amounts of boronic acid and base (Na<sub>2</sub>CO<sub>3</sub>, 2.5 equiv, 4-Tol-B(OH)<sub>2</sub>, 2.0 equiv, 1 h, 26% conversion; Na<sub>2</sub>CO<sub>3</sub>, 5.0 equiv, 4-Tol-B(OH)<sub>2</sub>, 4.0 equiv, 1 h, 31% conversion), the increase is much lower than previously observed using N-acylglutarimides (54% to >98%).<sup>9b</sup>

(5) A turnover number (TON) of 130 was determined under these conditions at 0.25 mol% Pd loading.

Overall, these mechanistic findings suggest that metal insertion may be the rate limiting step in the coupling.

In conclusion, we have developed a general and operationally-simple method for Suzuki-Miyaura cross-coupling of N-acylsuccinimides. The key features of our protocol include high efficiency, air-stable and commercially-available reagents and catalysts. The reaction employs new cross-coupling reagents, N-acylsuccinimides, which are activated for selective metal insertion by half-twist of the amide bond. Besides the great potential applications of N-acylsuccinimides in a wide range of transition-metal catalyzed cross-couplings, these reagents could also inspire future research to address tunable distortion of the amide bond. Further studies on the development of cross-coupling reactions of amides, including aliphatic amides, and detailed

## COMMUNICATION


## Organic &amp; Biomolecular Chemistry


mechanistic investigations are underway and will be reported in due course.

Rutgers University and the NSF (CAREER CHE-1650766) are acknowledged for support. The 500 MHz spectrometer was supported by the NSF-MRI grant (CHE-1229030).

## Notes and references

- Reviews: (a) R. Takise, K. Muto and J. Yamaguchi, *Chem. Soc. Rev.*, 2017, DOI: 10.1039/c7cs00182g; (b) G. Meng, S. Shi and M. Szostak, *Synlett*, 2016, **27**, 2530; (c) C. Liu and M. Szostak, *Chem. Eur. J.*, 2017, **23**, 7157; (d) J. E. Dander and N. K. Garg, *ACS Catal.*, 2017, **7**, 1413; (e) Y. Gao, C. L. Ji and X. Hong, *Sci. China Chem.* 2017, DOI: 10.1007/s11426-017-9025-1.
- General reviews on cross-coupling: (a) A. de Meijere, S. Bräse and M. Oestreich, *Metal-Catalyzed Cross-Coupling Reactions and More*, Wiley, 2014; (b) G. Molander, J. P. Wolfe and M. Larhed, *Science of Synthesis: Cross-Coupling and Heck-Type Reactions*, Thieme, 2013; For a review on electrophilic activation of amides: (c) D. Kaiser and N. Maulide, *J. Org. Chem.*, 2016, **81**, 4421; (d) For an excellent overview of amide cross-coupling, see: S. A. Ruider and N. Maulide, *Angew. Chem. Int. Ed.*, 2015, **54**, 13856; (e) For representative tandem amide coupling, see: J. A. Walker, K. L. Vickerman, J. N. Humke and L. M. Stanley, *J. Am. Chem. Soc.* **2017**, **139**, 10228, and references cited therein.
- For representative acyl-couplings, see: (a) L. Hie, N. F. F. Nathel, T. K. Shah, E. L. Baker, X. Hong, Y. F. Yang, P. Liu, K. N. Houk and N. K. Garg, *Nature*, 2015, **524**, 79; (b) P. Lei, G. Meng and M. Szostak, *ACS Catal.*, 2017, **7**, 1960; (c) G. Meng, P. Lei and M. Szostak, *Org. Lett.*, 2017, **19**, 2158.
- For representative decarbonylative couplings, see: (a) G. Meng and M. Szostak, *Angew. Chem. Int. Ed.*, 2015, **54**, 14518; (b) S. Shi, G. Meng and M. Szostak, *Angew. Chem. Int. Ed.*, 2016, **55**, 6959; (c) A. Dey, S. Sasmai, K. Seth, G. K. Lahiri and D. Maiti, *ACS Catal.*, 2017, **7**, 433; (d) H. Yue, L. Guo, H. H. Liao, Y. Cai, C. Zhu and M. Rueping, *Angew. Chem. Int. Ed.*, 2017, **56**, 4282; (e) H. Yue, L. Guo, S. C. Lee, X. Liu and M. Rueping, *Angew. Chem. Int. Ed.*, 2017, **56**, 3972.
- For studies on the effect of amide bond geometry on cross-coupling, see: (a) V. Pace, W. Holzer, G. Meng, S. Shi, R. Lalancette, R. Szostak and M. Szostak, *Chem. Eur. J.*, 2016, **22**, 14494; (b) R. Szostak, S. Shi, G. Meng, R. Lalancette and M. Szostak, *J. Org. Chem.*, 2016, **81**, 8091.
- For the lead reference on electronic activation, see: (a) R. Szostak, G. Meng and M. Szostak, *M., J. Org. Chem.*, 2017, **82**, 6373; (b) See also refs. 16 and 17.
- For recent developments of new catalytic systems in amide bond cross-coupling, see: Pd-NHC: (a) Ref. 3b; (b) P. Lei, G. Meng, S. Shi, Y. Ling, J. An, R. Szostak and M. Szostak, *Chem. Sci.*, 2017, **8**, 6252; (c) P. Lei, G. Meng, Y. Ling, J. An and M. Szostak, *J. Org. Chem.*, 2017, **82**, 6638; Ni-NHC: (d) L. Hie, E. L. Baker, S. M. Anthony, J. N. Desrosiers, C. Senanayake and N. K. Garg, *Angew. Chem. Int. Ed.*, 2016, **55**, 15129; (e) J. E. Dander, E. L. Baker and N. K. Garg, *Chem. Sci.*, 2017, **8**, 6433.
- A. Greenberg, C. M. Breneman and J. F. Liebman, *The Amide Linkage: Structural Significance in Chemistry, Biochemistry, and Materials Science*, Wiley, 2000.
- (a) G. Meng and M. Szostak, *Org. Lett.*, 2015, **17**, 4364; (b) G. Meng and M. Szostak, *Org. Biomol. Chem.*, 2016, **14**, 5690.
- For selected examples, see: (a) Ref. 4a-b, 4d-e, 9; (b) S. Shi and M. Szostak, *Org. Lett.*, 2017, **19**, 3095; (c) C. Liu and M. Szostak, *Angew. Chem. Int. Ed.*, 2017, DOI: 10.1002/anie.201707102; (d) W. Srimontree, A. Chatupheeraphat, H. H. Liao and M. Rueping, *Org. Lett.*, 2017, **19**, 3091; (e) A. Chatupheeraphat, H. H. Liao, S. C. Lee and M. Rueping, *Org. Lett.*, 2017, **19**, 4255; (f) S. Ni, W. Zhang, H. Mei, J. Han and Y. Pan, *Org. Lett.*, 2017, **19**, 2536.
- N. A. Weires, E. L. Baker and N. K. Garg, *Nat. Chem.*, 2016, **8**, 75.
- X. Li and G. Zou, *Chem. Commun.*, 2015, **51**, 5089.
- G. Meng, S. Shi and M. Szostak, *ACS Catal.*, 2016, **6**, 7335.
- (a) C. Liu, G. Meng, Y. Liu, R. Liu, R. Lalancette, R. Szostak and M. Szostak, *Org. Lett.*, 2016, **18**, 4194; (b) H. Wu, M. Cui, J. Jian and Z. Zheng, *Adv. Synth. Catal.*, 2016, **358**, 3876; (c) For the seminal study on using N-bromosaccharin, see: B. Zajc, *Synth. Commun.*, 1999, **29**, 1779.
- C. Liu, Y. Liu, R. Liu, R. Lalancette, R. Szostak and M. Szostak, *Org. Lett.*, 2017, **19**, 1434.
- G. Meng, R. Szostak and M. Szostak, *Org. Lett.*, 2017, **19**, 3596.
- G. Meng, R. Lalancette, R. Szostak and M. Szostak, *Org. Lett.*, 2017, **19**, 4656.
- S. Shi and M. Szostak, *Synthesis*, 2017, **49**, 3602.
- Reviews on acyl-metal intermediates: (a) L. J. Gooßen, N. Rodriguez and K. Gooßen, *Angew. Chem. Int. Ed.*, 2008, **47**, 3100; (b) W. Dzik, P. Lange and L. Gooßen, *Chem. Sci.*, 2012, **3**, 2671; (c) For a recent advance in generating acyl-metal intermediates, see: S. D. Friis, A. T. Lindhardt and T. Skrydstrup, *Acc. Chem. Res.*, 2016, **49**, 594.
- (a) For an example of using N-acylsuccinimides in Sml<sub>2</sub>-mediated reductive coupling, see: R. H. Taaning, K. B. Lindsay, B. Schiøtt, K. Daabjerg and T. Skrydstrup, *J. Am. Chem. Soc.*, 2009, **131**, 10253. (b) For a pertinent review, see: P. Ebran, C. M. Jensen, S. A. Johannesen, J. Karaffa, K. B. Lindsay, R. Taaning and T. Skrydstrup, *Org. Biomol. Chem.*, 2006, **4**, 3553.
- (a) For an elegant study on using N-acylsuccinimides as acylating reagents, see: C. A. Goodman, J. B. Eagles, L. Rudahindwa, C. G. Hamaker and S. R. Hitchcock, *Synth. Commun.*, 2013, **43**, 2155; (b) For a relevant use of N-acylsuccinimides in Ni/Ir-catalyzed acyl transfer, see: J. Amani, R.; Alam, S.; Badir and G. A. Molander, *Org. Lett.*, 2017, **19**, 2426.
- M. Cui, Z. Chen, T. Liu, H. Wang and Z. Zeng, *Tetrahedron Lett.*, 2017, DOI: 10.1016/j.tetlet.2017.08.044. Online publication date: 31 August 2017.
- Note that the reference 22 incorrectly assigns the effect of acid coordination to the amide bond in N-acylsuccinimides and N-acylglutarimides. It has been conclusively demonstrated that these amides favour O-protonation,<sup>5a,b</sup> which is in contrast to bridged lactams, which favour N-protonation: R. Szostak, J. Aubé and M. Szostak, *Chem. Commun.*, 2015, **51**, 6395.
- The following features additionally demonstrate the high activity of the present catalytic system: (1) using the alternative system,<sup>22</sup> the yield drops significantly upon lowering the Pd-loading to 3 mol%, which is a standard protocol in the present method; (2) the present method tolerates alkyl N-acylsuccinimide amides. Alkyl amides, in general, are challenging substrates for cross-coupling. See the following references that address this point: 7d and 7e.





The palladium-catalyzed Suzuki-Miyaura cross-coupling of *N*-acylsuccinimides as versatile acyl-transfer reagents via selective amide N–C bond cleavage is reported.