


Palladium-Catalyzed Hydroxylation of Aryl Halides under Ambient Conditions**

Alexey G. Sergeev, Thomas Schulz, Christian Torborg, Anke Spannenberg, Helfried Neumann, and Matthias Beller*

Phenols are found in various natural products ranging from coal tar containing simple phenols to red wine possessing numerous colored polyphenols with antioxidant properties.^[1,2] In organic synthesis the development of mild, general, and efficient methods for the preparation of phenols still constitutes a significant challenge. Considering the availability of starting materials, the direct nucleophilic substitution of a halogen atom in aryl halides is an appealing approach to the synthesis of substituted phenols.^[3,4] However, reactions of non-activated substrates typically proceed under harsh reaction conditions (200–350 °C).^[1] Hence, the development of a milder catalytic phenol synthesis through a two-step coupling procedure by Hartwig and co-workers was an important advancement.^[5] Recently, the direct hydroxylation of aryl halides was developed by Buchwald and co-workers as well as by Chan and co-workers using palladium catalysts with bulky tri-*tert*-butylphosphine and biphenylphosphines (**L1**, **L2**).^[6,7]




Among the various catalytic systems studied, palladium complexes based on ligands **L1** and **L2** allowed C–O coupling of both activated and non-activated aryl halides at 80–100 °C.^[7] Inspired by the efficiency of palladium/biphenylphosphine catalysts, we recently demonstrated the hydroxylation of aryl halides in the presence of imidazole-based ligands.^[8] Although progress in palladium-catalyzed C–O bond-forming reactions made it possible to arylate alcohols under milder conditions,^[9,10] the coupling using water remained problematic.^[4] Indeed, no examples of phenol

syntheses from non-activated aryl halides at ambient temperature have been described to date,^[11] and the mechanistic understanding of this coupling reaction is limited.

Herein we report the first room temperature palladium-catalyzed hydroxylation of aryl chlorides and bromides as deduced from studies of the elementary steps of the catalytic cycle. In the presence of a novel palladium precursor and the imidazolyphosphine ligand **L3** (Ad = adamantyl), a variety of phenols can be obtained from aryl halides in excellent yield.

Initially, we were interested in the structure and reactivity of palladium complexes of the bulky ligand **L3**, which was previously involved in the hydroxylation of aryl halides. To prepare the corresponding oxidative addition complex, bromomesitylene was reacted with different palladium sources and ligand **L3** in THF at room temperature. Notably, only [Pd(cod)(CH₂SiMe₃)₂] allowed isolation of the desired oxidative addition product in pure form. The oxidative addition of bromomesitylene in the presence of [Pd(cod)(CH₂SiMe₃)₂] and **L3** took place within 10 minutes at room temperature to give complex **1** in 65% yield (Scheme 1). In contrast, the



Scheme 1. Synthesis of complex **1**. cod = 1,5-cyclooctadiene.

reaction with [Pd₂(dba)₃] (dba = dibenzylideneacetone) proceeded more slowly (14 h) to give the oxidative addition complex in moderate yield (44%) upon isolation.^[12] Recent attempts to synthesize related [Pd(Ar)(X)(L)] (X = halogen) complexes bearing 2-dialkylphosphino-2,4,6-tri-*iso*-propylbiphenyl ligands (X-Phos-type phosphines including **L1** and **L2**) were not successful.^[13,14] However, the use of a ligand having methoxy groups introduced into the 3,6-positions of X-Phos allowed the synthesis of the stable oxidative addition complexes.^[14] Thus, introduction of additional donor atoms may play a profound role in the improved stability of the oxidative addition intermediate.

The crystal structure of the oxidative addition complex **1** is shown in Figure 1.^[15] Complex **1** contains a single phosphine ligand coordinated to the metal through both the phosphorous and nitrogen atoms, thereby forming a four-membered P,N-chelate ring.^[16] The additional coordination of the nitro-

[*] Dr. A. G. Sergeev, Dipl.-Chem. T. Schulz, Dipl.-Chem. C. Torborg, Dr. A. Spannenberg, Dr. H. Neumann, Prof. Dr. M. Beller
 Leibniz-Institut für Katalyse e.V.
 Albert-Einstein-Strasse 29a, 18059 Rostock (Germany)
 Fax: (+49) 381-1281-5000
 E-mail: matthias.beller@catalysis.de

[**] We thank Dipl.-Ing. A. Koch, S. Leiminger, and A. Kammer for excellent analytical support. We gratefully acknowledge Evonik (formerly Degussa) for financial support and the precious gift of different chemicals.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.200902148>.





Figure 1. Crystal structure and important through-space interactions in NOESY NMR spectrum of complex **1** (see the Supporting Information for details). The thermal ellipsoids are set at 30% probability.

gen atom “freezes” the rather sterically unfavorable conformation of the coordinated phosphine with bulky diisopropylphenyl and di-1-adamantylphosphino groups facing each other.

Thus, one side of the complex is sterically overcrowded with the phosphine, whereas the opposite side is relatively open for nucleophilic attack. As determined by NOESY NMR experiments **1** exists in a single form in solution,^[17] which corresponds to that found in the solid state (Figure 1).^[18]

Next, we investigated the preparation of the corresponding hydroxo complex “[Pd(Ar)(OH)(L3)]” (**2**), which is the key catalytic intermediate for the formation of phenol. Thus, **1** was reacted with an excess of *n*Bu₄NOH·30H₂O in THF at room temperature. The starting complex **1** was completely consumed within less than 20 minutes. To our surprise instead of the desired hydroxo complex **2**, 2,4,6-trimethylphenol was formed in 75% yield (Scheme 2). Apparently, complex **2** immediately underwent C–O bond-forming reductive elimi-



Scheme 2. Reaction of **1** with tetra-*n*-butyl ammonium hydroxide.

nation at room temperature. To the best of our knowledge such direct reductive elimination of phenols from palladium has not been reported until now.^[19] It is this step that is considered as the “bottle-neck” of the palladium-catalyzed hydroxylation and which is responsible for rather high reaction temperatures ($\geq 80^\circ\text{C}$).^[6,7]

Apparently, the combination of [Pd(cod)(CH₂SiMe₃)₂]/**L3** allows all steps of the catalytic cycle to proceed at room temperature within minutes. These findings encouraged us to study the corresponding catalytic reaction in the presence of [Pd(cod)(CH₂SiMe₃)₂] and other palladium precursors (Table 1). To our delight the reaction of bromomesitylene with an excess of CsOH·H₂O in the presence of [Pd(cod)(CH₂SiMe₃)₂] and **L3** in THF gave mesitol in nearly quantitative yield (Table 1, entry 2). The oxidative addition complex **1** as catalyst led to the product in identical yield

Table 1: Variation of palladium sources and ligands in the hydroxylation of bromomesitylene.^[a]

Entry	Ligand	“Pd” source	Yield [%] ^[b]
1	–	1	99
2	L3	[Pd(cod)(CH ₂ SiMe ₃) ₂]	98 (88) ^[c]
3	L3	[Pd ₂ (dba) ₃]	76 (44) ^[c]
4	L3	[Pd(allyl)(Cp)]	37
5	L3	[Pd(TMEDA)(Me) ₂]	35
6	L3	Pd(OAc) ₂	30
7	L1	[Pd(cod)(CH ₂ SiMe ₃) ₂]	1
8	L2	[Pd(cod)(CH ₂ SiMe ₃) ₂]	0
9	PtBu ₃	[Pd(cod)(CH ₂ SiMe ₃) ₂]	0
10	Q-Phos	[Pd(cod)(CH ₂ SiMe ₃) ₂]	0
11	L4	[Pd(cod)(CH ₂ SiMe ₃) ₂]	11

[a] Reaction conditions: bromomesitylene (1 mmol), CsOH·H₂O (3 mmol), [Pd(cod)(CH₂SiMe₃)₂] (2 mol%), phosphine ligand (3 mol%), hexadecane (100 μL), THF (2 mL), 24 $^\circ\text{C}$, 20 h. [b] Yields determined by using GC methods. [c] In parentheses: 1 mol% of Pd and 1.5 mol% of a ligand was used. Cp = cyclopentadiene, TMEDA = *N,N,N',N'*-tetramethylethylenediamine.

(Table 1, entry 1). Notably, all other palladium precursors tested with **L3** are considerably less efficient (Table 1, entries 3–6). The advantage of [Pd(cod)(CH₂SiMe₃)₂] in the room temperature hydroxylation is explained by the efficient generation of the corresponding palladium(0) complex (PdL) by rapid substitution of 1,5-cyclooctadiene for the bulky phosphine ligand L and a subsequent fast reductive elimination of bis(trimethylsilyl)ethane.^[20] As a result, the oxidative addition occurs fast at room temperature. Inferior results with the common palladium(0) precursor, [Pd₂(dba)₃] in the catalytic reaction (Table 1, entry 3) are attributed to a much slower oxidative addition of the aryl bromide (see below). This deceleration should be caused by the stronger coordination of the electron-poor olefin (dibenzylidenacetone) to the palladium center.^[21] Similar unfavorable effects of dba in palladium-catalyzed reactions are well documented in the literature.^[22]

Notably, despite applications of [Pd(cod)(CH₂SiMe₃)₂] (R = Me, Ph) in organometallic synthesis,^[14,20] these complexes have never been used in catalytic reactions.^[23] The ability of [Pd(cod)(CH₂SiMe₃)₂] to generate catalytically active species in situ under mild reaction conditions and in a one-step synthesis from commercially available precursor^[20] make this complex an appealing tool for catalysis of cross-coupling reactions at room temperature.^[24]

Furthermore, the palladium(0) precursor was tested with other ligands, which were successfully employed for hydroxylation of aryl halides at high temperature (**L1**, **L2**, and PtBu₃). For comparison, we also employed Q-Phos which is the unique ligand for arylation of *t*BuOH and *t*BuMe₂SiOH at room temperature.^[10] However, none of these tests led to the product at room temperature, demonstrating the superiority of **L3** for this reaction (Table 1, entries 7–10).

We suppose that the improved activity of the **L3**-based catalyst in the room temperature hydroxylation stems from a

favorable combination of the steric bulk of the biphenyl-type backbone and alkyl substituents bound to the phosphorous center, as well as the ability to coordinate to the metal through the sp^2 -nitrogen atom. The favorable effect of increased steric bulk of biphenyl-type ligands has already been highlighted.^[8,25] Indeed, in the hydroxylation at elevated temperatures (80–100 °C), the efficiency of the catalysts based on bulky biphenyl ligands (**L1**, **L2**), and their arylimidazolyl congeners with a similar backbone is rather close.^[7,8] However, in the room temperature reaction the novel palladium precursor and imidazolyl ligand **L3** exhibits an enormous difference in efficiency relative to reaction runs with the biphenyl phosphines **L1** and **L2** (Table 1, entries 2 and 7–8).

The main reason for this different catalytic behavior is seen in the ability of **L3** to coordinate to the palladium through both the P and N atoms (Figure 1). In agreement with this assumption, the pyrrole-based ligand **L4** which has the same framework as **L3**, but contains no second nitrogen atom available for coordination, provided only 11 % yield of phenol versus 98 % in the presence of **L3** (Table 1, entries 11 and 2).

Next, we optimized the reaction conditions for room temperature hydroxylation of bromomesitylene (Table 2). Among the various hydroxides tested, the best results were obtained in the presence of $CsOH \cdot H_2O$ (Table 2, entry 3). The mediocre performance of $nBu_4NOH \cdot 30H_2O$ in the catalytic reaction (Table 2, entry 5) compared to the reaction using stoichiometric amounts of $nBu_4NOH \cdot 30H_2O$ (Scheme 2) is attributed to the formation of a biphasic mixture resulting from the high water content in the commercial hydroxide.

Table 2: Variation of hydroxides in the hydroxylation of bromomesitylene.^[a]

Entry	Hydroxide	Conversion [%] ^[b]	Yield [%] ^[b]
1	LiOH	7	0
2	KOH	37	27
3	$CsOH \cdot H_2O$	100	98
4	$Me_4NOH \cdot 5H_2O$	98	87
5	$nBu_4NOH \cdot 30H_2O$	41	36

[a] Reaction conditions: bromomesitylene (1 mmol), base (3 mmol), $[Pd(cod)(CH_2SiMe_3)_2]$ (2 mol%), **L3** (3 mol%), hexadecane (100 μ L), THF (2 mL), 24 °C, 20 h. [b] Determined by GC methods.

Finally, we applied the optimized reaction conditions to the hydroxylation of various aryl halides at ambient conditions. As shown in Table 3, different aryl bromides and even aryl chlorides are transformed into the corresponding phenols in good to excellent yields (isolated; 67–99 %). In all cases, aryl chlorides were as active as the corresponding bromides and no noticeable amounts of diphenyl ethers were detected. Aryl halides containing electron-withdrawing substituents in the *ortho* and *para* positions reacted smoothly (Table 3, entries 10–16). Functional groups such as cyano, nitro, and

Table 3: Palladium-catalyzed hydroxylation of aryl halides at room temperature.^[a]

Entry	Aryl Halide	Product	"Pd" [mol %]	Yield [%] ^[b]
1			1 2	n.d. (88) 93 (98)
2			2	96
3		X = Br	2	92
4		X = Cl	2	92
5			4	70 ^[c]
6			2	72 (97) ^[c,d]
7		X = Br	2	85
8		X = Cl	2	75
9			2	87
10			2	94
11			4	88
12			1	99
13		X = Br	2	83 (89) ^[d]
14		X = Cl	2	73 (98) ^[d]
15			4	67 (72) ^[d]
16			2	95
17			2	99

[a] Reaction conditions: aryl halide (1 mmol), $CsOH \cdot H_2O$ (3 mmol), $[Pd(cod)(CH_2SiMe_3)_2]$ (1–4 mol%), **L3** (1.5–6 mol%), hexadecane (100 μ L), THF (2 mL), 24 °C, 20 h; reaction time is not optimized. [b] Yields for isolated product, and yields given in parentheses were determined by using GC methods. The reported yields refer to independent runs. [c] The reaction was carried out at 50 °C. [d] Volatile compound.

keto substituents are well tolerated. Moreover, the room temperature hydroxylation has a broader functional group scope compared to the reaction run at elevated temperature. For example, at 100 °C, we have observed only limited tolerance to nitriles.^[26] In contrast, the present mild protocol enables preservation of nitrile group in all cases. Indeed, hydroxylation of 4-bromo-3-methylbenzonitrile and 4-bromobenzonitrile gave phenols in 88 % and 99 % yields,

respectively, using lower catalyst loadings (Table 3, entries 11 and 12). Previously, all attempts to hydroxylate aryl halides possessing a CF₃ substituent at elevated temperatures failed because of the hydrolysis of the CF₃ group. Under the present set of conditions both *para*- and *ortho*-aryl halides possessing CF₃ groups were coupled successfully at room temperature in good yields (Table 3, entries 13–15). In addition, a heterocyclic derivative, such as 4-chloro-2-methylquinoline, is smoothly transformed into the corresponding phenol in quantitative yield (Table 3, entry 17). With regard to limitations of the present method, we found that CO₂H and OH groups are not tolerated presumably because of the formation poorly soluble salts.

The novel protocol was also successfully applied for the hydroxylation of various non-activated aryl bromides including bromobenzene, *ortho*- and *para*-substituted halobenzenes, and 1-halonaphthalenes to give substituted phenols in 70–93% yields (Table 3, entries 1–9). Bromo- and chloronaphthalenes smoothly reacted at room temperature affording naphthols in 75–87% yields (Table 3, entries 7–9). Bromobenzene and 4-bromotoluene were somewhat less reactive and required a reaction temperature of 50°C (Table 3, entries 5–6). The introduction of one or two methyl groups in *ortho* positions of halobenzenes increased the reactivity of the halides, allowing the reaction again to proceed at room temperature with excellent yields (Table 3, entries 1–4). For example, bromomesitylene and 2-bromotoluene furnished phenols in 93% and 92% yields, respectively. Hydroxylation of similar chloro derivatives also proceeded under the same conditions (Table 3, entries 2 and 4). Acceleration of the reaction by *ortho* substituents in haloarenes, as well as the similar reactivity of aryl bromides and aryl chlorides, makes it likely that reductive elimination is the rate limiting step for this hydroxylation reaction. Indeed, it is known that carbon–heteroatom bond-forming reductive elimination from aryl-palladium complexes is accelerated by *ortho* substituents in aryl halides.^[27] However, we found that the further increase in the steric bulk of the *ortho* substituents, for example, from methyl to *iso*-propyl decelerates the reaction. Hence, room temperature hydroxylation of 2-bromo-*iso*-propylbenzene proceeded only to 10% conversion, whereas the reaction of 2,4,6-tri-*iso*-propylbromobenzene did not occur at all and led to decomposition of the catalyst.

This effect of the different *ortho* substituents on the rate of the hydroxylation may imply a change of the rate-determining step.

In summary, we demonstrated for the first time that all steps of the catalytic cycle of the palladium-catalyzed hydroxylation of aryl chlorides and bromides can proceed at room temperature. Based on these findings, the catalytic synthesis of phenols occurs under unprecedented mild conditions. The key to the success is the combination of the novel palladium precursor [Pd(cod)(CH₂SiMe₃)₂] and the imidazole-based ligand **L3**, which should be useful for numerous other catalytic coupling reactions as well.

Received: April 21, 2009

Revised: August 7, 2009

Published online: September 8, 2009

Keywords: homogeneous catalysis · hydrolysis · hydroxylation · palladium · phenols

- [1] a) *Patai Series: The Chemistry of Functional Group. The Chemistry of Phenols* (Ed.: Z. Rappoport), Wiley, Chichester, **2003**; b) J. H. P. Tyman, *Studies in Organic Chemistry* 52. *Synthetic and Natural Phenols*, Elsevier, Amsterdam, **1996**.
- [2] For polyphenols in wine, see: G. J. Soleas, E. P. Diamandis, D. M. Goldberg, *J. Clin. Lab. Anal.* **1997**, *11*, 287–313.
- [3] For typical methods for the preparation of phenols, see: reference [1] and D. A. Whiting in *Comprehensive Organic Chemistry. The Synthesis and Reactions of Organic Compounds, Vol. 1* (Eds.: D. Barton, W. D. Ollis), Pergamon Press, Oxford, **1979**, pp. 717–730.
- [4] For a highlight of the palladium-catalyzed synthesis of phenols, see: M. C. Willis, *Angew. Chem.* **2007**, *119*, 3470–3472; *Angew. Chem. Int. Ed.* **2007**, *46*, 3402–3404.
- [5] S. Mann, C. Incarvito, A. L. Rheingold, J. F. Hartwig, *J. Am. Chem. Soc.* **1999**, *121*, 3224–3225.
- [6] G. Chen, A. S. C. Chan, F. Y. Kwong, *Tetrahedron Lett.* **2007**, *48*, 473–476.
- [7] K. W. Anderson, T. Ikawa, R. E. Tundel, S. L. Buchwald, *J. Am. Chem. Soc.* **2006**, *128*, 10694–10695.
- [8] T. Schulz, C. Torborg, B. Schäffner, J. Huang, A. Zapf, R. Kadyrov, A. Börner, M. Beller, *Angew. Chem.* **2009**, *121*, 936–939; *Angew. Chem. Int. Ed.* **2009**, *48*, 918–921.
- [9] For overview of palladium-catalyzed arylation of phenols and alcohols, see: a) J. F. Hartwig in *Handbook of Organopalladium Chemistry for Organic Synthesis, Vol. 1* (Eds.: E.-i. Negishi, A. de Meijere), Wiley, New York, **2002**, pp. 1097–1106; b) A. Zapf, M. Beller, T. Riermeier in *Transition Metals for Organic synthesis. Building Blocks and Fine Chemicals, Vol. 1* (Eds.: M. Beller, C. Bolm), Wiley, Weinheim, **2004**, pp. 231–256.
- [10] For room temperature arylation of alcohols using Pd/Q-Phos, see: a) Q. Shelby, N. Kataoka, G. Mann, J. Hartwig, *J. Am. Chem. Soc.* **2000**, *122*, 10718–10719; b) N. Kataoka, Q. Shelby, J. Stambuli, J. F. Hartwig, *J. Org. Chem.* **2002**, *67*, 5553–5566.
- [11] Hydroxylation of activated aryl halides such as *ortho*-halonitrobenzenes was carried out in the presence of Pd/PtBu₃ at 25–50°C (see reference [6]).
- [12] In addition to the main product some minor side-products were formed (5–10%), which could not be removed by recrystallization. Reaction with [Pd(TMEDA)(Me)₂] gave a mixture of [Pd(**L3**)(2,4,6-Me₃C₆H₂)(Br)] (**1**) and [Pd(TMEDA)(2,4,6-Me₃C₆H₂)(Br)] in the 2.57:1 ratio.
- [13] T. E. Barder, M. R. Biscoe, S. L. Buchwald, *Organometallics* **2007**, *26*, 2183–2193.
- [14] B. P. Fors, D. A. Watson, M. R. Biscoe, S. L. Buchwald, *J. Am. Chem. Soc.* **2008**, *130*, 13552–13554.
- [15] CCDC 728525 (**1**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [16] Four-membered P,N-chelates were also observed in palladium and platinum complexes of 2-(dialkylphosphino)-1-methyl-1H-imidazoles: a) D. B. Grotjahn, Y. Gong, L. N. Zakharov, J. A. Golen, A. L. Rheingold, *J. Am. Chem. Soc.* **2006**, *128*, 438–453; b) D. B. Grotjahn, Y. Gong, A. G. DiPasquale, L. N. Zakharov, A. L. Rheingold, *Organometallics* **2006**, *25*, 5693–5695.
- [17] Complex **1** exhibits no dynamic behavior in the temperature range of 25–(–80)°C as evidenced by the ¹H and ³¹P{¹H} NMR spectra in [D₈]THF (see the Supporting Information).
- [18] In contrast to **1**, oxidative addition palladium complexes bearing related biphenyl ligands show equilibrium between two forms, where one of them exhibits interaction of palladium atom and second arene ring of the biphenyl ligand, see reference [14] and

- M. Yamashita, I. Takamiya, K. Jin, K. Nozaki, *J. Organomet. Chem.* **2006**, 691, 3189–3195.
- [19] For studies of using stoichiometric amounts of arylpalladium alkoxide complexes for reductive elimination of aryl ethers, see: a) G. Mann, C. Incarvito, A. L. Reingold, J. F. Hartwig, *J. Am. Chem. Soc.* **1999**, 121, 3224–3225; b) G. Mann, Q. Shelby, A. H. Roy, J. F. Hartwig, *Organometallics* **2003**, 22, 2775–2789.
- [20] Y. Pan, G. B. Young, *J. Organomet. Chem.* **1999**, 577, 257–264.
- [21] Indeed, dibenzylideneacetone is more strongly bound to Pd⁰ compared to 1,5-cyclooctadiene. Palladium (0) complexes of dibenzylideneacetone ([Pd(dba)₂] and [Pd₂(dba)₃]) are stable and commercially available, whereas the corresponding complex with 1,5-cyclooctadiene ([Pd(cod)₂]) decomposes above –30 °C: a) M. Green, J. A. K. Howard, J. L. Spencer, F. G. A. Stone, *J. Chem. Soc. Chem. Commun.* **1975**, 449–451; b) E. Sturm, H. Reinhard, K.-J. Kreutz (Air Products & Chemicals Inc.), DE2555374A1, **1975**.
- [22] See for example: a) C. Amatore, A. Jutand, *Coord. Chem. Rev.* **1998**, 178–180, 511–528; b) C. Amatore, A. Jutand, F. Khalil, M. A. M'Barki, L. Mottier, *Organometallics* **1993**, 12, 3168–3178.
- [23] The reason that these complexes were not previously used as palladium precursors can hardly be explained by their limited thermal stability or sensitivity to air, since “as dry solids, these complexes survive for several hours at ambient temperature in air, without evidence of decomposition. At –20 °C under argon, they can be stored indefinitely” (see reference [20]).
- [24] Recently, Buchwald et al. reported an alternative phosphine ligated palladium pre-catalyst which forms PdL species at low temperature (–40 °C). However, multistep synthesis is required for the each of these phosphine-containing precursors: M. R. Biscoe, B. P. Fors, S. L. Buchwald, *J. Am. Chem. Soc.* **2008**, 130, 6686–6687.
- [25] R. Martin, S. L. Buchwald, *Acc. Chem. Res.* **2008**, 41, 1461–1473.
- [26] Whereas the hydroxylation of 4-bromo-3-methylbenzotrile using 8 mol% of palladium afforded the phenol in 68% yield, the corresponding reaction with less reactive 4-bromobenzotrile gave no desired product; see reference [8].
- [27] J. F. Hartwig, *Acc. Chem. Res.* **1998**, 31, 852–860.
-