

## Original article

Silica gel catalyzed  $\alpha$ -bromination of ketones using *N*-bromosuccinimide: An easy and rapid methodBodireddy Mohan Reddy<sup>a</sup>, Velpula Venkata Ramana Kumar<sup>a</sup>, Nallagonda Chinna Gangi Reddy<sup>a,\*</sup>, Siripragada Mahender Rao<sup>b</sup><sup>a</sup> Department of Chemistry, School of Physical Sciences, Yogi Vemana University, Kadapa 516 003, India<sup>b</sup> Research & Development (API) Centre, Panacea Biotech Ltd., B-1 Extn./A-27, Mohan Co-operative Industrial Estate, New Delhi 110 044, India

## ARTICLE INFO

## Article history:

Received 28 April 2013

Received in revised form 29 July 2013

Accepted 30 August 2013

Available online 5 November 2013

## Keywords:

Heterogeneous catalysis

 $\alpha$ -Bromination*N*-bromosuccinimide (NBS)

Silica gel

Ketones

## ABSTRACT

An easy and rapid method for the  $\alpha$ -bromination of ketones using *N*-bromosuccinimide (NBS) catalyzed by silica gel in methanol under reflux conditions was developed. The expected products were formed in excellent isolated yields within a short period of time (5–20 min). Major advantages of the present procedure include use of inexpensive and readily available catalyst, exclusion of pre- and post-chemical treatment of catalyst and use of methanol as solvent instead of ethers and chlorinated solvents.

© 2013 Nallagonda Chinna Gangi Reddy. Published by Elsevier B.V. on behalf of Chinese Chemical Society. All rights reserved.

## 1. Introduction

The  $\alpha$ -bromination of ketones has attracted much attention because the resulting  $\alpha$ -bromination of ketones are the key intermediates in organic synthesis [1,2]. Accordingly, a number of protocols have been developed which include cupric bromide [3], dioxane dibromide [4], tetra-butyl ammonium tribromide [5], H<sub>2</sub>O<sub>2</sub>-HBr [6], bromodimethyl sulfoniumbromide [7], ethylene bis(*N*-methyl imidazolium) ditribromide [8], trihaloisocyanuric acids [9], pyridinium bromochromate [10] and NH<sub>4</sub>Br-oxone [11]. Moreover, owing to user friendliness and availability, *N*-bromosuccinimide has been extensively used as brominating agent: transformations can be modulated under various reaction conditions, for example: (i) using different types of catalysts such as Mg(ClO<sub>4</sub>)<sub>2</sub> [12], NH<sub>4</sub>OAc [13], Amberlyst-15 [14], silica-supported NaHSO<sub>4</sub> [15], silica supported NaHCO<sub>3</sub> [16], sulfonic acid functionalized silica [17] and FeCl<sub>3</sub> [18]; (ii) ionic liquids [19], (iii) photochemical bromination [20]; (iv) sonochemical bromination [21] and (v) solvent free reaction conditions (SFRC) [22].

An extensive literature search revealed that bromination with NBS depends on the catalyst used and this encouraged researchers to develop a number of catalysts to make the bromination process


easier and more efficient. Silica gel is known as a good catalyst in many synthetic applications [23,24] and also for ring bromination [25–27]; However, no reports were found on the investigation of silica gel as catalyst for the  $\alpha$ -bromination of ketones using NBS. As a part of continuing agenda to explore the use of inorganic materials in the synthesis of organic compounds, we found that silica gel is an efficient catalyst for the  $\alpha$ -bromination of ketones when *N*-bromosuccinimide in methanol is used under reflux conditions (Scheme 1).

## 2. Experimental

Ketones were purchased from Acros, Merck and S.D. Fine Chemicals Ltd., Mumbai, India. *N*-bromosuccinimide and silica gel (60–120 mesh) were procured from Merck, India. Methanol (99.0%) and ethyl acetate were purchased from Finar Chemicals Ltd., India. Double distilled water was used for work-up. Melting points were determined in open capillaries on a Mel-Temp apparatus and are uncorrected. The <sup>1</sup>H NMR spectra were recorded on a Varian 300 MHz. Chemical shifts were expressed in parts per million (ppm). Splitting patterns describe apparent multiplicities and are designated as s (singlet), d (doublet), t (triplet), q (quartet), m (multiplet) or br (broad). Mass spectra (MS) were recorded on Agilent, model-6410, triple quad LCMS. Thin-layer chromatography was performed using 0.25 mm Merck silica gel plates (60F-254) and visualized with UV light. Column chromatography was performed on silica gel of 60–120 mesh, Merck).

\* Corresponding author. C.G.R. Nallagoudu.

E-mail address: [ncgreddy@yogivemanauniversity.ac.in](mailto:ncgreddy@yogivemanauniversity.ac.in) (N. Chinna. Gangi Reddy).



**Scheme 1.** The  $\alpha$ -bromination of ketones using NBS in presence of silica gel in methanol.

Column chromatographic grade silica gel (60–120 mesh) supplied by Merck, India was employed in the present investigation. The pH value of 10% aqueous suspension of silica gel is  $\sim$ 6.5–7.5. Silica gel catalyst was used as it is, without any activation or any further treatment or chemical modifications.

### 2.1. Typical experimental procedure

The  $\alpha$ -bromination reaction was carried out using acetophenone (1200 mg, 10 mmol), N-bromosuccinimide (2136 mg, 12 mmol), 10% (w/w) silica gel (120 mg) in 10 mL of methanol at reflux conditions until the disappearance of the substrate. (Note: 2136 mg of N-bromosuccinimide was added portion wise *i.e.* 356 mg for each time in six portions). The progress of the reaction was monitored by TLC. The reaction mass was filtered after the completion of the reaction as per TLC and the catalyst was collected for reuse. The filtrate was concentrated under vacuum. Double distilled water was added to the reaction mixture and quenched with aqueous sodium thiosulfate and the product extracted with dichloromethane (Caution: Severe burning sensation of eyes was observed during the work-up process). The layers were separated and the organic layer was collected and washed thrice with distilled water ( $3 \times 50$  mL). The collected organic layer was dried over anhydrous  $\text{Na}_2\text{SO}_4$ , filtered and concentrated. The obtained crude product was purified by column chromatography over silica gel (60–120 mesh) using *n*-hexane–EtOAc (99:1 ratio).

With the aim of studying the recycling of the catalyst, the isolated catalyst was washed with ethyl acetate (5 mL) after its filtration from the reaction medium, collected and dried in vacuum at 70 °C to a constant weight. Subsequently it was reused for the  $\alpha$ -bromination of acetophenone and achieved 95%, 86% and 83% yields of product (**2a**) for first, second and third reuse of catalyst respectively. All products gave spectroscopic data in agreement with the literature [15,21,27–30].

The method is also very practical for scale up in process development. We attempted large scale (100 gram scale) synthesis of 2-bromo-1-phenylethanone **2a** and obtained fruitful results with isolated yields ranging from 93% to 96%.

### 2.2. General experimental procedure for large scale (100 g scale) preparation

The  $\alpha$ -bromination reaction was carried out using acetophenone (100 g, 0.832 mol), N-bromosuccinimide (213.6 g, 1.2 mol), 10% (w/w) silica gel (10 g) in 1 L of methanol under reflux conditions until the disappearance of the substrate (Note: 213.6 g of N-bromosuccinimide was added portion wise *i.e.* 35.6 g for each time in six portions). Excess N-bromosuccinimide (+0.1 mol) was added to complete the reaction as per TLC. The reaction mass was filtered, after the completion of the reaction as per TLC and the catalyst was collected for reuse. The solvent was removed and double distilled water was added to the reaction mixture and was quenched with aqueous sodium thiosulfate. The product was extracted with dichloromethane. (Caution: Severe burning sensation of eyes was observed during the work-up process). The layers

were separated and the organic layer was collected and it was washed with distilled water for three times ( $3 \times 1$  L). The collected organic layer was dried over anhydrous  $\text{Na}_2\text{SO}_4$  and filtered, concentrated and purified by recrystallization using *n*-hexane (300 mL). The resulting yield of pure  $\alpha$ -brominated ketone, *i.e.* 2-bromo-1-phenylethanone (**2a**) was 94%. The same experiment was repeated twice and the yields were 93% and 96%, respectively. The reproducibility of the protocol was thus confirmed with the consistent isolated yields of desired product 2-bromo-1-phenylethanone (**2a**).

## 3. Results and discussion

To study the effect of solvents and temperature on the  $\alpha$ -bromination of ketones, we carried out a reaction with acetophenone using N-bromosuccinimide in methanol in presence of silica gel at room temperature. However a low yield (56%, Table 1 entry 1) of the product 2-bromo-1-phenylethanone (**2a**) was obtained after 4 h. When the same reaction was carried out under reflux conditions (Table 1, entry 1), the yield was 95% within 18 min. Other solvents such as  $\text{Et}_2\text{O}$ , THF,  $\text{CH}_3\text{CN}$ ,  $\text{CH}_2\text{Cl}_2$  and  $\text{CHCl}_3$  were also studied, but lower yields of the desired products were obtained (10–35%) as shown in Table 1. With  $\text{Et}_2\text{O}$ , THF, 1,4-dioxane and acetonitrile, lower yields of the product **2a** was formed, along with very low amounts of the  $\alpha$ ,  $\alpha'$ -dibrominated

**Table 1**  
Effect of solvent and temperature on the formation of  $\alpha$ -brominated product (**2a**).<sup>a</sup>

Entry	Solvent	Time (min)	Major product	Yield (%) <sup>b</sup>	Yield (%) <sup>c</sup>	Selectivity ( <b>2a:3a:4a</b> )
1	MeOH	18	<b>2a</b>	95	56	95:03:00
2	$\text{Et}_2\text{O}$	15	<b>4a</b>	24	18	24:04:50
3	THF	18	<b>4a</b>	18	15	18:03:45
4	1,4-Dioxane	28	<b>4a</b>	16	14	16:05:42
5	$\text{CH}_3\text{CN}$	180	<b>2a</b>	35	22	35:14:28
6	$\text{CH}_2\text{Cl}_2$	1500	<b>2a</b>	12	8	12:01:00
7	$\text{CHCl}_3$	1500	<b>2a</b>	14	7	14:04:00
8 <sup>d</sup>	MeOH	1500	–	0	0	0

<sup>a</sup> Reactions were conducted with acetophenone (10 mmol); NBS (12 mmol), 10% (w/w) silica gel, solvent (10 mL) under reflux conditions.

<sup>b</sup> Isolated yields under reflux condition.

<sup>c</sup> Isolated yields at room temperature.

<sup>d</sup> Reaction carried out without catalyst.

**Table 2**  
Role of catalyst load on the formation of  $\alpha$ -brominated product (**2a**).<sup>a</sup>

Entry	Catalyst (w/w %)	Time (min)	Major Product	Yield (%) <sup>b</sup>	Selectivity (%) ( <b>2a:3a</b> )
1	1	90	<b>2a</b>	41	41:03
2	5	60	<b>2a</b>	65	65:02
3	10	18	<b>2a</b>	95	95:03
4	15	20	<b>2a</b>	82	82:14
5	20	15	<b>2a</b>	74	74:18
6	30	11	<b>2a</b>	67	67:21

<sup>a</sup> Reactions were conducted with acetophenone (10 mmol); NBS (12 mmol), 10% (w/w) silica gel, methanol (10 mL) under reflux conditions.

<sup>b</sup> Isolated yields.

product (**3a**) and interestingly, the nuclear brominated product **4a** was formed as the major product (Table 1, entries 2–4). In chlorinated solvents, the formation of the desired product **2a** was very low. This may be attributed to the lack of solubility of NBS in these solvents (Table 1, entries 6–7). The reaction did not proceed in the absence of catalyst (Table 1, entry 8). Solvent screening experiments revealed that methanol was the best solvent for the silica gel catalyzed  $\alpha$ -bromination. For further studies, methanol was used as a solvent to optimize other reaction parameters.


We studied the time course of the  $\alpha$ -bromination reaction and subsequent effect on yield of the product (**2a**) using acetophenone as a model substrate in the presence of different weight percentage (w/w) of silica gel in methanol under reflux conditions with the acquired results displayed in Table 2. When 1% of catalyst was

used, 41% yield of the product (**2a**) was obtained in 90 min. Upon use of 5% of catalyst, a 65% yield was observed in 60 min and with 10% of catalyst, the obtained yield reached the maximum of 95% in 18 min. On the other hand with 15, 20 and 30% of catalyst, 82, 74 and 67% yields were obtained, respectively and the reaction was completed within 10–15 min. Low catalyst load resulted in the existence of high percentage of unreacted substrate (Table 2, entries 1 and 2) even with long reaction times. In contrast, an increase of catalyst load caused a decrease in the isolated yield of the desired product **2a** and a subsequent increase of  $\alpha,\alpha'$ -dibrominated product (**3a**) formation (Table 2, entries 4–6), but the reaction was rapidly completed.

To further examine the selectivity of this silica gel catalyzed  $\alpha$ -bromination reaction, different substituted acetophenones,

**Table 3**

The  $\alpha$ -bromination of various ketones using NBS in presence of silica gel.<sup>a</sup>



R = Acyclic or cyclic alkyl groups, Aryl, Naphthyl

Entry	Substrate	Product	Time (min)	Yield(%) <sup>b</sup>	Entry	Substrate	Product	Time (min)	Yield (%) <sup>b</sup>
1			18	95 [28]	12			15	47 [28]
2			20	71 [28]	13			18	45 [29]
3			17	90 [28]	14			21	82 [28]
4			15	93 [28]	15			18	87 [28]
5			17	90 [28]	16			12	76 [15]
6			16	88 [28]	17			10	22 [15]
7			14	87 [28]	18			10	14 [30]
8			22	80 [28]	19			10	83 [30]
9			19	91 [28]	20			7	85 [22]
10			13	64 [28]	21			5	89 [22]
11			17	42 [28]	22			6	91 [31]

<sup>a</sup> Reactions were conducted with ketone (**1**) (10 mmol); NBS (12 mmol), 10% (w/w) silica gel, methanol (10 mL) under reflux conditions.

<sup>b</sup> Isolated yields.

acenaphthones, acyclic and cyclic ketones were employed. In the  $\alpha$ -bromination reaction of acetophenone, a high yield (95%) of product (**2a**) was observed (Table 3, entry 1). The  $\alpha$ -bromination of substituted acetophenones gave moderate to high yields, regardless of the electronic influence of substituents (Table 3, entries 2–13), with the exception of *m*-nitroacetophenone, *p*-nitroacetophenone and *p*-cyanoacetophenone (Table 3, entries 11–13) wherein low yields of products **2k**, **2l** and **2m** were obtained respectively. This may be due to the presence of strong electron withdrawing groups on the aromatic ring. The products, **2k**, **2l** and **2m** were stable during reaction conditions as per TLC, but showed only moderate stability under the work-up process. However, after isolation, compounds (**2k-m**) were preserved under cooling (0–5 °C) to prevent decomposition. Compounds (**2k-m**) were soluble both in methanol and dichloromethane. Acenaphthones without substitution gave the corresponding products **2n** and **2o** in 82 and 87% yield (Table 3, entries 14 and 15), respectively. The  $\alpha$ -bromination of acyclic and cyclic ketones provided good yields of products within a short period of time (Table 3, entries 16–22).

#### 4. Conclusion

In summary, a new method was developed using silica gel as a catalyst for the  $\alpha$ -bromination of ketones (acyclic, cyclic and aralkyl ketones) in methanol utilizing NBS under reflux conditions. Thus, our new bromination reaction proceeded smoothly and afforded moderate to excellent isolated yields under optimized reaction conditions. The catalyst was efficiently recovered and reused for 3 times without loss of activity. Major advantages of the present protocol include shorter reaction times, user friendly process, simple work-up and exclusion of pre- and post-chemical treatment of catalyst, improved quality and appearance of the isolated products. In addition, the present method is highly useful for large scale ( $\geq 100$  g scale) applications both in chemical and pharmaceutical industries.

#### Acknowledgment

We express our thanks to Council of Scientific and Industrial Research (CSIR), New Delhi, Government of India for the financial support under major research project (No. 01 (2391)/10/EMR-II).

#### References

- [1] R.C. Larock, *Comprehensive Organic Transformations*, 2nd ed., Wiley-VCH, New York, 1999, p. 715.
- [2] For review see: A.W. Erian, S.M. Sherif, H.M. Gaber, The chemistry of  $\alpha$ -haloketones and their utility in heterocyclic synthesis, *Molecules* 8 (2003) 93–865.
- [3] L.C. King, G.K. Ostrum, Selective bromination with copper (II) bromide, *J. Org. Chem.* 29 (1964) 3459–3461.
- [4] S.J. Pasaribu, L.K. Williams, Selective bromination of substituted acetophenones with dioxan dibromide, *Aust. J. Chem.* 26 (1973) 1327–1331.
- [5] S. Kajigaeshi, T. Kakinami, T. Okamoto, S. Fujisaki, Synthesis of bromoacetyl derivatives by use of tetrabutylammonium tribromide, *Bull. Chem. Soc. Jpn.* 60 (1987) 1159–1160.
- [6] J. Ju, Y.J. Li, J.R. Gao, et al., High selectively oxidative bromination of toluene derivatives by the  $H_2O_2$ -HBr system, *Chin. Chem. Lett.* 22 (2011) 382–384.
- [7] A.T. Khan, M.A. Ali, P. Goswami, L.H. Choudhury, A mild and regioselective method for  $\alpha$ -bromination of  $\beta$ -Keto esters and 1, 3-diketones using bromodimethylsulfonium bromide (BDMS), *J. Org. Chem.* 71 (2006) 8961–8963.

- [8] R. Hosseinzadeh, M. Tajbakhsh, M. Mohadjerani, Z. Lasemi, Ethylenebis(*N*-methylimidazolium) ditribromide (EBMIDTB): an efficient reagent for the monobromination of 1,3-diketones and  $\beta$ -ketoesters, *Monatsh. Chem.* 140 (2009) 57–60.
- [9] G.F. Mendonça, H.C. Sindra, L.S. de Almeida, P.M. Esteves, M.C.S. de Mattos, Trihaloisocyanuric acids as convenient reagents for regioselective halogenation of  $\beta$ -dicarbonyl compounds, *Tetrahedron Lett.* 50 (2009) 473–475.
- [10] Y. Sarrafi, M. Sadatshahabi, K. Alimohammadi, A mild, simple and efficient method for selective  $\alpha$ -monobromination of 1,3-diketones and  $\beta$ -keto-esters using pyridinium bromochromate, *Chin. Chem. Lett.* 20 (2009) 393–396.
- [11] M. Arun Kumar, C.N. Rohitha, M. Mahender Reddy, P. Swamy, N. Narender, Oxidative bromination of ketones using ammonium bromide and oxone, *Tetrahedron Lett.* 53 (2012) 191–195.
- [12] D. Yang, Y. Yan, B. Lui, Mild  $\alpha$ -halogenation reactions of 1,3-dicarbonyl compounds catalyzed by Lewis acids, *J. Org. Chem.* 67 (2002) 7429–7431.
- [13] K. Tanemura, T. Suzuki, Y. Nishida, K. Satsumabayashi, T. Horaguchi, A mild and efficient procedure for  $\alpha$ -bromination of ketones using *N*-bromosuccinimide catalysed by ammonium acetate, *Chem. Commun.* (2004) 470–471.
- [14] H.M. Meshram, P.N. Reddy, K. Sadashiv, J.S. Yadav, Amberlyst-15<sup>®</sup>-promoted efficient 2-halogenation of 1,3-keto-esters and cyclic ketones using *N*-halosuccinimides, *Tetrahedron Lett.* 46 (2005) 623–626.
- [15] B. Das, K. Venkateswarlu, G. Mahender, I. Mahender, A simple and efficient method for  $\alpha$ -bromination of carbonyl compounds using *N*-bromosuccinimide in the presence of silica-supported sodium hydrogen sulfate as a heterogeneous catalyst, *Tetrahedron Lett.* 46 (2005) 3041–3044.
- [16] A. Rahman, S.B. Jonnalagadda, Simple and efficient system for the  $\alpha$ -bromination of a  $\beta$ -ketoester by using *N*-bromosuccinimide in the presence of silica supported  $NaHCO_3$  as the heterogeneous catalyst: an environmentally benevolent approach, *Synth. Commun.* 42 (2012) 1091–1100.
- [17] B. Das, K. Venkateswarlu, H. Holla, M. Krishnaiah, Sulfonic acid functionalized silica: a remarkably efficient heterogeneous reusable catalyst for  $\alpha$ -monobromination of carbonyl compounds using *N*-bromosuccinimide, *J. Mol. Catal.* 253 (2006) 107–111.
- [18] H. Jin, Z.D. Huang, C.X. Kuang, X.K. Wang, Iron-catalyzed bromination of aryl azides by *N*-bromosuccinimide: efficient method for the synthesis of brominated aryl azides, *Chin. Chem. Lett.* 22 (2011) 310–313.
- [19] H.M. Meshram, P.N. Reddy, K. Vishnu, K. Sadashiv, J.S. Yadav, A green approach for efficient  $\alpha$ -halogenation of  $\beta$ -dicarbonyl compounds and cyclic ketones using *N*-halosuccinimides in ionic liquids, *Tetrahedron Lett.* 47 (2006) 991–995.
- [20] S.S. Arbu, S.B. Waghmode, A.V. Ramaswamy, Photochemical  $\alpha$ -bromination of ketones using *N*-bromosuccinimide: a simple, mild and efficient method, *Tetrahedron Lett.* 48 (2007) 1411–1415.
- [21] M.V. Adhikari, S.D. Samant, Sonochemical bromination of acetophenones using *p*-toluenesulfonic acid – *N*-bromosuccinimide, *Ultrason. Sonochem.* 9 (2002) 107–111.
- [22] I. Pravst, M. Zupan, S. Stavber, Halogenation of ketones with *N*-halosuccinimides under solvent-free reaction conditions, *Tetrahedron* 64 (2008) 5191–5199.
- [23] M. Hashmat Ali, M. McDermott, Oxidation of thiols to disulfides with molecular bromine on hydrated silica gel support, *Tetrahedron Lett.* 43 (2002) 6271–6273.
- [24] E. Selcuk, S. Mehmet Emin, S. Ertan, D. Arif, Bromination of 2,3-dihydrobenzobenzene and synthesis of its mono- and dibromide derivatives: unexpected Wagner-Meerwein rearrangement on silica gel, *Turk. J. Chem.* 35 (2011) 587–598.
- [25] R.K. Sharma, C. Sharma, Oxidative bromination reaction using  $Cu^{2+}$ -perfluorophthalocyanine-immobilized silica gel catalyst under mild reaction conditions, *Tetrahedron Lett.* 51 (2010) 4415–4418.
- [26] K. Smith, M. James, A.G. Mistry, M.R. Bye, D.J. Faulkner, A new method for bromination and polybromination of carbazoles,  $\beta$ -carboline and iminodibenzyls by using *N*-bromosuccinimide and silica gel, *Tetrahedron* 48 (1992) 7479–7488.
- [27] H. Alinezhad, M. Tajbakhsh, M. Zare, One-pot regioselective synthesis of 4-bromopyrazole derivatives under solvent free conditions, *J. Mex. Chem. Soc.* 55 (2011) 238–241.
- [28] R.B. Mohan, N.C. Gangi Reddy, Regioselective  $\alpha$ -bromination of aralkylketones using *N*-bromosuccinimide in presence of Montmorillonite K-10 clay: a simple and efficient method, *Synth. Commun.* 43 (2013) 2603–2614.
- [29] A. Tsuruoka, Y. Kaku, H. Kakinuma, et al., Synthesis and antifungal activity of novel thiazole-containing triazole antifungals. II. Optically active ER-30346 and its derivatives, *Chem. Pharm. Bull.* 46 (1998) 623–630.
- [30] W. Peter, B. Joachim, Terpenes and terpene derivatives, synthesis and olfactive properties of (–)- and *rac*-siphiperfol-5-en-3-ol and of some tris-nor derivatives, *Liebigs Ann. Chem.* 1992 (1992) 669–678.
- [31] R.D. Patil, G. Joshi, S. Adimurthy, B.C. Ranu, Facile one-pot synthesis of  $\alpha$ -bromoketones from olefins using bromide/bromate couple as a nonhazardous brominating agent, *Tetrahedron Lett.* 50 (2009) 2529–2532.