

Communication

Highly Selective gem-Difluoroallylation of Organoborons with Bromodifluoromethylated Alkenes Catalyzed by Palladium

Qiao-Qiao Min, Zengsheng Yin, Zhang Feng, Wen-Hao Guo, and Xingang Zhang

J. Am. Chem. Soc., **Just Accepted Manuscript** • Publication Date (Web): 07 Jan 2014

Downloaded from <http://pubs.acs.org> on January 8, 2014

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



Highly Selective *gem*-Difluoroallylation of Organoborons with Bromodifluoromethylated Alkenes Catalyzed by Palladium

Qiao-Qiao Min, Zengsheng Yin, Zhang Feng, Wen-Hao Guo, and Xingang Zhang*

Key Laboratory of Organofluorine Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 345 Lingling Road, Shanghai 200032, China

Supporting Information Placeholder


ABSTRACT: A first example of Pd-catalyzed *gem*-difluoroallylation of organoborons using 3-bromo-3,3-difluoropropene (BDFP) in high efficiency with high α/γ -substitution regioselectivity has been developed. The reaction can also be extended to substituted BDFPs, and has advantages of low catalyst loading (0.8 mol % to 0.01 mol %), broad substrate scope, and excellent functional group compatibility, thus providing a facile route for practical application in drug discovery and development.

The importance of fluorinated compounds in agrochemicals, pharmaceuticals, and materials science¹ has triggered an explosion of research efforts in developing new and efficient methods to introduce fluorinated functional groups into organic molecules. Over the past few years, although considerable progresses have been achieved in fluorination and trifluoromethylation of organic substrates,² to date, strategies for introduction of a difluoromethylene (CF₂) group into organic molecules have been less explored³ despite the unique properties and important biological activity of compounds containing CF₂.⁴ For instance, CF₂ is usually considered as a bioisostere of the oxygen or a carbonyl group,⁵ which leads to increased dipole moments, enhanced acidity of its neighboring group and conformational changes.⁶ In particular, the introduction of a CF₂ onto the aromatic rings can dramatically improve the metabolic stability and oral bioavailability of biological active compounds.⁷ Generally, such a structural moiety is achieved by conversion of carbonyl group with DAST or Deoxofluor.⁸ But the intrinsic limitations of this method, such as the important functional groups incompatibility and the use of expensive and toxic fluorinated reagents, significantly restrict its widespread synthetic applications. To address these issues, one alternative approach is to directly introduce a functionalized CF₂ group into the organic molecules catalyzed by transition-metal. However, these difluoroalkylation processes still remain challenging, and only a handful of examples have been developed so far.^{9,10}

As part of ongoing study in transition-metal-catalyzed reactions for introduction of fluorinated functional groups into organic molecules,¹¹ herein we demonstrate the feasibility of Pd-catalyzed *gem*-difluoroallylation. The *gem*-difluoroallyl group (CF₂CH=CH₂) is extremely appealing due to the versatile synthetic utility of the carbon-carbon double bond. However, synthetical-

ly useful methods to access such valuable structures are rare.¹² We chose 3-bromo-3,3-difluoropropene (BDFP) as the starting reagent for *gem*-difluoroallylation because it is readily and commercially available,¹³ and a transition-metal-catalyzed cross-coupling would thus provide a cost-efficient access to *gem*-difluoroallylated structures that can be applied in the synthesis of bioactive compounds. In this study, we focused our research on addressing three crucial issues: (1) regiochemical selectivity, *i.e.*, α -substitution *vs* γ -substitution (Scheme 1a), (2) efficient catalytic system that can suppress undesired defluorination side reaction resulted from *gem*-difluoroallylated products¹⁴ (Scheme 1b), and (3) efficient and practical processes with broad substrate scope. As a result, we disclose the first example of highly selective Pd-catalyzed *gem*-difluoroallylation of organoborons with BDFP. The reaction can also be extended to substituted BDFPs. The notable advantages of this reaction are its high efficiency and regioselectivity (α/γ up to >37:1), low catalyst loading (0.8 mol % to 0.01 mol % Pd), applicability for large scale production (10 gram), broad substrate scope, and excellent functional group compatibility.


Scheme 1. Transition-Metal-Catalyzed Cross-Coupling Between 3-Bromo-3,3-difluoropropene and Organometallics.



We began our study with the reaction of air stable 4-(*tert*-butyl)phenylboronic acid **1a** and 3-bromo-3,3-difluoropropene **2** in the presence of Pd-catalyst (Table 1). After screening the reaction conditions, we found a 77% yield of a mixture of **3a** and **4a** with α -substitution **3a** as major product (**3a/4a** = 6.2:1) was obtained when the reaction was carried out with **1a** (1 mmol), **2** (2 equiv), Cs₂CO₃ (1.5 equiv), Pd(OAc)₂ (5 mol%), and PPh₃ (10 mol%) in dioxane at 80 °C (Table 1, entry 1). Notably, no defluorination of **3a** was observed under these reaction conditions. This is in sharp contrast to previous results, in which the defluorination of *gem*-difluoroallylated compounds frequently occurred.¹⁴ The reaction was also found very sensitive to the bases (Table 1, en-

tries 2-5) and solvents. K_2CO_3 and dioxane were the optimal choice. Other bases and solvents led to either lower yields or poor regioselectivities. No product was obtained when polar solvents, such as DMF, DMSO, and CH_3CN , were used due to the formation of uncertain byproducts from BDFP (see Supporting Information). To our delight, a low catalyst loading of $Pd(PPh_3)_4$ (0.2 mol %) showed a beneficial effect on both reaction efficiency and regioselectivity (94% yield, $3a/4a = 10:1$) (Table 1, entry 8). Switching $Pd(PPh_3)_4$ to $Pd_2(dba)_3$ dramatically increased the α/γ ($3a/4a$) regioselectivity ($3a/4a = 20:1$) (Table 1, entry 9). Importantly, even the amount of catalyst loading was reduced to 0.05 mol % in combination of 1.2 equiv of **2**, high yield of **3a** was still afforded while maintaining high levels of regioselectivity (Table 1, entry 11). Finally, the optimal reaction conditions were identified by increasing the amount of K_2CO_3 to 3.0 equiv with utilization of 0.48 equiv of H_2O (Table 1, entry 12), providing **3a** in high yield with excellent regioselectivity ($3a/4a = 20:1$) (for details see Supporting Information). It is noteworthy that this reaction could be successfully conducted under air in wet solvent, thus featuring an operational simplicity advantage of the present process. It should be mentioned that even BDFP **2** was treated with 2.5 equiv of **1a** under optimized reaction conditions (Table 1, entry 12), only 5% yield (determined by ^{19}F NMR) of defluorinated side product fluoroolefin was observed, and 90% yield of desired product **3a** still could be obtained with high α/γ regioselectivity ($3a/4a > 20:1$) (for details see Supporting Information).

Table 1. Representative Results for Optimization of Pd-Catalyzed Cross-Coupling between (4-(*tert*-Butyl)phenyl)boronic Acid **1a and 3-Bromo-3,3-difluoropropene **2**.^a**




entry	[Pd] (mol %)	Base (equiv)	3a+4a Yield (%), 3a/4a ^b
1 ^c	$Pd(OAc)_2$ (5)	CS_2CO_3 (1.5)	77, 6.2:1
2 ^c	$Pd(OAc)_2$ (5)	K_2CO_3 (1.5)	78, 6.4:1
3 ^c	$Pd(OAc)_2$ (5)	Na_2CO_3 (1.5)	26, 5.9:1
4 ^c	$Pd(OAc)_2$ (5)	KOAc (1.5)	11, only 3a
5 ^c	$Pd(OAc)_2$ (5)	K_3PO_4 (1.5)	28, 5.9:1
6	$Pd(PPh_3)_4$ (5)	K_2CO_3 (1.5)	75, 3.2:1
7	$Pd(PPh_3)_4$ (0.4)	K_2CO_3 (1.5)	97, 9.4:1
8	$Pd(PPh_3)_4$ (0.2)	K_2CO_3 (1.5)	94, 10:1
9	$Pd_2(dba)_3$ (0.4)	K_2CO_3 (1.5)	89, 20:1
10 ^d	$Pd_2(dba)_3$ (0.1)	K_2CO_3 (1.5)	77, 20:1
11 ^d	$Pd_2(dba)_3$ (0.05)	K_2CO_3 (1.5)	73, 20:1
12 ^e	$Pd_2(dba)_3$ (0.4)	K_2CO_3 (3.0)	92 (93), 20:1

^aReaction conditions (unless otherwise specified): **1a** (1.0 mmol, 1.0 equiv), **2** (1.5 equiv), dioxane (5 mL), 80 °C, 24 h under N_2 . ^bDetermined by ^{19}F NMR using trifluoromethylbenzene as internal standard and number in parenthesis is isolated yield. The ratio of **3a/4a** was determined by ^{19}F NMR before working up. ^cUsing 2.0 equiv of **2**, 10 mol % PPh_3 . ^dUsing 1.2 equiv of **2**. ^eUsing 0.48 equiv of H_2O , and reaction was conducted under air.

To ascertain the substrate scope of this method, a wide range of aryl boronic acids were then examined (Table 2). Generally, aro-

matic boronic acids bearing either electron-rich or electron-deficient substituents all led to **3** in high yields with high to excellent α/γ regioselectivities (**3a-m**). Importantly, in many cases the $Pd_2(dba)_3$ loading can be reduced to 0.2 mol % without compromising the reaction efficiency and α/γ regioselectivities (**3g-i**, **3k-m**). However, in the case of 3-nitrophenyl boronic acid, a dramatically decreased yield (44% determined by ^{19}F NMR) of **3j** was observed. Further improvement of the reaction efficiency by employing $Pd(PPh_3)_4$ (0.4 mol %) in conjunction with catalytic amount of CuI (6 mol %) provided **3j** in good yield (70%) with high regioselectivity (**3j/4j** = 13:1). The steric effect of the aromatic boronic acids did not interfere with the reaction. The phenyl substituent at the *para*, *meta*, and *ortho* position of the phenyl boronic acid afforded the corresponding products **3** in high efficiency with high α/γ regioselectivity (**3b-d**). Notably, even sterically hindered mesitylboronic acid previously demonstrated to be a challenging substrate for fluoroalkylation was also a competent coupling partner (**3f**), thus demonstrating the generality of the present reaction.

Table 2. *gem*-Difluoroallylation of Organoboronic Acids **1 with 3-Bromo-3,3-difluoropropene **2**.^a**



3a 93% (20:1)	3b 85% (16:1)	3c 87% (16:1)	3d 83% (15:1)	3e 92% (18:1)
3f ^b 78% (17:1)	3g ^c 87% (28:1)	3h ^c 71% (24:1)	3i ^c 82% (16:1)	3j ^{d,e} 70% (13:1)
3k ^e 78% (13:1)	3l ^e 60% (10:1)	3m ^e 80% (15:1)	3n ^b 68% (19:1)	3o ^c 82% (28:1)
3p ^b 90% (20:1)	3q 67% (12:1)	3r 74% (20:1)	3s 93% (17:1)	3t 60% (8:1)
3u ^c 90% (20:1)	3v ^f 72% (8:1)	3w ^b 92% (26:1)	3x ^b 72% (15:1)	


^aReaction conditions (unless otherwise specified): **1** (1 mmol), **2** (1.5 mmol), dioxane (5 mL) under air for 24 h. All reported yields are combined isolated yields of **3** and **4**. The ratio of **3/4** in parentheses was determined by ^{19}F NMR before working up. ^b1.5 mmol of **1** and 1.0 mmol of **2** were used. ^c0.2 mol % of $Pd_2(dba)_3$ was used. ^d0.4 mol % of $Pd(PPh_3)_4$ and 6 mol % of CuI were used. ^e2.0 mmol of **2** was used. ^f0.2 mol % of $Pd(PPh_3)_4$ was used.

A variety of versatile functional groups (nitro, ester, aldehyde, alcohol, thioether, and vinyl) were quite well-tolerated (**3j-p**).

Most remarkably, the successfully selective formation of **3q-r** with intact bromide provided a good platform for further functionalization. Heteroaromatic rings are a prominent structural motif found in numerous pharmaceuticals and agrochemicals. Boronic acids derived from dibenzo[*b,d*]thiophene, carbazole, and dibenzo[*b,d*]furan all underwent *gem*-difluoroallylation smoothly (**3t-v**). Most importantly, *N*-containing heterocycle carbazole furnished its *gem*-difluoroallylated product **3u** in 90% yield with high regioselectivity (**3u/4u** = 20:1). However, pyridine-containing boronic acids were not suitable substrates. Furthermore, the reaction was not restricted to the (hetero)aryl boronic acids, vinyl boronic acids were also suitable substrates, and provided linear and branched *gem*-difluoromethylenated dienes in high efficiency with high to excellent α/γ regioselectivities (**3w-x**).

In spite of substantial progress in fluoroalkylation of organic molecules, efficient method that enables low catalyst loading to practically synthesize fluorinated compounds has received rare attention. In this context, grams scale synthesis of *gem*-difluoroallylated arenes **3** with utilization of 0.1 mol % to 0.01 mol % of Pd catalyst was conducted to demonstrate the synthetic utility of the protocol further (Scheme 2). As shown in Scheme 2a, gram-scale synthesis of **3o** and **3r** in the presence of 0.05 mol % Pd₂(dba)₃ proceeded smoothly with high α/γ regioselectivity, thus offering a reliable and practical access to highly functionalized difluoromethylenated structure. Most excitingly, a 10-gram-scale reaction with Pd-catalyst loading as low as 0.01 mol % afforded **3a** in 80% yield with high regioselectivity (α/γ = 10:1) (Scheme 2b). To the best of our knowledge, this is the first example of transition-metal-catalyzed fluoroalkylation of organic molecules by using as low as 0.01 mol % catalyst in a practical manner.


Scheme 2. Grams-Scale Synthesis of *gem*-Difluoroallylated Arenes with Low Catalyst Loading.^a



^aAll reported yields are combined isolated yields of **3** and **4**, and the number in parentheses is the ratio of **3/4**.

The substituted BDFPs¹⁵ were also applicable to the reaction. As shown in Table 3, branched BDFP derivatives including phenyl and alkyl-substituted substrates underwent the *gem*-difluoroallylation in high efficiency with excellent α/γ regioselectivity (**6a-g**). Again, important functional groups were compatible with the reaction conditions (**6c-e**, **6g**). It was found that the linear substituted BDFPs were more reactive than BDFP, and resulted in some undesired defluorinated compounds and other uncertain byproducts. To our delight, decreasing the reaction temperature and using Pd(PPh₃)₄ (0.4 mol %) as a catalyst could afford *gem*-difluoroallylated products in moderate yields (**6h-i**). Importantly, no branched products (γ -substitution) were observed during the reaction. We reasoned that this is probably due to the steric effect of the substituents. It should be pointed out that bisubstituted BDFP was also a suitable substrate, providing **6j** in synthetically useful yield. This is noteworthy, as it is difficult to prepare such fluorinated molecule otherwise.


Table 3. *gem*-Difluoroallylation of Organoboronic Acids 1 with Substituted BDFPs 5.^a



^aReaction conditions (unless otherwise specified): **1** (1.3 mmol), **5** (1.0 mmol), dioxane (5 mL) under air for 10 h. All reported yields are isolated yields, the number in parentheses is the ratio of α/γ . ^b0.4 mol % of Pd(PPh₃)₄ and 3.0 mmol of KOH were used. ^c1.0 mmol of **1** and 1.3 mmol of **5** were used.

The *gem*-difluoroallylation of organoboronic acids can also be extended to organoborates without affecting the reaction efficiency (Scheme 3). High yield of **3a** with a slightly decreased α/γ regioselectivity (**3a/4a** = 13:1) was obtained by reaction of arylborate **7** with BDFP **2** (Scheme 3a). Furthermore, aryl potassium trifluoroborate salt **8** also underwent the reaction smoothly in high yield (90%) with excellent regioselectivity (**3a/4a** > 20:1) (Scheme 3b) (for details see Supporting Information). However, the Burke's MIDA esters afforded poor yields.

Scheme 3. *gem*-Difluoroallylation of Organoborates and Potassium Trifluoroborate Salt.



^aCombined isolated yield of α and γ substitutions, and the number in parentheses is the ratio of α/γ .

The usefulness of this protocol can also be featured by the late-stage *gem*-difluoroallylation of bioactive natural product. This offered a unique and highly valuable opportunity for drug discovery and development, because the unique structure of *gem*-difluoroallyl group not only provides a platform to use CF₂ as a bioisostere of the oxygen or a carbonyl group,⁵ but also makes it possible to further modify the bioactive compounds through transformation of double bond. As shown in Scheme 3c, treatment of arylborate **9** derived from estrone with BDFP on a gram-scale afforded *gem*-difluoroallylated compound **10** in 88% yield with high regioselectivity (α/γ = 15:1). Subsequently, dihydroxylation of **10** afforded modified bioactive compound **11** in high efficiency. While previous preparation of aryl *gem*-difluoroalkyls required electrochemical fluorination, and only low yield of desired product was provided,¹⁶ thus highlighting the advantages of the pre-

sent process. In addition, the successful use of aromatic pinacol esters as the coupling partners also provides the possibility for sequential C-H borylation/*gem*-difluoroallylation reactions. This is because aromatic pinacol esters are readily available through Ir-catalyzed C-H borylation.¹⁷

In conclusion, we have demonstrated a first example of Pd-catalyzed *gem*-difluoroallylation of organoborons with 3-bromo-3,3-difluoropropene. The reaction allowed *gem*-difluoroallylation of a wide range of organoborons including (hetero)aryl and vinyl boronic acids, borates, and potassium trifluoroborate salt with low catalyst loading under mild and operationally simple conditions. This reaction can also be extended to substituted BDFPs. Application of the method led to modified fluorinated bioactive compounds in a highly efficient and practical manner. Because of the unique structure of *gem*-difluoroallyl group and the advantages of this protocol, such as practicality with low catalyst loading, high regioselectivity and excellent functional group compatibility, we believed that this protocol should be useful for drug discovery and development. Further studies to uncover the mechanism as well as other derivative reactions are now in progress in our laboratory.

ASSOCIATED CONTENT

Supporting Information

Detailed experimental procedures, and characterization data for new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

* xgzhang@mail.sioc.ac.cn

Notes

The authors declare no competing financial interests.

ACKNOWLEDGMENT

This work was financially supported by the National Basic Research Program of China (973 Program) (No. 2012CB821600), the National Natural Science Foundation of China (Nos 21172242 and 21332010), and SIOC.

REFERENCES

- (1) For recent reviews, see: (a) Smart, B. E. *J. Fluorine Chem.* **2001**, *109*, 3. (b) Maienfisch, P.; Hall, R. G. *Chimia.* **2004**, *58*, 93. (c) Special issue on "Fluorine in the Life Sciences", *ChemBioChem* **2004**, *5*, 557. (d) Müller, K.; Faeh, C.; Diederich, F. *Science* **2007**, *317*, 1881. (e) Purser, S.; Moore, P. R.; Swallow, S.; Gouverneur, V. *Chem. Soc. Rev.* **2008**, *37*, 320.
- (2) For selected reviews, see: (a) Furuya, T.; Kamlet, A. S.; Ritter, T. *Nature* **2011**, *473*, 470. (b) Tomashenko, O. A.; Grushin, V. V. *Chem. Rev.* **2011**, *111*, 4475. (c) Hollingworth, C.; Gouverneur, V. *Chem. Commun.* **2012**, *48*, 2929. (d) Besset, T.; Schneider, C.; Cahard, D. *Angew. Chem., Int. Ed.* **2012**, *51*, 5048. (e) Qing, F.-L. *Chin. J. Org. Chem.* **2012**, *32*, 815. (f) Wang, X.; Zhang, Y.; Wang, J. *Sci. Sin. Chim.* **2012**, *42*, 1417.
- (3) For selected examples: (a) Zhang, X.; Xia, H.; Dong, X.; Jin, J.; Meng, W.-D.; Qing, F.-L. *J. Org. Chem.* **2003**, *68*, 9026. (b) Hu, J.; Zhang, W.; Wang, F. *Chem. Commun.* **2009**, 7465. (c) Fujikawa, K.; Fujioka, Y.; Kobayashi, A.; Amii, H. *Org. Lett.* **2011**, *13*, 5560. (d) Fier, P. S.; Hartwig, J. F. *J. Am. Chem. Soc.* **2012**, *134*, 5524. (e) Prakash, G. K. S.; Ganesh, S. K.; Jones, J.-P.; Kulkarni, A.; Masood, K.; Swabeck, J. K.; Olah, G. A. *Angew. Chem., Int. Ed.* **2012**, *51*, 12090. (f) Fujiwara, Y.; Dixon, J. A.; Rodriguez, R. A.; Baxter, R. D.; Dixon, D. D.; Collins, M. R.; Blackmond, D. G.; Baran, P. S. *J. Am. Chem. Soc.* **2012**, *134*, 1494.

(g) Zhou, Q.; Ruffoni, A.; Gianatassio, R.; Fujiwara, Y. Sella, E.; Shabat, D.; Baran, P. S. *Angew. Chem. Int. Ed.* **2013**, *52*, 3949.

(4) (a) Hertel, L. W.; Kroin, J. S.; Missner, J. W.; Tustin, J. M. *J. Org. Chem.* **1988**, *53*, 2406. (b) Akahoshi, F.; Ashimori, A.; Sakashita, H.; Yoshimura, T.; Eda, M.; Imada, T.; Nakajima, M.; Mitsutomi, N.; Kuwahara, S.; Ohtsuka, T.; Fukaya, C.; Miyazaki, M.; Nakamura, N. *J. Med. Chem.* **2001**, *44*, 1297. (c) Makarov, A. A.; Tsvetkov, P. O.; Villard, C.; Esquieu, D.; Pourroy, B.; Fahy, J.; Braguer, D.; Peyrot, V.; Lafitte, D. *Biochemistry* **2007**, *46*, 14899.

(5) (a) Blackburn, C. M.; England, D. A.; Kolkman, F. *J. Chem. Soc., Chem. Commun.* **1981**, 930. (b) Blackburn, G. M.; Kent, D. E.; Kolkman, F. *J. Chem. Soc. Perkin Trans. 1* **1984**, 1119. (c) Kitazume, T.; Kamazaki, T. *Experimental Methods in Organic Fluorine Chemistry*; Gordon and Breach Science: Tokyo, 1998.

(6) O'Hagan, D. *Chem. Soc. Rev.* **2008**, *37*, 308.

(7) Burgey, C. S.; Robinson, K. A.; Lyle, T. A.; Sanderson, P. E. J.; Dale, Lewis, S.; Lucas, B. J.; Krueger, J. A.; Singh, R.; Miller-Stein, C.; White, R. B.; Wong, B.; Lyle, E. A.; Williams, P. D.; Coburn, C. A.; Dorsey, B. D.; Barrow, J. C.; Stranieri, M. T.; Holahan, M. A.; Sitko, G. R.; Cook, J. J.; McMaster, D. R.; McDonough, C. M.; Sanders, W. G.; Wallace, A. A.; Clayton, F. C.; Bohn, D.; Leonard, Y. M.; Detwiler, Jr., T. J.; Lynch, Jr., J. J.; Yan, Y.; Chen, Z.; Kuo, L.; Gardell, S. J.; Shafer, J. A.; Vacca, J. P. *J. Med. Chem.* **2003**, *46*, 461.

(8) (a) Markovski, L. N.; Pahinnik, V. E.; Kirsanov, A. V. *Synthesis* **1973**, 787. (b) Middleton, W. J. *J. Org. Chem.* **1975**, *40*, 574. (c) Lal, G. S.; Pez, G. P.; Pesaresi, R. J.; Prozonc, F. M.; Cheng, H. *J. Org. Chem.* **1999**, *64*, 7048.

(9) For transition metal catalyzed difluoroalkylation, see: (a) Schwaebe, M. K.; McCarthy, J. R.; Whitten, J. P. *Tetrahedron Lett.* **2000**, *41*, 791. (b) Feng, Z.; Chen, F.; Zhang, X. *Org. Lett.* **2012**, *14*, 1938. (c) Belhomme, M.-C.; Poisson, T.; Pannecouke, X. *Org. Lett.* **2013**, *15*, 3428, and ref 3c.

(10) For transition metal mediated difluoroalkylation, see: (a) Taguchi, T.; Kitagawa, O.; Morikawa, T.; Nishiwaki, T.; Uehara, H.; Endo, H.; Kobayashi, Y. *Tetrahedron Lett.* **1986**, *27*, 6103. (b) Sato, K.; Omote, M.; Ando, A.; Kumadaki, I. *J. Fluorine Chem.* **2004**, *125*, 509. (c) Qiu, W.; Burton, D. J. *Tetrahedron Lett.* **1996**, *37*, 2745. (d) Yokomatsu, T.; Murano, T.; Suemune, K.; Shibuya, S. *Tetrahedron* **1997**, *53*, 815, and ref 3d, e.

(11) (a) He, C.-Y.; Fan, S.; Zhang, X. *J. Am. Chem. Soc.* **2010**, *132*, 12850. (b) Zhang, X.; Fan, S.; He, C.-Y.; Wan, X.; Min, Q.-Q.; Yang, J.; Jiang, Z.-X. *J. Am. Chem. Soc.* **2010**, *132*, 4506. (c) Fan, S.; Chen, F.; Zhang, X. *Angew. Chem., Int. Ed.* **2011**, *50*, 5918. (d) Chen, Z.; He, C.-Y.; Yin, Z.; Chen, L.; Zhang, X. *Angew. Chem., Int. Ed.* **2013**, *52*, 5813, and ref 9b.

(12) (a) Hasek, W. R.; Smith, W. C.; Engelhardt, V. A. *J. Am. Chem. Soc.* **1960**, *82*, 543. (b) Shi, G.-q.; Cai, W.-I. *Synlett.* **1996**, *4*, 371. (c) Bergeron, M.; Johnson, T.; Paquin, J.-F. *Angew. Chem., Int. Ed.* **2011**, *50*, 11112, and ref 9c.


(13) 3-Bromo-3,3-difluoropropene (BDFP) can also be easily prepared from CF₂Br₂ and ethylene, see: (a) Tarrant, P.; Lovelace, M. *J. Am. Chem. Soc.* **1954**, *76*, 3466. (b) Seyferth, D.; Simon, R. M.; Sepelak, D. J.; Klein, H. A. *J. Org. Chem.* **1980**, *45*, 2273.

(14) (a) Shi, G.-q.; Huang, X.-h.; Zhang, F.-J. *Tetrahedron Lett.* **1995**, *36*, 6305. (b) Kirihaara, M.; Takuwa, T.; Okumura, M.; Wakikawa, H.; Takahata, H.; Momose, T.; Takeuchi, Y.; Nemoto, H. *Chem. Pharm. Bull.* **2000**, *48*, 885. (c) Hazari, A.; Gouverneur, V.; Brown, J. M. *Angew. Chem., Int. Ed.* **2009**, *48*, 1296. (d) Pigeon, X.; Bergeron, M.; Barabe, F.; Dube, P.; Frost, H. N.; Paquin, J.-F. *Angew. Chem., Int. Ed.* **2010**, *49*, 1123, and ref 12c.

(15) The substituted BDFP **5** are not commercially available and were prepared according to the literature, see: (a) Nihei, T.; Iwai, N.; Matsuda, T.; Kitazume, T. *J. Org. Chem.* **2005**, *70*, 5912. (b) Gonzalez, J.; Foti, C. J.; Elsheimer, S. *J. Org. Chem.* **1991**, *56*, 4322. (c) Elsheimer, S.; Foti, C. J.; Bartberger, M. D. *J. Org. Chem.* **1996**, *61*, 6252.

(16) Nagura, H.; Murakami, S.; Fuchigami, T. *Tetrahedron* **2007**, *63*, 10237.

(17) Mkhallid, I. A. I.; Barnard, J. H.; Marder, T. B.; Murphy, J. M.; Hartwig, J. F. *Chem. Rev.* **2010**, *110*, 890.

Highly Selective *gem*-Difluoroallylation of Organoborons with Bromodifluoromethylated Alkenes Catalyzed by Palladium

- ✓ High Efficiency and Regioselectivity
- ✓ Low Catalyst Loading
- ✓ Broad Substrate Scope
- ✓ Excellent Functional Group Compatibility (Br, OH...)