


This article was downloaded by: [University of Bristol]

On: 26 January 2015, At: 11:50

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954

Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

Reactions of 3-substituted 5-Arylmethylene-1,3-Thiazolidin-2,4-diones with Azide and Cyanide Ions

Ahmed Said Ahmed Youssef ^a

^a Chemistry Department , Ain Shams University ,
Abbassia, Cairo, Egypt

Published online: 29 Aug 2007.

To cite this article: Ahmed Said Ahmed Youssef (2007) Reactions of 3-substituted 5-Arylmethylene-1,3-Thiazolidin-2,4-diones with Azide and Cyanide Ions, Phosphorus, Sulfur, and Silicon and the Related Elements, 182:6, 1335-1349, DOI: [10.1080/10426500601160967](https://doi.org/10.1080/10426500601160967)

To link to this article: <http://dx.doi.org/10.1080/10426500601160967>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or

indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Reactions of 3-substituted 5-Arylmethylene-1,3-Thiazolidin-2,4-diones with Azide and Cyanide Ions

Ahmed Said Ahmed Youssef

Chemistry Department, Ain Shams University, Abbassia, Cairo, Egypt

3-substituted-5-arylmethylene-1,3-thiazolidine-2,4-diones 3a–e react- ed with sodium azide to afford pyrazolinone derivatives 8a–d as well as 5-(4-chlorophenylmethylene)-1,3-thiazolidine-2,4-dione 7a in the case of 3d and 5-(4-methoxyphenylmethylene)-1,3-thiazolidin-2,4-dione 7b as a sole product in the case of 3e. Reactions of potassium cyanide with 3b–d gave butyronitrile derivative 9 as well as 7a and a water soluble adduct that, upon treatment with chloroacetic acid, yielded 10 in the case of 3c and 3d. Also, the treatment of 3e with potassium cyanide afforded a mixture of 7b, but-3-enonitrile derivative 11 and 10. Structures of all products were elucidated by microanalytical and spectral data.

Keywords 5-arylmethylene-1,3-thiazolidin-2,4-diones; 2-oxo-but-3-enonitrile; butyronitrile derivative; pyridazine derivatives; pyrazolinone derivatives

INTRODUCTION


It has been reported^{1–4} that 3-substituted-5-arylmethylene-1,3-thiazolidin-2,4-diones react with hydrazines, morpholine, benzylamine, and piperidine to afford pyrazolinones, thiolo-propenamides, arylamino-carboxamide, and N-phenylacrylamide derivatives, respectively. However, the 3-unsubstituted derivatives gave, upon treatment with hydrazine hydrate and piperidine, the corresponding triazinone^{3,5} and acrylamide derivatives.⁶ The latent ability of this ring system to cleave at 1,2- as well as 3,4- bonds encouraged us to synthesize some 3-substituted 5-arylmethylene-1,3-thiazolidin-2,4-diones to investigate the behavior of this ring toward azide and cyanide ions that had previous not been studied.

Received September 11, 2006; accepted November 18, 2006.

Address correspondence to Ahmed Said Ahmed Youssef, Chemistry Department, Faculty of Science, Ain Shams University, P.O. Box 11566, Abbassia, Cairo, Egypt. E-mail: ahmedy84@hotmail.com

RESULTS AND DISCUSSION

The starting materials (**3a,b**) were synthesized by condensing 3-phenyl-1,3-thiazolidin-2,4-dione (**1b**) with the appropriate aldehyde using sodium acetate-acetic acid.⁶ However, (**3c-e**) were prepared following the method of Lo et al.⁷ by treating the potassium salt of the respective *N*-unsubstituted analogues obtained from 1,3-thiazolidin-2,4-dione **1a** and the appropriate aldehyde with allyl or isopropyl bromide in dimethylformamide and were shown to be mixtures of the (*E*) and (*Z*)-isomers,³ as represented in Scheme 1. In the present study, the (*E,Z*)-mixtures were used without separation.




SCHEME 1

Unfortunately, attempts to prepare new 5-arylidene-1,3-thiazolidin-2,4-diones through the condensation of (**1a,b**) with benzoin (**4a**), 4-methoxybenzoin (**4b**), and benzoylacetone (**4c**) gave a poor yield of adducts.


Thus reacting equimolar amounts of 1,3-thiazolidin-2,4-dione (**1a**) with benzoin (**4a**), 4-methoxybenzoin (**4b**), or benzoylacetone (**4c**) in refluxing toluene in the presence of ammonium acetate as a base afforded a mixture of **5a** (major), and **6a** (minor), **5b**, or **6b**, respectively. Similar treatment of 3-phenyl-1,3-thiazolidin-2,4-dione (**1b**) with **4a** gave a mixture of **5a** (major) and **6c** (minor).

The structures of compounds **5** and **6** were elucidated from their analytical and spectral data. The infrared spectra of **5** showed absorption bands corresponding to C=N and/or C=C. However, those of compounds **6** exhibited stretching absorptions assignable to NH and/or OH as well as two strong absorptions, which were characteristic of C=O



SCHEME 2


in the respective systems. The ¹H NMR spectra of compounds **5** and **6a,c** were in accord with the proposed structures. Compound **5b** exhibited a singlet signal for methoxy protons. However, compounds **6a,c** revealed exchangeable singlets for OH and/or NH protons as well as singlet signals for methine protons. The downfield value for the OH proton signal of compounds **6a,c** suggested the existence of chelation with the 4-oxo group of the heteroring. Further elucidation was gained from EI-MS spectra that showed peaks for compounds **5**. However, the EI-MS spectra of compounds **6a,b** did not show the molecular ion peaks; instead they showed (M⁺ - ^{CO}Δ_{NH}) peak for compound **6a** and (M⁺ - HNCO - CH=C(OH)CH₃) peak in the case of **6b**, beside some of abundant peaks.




SCHEME 3

The formation of compounds **5** is shown in Scheme 3.

The treatment of (*E,Z*)-3-substituted-5-arylmethylene-1,3-thiazolidin-2,4-diones **3a-d** with sodium azide (3 equiv.) in refluxing dimethyl-



- a, R = Ph; Ar = 4-MeOC₆H₄-
 b, R = Ph; Ar = 4-ClC₆H₄-
 c, R = -CH₂-CH=CH₂; Ar = 4-ClC₆H₄-
 d, R = (CH₃)₂CH-; Ar = 4-ClC₆H₄-



- a, Ar = 4-ClC₆H₄-
 b, Ar = 4-MeOC₆H₄-

SCHEME 4


formamide afforded the pyrazolinone derivatives **8a–d** (major) as well as 5-(4-chlorophenylmethylene)-1,3-thiazolidin-2,4-dione (**7a**) in a minor amount in the case of **3d**. Similar treatment of **3e** with sodium azide yielded 5-(4-methoxyphenylmethylene)-1,3-thiazolidin-2,4-dione as sole products.

The structure of compounds **8a–d** was substantiated from their analytical and spectral data. Their infrared spectra showed stretching absorptions assignable to OH, C=O, and C=C groups of pyrazolinones.⁸ The ¹HNMR spectra of **8** were in good agreement with the suggested structure. Further elucidation was gained from their EI-MS where they did not show the molecular ion peak; instead a peak corresponding to M⁺ -16. The fragmentation pathway of compounds **8** is shown in Scheme 5.

The structure of compounds **7a** and **7b** was rigidly confirmed by mixture m.p. with authentic samples.⁷


The formation of compounds **7** and **8** (cf. Scheme 6) could be rationalized on the basis of an attack of the azide ion at the allyl or isopropyl group to afford the anion A. Protonation of this anion yielded adducts **7a** and **7b** (route a). The thiazolidine ring opening through an attack of the azide ion at the 2-oxo- group followed by Curtius rearrangement gave the isocyanate intermediate (B). Successive hydrolysis, intramolecular cyclization followed by elimination of hydrogen sulfide, afforded the pyrazolinone salt C. Acidification of (C) gave compounds **8** (route b).

However, treating (E,Z)-3-phenyl-5-(4-chlorophenylmethylene)-(**3b**)-, (E,Z)-3-allyl-5-(4-chlorophenylmethylene)-(**3c**)-, and (E,Z)-3-isopropyl-5-(4-chlorophenylmethylene)-(**3d**)-1,3-thiazolidin-2,4-diones with potassium cyanide in a boiling mixture of methanol and dioxane afforded 4-cyano-4-(4-chlorophenyl)-2-oxobutyronitrile (**9**) as well as (E,Z)-5-(4-chlorophenylmethylene)-1,3-thiazolidin-2,4-dione (**7a**) and a water soluble adduct that, upon treating its methanolic



SCHEME 5

solution with chloroacetic acid, gave ammonium chloroacetate (**10**) in the case of **3c** and **3d**. Similar treatment of (*E,Z*)-3-allyl-5-(4-methoxyphenylmethylene)-1,3-thiazolidin-2,4-dione (**3e**) with potassium cyanide gave a mixture of (*E,Z*)-5-(4-methoxyphenylmethylene)-1,3-thiazolidin-2,4-dione (**7b**), but-3-enonitrile derivative **11** and a water soluble adduct that yielded compound **10** upon treating its methanolic solution with chloroacetic acid.






FIGURE 6




FIGURE 7

The structures of compounds **9** and **11** were substantiated from their analytical and spectral data. Their IR spectra exhibited stretching absorptions for CN and CO groups beside an additional absorption for the SH group in the case of compound **11**.

The 1H NMR spectrum of compound **9** was consistent with the suggested structure as it showed from low to high field the signals of aromatic, methine, and methylene protons. Its EI-MS spectrum did not show the molecular ion peak, but it showed a peak at m/e 190 that corresponds to $(M^+ - CO)$. The fragmentation pathway is represented in Scheme 7.

The 1HNMR spectrum of **11** revealed the existence of two closely spaced singlets for protons of two methoxy groups, an exchangeable singlet for the SH proton, as well as two doublet signals for aromatic protons. Moreover, the structure of **11** got further support from its EI-MS as it showed the molecular ion peak beside some abundant peaks. The fragmentation pattern is shown in Scheme 8.

The structures of compounds **7a** and **7b** were rigidly confirmed by m.p. comparison with those reported.⁷ However, the structure of **10** was confirmed by mixture m.p. with authentic specimen.

The overall picture for the mode of action of the cyanide ion on 3-aryl and 3-alkyl substituted-5-arylmethylene-1,3-thiazolidin-2,4-diones **3b-e** is easily visualized as represented by type (A) and type (B) (cf. Scheme 9). Type (A) involves an initial attack by the cyanide ion at the allyl or isopropyl group to afford the anion (A). Protonation of this anion yielded the adducts **7a** and **7b**. Thiazolidine ring opening of this anion at the 1,2-bond gave the thiolate anion (B), which underwent 3,4-bond cleavage to give isocyanate (C) and thirane (D). The




FIGURE 8

addition of water to (C) gave the carbamate (E), which, on treatment of its methanolic solution with chloroacetic acid, afforded **10**. A thirane ring opening through attack by the CN^- ion at the 2-oxo group followed by the addition of the solvent molecule (MeOH)⁹ at the β -carbon of the exocyclic double bond gave the methoxy adduct (F). Oxidation of the methoxy adduct (F) afforded (G), which, upon protonation, yielded **11**.

Type (B) involves the thiazolidine ring opening through successive cleavage at the 1,2- as well as the 3,4-bonds to give the isocyanate (H) and the thirane (D). The thirane ring opening through attack by




FIGURE 9

the cyanide ion at the 2-oxo group followed by reaction of the formed thiolate ion with the isocyanate (H) gave (I). Cleavage of (I) yielded the thiocarbamate (J) and the acetylenic ketone (K). Attack by another cyanide ion at the β -acetylenic carbon of (K) followed by reduction afforded **9**.

CONCLUSION

It is inferred that reactions of 3-substituted-5-arylmethylene-1,3-thiazolidine-2,4-diones with azide and cyanide ions proceed through 1,2- as well as 3,4- bond cleavage, as reported¹⁻⁴ before with hydrazines, morpholine, benzyl amine, and piperidine, and not through an attack at the β -carbon of the exocyclic double bond as reported¹⁰ in the reaction of 4-methoxyphenylazide with 3-substituted-5-arylmethylene-4-oxo-2-thioxo-1,3-thiazolidine.

EXPERIMENTAL

All melting points are uncorrected. IR spectra were measured on a Unicam SP1200 spectrometer as KBr discs. Unless otherwise stated, the ¹HNMR spectra were measured in CDCl₃ solution on Varian Gemini 200 MHz or Bruker ES 200 MHz instruments with chemical shift (δ) expressed in ppm downfield from Me₄Si. Mass spectra were recorded on Shimadzu GC-MS-QP 1000 Ex or Finnigan GCQ instruments operating at 70 eV. Column chromatography and TLC^{10,11} were run using Silica Gel Voein, activity III/30 mm according to Brockmann and Schodder and TLC aluminium sheets Silica Gel 60 F₂₅₄ (Merck).

Reactions of (1a,b) with (4a-c)

A solution of (1a) or (1b) (10 mmoles) and (4a-c) (10 mmoles) in 25 mL toluene in the presence of ammonium acetate (2.31 g, 30 mmoles) was refluxed for 15 h. The reaction mixture was cooled and then poured into cold water. The toluene layer was dried over anhydrous calcium chloride, concentrated, and left to stand over night at room temperature. The precipitated solid was recrystallized from a proper solvent to give the adducts **5b** and **6b** in the case of reactions of **1a** with **4b** and **4c**. The adduct **5a** was isolated from the reaction of **1a** with **4a**, and the residual oil was chromatographed over silica gel. Elution with petroleum ether (b.p. 40-60)/ether (4 : 1 V/V) gave **6a**. The precipitated solid in the case of the reaction of **1b** with **4a** was fractionally crystallized to give first **6c** as yellow crystals (ethanol). The insoluble part in boiling ethanol was recrystallized from benzene to give **5a** as white crystals.

2,3,5,6-Tetraphenyl-1,4-diazine (5a)

1.5 g (30%) white crystals (benzene), m.p. 241–243 °C; IR: ν_{\max} = 3080 (aryl-H), 1615 (C=C and/or C=N), 760, 690 cm^{-1} . $^1\text{HNMR}$: δ 7.31–7.68 (m, 20, ArH). EI-MS m/z (%) 384 (M^+ , 77), 383 (68), 179 (14), 178 (base), 177 (20), 176 (35), 152 (20), 151 (14), 103 (13), 77 (10), 76 (13), 51 (11). Anal. calcd. for $\text{C}_{28}\text{H}_{20}\text{N}_2$: C, 87.47; H, 5.24; N, 7.29. Found : C, 87.21; H, 5.49; N, 7.10%.

2,6-Diphenyl-3,5-di-(4-methoxyphenyl)-1,4-diazine (5b)

1.1 g (25%), white crystals (methanol-benzene), m.p. 190–192 °C; IR: ν_{\max} = 3020 (aryl-H), 2920, 2830 (alkyl-H), 1610 (C=C and/or C=N), 835, 760, 700 cm^{-1} . $^1\text{HNMR}$: δ 3.83 (s, 6, 2OCH_3), 6.83–7.69 (m, 18, ArH). EI-MS m/z (%) 444 (M^+ , base), 443 (53), 208 (31), 193 (54), 165 (50), 164 (17), 163 (11), 139 (11), 104 (25). Anal. calcd. for $\text{C}_{30}\text{H}_{24}\text{N}_2\text{O}_2$: C, 81.06; H, 5.44; N, 6.30. Found: C, 81.22; H, 5.11; N, 6.45%.

5-(1,2-Diphenyl-2-hydroxy)ethylene-1,3-thiazolidin-2,4-dione (6a)

0.37 g (12%), yellow crystals (ethanol), m.p. 314–316 °C; IR: ν_{\max} = 3340, 3130 (OH and/or NH), 3055 (aryl-H), 2930, 2860 (alkyl-H), 1740, 1690 (C=O), 765, 690 cm^{-1} . $^1\text{HNMR}$: δ 4.02 (s, 1, CH), 7.24–7.42 (m, 10, ArH), 10.01 (br.s, 1, OH exchangeable), 10.15 (br.s, 1, NH exchangeable). EI-MS m/z (%) 311 (M^+ , missed), 236 ($\text{M}^+ - \overset{\text{O}}{\text{N}}$, 88), 165 (16), 105 (63), 104 (base), 103 (16), 77 (51), 76 (12), 51 (35). Anal. calcd. for $\text{C}_{17}\text{H}_{13}\text{NO}_3\text{S}$: C, 65.58; H, 4.21; N, 4.49. Found: C, 65.41; H, 4.46; N, 4.19%.

5-(1-Phenyl-3-hydroxy)-but-2-en-1-ylene-1,3-thiazolidin-2,4-dione (6b)

0.39 g (15%), pale yellow crystals (ethanol), m.p. 140–142 °C; IR: ν_{\max} = 3314, 3160 (OH and/or NH), 3060 (aryl-H), 2983, 2875 (alkyl-H), 1743, 1635 (C=O), 1595 (C=C), 641, 754 cm^{-1} . EI-MS m/z (%) 261 (M^+ , missed), 161 ($\text{M}^+ - \text{HNCO}-\text{CH}=\text{C}(\text{OH})\text{CH}_3$, 38), 160 (88), 84 (base), 77 (33), 51 (29). Anal. calcd. for $\text{C}_{13}\text{H}_{11}\text{NO}_3\text{S}$: C, 59.76; H, 4.24; N, 5.36. Found: C, 59.42; H, 4.39; N, 5.29%.

3-Phenyl-5-(1,2-diphenyl-2-hydroxy)ethylene-1,3-thiazolidin-2,4-dione (6c)

0.77 g (20%), yellow crystals (ethanol), m.p. 138–140°C, IR: ν_{\max} = 3380, 3210 (OH and/or NH), 3070 (aryl-H), 2985, 2874 (alkyl-H), 1728, 1676 (C=O), 752, 692 cm^{-1} . $^1\text{H NMR}$: δ 4.1 (s, 1, CH) 7.18–7.51 (m, 15, ArH), 10.12 (br.s, 1, OH exchangeable) Anal. calcd. for $\text{C}_{23}\text{H}_{17}\text{NO}_3\text{S}$: C, 71.29; H, 4.42; N, 3.62. Found: C, 71.48; H, 4.23; N, 3.79%.

Reactions of 3-Substituted-5-arylmethylene-1,3-thiazolidin-2,4-diones (3a–e) with Sodium Azide

A mixture of (3a–e) (10 mmoles) and sodium azide (1.95 g, 30 mmoles) was refluxed in dimethylformamide (20 mL) for 20 h. The color of the reaction mixture changed from yellow to brown to orange, and then to yellow again. The precipitated solid during the reaction was filtered off; part of it was dissolved in water and then acidified with ice cold HCl, where no precipitate was obtained. This solid was not identified due to its very poor yield and the difficulty of doing detailed characterization as it is insoluble in all organic solvents and its m.p. is $>300^\circ\text{C}$. The reaction mixture was cooled, concentrated, and then poured into ice cold water. The precipitated solid was recrystallized from a proper solvent to give (10–15%) of unreacted starting materials 3a–e. The filtrate was acidified with ice cold HCl. The solid obtained was fractionally crystallized to afford 7a and 7b in the case of 3d and 3e (identical m.p. with the reported⁷), and 8a–d, respectively.

E,Z-5-(4-Chlorophenylmethylene)-1,3-thiazolidin-2,4-dione (E,Z-7a)

0.6 g (10%); yellow crystals (ethanol-benzene), m.p. 224–226°C, Lit.⁷, m.p. 223–225°C.

E,Z-5-(4-Methoxyphenylmethylene)-1,3-thiazolidin-2,4-dione (E,Z-7b)

0.7 g (12%), yellow crystals (ethanol), m.p. 208–210°C, Lit.,⁷ m.p. 212–214°C.

1-Carboxy-2-phenyl-3-oxo-5-(4-methoxyphenyl)-2,3-dihydropyrazole (8a)

1.55 g (50%), colorless crystals (ethanol-benzene), m.p. 152–154°C; IR: ν_{\max} = 3400, 3150 (br.) (OH), 1665 (C=O), 1600 (C=C), 1550 (pyrazole)

ring), 830, 690 cm^{-1} . $^1\text{HNMR}$: δ 3.77 (s, OMe), 6.89–7.97 (m, 10, ArH + CH=), 9.1 (br.s, 1, OH exchangeable). EI-MS m/z (%) 310 (M^+ , missed), 294 ($\text{M}^+ - 16$, 23), 202 (26), 146 (15), 135 (24), 134 (19), 119 (25), 93 (base), 77 (20), 76 (15), 65 (16), 51 (17). Anal. calcd. for $\text{C}_{17}\text{H}_{14}\text{N}_2\text{O}_4$: C, 65.80; H, 4.55; N, 9.03. Found: C, 65.97; H, 4.35; N, 8.89%.

1-Carboxy-2-phenyl-3-oxo-5-(4-chlorophenyl)-2,3-dihydropyrazole (8b)

1.73 g (55%), colorless crystals (xylene), m.p. 208–210°C; IR: ν_{max} = 3389, 3196 (br.) (OH), 1671 (C=O), 1601 (C=C), 1548 (pyrazole ring), 829, 747, 684 cm^{-1} . $^1\text{HNMR}$: δ 7.37–7.98 (m, 10, ArH + CH=), 8.92 (br.s, 1, OH exchangeable). EI-MS m/z (%) 314 (M^+ , missed), 300 ($\text{M}^+ + 2 - 16$, 5), 298 ($\text{M}^+ - 16$, 15), 206 (19), 139 (12), 123 (15), 93 (base) 77 (12), 65 (12), 51 (10). Anal. calcd. for $\text{C}_{16}\text{H}_{11}\text{ClN}_2\text{O}_3$: C, 61.06; H, 3.52; N, 8.90. Found: C, 61.20; H, 3.39; N, 8.74%.

1-Carboxy-2-allyl-3-oxo-5-(4-chlorophenyl)-2,3-dihydropyrazole (8c)

1.33 g (48%), metallic needles (ethanol), m.p. 130–132°C; IR: ν_{max} = 3400, 3118 (br.) (OH), 2980, 2906 (alkyl-H), 1668 (C=O), 1637, 1612 (C=C), 1559 (pyrazole ring), 824 cm^{-1} . $^1\text{HNMR}$: δ 4.08 (m, 2, N-CH₂), 5.22 (m, 2, CH₂=CH-), 5.91 (m, 1, CH₂=CH-), 7.28–7.87 (m, 5, ArH + CH=), 13.05 (br.s, 1, OH exchangeable). EI-MS m/z (%) 278 (M^+ , missed), 264 ($\text{M}^+ + 2 - 16$, 13), 262 ($\text{M}^+ - 16$, 37), 247 (14), 208 (16), 206 (46), 138 (14), 123 (19), 114 (10), 57 (81), 56 (base). Anal. calcd. for $\text{C}_{13}\text{H}_{11}\text{ClN}_2\text{O}_3$: C, 56.03; H, 3.98; N, 10.05. Found: C, 55.84; H, 4.12; N, 10.21%.

1-Carboxy-2-isopropyl-3-oxo-5-(4-chlorophenyl)-2,3-dihydropyrazole (8d)

1.49 g (53%), colorless crystals (benzene-ethanol), m.p. 145–147°C; IR: ν_{max} = 3406, 3104 (br.) (OH), 3050 (aryl-H), 2948, 2899 (alkyl-H), 1669 (C=O), 1550 (pyrazole ring), 820 cm^{-1} . $^1\text{HNMR}$: δ 1.26 (d, 2CH₃, J = 6.5 Hz), 4.26 (m, 1, CH(CH₃)₂, J = 6.6 Hz), 7.25–7.79 (m, 5, ArH + CH=), 12.41 (br.s, 1, OH exchangeable). EI-MS m/z (%) 280 (M^+ , missed), 266 ($\text{M}^+ + 2 - 16$, 5), 264 ($\text{M}^+ - 16$, 16), 249 (16), 208 (14), 206 (43), 138 (13), 123 (19), 58 (base). Anal. calcd. for $\text{C}_{13}\text{H}_{13}\text{ClN}_2\text{O}_3$: C, 55.62; H, 4.67; N, 9.98. Found: C, 55.41; H, 4.81; N, 9.66%.

Reactions of (3b–e) with Potassium Cyanide

A solution of (25 mL) dioxane containing **3a–e** (10 mmoles) and potassium cyanide (1.95 g, 30 mmoles) was heated to 90°C for 25 min. During the reflux, 30 mL of methanol was added in five equal portions. The mixture was refluxed for 4 h. The precipitated solid during heating was filtered off and dissolved in water. Acidification with ice cold HCl was accompanied by effervescence without precipitation of a solid. Treatment of a methanolic solution of this solid with chloroacetic acid afforded **10**. The reaction mixture was concentrated and left to stand at room temperature for 2 h. The precipitated solid was filtered off, dissolved in water, and acidified with ice cold HCl to give a solid. Crystallization from ethanol–benzene afforded **7a** in the case of **3c** and **3d**, in a yield of 8%, m.p. 225–227°C (identical with the reported⁷), and **7b** as an E/Z mixture in a ratio of 35:65 in the case of **3e**, with a yield of 10%, m.p. 208–210°C (identical with the reported⁷). On leaving the mother liquor over night at room temperature, afforded compound **9** in the case of **3b**, **3c**, & **3d**, while **11** was obtained in the case of **3e**.

Ammonium Chloroacetate (10)

0.12 g (11%), white crystals (methanol), m.p. 137–139°C (undepressed on an admixture with an authentic sample prepared by passing ammonia gas over a methanolic solution of chloroacetic acid); IR: ν_{\max} = 3401 (NH), 2962, 2942, 2855 (alkyl-H), 1648 cm^{-1} (C=O). ¹HNMR (DMSO-*d*₆): δ 4.04 (s, 2, CH₂), 4.83 (br.s, 4, NH₄ exchangeable). EI-MS *m/z* (%) 111 (M⁺, missed), 96 (M⁺ + 2 -NH₃, 1.4), 94 (M⁺ -NH₃, 4.2), 79 (ClCH₂-C≡O⁺, 1.3), 78 (ClCH=C=O, 0.5), 77 (ClCH₂C≡O⁺, 4.3), 76 (ClCH=C=O, 1), 58 (M⁺ -NH₃-HCl, 1), 50 (M⁺ -NH₃-CO₂, base).

4-Cyano-4-(4-chlorophenyl)-2-oxobutyronitrile (9)

0.87–1.31 g (40–60%), creamy crystals (ethanol), m.p. 87–89°C; IR: ν_{\max} = 3050 (aryl-H), 2973, 2945 (alkyl-H), 2247 (CN), 1657 (C=O), 825 cm^{-1} . ¹HNMR: δ 2.97 (m, 2, CH₂, J = 6.4 Hz), 4.15 (t, 1, CH, J = 6.8 Hz), 7.37 (d, 2, H_a, H_a', J_o = 6.8 Hz), 7.46 (d, 2, H_b, H_b', J_o = 6.8 Hz). EI-MS *m/z* (%) 218 (M⁺, missed), 192 (M⁺ + 2 -CO, 7), 190 (M⁺ -CO, 19), 152 (M⁺ + 2 -CH₂COCN, 36), 150 (M⁺ -CH₂COCN, base), 115 (11), 88 (11), 75 (22), 74 (14), 63 (20), 62 (13), 61 (11), 51 (21), 50 (34). Anal. calcd. for C₁₁H₇ClN₂O: C, 60.43; H, 3.23; N, 12.81. Found: C, 60.25; H, 3.31; N, 12.56%.

3-Mercapto-4-methoxy-4-(4-methoxyphenyl)-2-oxo-but-3-enonitrile (11)

1.2 g (48%), pale brown crystals (ethanol), m.p. 279–281°C; IR: ν_{\max} = 3001 (aryl-H), 2929, 2840 (alkyl-H), 2423 (SH), 2056 (CN), 1673 (C=O), 1607 (C=C), 817 cm^{-1} . $^1\text{HNMR}$ (DMSO- d_6): δ 3.76, 3.77 (two singlets, 2OCH₃), 2.49 (br.s, 1, SH exchangeable), 7.01 (d, 2, H_a, H_{a'}, J₀ = 8.8 Hz), 7.47 (d, 2, H_b, H_{b'}, J₀ = 8.6 Hz). EI-MS m/z (%) 251 (M⁺ + 2, 6), 249 (M⁺, 68), 237 (2), 235 (30), 216 (17), 191 (13), 174 (18), 173 (26), 165 (11), 164 (base), 163 (49), 149 (69), 148 (18), 145 (25), 121 (22), 120 (22), 119 (12), 94 (12), 89 (14), 82 (10), 77 (25), 69 (13), 63 (14), 62 (13), 51 (17), 50 (11). Anal. calcd. for C₁₂H₁₁NO₃S: C, 57.82; H, 4.45; N, 5.62. Found: C, 57.52; H, 4.62; N, 5.74%.

REFERENCES

- [1] M. T. Omar, M. M. Habashy, A. M. Youssef, and F. A. Sherif, *J. Prakt. Chemie.*, **331**, 393 (1989).
- [2] M. T. Omar and F. A. Sherif, *Synthesis*, **9**, 742 (1981).
- [3] A. R. A. Raouf, M. T. Omar, and M. M. El-Attal, *Acta. Chimica. Sci. (Hung.)*, **87**, 187 (1975).
- [4] A. R. A. Raouf, M. T. Omar, and M. M. El-Attal, *Acta. Chimica. Sci. (Hung.)*, **83**, 367 (1975).
- [5] M.T. Omar, M. M. Habashy, and A. El-Khamry, *Aust. J. Chem.*, **33**, 619 (1980).
- [6] E. Campaigne and R. E. Cline, *J. Org. Chem.*, **21**, 32 (1956).
- [7] C. P. Lo, E. Y. Shorpsshire, and W. J. Croxall, *J. Am. Chem. Soc.*, **75**, 4845 (1953).
- [8] J. Elguero, C. Marzin, A. R. Katritzky, and P. Linda, *The Tautomerism of Heterocycles* (Academic Press, New York, 1976).
- [9] D. Papa, E. Schwenk, F. Villiani, and E. Klingsberg, *J. Am. Chem. Soc.*, **70**, 3555 (1948).
- [10] M. T. Omar, A. S. A. Youssef, and K. A. Kandeel, *Phosphorus, Sulfur and Silicon*, **162**, 25 (2000).
- [11] K. A. Kandeel and A. S. A. Youssef, *Molecules*, **6**, 510 (2001).