

Direct Conversion of Primary Alcohols to 1,2-Amino Alcohols: Enantioselective Iridium-Catalyzed Carbonyl Reductive Coupling of Phthalimido-Allene via Hydrogen Auto-Transfer

Kim Spielmann, Ming Xiang,¹ Leyah A. Schwartz, and Michael J. Krische*²

Department of Chemistry, University of Texas at Austin, Austin, Texas 78712, United States

S Supporting Information

ABSTRACT: The first catalytic enantioselective carbonyl (α -amino)allylations are described. Phthalimido-allene **1** and primary alcohols **2a–2z**, **2a'–2c'** engage in hydrogen auto-transfer-mediated carbonyl reductive coupling by way of (α -amino)allyliridium–aldehyde pairs to form vicinal amino alcohols **3a–3z**, **3a'–3c'** with high levels of regio-, *anti*-diastereo-, and enantioselectivity. Reaction progress kinetic analysis and isotopic labeling studies corroborate a catalytic cycle involving turnover-limiting alcohol dehydrogenation followed by rapid allene hydro-metalation.

Asymmetric carbonyl addition ranks foremost among methods used for the convergent construction of enantiomerically enriched alcohols.¹ Data mining of patents from the pharmaceutical industry reveals that carbonyl addition (alongside Suzuki coupling) remains one of the most frequently utilized methods for C–C bond formation.² The vast majority of carbonyl addition reactions rely on the use of preformed carbanions, which can be moisture sensitive, unsafe, and often require multi-step preparation and cryogenic conditions. Metal-catalyzed carbonyl reductive coupling of π -unsaturated pronucleophiles has emerged as an alternative to the use of stoichiometric carbanions.³ However, many of the terminal reductants utilized in such processes (e.g., Mn, Zn, Et₃B, Et₂Zn) are as problematic as the premetalated reagents they replace. Carbonyl reductive coupling via hydrogen auto-transfer does not require an exogenous reductant, as alcohol reactants serve dually as reductant and carbonyl proelectrophile.⁴

Based on this concept and motivated by the prevalence (>40%) of chiral amines (including vicinal amino alcohols) in FDA-approved drugs,^{5a,b} a catalytic enantioselective carbonyl (α -amino)allylation was sought.^{6–8} In 1993, Barrett reported a boron reagent for asymmetric carbonyl (α -amino)allylation.⁷ Remarkably, after more than 25 years, corresponding catalytic enantioselective processes have remained elusive, and the only related catalytic transformation to have appeared is the 2-azadiene-ketone reductive coupling reported by Malcolmson.⁹ Here, we disclose that phthalimido-allene **1**, a tractable crystalline solid (mp = 79–81 °C), participates in catalytic enantioselective carbonyl reductive coupling via hydrogen auto-transfer to deliver vicinal amino alcohols with high levels of regio-, *anti*-diastereo-, and enantiocontrol (Figure 1). This work represents a rare example of the use of allene pronucleophiles in enantioselective carbonyl reductive coupling.¹⁰





Figure 1. Selected enantioselective methods for convergent construction of vicinal amino alcohols via classical and metal-catalyzed carbonyl addition.

Phthalimido allene **1** is readily prepared through base-catalyzed isomerization of commercially available *N*-propargyl phthalimide.¹¹ Guided by seminal findings from our labo-

Received: August 12, 2019


Table 1. Selected Optimization Experiments in the Enantioselective Iridium-Catalyzed (α -Amino)allylation of Phthalimido-Allene **1 with Alcohol **2a**^a**



^aYields are of material isolated by silica gel chromatography. Diastereoselectivities were determined by ¹H NMR of crude reaction mixtures. Enantioselectivities were determined by chiral stationary phase high performance liquid chromatography (HPLC) analysis. See [Supporting Information](#) for experimental details.

ratory,¹² it was posited that hydrogen transfer from primary alcohols to allenimide **1** would generate transient (phthalimido)allyliridium–aldehyde pairs that combine by way of closed six-centered transition structures to furnish *anti*-vicinal amino alcohols. The feasibility of this transformation was rendered uncertain by competing conventional transfer hydrogenation of allene **1** in response to the steric demand of the phthalimide moiety, which may retard the rate of aldehyde addition. An assay of diverse chiral ruthenium and iridium complexes was undertaken, and a promising result was obtained using the cyclometalated π -allyliridium complex modified by 3-nitrobenzoic acid and (*R*)-SEGPHOS, Ir-I, which delivered the desired amino alcohol **3a** in 10% yield and 40% ee with >20:1 *anti*-diastereoselectivity (Table 1). Enantioselectivity improved using the more Lewis acidic 4-cyano-3-nitro-*C,O*-benzoate, Ir-II, but the isolated yield of **3a** remained modest due to low conversion. Similar trends were observed with the corresponding catalysts based on (*R*)-BINAP, Ir-III and Ir-IV, but with a small increase in enantioselectivity. A pronounced improvement in both conversion and enantioselectivity was observed upon use of Ir-V, which incorporates (*R*)-H₈-BINAP.¹³ Use of the (*R*-

Table 2. Diastereo- and Enantioselective Iridium-Catalyzed Hydrohydroxyalkylation of Phthalimido-Allene **1 with Alcohols **2a–2z**, **2a'–2c'** To Form 1,2-Amino Alcohols **3a–3z**, **3a'–3c'**^a**



^aYields are of material isolated by silica gel chromatography. Diastereoselectivities were determined by ¹H NMR of crude reaction mixtures. Enantioselectivities were determined by chiral stationary phase HPLC analysis. Standard conditions: 0.2 mmol scale, 48 h. See [Supporting Information](#) for experimental details. ^b72 h, ^cIr-V, ^dIr-VI (7.5 mol%).





Figure 2. Product formation as monitored by ^1H NMR analysis in reactions conducted using the “different excess” protocol: $[\text{Ir}] = 0.01\text{ M}$ and $[\text{KH}_2\text{PO}_4] = 0.2\text{ M}$, with (left) $[2\text{a}]_0 = 0.2\text{ M}$, $[1]_0 =$ as noted and (middle) $[1]_0 = 0.3\text{ M}$, $[2\text{a}]_0 =$ as noted. (Right) Product formation as monitored by NMR analysis in reactions with varying catalyst loading: $[1] = 0.3\text{ M}$, $[2\text{a}] = 0.2\text{ M}$, $[\text{KH}_2\text{PO}_4] = 0.2\text{ M}$, and $[\text{cat.}] =$ as noted.

H_8 -BINAP iridium complex bound by 3,4-dinitro-*C,O*-benzoic acid, Ir-VI, provided still higher levels of enantioselectivity. Finally, introduction of monobasic potassium phosphate led to higher conversion, allowing **3a** to be formed in 80% yield, 96% ee with complete *anti*-diastereoselectivity (Table 1). As borne out by single-crystal X-ray diffraction analysis of Ir-VI (see Supporting Information), the dihedral angle between the tetralin rings of (*R*)- H_8 -BINAP (ca. 86°) is significantly larger than the dihedral angle between the naphthalene rings of BINAP (ca. 75°) or SEGPHOS (ca. 72°),¹⁴ which may better accommodate the sterically demanding phthalimide moiety to facilitate alkoxide exchange at the metal center.


Reaction scope was evaluated by applying optimal conditions identified for the (α -amino)allylation of 2-phenylethanol **2a** to diverse alcohols **2b–2z**, **2a'–2c'** (Table 2). All vicinal amino alcohols **3a–3z**, **3a'–3c'** were formed in good yield with excellent levels of diastereo- and enantioselectivity. The (α -amino)allylations of *N*-Boc-ethanolamine **2j**, *N*-Boc-propanolamine **2k**, and trifluorobutanol **2m**, which are commercially available, are significant, as the corresponding aldehydes are not available for purchase and are relatively unstable. Modification of the heteroaryl-containing alcohols **2c–2i**, **2t**, and **2u**, which includes perphenazine **2g**, an FDA approved drug, establishes the feasibility of utilizing this method for late-stage functionalization.¹⁵ Due to a pronounced kinetic bias for primary alcohol dehydrogenation,¹⁶ free secondary hydroxyl groups are tolerated, as illustrated in the site-selective formation of (*R*)-butane diol adducts **3b'** and **3c'**, which occur with complete levels of catalyst-directed diastereoselectivity. Using this first generation catalytic system, benzylic alcohols are converted to the amino alcohols in high yield but lower enantioselectivities are observed. As demonstrated by the conversion of *dehydro-2l* to amino alcohol **3l**, the reactions can also be conducted from the aldehyde oxidation level using 2-propanol as terminal reductant (eq 1). Given the frequent appearance of morpholines as substructures in pharmaceutical ingredients,¹⁷ compound **3a** was converted to the morpholine **5a** (eq 2).¹⁸ To further demonstrate utility of amino alcohols **3a–3z**, **3a'–3c'**, adduct **3m** was subjected to alkene oxidative cleavage to provide the nonproteinogenic amino acid derivative **6m** (eq 3).¹⁹

Reaction progress kinetic analysis (RPKA) using the “different excess” protocol was used to gain mechanistic insight (Figure 2).²⁰ The kinetic order of reactants varied over time; therefore, general trends were evaluated. Doubling the initial concentration of allene **1** slightly decreases the rate of product formation. This data suggests allene hydrometalation is rapid, allene **1** is not involved in the turnover-limiting step and, at higher concentrations, allene **1** inhibits the rate of product formation (Figure 2, left). Doubling the initial concentration of



alcohol **2a** results in a slight increase in the rate of product formation, signifying a positive order in alcohol **2a** (Figure 2, middle). Increasing the loading of iridium catalyst, (*R*)-Ir-VI, results in a dramatic increase in the rate of product formation, demonstrating the reaction is positive order in catalyst (Figure 2, right). Separate experiments using the “same excess” protocol reveal significant catalyst deactivation that is contributed to by product inhibition.²¹ Additionally, introduction of aldehyde *dehydro-2a* (10 mol%) inhibits product formation, suggesting carbonyl addition may not be turnover limiting.²¹

Deuterium labeling studies provide additional information on the reaction mechanism (eqs 4–6).²² Exposure of allene **1** to *deuterio-2a* under standard reaction conditions delivers *deuterio-3a* (eq 4). Deuterium is completely retained at the carbinol position, suggesting *deuterio-3a* is inert with respect to dehydrogenation. Incorporation of deuterium at both the internal and terminal vinylic positions corroborates reversible allene hydrometalation with incomplete regioselectivity. In a competition kinetics experiment, allene **1** was exposed to equimolar quantities of alcohol **2a** and *deuterio-2a* (eq 5). The observed levels of deuterium incorporation at the carbinol position of *deuterio-3a* are consistent with a normal primary kinetic isotope effect ($k_{\text{H}}/k_{\text{D}} \approx 2.3$). Evaluation of the initial rates for the reaction of both **2a** and *deuterio-2a* also reveals a primary kinetic isotope effect ($k_{\text{H}}/k_{\text{D}} \approx 1.5$) (Figure 3). Along with the reaction orders suggested from the RPKA experiments, this kinetic isotope effect data was consistent with two scenarios: (1) reversible alcohol dehydrogenation followed by rate-



determining carbonyl addition, or (2) rate-determining alcohol dehydrogenation.²² To determine which of these processes is operative an additional experiment was undertaken (eq 6). When phthalimido-allene **1** is exposed to equimolar quantities of *deuterio-2a* and *dehydro-2l* under standard conditions, hydrogen–deuterium exchange is not observed at the carbinol position of *deuterio-3a* and *dehydro-3l*, suggesting alcohol–aldehyde redox equilibration does not occur in advance of carbonyl addition. Hence, the collective data implicate turnover-limiting alcohol dehydrogenation followed by rapid allene hydrometalation.





Figure 3. Initial rates study: $[1]_0 = 0.3$ M; $[2a]_0$ or $[deuterio-2a]_0 = 0.2$ M; $[cat.] = 0.01$ M.

Based on the kinetic and isotopic labeling studies, the indicated catalytic mechanism is proposed (Scheme 1). Entry into the catalytic cycle occurs through protonolysis of the allyliridium complex (R)-Ir-VI by the reactant alcohol. The resulting iridium alkoxide **I** undergoes irreversible dehydrogenation to form the iridium hydride **II**, which is rapidly consumed by reversible allene hydrometalation. Due to the steric demand of the phthalimide moiety, the (*Z*)- σ -(amino)allyliridium complex **IIIa** is anticipated to be the kinetic product of allene hydrometalation. Isomerization to the thermodynamically preferred (*E*)- σ -allyliridium complex **IIIb** is followed by aldehyde coordination and carbonyl addition through a closed chair-like transition structure to form iridium(III) alkoxide **IV**.

Scheme 1. General Catalytic Mechanism As Corroborated by Kinetic and Isotopic Labeling Studies



Exchange with the primary alcohol reactant releases product and regenerates iridium alkoxide **I** to close the catalytic cycle.

In summary, we report a catalytic method for the direct conversion of primary alcohols to vicinal amino alcohols that occurs with high levels of regio-, *anti*-diastereo-, and enantioselectivity. This hydrogen auto-transfer process exploits the tractable, crystalline phthalimido-allene **1** as pronucleophile and represents the first protocol for catalytic enantioselective carbonyl (α -amino)allylation. More broadly, this work contributes to an evolution from use of traditional carbonyl addition methods that exploit preformed carbanions to byproduct-free catalytic carbonyl reductive couplings, where alcohol proelectrophiles and π -unsaturated pronucleophiles combine by way of transient organometallics.⁴

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.9b08715.

- Experimental procedures and spectral data (PDF)
- X-ray diffraction data for compound **3a** (CIF)
- X-ray diffraction data for compound **3v** (CIF)
- X-ray diffraction data for the (R)-H₈-BINAP-modified iridium complex Ir-VI (CIF)

AUTHOR INFORMATION

Corresponding Author

*mkrische@mail.utexas.edu

ORCID

Ming Xiang: 0000-0002-2656-8804

Michael J. Krische: 0000-0001-8418-9709

Author Contributions

All authors contributed equally.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

The Robert A. Welch Foundation (F-0038), the NIH-NIGMS (RO1-GM069445, 1 S10 OD021508-01). We thank Professor Kami Hull for helpful discussions of the kinetic data.

REFERENCES

- (1) For selected reviews on carbonyl addition chemistry, see: (a) Noyori, R.; Kitamura, M. *Enantioselective Addition of Organo-*

metallic Reagents to Carbonyl Compounds: Chirality Transfer, Multiplication and Amplification. *Angew. Chem., Int. Ed. Engl.* **1991**, *30*, 49. (b) Soai, K.; Shibata, T. Alkylation of Carbonyl Groups. In *Comprehensive Asymmetric Catalysis I-III*; Jacobsen, E. N., Pfaltz, A., Yamamoto, H., Eds.; Springer-Verlag Berlin Heidelberg: Germany, 1999; Vol. 2, pp 911. (c) Pu, L.; Yu, H.-B. Catalytic Asymmetric Organozinc Additions to Carbonyl Compounds. *Chem. Rev.* **2001**, *101*, 757. (d) Trost, B. M.; Weiss, A. H. The Enantioselective Addition of Alkyne Nucleophiles to Carbonyl Groups. *Adv. Synth. Catal.* **2009**, *351*, 963. (e) *Comprehensive Organic Synthesis*, 2nd ed.; Knochel, P.; Molander, G. A., Eds.; Elsevier: Oxford, 2014; Vols. 1 and 2.

(2) Schneider, N.; Lowe, D. M.; Sayle, R. A.; Tarselli, M. A.; Landrum, G. A. Big Data from Pharmaceutical Patents: A Computational Analysis of Medicinal Chemists' Bread and Butter. *J. Med. Chem.* **2016**, *59*, 4385.

(3) For a recent review on metal-catalyzed carbonyl reductive coupling, see: Holmes, M.; Schwartz, L. A.; Krische, M. J. Intermolecular Metal-Catalyzed Reductive Coupling of Dienes, Allenes, and Enynes with Carbonyl Compounds and Imines. *Chem. Rev.* **2018**, *118*, 6026.

(4) For selected reviews on carbonyl reductive coupling via hydrogen auto-transfer, see: (a) Kim, S. W.; Zhang, W.; Krische, M. J. Catalytic Enantioselective Carbonyl Allylation and Propargylation via Alcohol-Mediated Hydrogen Transfer: Merging the Chemistry of Grignard and Sabatier. *Acc. Chem. Res.* **2017**, *50*, 2371. (b) Doerksen, R. S.; Meyer, C. C.; Krische, M. J. Feedstock Reagents in Metal-Catalyzed Carbonyl Reductive Coupling: Minimizing Preactivation for Efficiency in Target-Oriented Synthesis. *Angew. Chem., Int. Ed.* **2019**, DOI: 10.1002/anie.201905532.

(5) (a) Ghislieri, D.; Turner, N. J. Biocatalytic Approaches to the Synthesis of Enantiomerically Pure Chiral Amines. *Top. Catal.* **2014**, *57*, 284. (b) Breuer, M.; Ditrach, K.; Habicher, T.; Hauer, B.; Kessler, M.; Stürmer, R.; Zelinski, T. Industrial Methods for the Production of Optically Active Intermediates. *Angew. Chem., Int. Ed.* **2004**, *43*, 788. (c) Sehl, T.; Maugeri, Z.; Rother, D. Multi-step Synthesis Strategies towards 1,2-Amino Alcohols with Special Emphasis on Phenylpropanolamine. *J. Mol. Catal. B: Enzym.* **2015**, *114*, 65 and references cited therein.

(6) For racemic vicinal amino alcohols via metal-catalyzed carbonyl (α -amino)allylation, see: (a) Skucas, E.; Zbieg, J. R.; Krische, M. J. *anti*-Aminoallylation of Aldehydes via Ruthenium Catalyzed Transfer Hydrogenative Coupling of Sulfonamido-Allenes: 1,2-Aminoalcohols. *J. Am. Chem. Soc.* **2009**, *131*, 5054. (b) Zbieg, J. R.; McInturff, E. L.; Krische, M. J. Allenamide Hydro-Hydroxyalkylation: 1,2-Aminoalcohols via Ruthenium Catalyzed Carbonyl *anti*-Aminoallylation. *Org. Lett.* **2010**, *12*, 2514. (c) Zhang, W.; Chen, W.; Xiao, H.; Krische, M. J. Carbonyl *anti*-(α -Amino)allylation via Ruthenium Catalyzed Hydrogen Auto-Transfer: Use of an Acetylenic Pyrrole as an Allylmetal Pronucleophile. *Org. Lett.* **2017**, *19*, 4876. (d) Schwarz, J. L.; Schäfers, F.; Tlahuext-Aca, A.; Lückemeier, L.; Glorius, F. Diastereoselective Allylation of Aldehydes by Dual Photoredox and Chromium Catalysis. *J. Am. Chem. Soc.* **2018**, *140*, 12705.

(7) (a) Barrett, A. G. M.; Seefeld, M. A. B-[(*E*)-3-(Diphenylamino)-allyl]diisopinocampheylborane: An Excellent Reagent for The Stereoselective Synthesis of *anti*- β -Diphenylamino Alcohols. *J. Chem. Soc., Chem. Commun.* **1993**, 339. (b) Barrett, A. G. M.; Seefeld, M. A. The Use of B-[(*E*)-3-(Diphenylamino)allyl]diisopinocampheylborane as A Reagent for The Stereoselective Synthesis of *anti*- β -Diphenylamino Alcohols and *trans*-1-Diphenylamino-2-(1-hydroxylalkyl)-cyclopropanes. *Tetrahedron* **1993**, *49*, 7857. (c) Barrett, A. G. M.; Seefeld, M. A.; Williams, D. J. Convenient Asymmetric Synthesis of *anti*- β -Amino Alcohols: An X-Ray Crystallographic Study of (4*R*)-2,2-Dimethyl-4-[(2*S*)-(diphenylmethyleneamino)-(1*S*)-hydroxy-3-buten-1-yl]-1,3-dioxolane. *J. Chem. Soc., Chem. Commun.* **1994**, 1053. (d) Barrett, A. G. M.; Seefeld, M. A.; White, A. J. P.; Williams, D. J. Convenient Asymmetric Syntheses of *anti*- β -Amino Alcohols. *J. Org. Chem.* **1996**, *61*, 2677.

(8) For selected reviews on enantioselective carbonyl allylation, see: (a) Ramachandran, P. V. Pinane-based versatile "allyl" boranes. *Aldrichimica Acta* **2002**, *35*, 23. (b) Denmark, S. E.; Fu, J. Catalytic

Enantioselective Addition of Allylic Organometallic Reagents to Aldehydes and Ketones. *Chem. Rev.* **2003**, *103*, 2763. (c) Yu, C.-M.; Youn, J.; Jung, H.-K. Regulation of Stereoselectivity and Reactivity in the Inter- and Intramolecular Allylic Transfer Reactions. *Bull. Korean Chem. Soc.* **2006**, *27*, 463. (d) Marek, I.; Sklute, G. Creation of Quaternary Stereocenters in Carbonylallylation Reactions. *Chem. Commun.* **2007**, 1683. (e) Hall, D. G. Lewis and Brønsted Acid Catalyzed Allylboration of Carbonyl Compounds: From Discovery to Mechanism and Applications. *Synlett* **2007**, *11*, 1644. (f) Hargaden, G. C.; Guiry, P. J. The Development of the Asymmetric Nozaki-Hiyama-Kishi Reaction. *Adv. Synth. Catal.* **2007**, *349*, 2407. (g) Lachance, H.; Hall, D. G. Allylboration of Carbonyl Compounds. *Org. React.* **2009**, *73*, 1. (h) Yus, M.; González-Gómez, J. C.; Foubelo, F. Catalytic Enantioselective Allylation of Carbonyl Compounds and Imines. *Chem. Rev.* **2011**, *111*, 7774. (i) Yus, M.; Gonzalez-Gomez, J. C.; Foubelo, F. Diastereoselective Allylation of Carbonyl Compounds and Imines: Application to the Synthesis of Natural Products. *Chem. Rev.* **2013**, *113*, 5595. (j) Huo, H.-X.; Duvall, J. R.; Huang, M.-Y.; Hong, R. Catalytic Asymmetric Allylation of Carbonyl Compounds and Imines with Allylic Boronates. *Org. Chem. Front.* **2014**, *1*, 303. (k) Kumar, P.; Tripathi, D.; Sharma, B. M.; Dwivedi, N. Transition Metal Catalysis - A Unique Road Map in the Stereoselective Synthesis of 1,3-Polyols. *Org. Biomol. Chem.* **2017**, *15*, 733. (l) Spielmann, K.; Niel, G.; de Figueiredo, R. M.; Campagne, J. M. Catalytic Nucleophilic 'Unpoled' π -Allyl Reagents. *Chem. Soc. Rev.* **2018**, *47*, 1159.

(9) (a) Li, K.; Shao, X.; Tseng, L. M.; Malcolmson, S. J. 2-Azadienes as Reagents for Preparing Chiral Amines: Synthesis of 1,2-Amino Tertiary Alcohols by Cu-Catalyzed Enantioselective Reductive Couplings with Ketones. *J. Am. Chem. Soc.* **2018**, *140*, 598. (b) Shao, X.; Li, K.; Malcolmson, S. Enantioselective Synthesis of *anti*-1,2-Diamines by Cu-Catalyzed Reductive Couplings of Azadienes with Aldimines and Ketimines. *J. Am. Chem. Soc.* **2018**, *140*, 7083.

(10) Only five catalytic enantioselective allene-carbonyl reductive couplings have been reported, see: (a) Han, S. B.; Kim, I. S.; Han, H.; Krische, M. J. Enantioselective Carbonyl Reverse Prenylation from the Alcohol or Aldehyde Oxidation Level Employing 1,1-Dimethylallene as the Prenyl Donor. *J. Am. Chem. Soc.* **2009**, *131*, 6916. (b) Holmes, M.; Nguyen, K. D.; Schwartz, L. A.; Luong, T.; Krische, M. J. Enantioselective Formation of CF₃-Bearing All-Carbon Quaternary Stereocenters via C-H Functionalization of Methanol: Iridium Catalyzed Allene Hydrohydroxymethylation. *J. Am. Chem. Soc.* **2017**, *139*, 8114. (c) Tsai, E. Y.; Liu, R. Y.; Yang, Y.; Buchwald, S. L. A Region- and Enantioselective CuH-Catalyzed Ketone Allylation with Terminal Allenes. *J. Am. Chem. Soc.* **2018**, *140*, 2007. (d) Schwartz, L. A.; Holmes, M.; Brito, G. A.; Gonçalves, T. P.; Richardson, J.; Ruble, J. C.; Huang, K.-W.; Krische, M. J. Cyclometallated Iridium-PhanePhos Complexes Are Active Catalysts in Enantioselective Allene-Fluoral Reductive Coupling and Related Alcohol-Mediated Carbonyl Additions that Form Acyclic Quaternary Carbon Stereocenters. *J. Am. Chem. Soc.* **2019**, *141*, 2087. (e) Liu, R. Y.; Zhou, Y.; Yang, Y.; Buchwald, S. L. Enantioselective Allylation Using Allene, a Petroleum Cracking Byproduct. *J. Am. Chem. Soc.* **2019**, *141*, 2251.

(11) (a) Fernandez, I.; Monterde, M. I.; Plumet, J. On the Base-Induced Isomerization of Cyclic Propargylamides to Cyclic Allenamides. *Tetrahedron Lett.* **2005**, *46*, 6029. (b) Lindsay, V. N. G.; Fiset, D.; Gritsch, P. J.; Azzi, S.; Charette, A. B. Stereoselective Rh₂(S-IBAZ)₄-Catalyzed Cyclopropanation of Alkenes, Alkynes, and Allenes: Asymmetric Synthesis of Diacceptor Cyclopropylphosphonates and Alkylidene-cyclopropanes. *J. Am. Chem. Soc.* **2013**, *135*, 1463.

(12) Bower, J. F.; Skucas, E.; Patman, R. L.; Krische, M. J. Catalytic C-C Coupling via Transfer Hydrogenation: Reverse Prenylation, Crotylation and Allylation from the Alcohol or Aldehyde Oxidation Level. *J. Am. Chem. Soc.* **2007**, *129*, 15134.

(13) (a) Zhang, X.; Mashima, K.; Koyano, K.; Sayo, N.; Kumobayashi, H.; Akutagawa, S.; Takaya, H. Synthesis of Partially Hydrogenated BINAP Variants. *Tetrahedron Lett.* **1991**, *32*, 7283. (b) Zhang, X.; Taketomi, T.; Yoshizumi, T.; Kumobayashi, H.; Akutagawa, S.; Mashima, K.; Takaya, H. Asymmetric Hydrogenation of Cycloalkanes Catalyzed by BINAP-Ir(I)-Aminophosphine Systems. *J.*

Am. Chem. Soc. **1993**, *115*, 3318. (c) Zhang, X.; Mashima, K.; Koyano, K.; Sayo, N.; Kumobayashi, H.; Akutagawa, S.; Takaya, H. Synthesis of Partially Hydrogenated 2,2'-Bis(diphenylphosphanyl)-1,1'-binaphthyl (BINAP) Ligands and Their Application to Catalytic Asymmetric Hydrogenation. *J. Chem. Soc., Perkin Trans. 1* **1994**, *1*, 2309.

(14) For X-ray diffraction data on analogous π -allyliridium-C,*O*-benzoate complexes modified by (*R*)-SEGPHOS and (*R*)-BINAP, see CCDC929090 and CCDC712857, respectively.

(15) For selected reviews on late-stage functionalization of complex small molecules, see: (a) Cernak, T.; Dykstra, K. D.; Tyagarajan, S.; Vachal, P.; Krska, S. W. The Medicinal Chemist's Toolbox for Late Stage Functionalization of Drug-Like Molecules. *Chem. Soc. Rev.* **2016**, *45*, 546. (b) Shugrue, C. R.; Miller, S. J. Applications of Nonenzymatic Catalysts to the Alteration of Natural Products. *Chem. Rev.* **2017**, *117*, 11894. (c) Blakemore, D. C.; Castro, L.; Churcher, I.; Rees, D. C.; Thomas, A. W.; Wilson, D. M.; Wood, A. Organic Synthesis Provides Opportunities to Transform Drug Discovery. *Nat. Chem.* **2018**, *10*, 383. (d) Dominguez-Huerta, A.; Dai, X.-J.; Zhou, F.; Querard, P.; Qiu, Z.; Ung, S.; Liu, W.; Li, J.; Li, C.-J. Exploration of New Reaction Tools for Late-Stage Functionalization of Complex Chemicals. *Can. J. Chem.* **2019**, *97*, 67.

(16) Although primary alcohol dehydrogenation is more endothermic than secondary alcohol dehydrogenation, primary alcohol dehydrogenation is kinetically preferred due to the steric demand of the catalyst. Site-selective methods for the C–C coupling of diols and higher polyols are summarized in ref 4a.

(17) For selected reviews highlighting the recurrence of *N*-heterocycles among FDA approved drugs, see: (a) Taylor, R. D.; MacCoss, M.; Lawson, A. D. G. Rings in Drugs. *J. Med. Chem.* **2014**, *57*, 5845. (b) Vitaku, E.; Smith, D. T.; Njardarson, J. T. Analysis of the Structural Diversity, Substitution Patterns, and Frequency of Nitrogen Heterocycles among U.S. FDA Approved Pharmaceuticals. *J. Med. Chem.* **2014**, *57*, 10257.

(18) The synthesis of morpholine **5a** was conducted in analogy to the following procedures: (a) Alexander, R.; Balasundaram, A.; Batchelor, M.; Brookings, D.; Crépy, K.; Crabbe, T.; Deltent, M.-F.; Driessens, F.; Gill, A.; Harris, S.; Hutchinson, G.; Kulisa, C.; Merriman, M.; Mistry, P.; Parton, T.; Turner, J.; Whitcombe, I.; Wright, S. 4-(1,3-Thiazol-2-yl)morpholine Derivatives as Inhibitors of Phosphoinositide 3-Kinase. *Bioorg. Med. Chem. Lett.* **2008**, *18*, 4316. (b) Hu, P.; Hu, J.; Jiao, J.; Tong, X. Amine-Promoted Asymmetric (4 + 2) Annulations for the Enantioselective Synthesis of Tetrahydropyridines: A Traceless and Recoverable Auxiliary Strategy. *Angew. Chem., Int. Ed.* **2013**, *52*, 5319.

(19) Chen, Y. K.; Lurain, A. E.; Walsh, P. J. A General, Highly Enantioselective Method for the Synthesis of *d* and *l* α -Amino Acids and Allylic Amines. *J. Am. Chem. Soc.* **2002**, *124*, 12225.

(20) For a review on reaction progress kinetic analysis, see: Blackmond, D. G. Reaction Progress Kinetic Analysis: A Powerful Methodology for Mechanistic Studies of Complex Catalytic Reactions. *Angew. Chem., Int. Ed.* **2005**, *44*, 4302.

(21) This experiment is described in the [Supporting Information](#).

(22) An authoritative treatment of the execution and interpretation of kinetic isotope effect experiments in reactions involving C–H bond cleavage is reported. Our catalytic mechanism appears consistent with reaction coordinate 1, [Figure 1](#): Simmons, E. M.; Hartwig, J. F. On the Interpretation of Deuterium Kinetic Isotope Effects in C–H Bond Functionalizations by Transition-Metal Complexes. *Angew. Chem., Int. Ed.* **2012**, *51*, 3066.