

## Accepted Manuscript

Facile Azide Formation *via* Diazotransfer Reaction in a Copper Tube Flow Reactor

Koen Nuyts, Matthias Ceulemans, Tatjana N. Parac-Vogt, Geert Bultynck, Wim M. De Borggraeve

PII: S0040-4039(15)00306-8  
DOI: <http://dx.doi.org/10.1016/j.tetlet.2015.02.036>  
Reference: TETL 45904

To appear in: *Tetrahedron Letters*

Received Date: 21 January 2015  
Revised Date: 4 February 2015  
Accepted Date: 10 February 2015

Please cite this article as: Nuyts, K., Ceulemans, M., Parac-Vogt, T.N., Bultynck, G., De Borggraeve, W.M., Facile Azide Formation *via* Diazotransfer Reaction in a Copper Tube Flow Reactor, *Tetrahedron Letters* (2015), doi: <http://dx.doi.org/10.1016/j.tetlet.2015.02.036>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.




**Graphical Abstract**

To create your abstract, type over the instructions in the template box below.  
Fonts or abstract dimensions should not be changed or altered.

**Facile Azide Formation via Diazotransfer Reaction in a Copper Tube Flow Reactor**

Leave this area blank for abstract info.

Koen Nuyts, Matthias Ceulemans, Tatjana N. Parac-Vogt, Geert Bultynck, Wim M. De Borggraeve





## Facile Azide Formation *via* Diazotransfer Reaction in a Copper Tube Flow Reactor

Koen Nuyts<sup>a</sup>, Matthias Ceulemans<sup>a</sup>, Tatjana N. Parac-Vogt<sup>a</sup>, Geert Bultynck<sup>b</sup> and Wim M. De Borggraeve<sup>a\*</sup>

<sup>a</sup>Molecular Design and Synthesis, KU Leuven – University of Leuven, Celestijnenlaan 200F, 3001 Leuven, Belgium

<sup>b</sup>Laboratory of Molecular and Cellular Signaling, KU Leuven – University of Leuven, O&N I Herestraat 49, 3000 Leuven, Belgium

### ARTICLE INFO

#### Article history:

Received  
Received in revised form  
Accepted  
Available online

#### Keywords:

Diazotransfer reaction  
Flow chemistry  
Azides  
Copper reactor

### ABSTRACT

A copper tube flow reactor is used in the conversion of primary amines into organic azides using imidazole-1-sulfonyl azide hydrogen sulfate. The catalyst is generated *in situ* from the metallic copper. The reaction can be quenched in acidic environment or *via* a cycloaddition of the azides formed with an alkyne. The possibility to perform this azide-alkyne cycloaddition using the copper released from the reactor is demonstrated with the synthesis of both a 1,2,3-triazole derivative of benzylamine and of a more complex BODIPY-DOTA adduct.

2009 Elsevier Ltd. All rights reserved.

Flow chemistry is a useful tool for the synthetic chemist.<sup>1-4</sup> The ease with which flow reactors can be constructed by utilizing standard tubing, connector pieces and syringe pumps, the possibility to tightly control reaction time and reaction temperature, the small quantities of product required for optimization are great advantages. Furthermore, safety issues such as toxicity and explosive decomposition of reagents are mitigated by the closed system and small reaction volumes.<sup>1-5</sup> The latter argument shows great potential for flow chemistry in the synthesis of organic azide.

Organic azides are a class of compounds with potential highly toxic and explosive properties.<sup>6</sup> Despite these attributes they are valuable and versatile intermediates in organic synthesis.<sup>7</sup> Organic azides can be synthesized from primary amines *via* a diazotransfer reaction.<sup>8,9</sup> Triflyl azide is most commonly used as a diazotransfer reagent.<sup>10</sup> Its short shelf life and highly explosive nature in neat form render this product rather unattractive.<sup>11,12</sup> The introduction of imidazole-1-sulfonyl azide (**ISA**) and more recently its hydrogen sulfate salt (**ISA·H<sub>2</sub>SO<sub>4</sub>**, Scheme 1) as a stable, easy to prepare and manipulate reagent has given the diazotransfer reaction a whole new set of applications.<sup>13-19</sup>

- Scheme 1 -

Recently, the possibility to perform the diazotransfer reaction under continuous flow conditions has been demonstrated by Delville *et al.*<sup>20</sup> However, issues with solubility of some of our compounds prompted us to use a different solvent system and a different catalyst. While screening for appropriate catalysts in preparation of performing the diazotransfer reaction under

continuous flow conditions, it was found that it was possible to catalyze the reaction by using copper turnings. Oxidation of metallic copper under the influence of a sulfonyl azide results in the formation of a catalytic active Cu(II)-sulfonylamide complex.<sup>21,22</sup> This prompted us to investigate the possibility to perform this reaction under flow conditions using a copper tube reactor. The use of metallic copper as a source of an ionic catalyst system has previously been demonstrated, *e.g.* in the synthesis of 1,4-disubstituted 1,2,3-triazoles.<sup>23,24</sup> The use of copper flow reactors has already been successfully applied in the synthesis of 1,4-disubstituted 1,2,3-triazoles and several coupling reactions.<sup>25-40</sup> To the best of our knowledge, the application of copper tube flow reactors was not reported for the diazotransfer reaction.

- Figure 1 -

A general procedure was developed *via* the optimization of the synthesis of  $\gamma$ -azidobutyric acid (**2**) starting from  $\gamma$ -aminobutyric acid (**1**) and **ISA·H<sub>2</sub>SO<sub>4</sub>** (Scheme 1). A 500  $\mu$ L internal volume reactor was constructed using copper GC tubing (Figure 1). The reactor was connected to two syringes *via* a T-mixing piece. A 10:3:3 mixture of methanol:dichloromethane:water was chosen as the solvent since this monophasic mixture is reported to result in a minimal precipitation of salts from the diazotransfer reaction.<sup>7,41</sup> This mixture also proved to have better solvating properties for the reagents used. The amine and diazotransfer reagent were dissolved in separate stock solutions that were brought in two separate syringes. Stock solution A is 145 mM in reagent **1**. Stock solution B contains a solution with variable quantities of **ISA·H<sub>2</sub>SO<sub>4</sub>** (according to the number of equivalents needed).

\* Corresponding author. Tel.: +32-16-327693; fax: +32-16-327990; e-mail: wim.deborggraeve@chem.kuleuven.be

Base was added as a second component to solution A in a fixed ratio to **ISA·H<sub>2</sub>SO<sub>4</sub>** (Supplementary Material). When the base is premixed with **ISA·H<sub>2</sub>SO<sub>4</sub>**, premature decomposition of this reagent is observed in the syringe as formation of gas bubbles. An ice bath was used to immerse the T-mixer to prevent decomposition while combining the mixtures before bringing them into the copper tube flow reactor at room temperature. Upon leaving the reactor the reaction mixture was directly quenched by adding the reaction mixture to acetic anhydride. Reaction time and equivalents of **ISA·H<sub>2</sub>SO<sub>4</sub>** were optimized and monitored *via* LC/MS (Figure 2 and Figure 3).

- Figure 2 –
- Figure 3-

Using less than three equivalents of **ISA·H<sub>2</sub>SO<sub>4</sub>** relative to the amine results in incomplete conversion of the amine. At the same time it was observed that a precipitate forms in the reactor. By using three or more equivalents of **ISA·H<sub>2</sub>SO<sub>4</sub>** this problem is alleviated. A comparable rate of amine conversion is observed when three or four equivalents of **ISA·H<sub>2</sub>SO<sub>4</sub>**. However, overall formation of azide is lower in the case where four equivalents are used. Addition of five equivalents results in both lower conversion of the amine and less formation of the azide. At shorter retention times the amine does not completely convert to the azide. Longer retention times result in lower yields of the azide. This could be due to further reduction of the formed azides in the presence of copper.<sup>21, 42</sup> Generally, the amine is converted optimally when a retention time of the mixture in the reactor of 60-300 seconds is used. The scope and limitations were studied at room temperature using 3 equivalents of **ISA·H<sub>2</sub>SO<sub>4</sub>** and a retention time of 300 seconds (Table 1). The reaction was quenched by introducing the mixture to a solution of 10 equivalents of 5-chloro-1-pentyne and CuI or in a 3:1 solution of methanol and acetic acid, resulting in the formation of the triazole **xb** or azide **xc** respectively (Figure 4).

- Figure 4 –

For ease of purification the ethyl ester of  $\gamma$ -aminobutyric acid was used to determine the isolated yield and the acidic quench was replaced with a quench using a mixture of 10 equivalents of 5-chloro-1-pentyne and CuI yielding the corresponding 1,4-triazole *via* a copper(I) catalyzed azide-alkyne cycloaddition (Cu AAC). This quenches the reaction and yields a less volatile compound. Product **1b** was obtained in a 59% overall isolated yield over the two reaction steps. Conversion of benzylamine and Fmoc-Lys-OH proceeded with yields of 77% and 74% respectively. The diazotransfer to compound **4a** resulted in selective transformation of the alkyl amine, leaving the hetaryl amine intact. Product **4c** was obtained in a 65% yield. Conversion of aromatic amines proved to be more problematic due to the instability of their corresponding azides. To prevent this undesired degradation during purification, all the organic azides were converted to their respective 1,4-triazoles by introducing the reaction mixture to a 5-chloro-1-pentyne/CuI solution. The isolated yield of these compounds decreases from 44% for **5b** to 26% for **6b** to 7% for **7b**, with no conversion observed for **8a**. This decrease within the aromatic amines and compared to the alkyl amines corresponds to a decrease in nucleophilicity of the nitrogen atom. This confirms the need of a nucleophilic amine in the diazotransfer reaction as proposed in the mechanism of Nyffeler *et al* and the elucidated mechanism of Stevens *et al*.<sup>7, 41, 43</sup>

- Table 1-

After each run the increase of the internal volume of the copper tube reactor was checked by weighing the reactor. This increase due to the previously mentioned oxidation of Cu(0) amounted to less than 0,5% of the total internal volume and was deemed insignificant.

To verify the possibility of cascading the diazotransfer reaction with the Cu AAC using the copper ions formed in the reactor tube, the synthesis of compound **2b** was repeated in one continuous process. The azide of compound **2a** was introduced to a solution 1,5 equivalents of 5-chloro-1-pentyne and 1,5 equivalents of sodium ascorbate. The reaction mixture was left stirring overnight and compound **2b** was isolated in a 57% overall yield. The same procedure was used in the synthesis of compound **11** (Scheme 2). The azide of compound **9** could not be isolated in batch diazotransfer reactions. Hence, azide **9b** was synthesized *in situ* by performing the herein developed flow procedure in a 150  $\mu$ L copper tube flow reactor. The reaction mixture was introduced to a flask containing 1,5 equivalents of sodium ascorbate and 1,5 equivalents of **10**. The reaction was left stirring overnight and was purified *via* HPLC. The previously elusive BODIPY functionalized complex **11** was obtained in a 9% overall isolated yield starting from **9**.

- Scheme 2 –

In conclusion, a straightforward flow protocol is developed for the synthesis of azides *via* a diazotransfer reaction using a copper tube flow reactor. This procedure requires less diazotransfer reagent, a lower retention time and no addition of catalyst to any of the stock solutions compared to the procedure of Delville *et al*.<sup>20</sup> It was shown that the reaction proceeds with higher yields using alkyl amines compared to aryl amines. Diazotransfer reactions to aryl amines are more efficient if the amine is more nucleophilic. The possibility to perform the Cu AAC using the copper released from the system is demonstrated by the synthesis of compound **2b** and compound **11**. Once constructed, the copper tube flow reactor, the easily applicable procedure and the possibility of cascading the diazotransfer reaction with the Cu AAC puts a wide range of azides and 1,4-triazoles within easy reach of a wide variety of scientists.

## Acknowledgments

K.N. thanks the research council of KU Leuven and the department of chemistry of KU Leuven for funding (project OT/11/047, project OT/14/067).

## Supplementary Material

Experimental procedures, characterization data.

## References and notes

1. Baxendale, I. R.; Brocken, L.; Mallia, C. J. *Green Process. Synth.* **2013**, *2*, 211-230.
2. Kirschning, A.; Wladimier, S.; Mennecke, K. *Chem. Eur. J.* **2006**, *12*, 5972-5990.
3. Wegner, J.; Ceylan, S.; Kirschning, A. *Adv. Synth. Catal.* **2012**, *354*, 17-57.
4. Ley, S. V.; Fitzpatrick, D. E.; Ingham, R. J.; Myers, R. M. *Ang. Chem. Int. Ed.* **2015**, *54*, 2-18.
5. Hessel, V.; Kralisch, D.; Kockmann, N.; Noël, T.; Wang, Q. *ChemSusChem* **2013**, *6*, 746-489.
6. Bräse, S.; Gil, C.; Knepper, K.; Zimmermann, V. *Ang. Chem. Int. Ed.* **2005**, *44*, 5188-5240.

7. Nyffeler, P. T.; Liang, C.-H.; Koeller, K. M.; Wong, C.-H. *J. Am. Chem. Soc.* **2002**, *124*, 10773-10778.
8. Alper, P. B.; Hung, S.-C.; Wong, C.-H. *Tetrahedron Lett.* **1996**, *37*, 6029-60332.
9. Cavender, C. J.; Shiner, V. J. *J. Org. Chem.* **1972**, *37*, 3567-3569.
10. *Organic Azides - Synthesis and Applications*. John Wiley & Sons Ltd: Chippingham, Wiltshire, 2010.
11. Johansson, H.; Pedersen, D. S. *Eur. J. Org. Chem.* **2012**, 4267-4281.
12. Sminia, T. J.; Pedersen, D. S. *Synlett* **2012**, *23*, 2643-2646.
13. Bastian, A. A.; Warszawik, E. M.; Panduru, P.; Arenz, C.; Herrmann, A. *Chem. Eur. J.* **2013**, *19*, 9151-9154.
14. Castro, V.; Blanco-Canosa, J. B.; Rodriguez, H.; Albericio, F. *ACS Comb. Sci.* **2013**, *15*, 331-334.
15. Fischer, N.; Goddard-Borger, E. D.; Greiner, R.; Klapötke, T. M.; Skelton, B. W.; Stierstorfer, J. *J. Org. Chem.* **2012**, *77*, 1760-1764.
16. Goddard-Borger, E. D.; Stick, R. V. *Org. Lett.* **2007**, *9*, 3797-3800.
17. Hansen, M. B.; van Gurp, T. H. M.; van Hest, J. C. M.; Löwik, D. W. P. M. *Org. Lett.* **2012**, *14*, 2330-2333.
18. Schoffelen, S.; van Eldijk, M. B.; Rooijackers, B.; Raijmakers, R.; Heck, A. J. R.; van Hest, J. C. M. *Chem. Sci.* **2011**, *2*, 701-705.
19. van Dongen, S. F. M.; Teeuwen, R. L. M.; Nallani, M.; van Berkel, S. S.; Cornelissen, J. J. L. M.; Nolte, R. J. M.; van Hest, J. C. M. *Bioconjugate Chem.* **2008**, *20*, 20-23.
20. Delville, M. M. E.; Nieuwland, P. J.; Janssen, P.; Koch, K.; van Hest, J. C. M.; Rutjes, F. P. J. T. *Chem. Eng. J.* **2011**, *167*, 556-559.
21. Kwart, H.; Kahn, A. A. *J. Am. Chem. Soc.* **1967**, *89*, 1950-1951.
22. Kwart, H.; Kahn, A. A. *J. Am. Chem. Soc.* **1967**, *89*, 1951-1953.
23. Cintas, P.; Barge, A.; Tagliapietra, S.; Boffa, L.; Cravotto, G. *Nat. Protoc.* **2010**, *5*, 607-616.
24. Cravotto, G.; Fokin, V. V.; Garella, D.; Binello, A.; Boffa, L.; Barge, A. *J. Comb. Chem.* **2010**, *12*, 13-15.
25. Tan, L.-M.; Sem, Z.-Y.; Chong, W.-Y.; Hendra; W.L., K.; C.-L.K., L. *Org. Lett.* **2013**, *15*, 65-67.
26. Tu, N. P.; Hochlowski, J. E.; Djuric, S. W. *Mol. Divers.* **2012**, *16*, 53-58.
27. Zhan, P.; Russell, M. G.; Jamison, T. F. *Org. Process Res. Dev.* **2014**, *18*, 1567-1570.
28. Zhang, Y.; Jamison, T. F.; Patel, S.; Mainolfi, N. *Org. Lett.* **2011**, *13*, 280-283.
29. Bogdan, A. R.; Sach, N. W. *Adv. Synth. Catal.* **2009**, *351*, 849-854.
30. Ceylan, S.; Klände, T.; Vogt, C.; Friese, C.; Kirschning, A. *SYNLETT* **2010**, 2009-2013.
31. Smith, C. D.; Baxendale, I. R.; Lanners, S.; Hayward, J. J.; Smith, S. C.; Ley, S. V. *Org. Biomol. Chem.* **2007**, *5*, 1559-1561.
32. Baxendale, I. R.; Ley, S. V.; Mansfield, A. C.; Smith, C. D. *Ang. Chem. Int. Ed.* **2009**, *48*, 4017-4021.
33. Fuchs, M.; Goessler, W.; Pilger, C.; Kappe, C. O. *Adv. Synth. Catal.* **2010**, *352*, 323.
34. Bogdan, A. R.; James, K. *Chem. Eur. J.* **2010**, *16*, 14506-14512.
35. Bogdan, A. R.; James, K. *Org. Lett.* **2011**, *13*, 4060-4063.
36. Varas, A. C.; Noël, T.; Wang, Q.; Hessel, V. *ChemSusChem* **2012**, *5*, 1703-1707.
37. Borukhova, S.; Noël, T.; Metten, B.; de Vos, E.; Hessel, V. *ChemSusChem* **2013**, *6*, 2220-2225.
38. Borukhova, S.; Seeger, A. D.; Noël, T.; Wang, Q.; Busch, M.; Hessel, V. *ChemSusChem* **2015**, *8*, 504-512.
39. Ötvös, S. B.; Mándity, I. M.; Kiss, L.; Fülöp, F. *Chem. Asian J.* **2013**, *8*, 800-808.
40. Ötvös, S. B.; Hatoss, G.; Georgiádes, A.; Kovács, S.; Mándity, I. M.; Novák, Z.; Fülöp, F. *RSC Adv.* **2014**, *4*, 46666-46674.
41. Stevens, M. Y.; Sawant, R. T.; Odell, L. R. *J. Org. Chem.* **2014**, *79*, 4826-4831.
42. Ahammed, S.; Saha, A.; Ranu, C. *J. Org. Chem.* **2011**, *76*, 7235-7239.
43. Loos, P.; Ronco, C.; Riedric, M.; Arndt, H.-D. *Eur. J. Org. Chem.* **2013**, 3290-3315.

**Scheme 1.** Synthesis of  $\gamma$ -azidobutyric acid via a copper catalyzed diazotransfer reaction using imidazole-1-sulfonyl azide.

**Scheme 2.** Synthesis of the BODIPY functionalized complex **11** by performing a diazotransfer reaction on amine **9**, followed by a Cu AAC with alkyne **10**.


**Figure 1.** Set up of the copper tube flow reactor.

**Figure 2.** Influence of the number of equivalents of  $\text{ISA}\cdot\text{H}_2\text{SO}_4$  and the retention time in the reactor on the conversion of amine **1** relative to an internal standard.

**Figure 3.** Influence of the number of equivalents of  $\text{ISA}\cdot\text{H}_2\text{SO}_4$  and the retention time in the reactor on the formation of azide **2** relative to an internal standard.

**Figure 4.** The diazotransfer reaction of **xa** is either quenched by introducing the mixture to a solution of 5-chloro-1-pentyne and CuI forming triazole **xb**, or quenched by introducing the mixture to a 3:1 solution of methanol and acetic acid forming azide **xc**.

**Table 1.** Diazotransfer reaction at room temperature, 3 equivalents of  $\text{ISA}\cdot\text{H}_2\text{SO}_4$ , 300 seconds retention time in a 500  $\mu\text{L}$  copper tube flow reactor. Yields are isolated yields.

Entry	Amine	Product	Yield (%)
1	<b>1a</b> ethyl 4-aminobutyrate·HCl	<b>1b</b>	59
2	<b>2a</b> benzylamine	<b>2b</b>	77
3	<b>3a</b> Fmoc-Lys-OH·HCl	<b>3c</b>	74
4	<b>4a</b> 	<b>4c<sup>a</sup></b>	65
5	<b>5a</b> p-anisidine	<b>5b</b>	44
6	<b>6a</b> p-toluidine	<b>6b</b>	26
7	<b>7a</b> p-chloroaniline	<b>7b</b>	7
8	<b>8a</b> 4-aminobenzonitrile	<b>8b</b>	/

<sup>a</sup> Synthesis in a 150  $\mu\text{L}$  copper tube flow reactor, reaction was quenched by introducing the reaction mixture to a 3:1 solution of methanol:acetic acid.






Figure 2






Figure 3

