

Microwave-promoted Synthesis of Methoxylated Benzaldehydes from Natural *cis*-Phenylpropenes Using NaIO₄/OsO₄ (cat.)¹

Arun K. Sinha,* Bhupendra P. Joshi, and Ruchi Acharya

Natural Plant Products Division, Institute of Himalayan Bioresource Technology, Palampur (H.P.)-176061, India

(Received April 28, 2003; CL-030360)

Microwave-promoted rapid oxidation of toxic natural *cis*-phenylpropenes (**1a–1e**) into high valued methoxylated benzaldehydes (**2a–2e**) was achieved within 2 min in high yield (71–84%) by reacting with sodium metaperiodate in the presence of catalytic amount of osmium tetroxide (co-oxidant) and benzyltriethylammonium chloride (PTC).


A large number of naturally occurring methoxylated benzaldehydes² contribute significantly to the taste and flavour of many foods, drinks, perfumery³ and also used for the synthesis of many bioactive compounds.⁴ There are synthetic methods⁵ reported for the preparation of these important benzaldehydes including oxidation of phenylpropenes.⁶ However, most of the reported methods are associated with drawbacks such as low yields, long reaction time, maintenance of temperature, expensive reagents and overall environmentally hazardous reaction conditions. In this context, new emerging microwave technique⁷ coupled with potassium permanganate (KMnO₄)⁸ is proven to be high yielding oxidant as well as free from most of the above drawbacks, however, KMnO₄ in our hands converted β -asarone⁹ (**1a**), a natural toxic *cis*-phenylpropene,¹⁰ into asaronaldehyde (**2a**) in only 37% yield. In addition, KMnO₄ generated excessive fumes inside the microwave oven which once resulted in a spark.¹¹ Similarly, manganese dioxide/sulphanilic acid^{6,9} coupled in microwave provided **2a** in 21% yield only and most of the starting **1a** remained unreacted. All this finally prompted us to search for an alternative oxidant. Herein, we report a facile oxidative conversion of an abundantly available toxic methoxylated *cis*-phenylpropenes or a mixture of *cis*- and *trans*-phenylpropenes (**1**) into methoxylated benzaldehydes (**2**) in aqueous¹² media using sodium metaperiodate (NaIO₄)¹³ as an oxidant in the presence of osmium tetroxide (OsO₄) and benzyltriethylammonium chloride under microwave irradiation (Scheme 1).

Treatment of *cis*-isomer¹⁴ (**1a**) with a mixture of NaIO₄, OsO₄ (co-oxidant) and benzyltriethylammonium chloride in a 4:1 mixture of water and THF under microwave irradiation⁷

gave 2,4,5-trimethoxybenzaldehyde (**2a**) in 82% yield in 50 s. Similarly, α -asarone (*trans*-isomer, procured from Sigma) upon oxidation produced **2a** in 84% yield which clearly confirmed that *cis* or *trans* geometry of phenylpropene does not effect the yield of the oxidized product.¹⁴ As an extension to this method, crude *Acorus calamus* oil¹⁵ rich in *cis*- and *trans*-asarone is also directly oxidized under microwave to provide **2a** in 76% yield in 1 min while conventional method provides 57% yield of **2a** in 6 h. It is worthwhile to mention that the direct utilization of crude calamus oil for its high value added product **2a** is a huge economical gain for *A. calamus* oil obtained from tetraploid and hexaploid strains.^{9,16} Moreover, high price of co-oxidant OsO₄ did not become restrain in our process as 0.001 mole of OsO₄ per mole of phenylpropene was effective in oxidizing phenylpropenes in optimum yield. Finally, the above combina-

Table 1. Conversion of Phenylpropenes (**1**) into Benzaldehydes (**2**)

Entry	Phenylpropene (1)	Benzaldehyde (2)	Microwave irradiation Time (s)	Yield (%)
a)			50	82
b)			60	79
c)			90	78
d)			90	79
e)			120	71



(a) and (b) R₂=R₄=R₅=OMe; R₃=R₆=H (c) R₃=R₄=R₅=OMe; R₂=R₆=H

(d) R₃=R₄=OCH₂O; R₂=R₅=R₆=H (e) R₃=R₄=OMe; R₂=R₅=R₆=H (f) R₄=OMe; R₂=R₃=R₅=R₆=H

Scheme 1.

tion of oxidants¹³ was employed successfully on a series of other toxic¹⁰ methoxylated *cis*-phenylpropenes (**1b–1e**) towards formation of respective methoxylated benzaldehydes (**2b–2e**) in high yield¹⁷ as summarized in Table 1.

In conclusion, we report a mild, ecofriendly, rapid and efficient process for the preparation of high valued methoxylated benzaldehydes from toxic *cis*-phenylpropenes using NaIO₄/OsO₄ under microwave irradiation in aqueous¹² condition. Moreover, this process is capable of converting crude calamus oil containing 85% of toxic β -asarone along with 3% α -asarone into useful asaronaldehyde³ in 76% yield which makes this process attractive for industrial use.

One of the authors (BPJ) is indebted to CSIR, Delhi for the award of SRF. The authors gratefully acknowledge the Director of I. H. B. T., Palampur for his kind cooperation and encouragement.

References and Notes

- 1 IHBT Communication No. 0314.
- 2 J. B. Harborne and H. Baxter, in "Phytochemical Dictionary, A Handbook of Bioactive Compounds from Plants," Taylor & Francis Ltd., London (1993).
- 3 A. Steffen, in "Perfume and Flavor Chemicals (Aroma Chemicals)," Allured Publishing Corp., IL (1994), Vol. I & II; K. Bauer, D. Garbe, and H. Surburg, in "Common Fragrance & Flavor Materials: Preparation, Properties & Uses," 4th ed., Wiley-VCH Verlag, Weinheim (2001).
- 4 K. Tomioka, Y. Kuboto, and K. Koga, *Tetrahedron Lett.*, **22**, 2953 (1989); J. Demyttenaere, K. V. Syngel, A. P. Markusse, S. Vervisch, S. Debenedetti, and N. D. Kimpe, *Tetrahedron*, **58**, 2163 (2002).
- 5 I. M. Godfrey, M. V. Sargent, and J. A. Elix, *J. Chem. Soc., Perkin Trans. 1*, **1974**, 1353; S. D. Anthony and K. J. Charles, U. S. Patent 5910613 (1999); H. P. Markus, R. Roos, and J. L. A. Peters, Eur. Patent 0542348 (1993); N. Ito and Hasebe, Jpn. Patent 11279104A2 (1999); Interchem Corp., British Patent 1092615 (1967); T. Yokoyama and N. Yamagata, *Appl. Catal., A*, **221**, 227 (2001).
- 6 P. Z. Bezoukian, in "Perfumery and Flavoring Synthetics," Allured Publishing Corporation, IL, USA (1986); T. F. Nicholas, U. S. Patent 2794813 (1957); P. S. S. Essfncefabri, British Patent 774608; C. Chen, A. Potthast, T. Rosenau, T. Gratzl, J. S. Gratzl, A. G. Kirkman, D. Nagai, and T. Miyakoshi, *J. Mol. Catal. B: Enzym.*, **8**, 213 (2000).
- 7 L. Pelle, T. Jason, W. Bernard, and W. Jacob, *Tetrahedron*, **57**, 9225 (2001); M. Larhed and A. Hallberg, *Drug Discovery Today*, **6**, 406 (2001); A. Shaabani, A. Banzir, F. Teimouri, and D. G. Lee, *Tetrahedron Lett.*, **43**, 5165 (2002).
- 8 F. Rudolf and S. Tibor, *Chem. Ber.*, **39**, 1211 (1906); L. N. Thach, T. H. Anh, L. T. X. Thi, and L. Andre, *Chem. Abstr.*, **126**, 211877z (1997); L. N. Thach, T. H. Anh, and D. T. A. Thu, *Chem. Abstr.*, **130**, 110007t (1999).
- 9 Phenylpropenes^{2,3} exist in three isomeric forms namely *trans* (α), *cis* (β) and allyl (γ) isomers where *cis*-phenylpropenes (e.g. β -asarone) are recently found carcinogenic and toxic⁹ which prompted us to utilize *cis*-phenylpropenes or a mixture of *cis*- and *trans*-isomers as a cheaper and economical starting material. Among three isomers, *cis*-phenylpropene is generally present in high concentration e.g., β -asarone (*cis*-2,4,5-trimethoxyphenylpropene) is present upto 90% in tetraploid and hexaploid strains (distributed extensively in India, Pakistan, Japan and China) restricts the market potential of *A. calamus* oil.
- 10 E. C. Miller, A. B. Swanson, D. H. Phillips, T. L. Fletcher, A. Liem, and J. A. Miller, *Cancer Res.*, **43**, 1124 (1983); G. Abel, *Planta Med.*, **53**, 251 (1987); S. C. Kim, A. Liem, B. C. Stewart, and J. A. Miller, *Carcinogenesis*, **20**, 1303 (1999).
- 11 W. S. Trahanovsky, in "Oxidation in Organic Chemistry," Academic Press, New York (1982), p 186; M. W. Beck and M. E. Brown, *Thermochim. Acta*, **65**, 197 (1983).
- 12 C. J. Li, *Chem. Rev.*, **93**, 2023 (1993); C. J. Li, *Tetrahedron*, **52**, 5643 (1996).
- 13 G. Cainelli, M. Contento, F. Manescalchi, and L. Plessi, *Synthesis*, **1989**, 47; H. Tanaka, R. Kikuchi, M. Baba, and S. Torii, *Bull. Chem. Soc. Jpn.*, **68**, 2989 (1995); D. L. J. Clive, Y. Tao, Y. Bo, Y. Hu, N. Selvakumar, S. Sun, S. Daigneault, and Y. Wu, *Chem. Commun.*, **2000**, 1341.
- 14 General procedure for microwave-promoted oxidation of methoxylated *cis*-phenylpropenes (**1a–1e**) into phenylaldehyde (**2a–2e**): A mixture of *cis*-phenylpropene (0.015 mol), OsO₄ (0.004 g, 0.015 mmol), NaIO₄ (11.75 g, 0.055 mol) and benzyltriethylammonium chloride (0.08–0.1 g) were dissolved in H₂O–THF (10 mL, 4:1) and irradiated for 40 s to 2 min (Table 1) under microwave oven (2450 MHz, 900 Watts, Kenstar, India). The mixture was filtered and washed with dichloromethane. Evaporation of the solvent under reduced pressure gave a crude product, which was purified on silica gel column with hexane–ethyl acetate, 9:1 to 4:1 to afford **2a–2e**.
- 15 The hydrodistillation of rhizomes of *Acorus calamus* gave yellowish oil in 1.7% yield (w/w) with a presence of 3% of α -asarone and 85% β -asarone (**1a**) (by GC) and used directly for oxidation. A mixture of crude calamus oil (9.00 g), OsO₄ (0.08 to 0.002 g), NaIO₄ (50 to 55 g) and benzyltriethylammonium chloride (0.15 g) were dissolved in H₂O–THF (50–60 mL, 4:1) and irradiated for 1 min in parts which provided **2a** in 76% yield (based upon percentage of α - and β -asarone present in crude calamus oil).
- 16 A. K. Sinha, B. P. Joshi, and R. Dogra, *Nat. Prod. Lett.*, **15**, 439 (2001); A. K. Sinha, R. Acharya, and B. P. Joshi, *J. Nat. Prod.*, **65**, 764 (2002); A. K. Sinha, U. S. Patent 6528041 (2002); A. K. Sinha, B. P. Joshi, and R. Dogra, U. S. Patent 6544390 (2003).
- 17 The yield of liquid anisaldehyde (**2e**) is found lesser than other solid phenylaldehydes (**2a–2d**) because of volatility of **2e** during microwave irradiation reaction.¹⁴