

Complexes of Some Alkali-metal and Silver Salts with 1,4-Dioxan

By John C. Barnes* and (Miss) C. S. Duncan, Chemistry Department, The University, Dundee DD1 4HN

The 1,4-dioxan (dx) complexes $\text{NaClO}_4 \cdot 3\text{dx}$, $\text{NaBF}_4 \cdot 3\text{dx}$, and $\text{NaI} \cdot 3\text{dx}$ have the cubic structure of $\text{AgClO}_4 \cdot 3\text{dx}$. The metal ion is octahedrally co-ordinated by oxygen atoms from dioxan with Na-O distance 2.43–2.46 Å, identical to those found in $\text{Na}(\text{OH}_2)_6$ octahedral. Size and shape requirements limit the anions found in complexes of this structure. The heavier alkali-metal ions are too large for the structure to be stable. The lattice energy of the parent salt is not important in selecting which alkali-metal ions form complexes with dioxan. The lattice energy of the complex is dominated by metal-dioxan ion-dipole interaction. Temperatures, enthalpies, and entropies of decomposition are reported. $\text{AgClO}_4 \cdot 3\text{dx}$ decomposes to $\text{AgClO}_4 \cdot \text{dx}$ and thence to AgClO_4 , $\text{NaClO}_4 \cdot \text{dx}$, and $\text{NaBF}_4 \cdot 3\text{dx}$ decompose directly to the parent salt. $\text{NaI} \cdot 3\text{dx}$ decomposes to NaI at atmospheric pressure but $\text{NaI} \cdot 2\text{dx}$ and $\text{NaI} \cdot \text{dx}$ have been characterised *in vacuo*. Temperatures and enthalpies of phase changes in AgClO_4 , NaClO_4 , NaBF_4 , and $\text{NaClO}_4 \cdot 3\text{dx}$ are reported.

THERE has been a revival of interest in the complexes formed by alkali metals.¹ Work on the polyether complexes has shown that ether groups are good ligands for these metals.² A number of complexes of alkali metal salts with 1,4-dioxan (dx) have been reported (Table 1).

TABLE 1

Dioxan complexes of alkali metal salts^a

	Cl^- ^b	Br^- ^b	I^- ^b	ClO_4^- ^c	BF_4^- ^d	SCN^-
Li^+	1	1	2	—	—	1
Na^+	0	0	2,3	3	3	3
K^+	0	0	1	0	0	2
Rb^+	0 ^d	0 ^d	0 ^d	0	—	0
Cs^+	0 ^d	0 ^d	0 ^d	0	—	0

^a The Table lists values of n in $\text{MX}_n \cdot \text{ndx}$. A zero entry indicates a system for which no complex was obtained, a dash indicates a system which has not been investigated. ^b H. Rheinboldt, A. Luyken, and H. Schmittman, *J. prakt. Chem.*, 1937, **148**, 81; **149**, 30. ^c T. Gruhn and M. Gorman, *J. Inorg. Nuclear Chem.*, 1965, **27**, 482. ^d Present work. ^e M. Gorman, D. DeMattio, D. Doonan, and G. McDonald, *J. Chem. Educ.*, 1970, **47**, 466.

These complexes are of very low thermal stability, exerting significant vapour pressures of dioxan at 300 K. Although some can be prepared from aqueous solution, the crystalline complexes react with atmospheric moisture, usually with loss of dioxan. Only the system $\text{LiCl} \cdot \text{dx} \cdot \text{H}_2\text{O}$ has been investigated in detail; $\text{LiCl} \cdot \text{dx} \cdot \text{H}_2\text{O}$ occurred over much of the phase diagram but this compound was stable only in contact with its mother-liquor.³

Table 1 shows that a number of complexes have the stoichiometry $\text{NaX} \cdot 3\text{dx}$. There have been no previous studies of these compounds beyond reports of their preparation. Analogous sodium and silver(I) salts are often isostructural and the structure of $\text{AgClO}_4 \cdot 3\text{dx}$ is known,⁴ and shown in Figure 1. The unit cell is cubic, with $a = 7.67$ Å. Vapour pressure measurements⁵ showed that $\text{AgClO}_4 \cdot 3\text{dx}$ decomposed to $\text{AgClO}_4 \cdot \text{dx}$ and thence to AgClO_4 . The values of ΔH_{298} and ΔS_{298} obtained from these measurements have been criticised.⁶

RESULTS

$\text{NaClO}_4 \cdot 3\text{dx}$, $\text{NaBF}_4 \cdot 3\text{dx}$, and $\text{NaI} \cdot 3\text{dx}$ are all cubic with unit cell parameters very close to that of $\text{AgClO}_4 \cdot 3\text{dx}$

¹ M. R. Truter, *Chemistry in Britain*, 1971, 203.

² D. Bright and M. R. Truter, *Nature*, 1970, **225**, 176.

³ C. C. Lynch, *J. Phys. Chem.*, 1942, **46**, 366.

(Table 2). It seems safe to assume that all these compounds are isostructural.




FIGURE 1 Unit-cell of $\text{AgClO}_4 \cdot 3(1,4\text{-dioxan})$. The anion rotates freely and the dioxan molecules rotate about the O-O axes (Prosen and Trueblood, *Acta Cryst.*, 1956, **9**, 741)

The decomposition of these 1:3 complexes has been studied by differential enthalpic analysis (d.e.a.), by thermogravimetry and by vapour pressure measurements.

TABLE 2

Unit cell parameters of cubic $\text{MX}_n \cdot 3\text{dx}$ compounds

	a , Å
$\text{AgClO}_4 \cdot 3\text{dx}$	7.66 ± 0.02 (7.67 ± 0.01) [*]
$\text{NaI} \cdot 3\text{dx}$	7.59 ± 0.03
$\text{NaBF}_4 \cdot 3\text{dx}$	7.66 ± 0.01
$\text{NaClO}_4 \cdot 3\text{dx}$	7.66 ± 0.02

^{*} Single-crystal data from R. J. Prosen and K. N. Trueblood, *Acta Cryst.*, 1956, **9**, 741.

The latter offer a more reliable route than d.e.a. to the enthalpies of decomposition of compounds which exert a finite vapour pressure at ambient temperature since a partially decomposed sample will exert the same vapour pressure as a stoichiometric sample but will give a peak area in d.e.a. which is too small by an uncertain amount. This point is illustrated in Table 3 in which the decomposition data is summarised.

⁴ R. J. Prosen and K. N. Trueblood, *Acta Cryst.*, 1956, **9**, 741.

⁵ L. W. Baasch, *Spectrochim. Acta*, 1959, **17**, 726.

⁶ J. C. Barnes, *Inorg. Chem.*, 1972, in the press.

AgClO₄.3dx.—Two endothermic decomposition reactions corresponded to the formation of AgClO₄.dx and AgClO₄. AgClO₄ shows phase changes at 371, 398.6, and 423 K (Table 4). The last represents the change to the cubic modification. Only the last phase change is seen in AgClO₄ freshly decomposed from AgClO₄.3dx.

ment on NaI₂dx showed a phase change and two endothermic decomposition steps to NaI. V.p. measurements on samples of NaI₂dx prepared by driving reaction (2) to completion in the v.p. apparatus, were unsatisfactory. Non-linear Arrhenius plots were obtained. No convincing explanation for this behaviour is available at present.

TABLE 3
Thermal decomposition reactions ^a

Reagent	T ₁ /K	T ₂ /K	ΔW _{expt} %	ΔW _{Th} %	ΔH/kJ mol ⁻¹ (d.e.a.)	ΔH ₂₉₈ /kJ mol ⁻¹ (v.p.)	ΔS ₂₉₈ /JK ⁻¹ mol ⁻¹ (v.p.)	-ΔG ₂₉₈ /kJ mol ⁻¹ (v.p.)
AgClO ₄ .3dx	358	397	55.6 ± 0.5	56.04	39.2 ± 0.9	58.96 ± 0.61	246 ± 10	14.24 ± 0.21
AgClO ₄ .dx	400	420	28.1 ± 1.0	29.82	59.2 ± 2.1	92.09 ± 1.10	399 ± 12	26.87 ± 0.40
NaClO ₄ .3dx	343	398	67.3 ± 0.7	68.35	54.9 ± 1.6	62.42 ± 0.57	260 ± 12	14.90 ± 0.20
NaI ₃ dx	321	383	62.4 ± 0.5	63.81	54.5 ± 1.3 ^b	41.93 ± 0.42 ^c	174 ± 10 ^c	9.90 ± 0.15 ^c
NaBF ₄ .3dx	283	356	71.9 ± 0.9	70.65	35.6 ± 0.7	44.00 ± 0.51	176 ± 12	8.59 ± 0.15
NaI ₂ dx	<301 ^e	365	54.9 ± 0.3	54.04	67.7 ± 1.0 ^b	61.84 ± 0.61 ^d	313 ± 10	12.33 ± 0.15
		381						

^a D.e.a. experiments were carried out at atmospheric pressure under nitrogen flowing at 30 cm³ min⁻¹, heating at 8.33 K min⁻¹. T₁ was the minimum temperature at which the endotherm was detected. T₂ was the peak temperature of the endotherm. All thermodynamic quantities are expressed 'per mole of dioxan released'. ΔW represents weights loss to MX. ^b Decomposition to NaI. ^c Decomposition to NaI₂dx. ^d Decomposition to NaI₂dx. ^e Obscured by previous endotherm.

NaBF₄.3dx and NaClO₄.3dx.—D.e.a., t.g., and v.p. measurements with analysis, agree that these compounds decompose directly into NaBF₄ and NaClO₄ in one step.

Two endothermic phase changes occur during the decomposition of NaClO₄.3dx (Table 4). Both were readily


TABLE 4
Phase transitions observed by d.e.a.

Compound	T ₁ /K	T ₂ /K	ΔH/kJ mol ⁻¹
NaI ₂ dx	321	325.3	5.91 ± 0.11
NaClO ₄ .3dx	316.3	317.6	70.0 ± 0.04
	368.6	369.2	0.92 ± 0.21
NaClO ₄	576.3	579.1	1.46 ± 0.05
NaBF ₄	515.6	520.7	5.56 ± 0.25
AgClO ₄	371.4	373.1	1.22 ± 0.02
	398.6	402.1	0.51 ± 0.01
	423.5	429.5	3.70 ± 0.02


Experimental conditions as in Table 3.

reversible. The areas of these endotherms were proportional to the fraction (1 - α) of the sample remaining undecomposed when the phase change occurred, indicating that both phase changes are associated with NaClO₄.3dx and not with the decomposition products. (Some control of the relationship between α and T can be obtained by varying the heating rate in d.e.a.) Phase changes in NaBF₄.dx occur between 273 and 293 K and appeared to be unreproducible, possibly for instrumental reasons. NaBF₄ and NaClO₄ change from the anhydrite structure to cubic at 516 and 576 K respectively. These phase changes also occur in samples decomposed from the dioxan complexes (Table 4).

NaI₃dx.—This compound behaved differently under different experimental conditions. D.e.a. measurements showed only a single decomposition endotherm, with no phase changes. The product was found to be NaI. V.p. measurements showed an initial reaction



This reaction is followed by



for which ΔH is greater than for reaction (1), leading to a transition temperature at 340 K (Figure 2). D.e.a. measure-

DISCUSSION

In the AgClO₄.3dx structure the size of the unit-cell is determined by the silver-dioxan contacts. The Ag-O distance is 2.46 Å, which is typical of non-covalent Ag-O bonds. From the unit-cell parameters and the O-O distance in dioxan the Na-O distances in NaX₃dx range between 2.43 and 2.46 Å. These are in good agreement




FIGURE 2 Vapour pressure data for the decomposition of NaI_ndx; Triangles NaI₃dx; Circles NaI₂dx

with the values of 2.37–2.47 and 2.40–2.45 Å found in Na₂SO₄.10H₂O and in borax respectively.⁷ In both these compounds the sodium ions are at the centres of irregular octahedra of water molecules and thus in a similar environment to the octahedron of oxygen atoms in NaClO₄.3dx. The Ag-Cl distance in AgClO₄.3dx is 6.63 Å compared with 3.46 Å in cubic AgClO₄.⁷

The anion, situated at the body centre of a cubic array of metal ions, must be at least pseudo-spherical to allow the rotational freedom observed in AgClO₄.3dx.⁴

⁷ R. W. G. Wyckoff, 'Crystal Structures,' Interscience, New York, 1964, vols. 2 and 3.

NaNO_3 and NaBrO_3 do not form compounds of this structure. The effective radius of the anion must be large enough to prevent rattling but small enough to allow rotation unimpeded by the rotating methylene groups of the dioxan. The limits, in terms of ionic radii, are shown by NaBr and NaIO_4 , neither of which have been found to form compounds of this series. The values are


Br^-	I^-	BF_4^-	ClO_4^-	IO_4^-
1.96	2.16	2.28	2.33	2.48 Å

using Kaputinskii's thermochemical radii⁸ for the polyatomic ions.

None of the corresponding potassium salts form complexes of this type. Substitution of K^+ for Na^+ would give a unit-cell parameter of 8.40 Å, increasing the radius of the anion site by 0.5 Å. This would allow even IO_4^- to rattle. Ionic radii show that LiCl could form $\text{LiCl}\cdot 3\text{dx}$ but only a phase reported as $\text{LiCl}\cdot \text{dx}$ is known and that is of very low thermal stability.³ The formation of $\text{AgBF}_4\cdot \text{dx}$ and $\text{AgClO}_4\cdot \text{dx}$, for which sodium analogues have not been prepared may represent Ag^+ adopting the familiar linear two-co-ordinate geometry.

The contrast between the alkali metal complexes formed by 1,4-dioxan and those formed by thiourea can be explored by lattice energy arguments. Boeyens and co-workers^{9,10} have shown that thiourea complexes are formed only for salts with lattice energies below 670 kJ mol^{-1} . These complexes have lattice energies *ca.* 750 kJ mol^{-1} . Metal-thiourea interactions make a relatively small contribution to the overall lattice energy of the complex.

The enthalpy cycle shows that dioxan complexes are favoured only if the lattice energy of the complex exceeds the sum of the lattice energy of the parent salt and the heat of vaporisation of dioxan, that is if the heat of decomposition of the complex exceeds ΔH_{vap} (37.5 kJ mol^{-1}).¹¹ This condition is only satisfied for the $\text{MX}\cdot 3\text{dx}$ structure in the few cases where the packing is very efficient. Comparison of the lattice energies of $\text{MX}\cdot 3\text{dx}$



obtained by use of the enthalpy cycle with the calculated cohesive Madelung energies for M^+ and X^- arranged in the position which these ions occupy in $\text{MX}\cdot 3\text{dx}$ shows that this Madelung energy (*ca.* 370 kJ mol^{-1}) is only about 40% of the total lattice energy (Table 5). Allowing for Born repulsion the contribution from ion-dipole

⁸ A. P. Kaputinskii, *Quart. Rev.*, 1956, **10**, 283.

⁹ J. C. A. Boeyens and G. Gafner, *J. Chem. Phys.*, 1968, **49**, 2435.

¹⁰ J. C. A. Boeyens, *Acta Cryst.*, 1968, **B24**, 1191.

¹¹ J. L. Crenshaw, A. C. Cope, N. Finkelstein, and R. Rogan, *J. Amer. Chem. Soc.*, 1938, **60**, 2308.

TABLE 5
Lattice energies and Madelung energies^a

	Lattice energy (MX), kJ mol^{-1}	Lattice energy (MX, 3dx), kJ mol^{-1}	Madelung energy (MX, 3dx), kJ mol^{-1}
$\text{AgClO}_4\cdot 3\text{dx}$	798 ^b	1008	369
$\text{NaClO}_4\cdot 3\text{dx}$	732 ^b	918	369
$\text{NaI}\cdot 3\text{dx}$	699 ^c	865	372
$\text{NaBF}_4\cdot 3\text{dx}$			369

^a Madelung energy is used here for the attraction between anions and cations, spaced as required by the $\text{AgClO}_4\cdot 3\text{dx}$ structure, in the absence of the dioxan molecules. ^b D. F. C. Morris, *J. Inorg. Nuclear Chem.*, 1958, **6**, 295. ^c D. F. C. Morris, *Acta Cryst.*, 1956, **9**, 197.

interactions must be about 670 kJ mol^{-1} . The relative positions of dioxan, M^+ and X^- in the structure show that the dioxan will be strongly polarised by M^+ so as to increase the two opposed dipoles in the molecule (a similar effect is found in thiourea¹⁰).

The decrease in the cohesive energy for the ions and the ion-dipole interaction as r_m^+ increases must be sufficient to compensate for the decreasing lattice energy of MX and to prevent the formation of dioxan complexes of the heavier alkali metals.

Comparison with data for other dioxan complexes indicates that the low thermal stability of the $\text{MX}\cdot 3\text{dx}$ series arises largely from unfavourable entropy terms. The enthalpies of decomposition reported for dioxan complexes lie between 50 and 85 kJ mol^{-1} per dioxan lost¹¹⁻¹³ so that the values reported in Table 3 are not unduly low.

EXPERIMENTAL

The adducts were precipitated by the addition of freshly distilled dioxan to aqueous solutions of the salts at room temperature. The dry adducts were kept in an atmosphere saturated with dioxan vapour. They were characterised by thermogravimetry and by determination of the anion content by ion-exchange. Attempts were made to obtain the corresponding potassium complexes and complexes of other sodium salts, without success.

D.e.a. measurements were made with the Perkin-Elmer DSC 1B scanning calorimeter used previously.^{12,13} I.r. spectra were recorded on Nujol mulls on a Perkin-Elmer 521 instrument, but showed no features of interest. X-Ray photographs were obtained using $\text{Cu-K}\alpha$ radiation either photographically using a 5.71 cm diameter camera calibrated with NaCl or using a recording diffractometer. Since no low temperature or controlled atmosphere X-ray facilities were available it was necessary to obtain successive powder patterns and extrapolate to zero time to eliminate reflections due to decomposition products.

Vapour pressure measurements were made using an apparatus based on that used by earlier workers,¹¹ maintained in a thermostatted oil bath.

The collaboration of Professor J. Iball in providing the X-ray photographs is acknowledged. The S.R.C. are thanked for financial assistance.

[2/171 Received, 27th January, 1972]

¹² J. C. Barnes and C. S. Duncan, *J. Chem. Soc. (A)*, 1969, 1746.

¹³ J. C. Barnes and C. S. Duncan, *J. Chem. Soc. (A)*, 1970, 1442.