

Raney nickel to give $C^{11}H_3NH_2$. Possible one carbon species with this behavior are methylenimine, cyclodiazomethane and diaminomethane. One carbon species may be the precursor of both methane and methylamine. The mechanism of the primary process is not dealt with here but will be considered in the full paper dealing with these experiments, unreported data and work in progress on this system.

Acknowledgments.—The authors wish to acknowledge the invaluable assistance of Dr. C. Baker, head of the BNL 60" cyclotron group. We also wish to acknowledge the help of Drs. G. Stöcklin and D. Christman in carrying out some of the carbon-11 determinations.

(14) Institute of Pharmaceutical Chemistry, University of Rome. Visiting Associate Chemist, Brookhaven National Laboratory, 1961-1962.

CHEMISTRY DEPARTMENT
BROOKHAVEN NATIONAL LABORATORY
UPTON, L. I., NEW YORK

FULVIO CACACE¹⁴
ALFRED P. WOLF

RECEIVED MAY 2, 1962

STERIODS. CCII.¹ A NEW ROUTE TO 19-NOR STEROIDS

Sir:

The increasing importance of 19-nor steroids,^{2a} particularly in the field of oral contraception,^{2b} has made it attractive to investigate new routes to this class of compounds which do not proceed *via* Birch reduction³ of ring A aromatic precursors, since the latter reaction in particular is inconvenient for large scale operations.

Recently we⁴ and others⁵⁻⁷ described chemical methods for the direct oxygenation of the C-19 methyl group, a primary prerequisite for the subsequent conversion to 19-nor steroids. However, none of these approaches offered a facile synthesis of 19-nor- Δ^4 -3-ketones.

The conversion of several Δ^5 -3 β -alcohols into their corresponding 19-nor- Δ^5 -3-ketones by an efficient process⁸ is now described, which is exemplified by the conversion of pregnenolone acetate (I) into 19-norprogesterone (VIII) in an overall yield of 37%.

Addition of hypobromous acid (N-bromoacetamide and aqueous perchloric acid) to pregnenolone

acetate (I) in dioxane solution gave 5 α -bromopregnane-3 β ,6 β -diol-20-one 3-acetate (II) (m.p. 165-167°, $[\alpha]_D +6.5^\circ$.⁹ Found for $C_{29}H_{36}BrO_4$: C, 60.36; H, 7.81; Br, 17.79; O, 14.17). Treatment of II with 1.2 moles of dry lead tetraacetate in anhydrous benzene^{4,7} afforded 6 β ,19-oxido-5 α -bromopregnane-3 β -ol-20-one acetate (III) (m.p. 152-154°, $[\alpha]_D +58^\circ$, ν_{max}^{KBr} 907¹⁰ cm⁻¹. Found for $C_{28}H_{33}BrO_4$: C, 61.18; H, 7.31; Br, 17.91; O, 13.97). Mild alkaline hydrolysis of III gave the corresponding 3 β -alcohol IV (m.p. 179-180°, $[\alpha]_D +61^\circ$. Found for $C_{21}H_{31}BrO_3$: C, 61.52; H, 7.70; Br, 20.19) which was oxidized with chromium trioxide in aqueous acetic acid and then directly converted with zinc dust in isopropyl alcohol¹¹ (reflux 24 hours) into 19-hydroxy- Δ^5 pregnene-3,20-dione (V). Treatment of V with oxalic acid in ethanol afforded 19-hydroxyprogesterone (VI) (m.p. 169-171°, $[\alpha]_D +182^\circ$, λ_{max} 243 m μ , log ϵ 4.22. Found for $C_{21}H_{30}O_3$: C, 76.07; H, 9.11; O, 14.69). Oxidation of VI with 8 N chromic acid in acetone solution furnished the corresponding 19-carboxylic acid VII (m.p. 147-149°, $[\alpha]_D +94^\circ$, λ_{max} 243 m μ , log ϵ 4.19. Found for $C_{21}H_{28}O_4$: C, 73.43; H, 8.22; O, 18.33) smoothly converted by acidic methanol¹² to 19-norprogesterone¹³ (VIII) (m.p. 142-144°, $[\alpha]_D +141^\circ$).

In a similar manner dehydroisoandrosterone acetate (IX) gave 19-norandrostene-3,17-dione¹⁴ (XV) (m.p. 166-168°, $[\alpha]_D +127^\circ$, λ_{max} 239 m μ , log ϵ 4.23) *via* 5 α -bromoandrostane-3 β ,6 β -diol-17-one 3-acetate (X),^{4b,15} 6 β ,19-oxido-5 α -bromoandrostane-3 β -ol-17-one acetate (XI) (m.p. 187-188°, $[\alpha]_D +39^\circ$. Found for $C_{21}H_{29}BrO_4$: C, 59.24; H, 7.04; Br, 18.99), its corresponding 3 β -ol XII (m.p. 209-211°, $[\alpha]_D +38^\circ$. Found for $C_{19}H_{27}BrO_3$: C, 59.74; H, 7.15; Br, 20.71), 19-hydroxy- Δ^4 -androstene-3,17-dione¹⁶ (XIII) (m.p. 168-170°, λ_{max} 243 m μ , log ϵ 4.21) and the 19-carboxylic acid XIV¹² (m.p. 148-150°, $[\alpha]_D +242^\circ$, λ_{max} 244 m μ , log ϵ 4.18. Found for $C_{19}H_{24}O_4$: C, 72.13; H, 7.69; O, 20.27).

19-Nor-17 α -acetoxyprogesterone (XXIII) also was prepared from 17 α -hydroxypregnenolone 3,17-diacetate¹⁷ (XVI), by an alternate reaction sequence.

Reaction of 5 α -bromopregnane-3 β ,6 β ,17 α -triol-20-one 3,17-diacetate (XVII) (m.p. 184-186°, $[\alpha]_D -49^\circ$. Found for $C_{28}H_{37}BrO_6$: C, 58.41; H, 7.31; Br, 16.04; O, 18.23) with lead tetraacetate gave 6 β ,19-oxido-5 α -bromopregnane-3 β ,17 α -diol-20-one 3,17-diacetate (XVIII) (m.p. 202-208°,

(9) All rotations in chloroform solution and ultraviolet spectra in 95% ethanol.

(10) All compounds with the 6 β ,19-oxide bridge were characterized by a sharp medium intensity band between 903-910 cm⁻¹.

(11) In contrast to the 5 α -bromo-6,19-oxides, the corresponding 5 α -chloro-6,19-oxides [e.g., 5 α -chloro-6 β ,19-oxidoandrostane-3 β -ol-20-one acetate (m.p. 186-189°, $[\alpha]_D +53^\circ$)] were recovered unchanged after treatment with zinc in isopropyl alcohol under reflux.

(12) H. Hagiwara, S. Noguchi and M. Nishikawa, *Chem. Pharm. Bull. Japan*, **8**, 84 (1960).

(13) L. Miramontes, G. Rosenkranz and C. Djerassi, *J. Am. Chem. Soc.*, **73**, 3540 (1951).

(14) A. L. Wilds and N. A. Nelson, *J. Am. Chem. Soc.*, **75**, 5366 (1953).

(15) V. Grenville, D. K. Patel, V. Petrow, I. A. Stuart-Webb and D. M. Williamson, *J. Chem. Soc.*, 4105 (1957).

(16) A. S. Meyer, *Experientia*, **11**, 99 (1955), reports m.p. 168-170°.

(17) R. B. Turner, *J. Am. Chem. Soc.*, **75**, 3489 (1953).

- (1) Steroids CCI, P. Crabbé and J. Romo, *Ciencia*, in press.
- (2) (a) D. A. McGinty and C. Djerassi, *Ann. N. Y. Acad. Sci.*, **71**, 500 (1958); (b) G. Pincus and A. P. Merrill, "Control of Ovulation," Ed. by C. A. Villee, Pergamon Press, New York.
- (3) A. J. Birch and H. Smith, *Quart. Rev. (London)*, **12**, 17 (1958).
- (4) (a) A. Bowers, L. C. Ibáñez, M. E. Cabezas and H. J. Ringold, *Chem. and Ind. (London)*, 1299 (1960); (b) A. Bowers, E. Denot, L. C. Ibáñez, M. E. Cabezas and H. J. Ringold, *J. Org. Chem.*, **27**, 1862 (1962).
- (5) D. H. R. Barton, J. M. Beaton, L. E. Geller and M. M. Pechet, *J. Am. Chem. Soc.*, **82**, 2640 (1960); M. Akhtar and D. H. R. Barton, *ibid.*, **83**, 2213 (1961); D. H. R. Barton and J. M. Beaton, *ibid.*, **84**, 199 (1962).
- (6) J. S. Mills and V. Petrow, *Chem. and Ind. (London)*, 946 (1961).
- (7) Ch. Meystre, K. Heusler, J. Kalvoda, P. Wieland, G. Anner and A. Wettstein, *Experientia*, **17**, 475 (1961).
- (8) Recently G. Gardi and C. Pedrali, *Gazz. Chim. Ital.*, **91**, 1420 (1961), also M. Akhtar and D. H. R. Barton, *J. Am. Chem. Soc.*, **84**, 1496 (1962), described the synthesis of 19-nor steroids from Δ^5 -3 β -hydroxyandrostenes by processes utilizing an ultraviolet light irradiation procedure as the key step. Also cf. R. Kwok, T. Jen and M. Wolff, Abstracts 141st Meeting of the American Chemical Society, March 1962, p. 43-N.

$[\alpha]_D -1^\circ$. Found for $C_{25}H_{35}BrO_6$: C, 58.77; H, 6.79; Br, 15.55; O, 18.67) which was treated with zinc dust in isopropyl alcohol to afford Δ^5 -pregnene-3 β ,17 α ,19-triol-20-one 3,17-diacetate (XIX) (m.p. 228–229°, $[\alpha]_D -48^\circ$. Found for $C_{25}H_{36}O_6$: C, 69.10; H, 8.66; O, 22.36). Hydrolysis of XIX gave the 3 β -alcohol XX (m.p. 245–247°, $[\alpha]_D -63^\circ$. Found for $C_{23}H_{34}O_5$: C, 70.59; H, 8.53; O, 20.62) which underwent Oppenauer oxidation (10 min.) to furnish 19-hydroxy-17 α -acetoxyprogesterone (XXI) (m.p. 252–254°, $[\alpha]_D +72^\circ$, λ_{max} 242 m μ , $\log \epsilon$ 4.17. Found for $C_{23}H_{32}O_5$: C, 71.38; H, 8.36; O, 20.34). Oxidation of XXI to the 19-acid XXII (m.p. 166–168°, $[\alpha]_D +116^\circ$, λ_{max} 244 m μ , $\log \epsilon$ 4.18. Found for $C_{23}H_{30}O_6$: C, 68.72; H, 7.60; O, 23.56) and treatment with acid gave 19-nor-17 α -acetoxyprogesterone¹⁸ (XXIII) (m.p. 227–229°, $[\alpha]_D +18^\circ$, λ_{max} 239 m μ , $\log \epsilon$ 4.23).

(18) A. Zaffaroni, H. J. Ringold, G. Rosenkranz, F. Sondheimer, G. H. Thomas and C. Djerassi, *J. Am. Chem. Soc.*, **80**, 6110 (1958).

RESEARCH LABORATORIES
SYNTEX, S. A.
APARTADO POSTAL 2679
MÉXICO, D. F.


A. BOWERS
R. VILOTTI
J. A. EDWARDS
E. DENOT
O. HALPERN

RECEIVED JUNE 13, 1962

SYNTHESIS OF PATCHOULI ALCOHOL¹

Sir:

After structural studies on the tricyclic sesquiterpene patchouli alcohol (1)² were complete we initiated work directed toward a synthesis of this natural product and the related patchoulione (19), a substance with highly desirable olfactory properties. A synthesis verifying both structure and absolute configuration of 1 is now presented.



Addition of allylmagnesium chloride to (–)-homocamphor³ (prepared from (+)-camphor) gave the carbinol (3) (80%), m.p. 36–36.5°, $[\alpha]_D -20^\circ$ (all in $CHCl_3$) which on treatment with diborane⁴ followed by oxidation with Jones' reagent^{5,6} was converted to the spiroactone (4) (54%), m.p. 89–90°, $[\alpha]_D -45^\circ$, $\nu_{max}^{CCl_4}$ 1770 cm^{-1} . Dehydration in the presence of 1% $ZnCl_2$ in hot acetic anhydride-acetic acid solution (3:1) furnished the *meso*-ketone (5), (10%), m.p. 101–103°, λ_{max}^{EtOH}

(1) Terpenes XVII.

(2) G. Büchi, R. E. Erickson and N. Wakabayashi, *J. Am. Chem. Soc.*, **83**, 927 (1961).

(3) H. Favre and J.-C. Richer, *Can. J. Chem.*, **37**, 417 (1959); H. Rupe and C. Frey, *Helv. Chim. Acta*, **27**, 627 (1944).


(4) H. C. Brown and G. Zweifel, *J. Am. Chem. Soc.*, **81**, 247 (1959).

(5) A. Bowers, T. G. Halsall, E. R. H. Jones and A. J. Lemin, *J. Chem. Soc.*, 2548 (1953).

(6) Cf. oxidation of secondary boranes to ketones, R. Pappo, *J. Am. Chem. Soc.*, **81**, 1010 (1959); H. C. Brown and C. P. Garg, *ibid.*, **83**, 2951 (1961).

247 m μ (ϵ 14,500), $\nu_{max}^{CCl_4}$ 1665, 1615 cm^{-1} , n.m.r. (all in CCl_4) at 4.25 (1H singlet); 8.82 τ (6H singlet) and the desired cyclopentenone (9) (40%) m.p. 100°, λ_{max}^{EtOH} 246 m μ (ϵ 13,000), $\nu_{max}^{CCl_4}$ 1695, 1635 cm^{-1} , three methyl resonance peaks in the n.m.r. spectrum at 8.95, 9.05 and 9.15 τ . The latter ketone (9) was nearly completely racemized, indicating reversible isomerization of 8 and the *meso*-acid (2) prior to cyclization. This was avoided as follows: Hydroboration of 3 followed by oxidation with hydrogen peroxide gave the diol (6), m.p. 138–139°, $[\alpha]_D -25^\circ$ identical with the product available by reduction of 4 with lithium aluminum hydride which was monoacetylated, dehydrated with phosphorus oxychloride in pyridine and the resulting acetate reduced with lithium aluminum hydride to the unsaturated alcohol (7). Oxidation with chromium trioxide furnished the acid (8), m.p. 81–82°, n.m.r. signals at –2.2 (1H), 4.9 (1H broad), 9.0 (3H), 9.08 τ (6H) (over-all yield 3 to 8, 65%).

Treatment of its acid chloride with aluminum chloride in carbon disulfide afforded 9, m.p. 110°, $[\alpha]_D -149^\circ$ (41%).



Wittig condensation⁷ with triphenylphosphine-methylene in peroxide free tetrahydrofuran furnished an unstable liquid diene (10), λ_{max}^{EtOH} 256 m μ which on hydrogenation over Raney nickel W2 in ethyl acetate gave an olefin, over-all yield 9 to 11, 38%, $[\alpha]_D -42^\circ$. Identity with β -patchoulene (11) was established by comparison of infrared and n.m.r. spectra, optical rotation and retention time in the gas chromatogram.

The tricyclic skeleton of patchouli alcohol (1) had to be constructed from 11 using some version of the Wagner–Meerwein rearrangement. Direct isomerization was ruled out because the equilibrium between 11 and 20 is quantitatively in favor of 11. Epoxidation of 11 with peracetic acid produced the liquid epoxide (12) (99%), $[\alpha]_D -8^\circ$ which was isomerized with boron fluoride in ether solution (20°, 5–10 min.) to the unsaturated alcohol (13) (57%), m.p. 30–31°, $[\alpha]_D -14^\circ$,

(7) U. Schöllkopf, *Angew. Chem.*, **71**, 280 (1959).