


This article was downloaded by: [Georgetown University]

On: 15 May 2013, At: 03:05

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/lsyc20>

The Oxidation of Primary Alcohols to Methyl Esters and Diols to Lactones Using Trichloroisocyanuric Acid

Prof. Gene A. Hiegel^a & Cynthia B. Gilley^a

^a Department of Chemistry and Biochemistry, California State University, Fullerton, California, USA

Published online: 17 Aug 2006.

To cite this article: Prof. Gene A. Hiegel & Cynthia B. Gilley (2003): The Oxidation of Primary Alcohols to Methyl Esters and Diols to Lactones Using Trichloroisocyanuric Acid, *Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry*, 33:12, 2003-2009

To link to this article: <http://dx.doi.org/10.1081/SCC-120021026>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.



SYNTHETIC COMMUNICATIONS®

Vol. 33, No. 12, pp. 2003–2009, 2003

The Oxidation of Primary Alcohols to Methyl Esters and Diols to Lactones Using Trichloroisocyanuric Acid

Gene A. Hiegel* and Cynthia B. Gilley

Department of Chemistry and Biochemistry, California State University, Fullerton, California, USA

ABSTRACT

Primary alcohols and diols are easily oxidized to methyl esters by a solution of trichloroisocyanuric acid with methyl alcohol in dichloromethane. In addition, α,ω -diols are also readily oxidized into lactones by refluxing with trichloroisocyanuric acid and pyridine in dichloromethane.

Key Words: Primary alcohols; Methyl esters; Lactones; Diols; Oxidation; Synthesis; Preparation; Trichloroisocyanuric acid.

*Correspondence: Prof. Gene A. Hiegel, Department of Chemistry and Biochemistry, California State University, 800 N. State College Blvd., Fullerton, California 92834, USA; Fax: 714-278-5316; E-mail: ghiegel@fullerton.edu.

2003

DOI: 10.1081/SCC-120021026
Copyright © 2003 by Marcel Dekker, Inc.

0039-7911 (Print); 1532-2432 (Online)
www.dekker.com




2004

Hiegel and Gilley

Although a variety of oxidizing agents have been shown to successfully convert diols to lactones,^[1-8] there are few methods for the synthesis of methyl esters from primary alcohols. *N*-Iodosuccinimide,^[7] calcium hypochlorite,^[9] and tert-butyl hypochlorite^[10] have been shown to oxidize primary alcohols to methyl esters. We wish to report simple procedures for the synthesis of methyl esters from primary alcohols and for lactones from diols using trichloroisocyanuric acid (**1**) [1,3,5-trichloro-1,3,5-triazine-2,4,6-(1H,3H,5H)-trione] as the oxidizing agent.^[11]

Primary alcohols and diols are easily converted into the corresponding methyl esters in a solution of **1** and methyl alcohol in dichloromethane (Eq. (1)). The yield and purity of the isolated methyl esters are shown in Table 1.



A reasonable pathway for the transformation is shown in the Sch. 1. Previous studies have shown that the aldehyde to methyl ester conversion^[12] is significantly faster than the conversion of primary alcohols to aldehydes.^[13]

The primary alcohol to methyl ester conversion is an exothermic reaction, but an increase in reaction temperature results in the formation of byproducts. In addition, there is a variable induction period between the time the reagents are mixed and when the temperature spikes. Characteristically cyanuric acid begins to precipitate as the temperature begins to rise. The oxidation of primary alcohols to methyl esters using **1** was carried out successfully with methyl alcohol in dichloromethane, with methyl alcohol in acetonitrile, and in methyl alcohol. The solvent of choice for moderate or large-scale reactions is dichloromethane because the temperature is controlled internally by the low boiling point. If the temperature is kept at or below 40°C, methanol or acetonitrile can be used as solvents. Since external heat control can be problematic, the recommendation is to use dichloromethane as the solvent and to place the reaction flask in a cold water bath to facilitate heat dissipation. Since chlorine is also present in the reaction mixture, the reaction was protected from light to prevent radical chain chlorination reactions.

A complex mixture of products was formed during the oxidation of phenethyl alcohol and 3-phenyl-1-propanol. Once phenethyl alcohol is oxidized to phenylacetaldehyde, it is expected to have a higher enol content than most aldehydes, and therefore, more likely to form the α -chloro




Oxidation Using Trichloroisocyanuric Acid

2005







Table 1. The oxidation of alcohols to methyl esters.

Alcohol	Methyl ester	Yield (%)	GC purity (%)
		81 ^a	98.9
		81 ^a	99.9
		85 ^a	99.7
		75 ^a	99.9
		79 ^a	99.6
		39 ^b	87.5
		67 ^b	97.6
		96 ^b	98.5
		86 ^b	99.8

^aDistilled through a concentric tube column.^bPurified by flash chromatography.

Scheme 1.

**Table 2.** The oxidation of alcohols to lactones.

Alcohol	Lactone	Yield ^a (%)	GC purity (%)
		76	98.4
		72	97.4
		97	99.9

^aPurified by flash chromatography.

derivative. Other byproducts may also result from chlorination on the aromatic ring.^[11]

The C₄ and C₅ α,ω -diols can be converted into their corresponding lactones by refluxing with **1** and pyridine in dichloromethane. The yield and purity of the isolated lactones are shown in Table 2. Pyridine is added to absorb the hydrogen chloride formed during the oxidation. Experiments performed without pyridine gave low yields of lactone possibly due to polymer formation. The lactone formation reaction is believed to proceed through a cyclic hemiacetal, which would then be oxidized to the lactone. In contrast, the oxidation of 1,6-hexanediol gave only a trace amount of ϵ -caprolactone. However, 1,4-butanediol, 1,5-pentanediol, 1,6-hexanediol and 1,12-dodecanediol did oxidize to the corresponding diesters when methyl alcohol was present. The dimethyl succinate from the oxidation of 1,4-butanediol contained an unidentified impurity of 11.3% which may be 2-methoxytetrahydrofuran based on an ¹H-NMR peak at 3.23 δ .

EXPERIMENTAL

Analysis by gas chromatography was performed using a Hewlett-Packard 5890 Series II instrument with a 6 ft \times 1/8 in 10% Carbowax 20 M column. FT-IR spectra were recorded using a Perkin Elmer 1650. ¹H-NMR spectra were recorded on an Anasazi-modified Varian EFT

**Oxidation Using Trichloroisocyanuric Acid**

2007

90 MHz nuclear magnetic resonance spectrometer in CDCl_3 or CCl_4 . Flash chromatography was carried out with Merck 60 Å silica gel (230–400 mesh).

All reagents were used as received unless otherwise stated. TCICA was obtained from OMNI with a purity of 99% (88% available chlorine). 1,4-Butanediol and 1,6-butanediol were obtained from J.T. Baker. Phenethyl alcohol was reagent grade from Eastman. All other chemicals were obtained from Aldrich. 1-Nonanol was obtained from Aldrich and distilled prior to use. All products were purified by distillation or flash chromatography and were characterized by comparison with authentic samples using IR and $^1\text{H-NMR}$ spectra and GC retention time.

Oxidation of Benzyl Alcohol to Methyl Benzoate

In a 250-mL three-neck flask were placed a magnetic stir bar, 60 mL of anhydrous dichloromethane, 39.0 mL (963 mmol) of anhydrous methyl alcohol, and 22.46 g (96.6 mmol) of **1**. After approximately 10 min of stirring, 10.28 g (10 mL, 95.1 mmol) of benzyl alcohol was added. The flask was flushed with nitrogen, covered with foil to keep out light, and placed in a cold-water bath ($\sim 2^\circ\text{C}$ at start of the reaction). After approximately 24 h, excess TCICA was destroyed by the slow addition of saturated aqueous sodium hydrogen sulfite until a negative test with wet potassium iodide-starch test paper was achieved. The cyanuric acid precipitate was removed by vacuum filtration and the solid was washed with pentane. Most of the solvent was removed from the filtrate with a rotary evaporator and the residue was diluted with 60 mL of pentane. The pentane solution was washed with 1 N NaOH (20 mL), saturated NaCl (20 mL), and dried over anhydrous magnesium sulfate. After filtration and concentration, the crude product was vacuum distilled [b.p. $96.0\text{--}97.8^\circ\text{C}$ (26.4–30.3 torr)] through a concentric tube column to give 10.16 g (79%) of methyl benzoate. Analysis by GC showed a purity of 99.6%, and the retention time was identical to that of an authentic sample. The IR and $^1\text{H-NMR}$ spectra were identical to that of an authentic sample.

Oxidation of 1,4-Butanediol to γ -Butyrolactone

In a 50-mL three-neck flask were placed a magnetic stir bar, 30.0 mL of anhydrous dichloromethane, 0.4544 g (5.042 mmol) of 1,4-butanediol,



2008

Hiegel and Gilley

0.90 mL (11.2 mmol) of pyridine, and 1.2899 g (5.550 mmol) of **1**. The flask was flushed with nitrogen, covered with foil to keep out light, and refluxed using a temperature-controlled oil bath. After approximately 5 h, excess TCICA was destroyed by the slow addition of saturated aqueous sodium hydrogen sulfite until a negative test with wet potassium iodide-starch test paper was achieved. The cyanuric acid precipitate was removed by vacuum filtration and the solid was washed with dichloromethane. The solution was washed with 1 N HCl (15 mL). Dichloromethane (2 × 15 mL) was used to back-extract the HCl wash. The solution was then dried over anhydrous sodium sulfate. After filtration and concentration, the oily residue was purified by flash chromatography to give 0.3305 g (76%) of γ -butyrolactone. Analysis by GC showed a purity of 98.4%, and the retention time was identical to that of an authentic sample. The IR and ¹H-NMR spectra were identical to that of an authentic sample.

ACKNOWLEDGMENT

This work was supported in part by an NSF-REU Program grant, a CSUF Undergraduate Research and Creativity Award, and an award from the Rohm and Haas Company: Otto Haas Award for Technical Excellence (courtesy of Robert K. Barr).

REFERENCES

1. Shaabani, A.; Lee, D.G. Solvent free permanganate oxidations. *Tetrahedron Lett.* **2001**, *42* (34), 5833–5836.
2. Tanaka, H.; Kawakami, Y.; Goto, K.; Kuroboshi, M. An aqueous silica gel disperse electrolysis system. N-Oxyl-mediation electrooxidation of alcohols. *Tetrahedron Lett.* **2001**, *42* (3), 445–488.
3. Mukhopadhyay, S.; Chandalia, S.B. Production of γ -butyrolactone by liquid phase oxidation of 1,4-butanediol. *Indian J. Chem. Technol.* **1999**, *6* (4), 237–239.
4. Yang, G.; Chen, Z.; Zhang, S.; Chen, J. Study on polymer-supported tribromide oxidizing agent. *Lizi. Jiaohuan Yu Xifu* **1998**, *14* (6), 475–480; *Chem. Abstr.* *130*, 313445.
5. Yang, G.; Chen, Z.; Shi, C. Study on polymer-supported bromate ion oxidizer with sodium bisulfate. *Hubei Daxue Xuebao, Ziran Kexueban.* **1998**, *20* (3), 256–259; *Chem. Abstr.* *130*, 24638.



Oxidation Using Trichloroisocyanuric Acid

2009

6. Morimoto, T.; Hirano, M.; Iwasaki, K.; Ishikawa, T. Oxidation of diols with sodium bromite trihydrate in organic solvents in the presence of alumina. *Chem. Lett.* **1994**, (1), 53–54.
7. McDonald, C.E.; Holcomb, H.L.; Leathers, T.W.; Kennedy, K.E. The oxidation of primary alcohols to methyl esters and lactones using N-iodosuccinimide. *Microchem. J.* **1993**, 47 (1–2), 115–119.
8. Kondo, S.; Kawasoe, S.; Kunisada, H.; Yuki, Y. Convenient synthesis of lactones by the reaction of diols with N-halomides. *Synth. Commun.* **1995**, 25, 719–724; two examples using **1** are included.
9. McDonald, C.E.; Nice, L.E.; Shaw, A.W.; Nestor, N.B. Calcium hypochlorite-mediated oxidation of primary alcohols to methyl esters. *Tetrahedron Lett.* **1993**, 34 (17), 2741–2744.
10. Milovanovic, J.N.; Vasojevic, M.; Gojkovic, S. Oxidation of primary alcohols to methyl esters using tert-butyl hypochlorite, pyridine and methyl alcohol. *J. Chem. Soc. Perkin Trans. 2* **1991**, 1231–1233.
11. For synthetic applications of **1** see: Hiegel, G.A. Trichloroisocyanuric Acid. In *Encyclopedia of Reagents for Organic Synthesis*; Paquette, L.A., Ed.; Wiley: Chichester, Sussex, England, 1995; Vol. 7, 5072–5073.
12. Hiegel, G.A.; Bayne, C.D.; Donde, Y.; Tarmashiro, G.S.; Hilberath, L.A. The oxidation of aldehydes to methyl esters using trichloroisocyanuric acid. *Synth. Commun.* **1996**, 26, 2633–2639.
13. Unpublished results by Katherine A. Alvarado and Klaas Schildknecht, The oxidation of primary alcohols with trichloroisocyanuric acid.

Received in the USA October 17, 2002



MARCEL DEKKER, INC. • 270 MADISON AVENUE • NEW YORK, NY 10016

©2003 Marcel Dekker, Inc. All rights reserved. This material may not be used or reproduced in any form without the express written permission of Marcel Dekker, Inc.