


This article was downloaded by: [Moskow State Univ Bibliote]

On: 18 February 2014, At: 07:05

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/Incn20>

Synthesis of 2',3'-Dideoxy-2'-fluoro-3'-thioarabinothymidine and Its 3'-Phosphoramidite Derivative

Mohamed I. Elzagheid^a, Anna Lisa Tedeschi^a & Masad J. Damha^{a,b}

^a Department of Chemistry, McGill University, Montreal, Quebec, Canada

^b McGill University, Otto Maass Chemistry Building 801 Sherbrooke Street W, H3A 2K6, Montreal, QC, Canada

Published online: 31 Aug 2006.

To cite this article: Mohamed I. Elzagheid, Anna Lisa Tedeschi & Masad J. Damha (2003) Synthesis of 2',3'-Dideoxy-2'-fluoro-3'-thioarabinothymidine and Its 3'-Phosphoramidite Derivative, *Nucleosides, Nucleotides and Nucleic Acids*, 22:5-8, 1343-1346, DOI: [10.1081/NCN-120022961](https://doi.org/10.1081/NCN-120022961)

To link to this article: <http://dx.doi.org/10.1081/NCN-120022961>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Synthesis of 2',3'-Dideoxy-2'-fluoro-3'-thioarabinothymidine and Its 3'-Phosphoramidite Derivative

Mohamed I. Elzagheid, Anna Lisa Tedeschi, and Masad J. Damha*

Department of Chemistry, McGill University, Montreal,
Quebec, Canada

ABSTRACT

An efficient method for the synthesis of 5'-O-monomethoxytrityl-2',3'-dideoxy-2'-fluoro-3'-thioarabinothymidine [$^{5'}\text{-MMT}_{\text{araF-T}_{3'\text{SH}}}$ (**5**)] and its 3'-phosphoramidite derivative (**6**) suitable for automated incorporation into oligonucleotides, is demonstrated. A key step in the synthesis involves reaction of 5'-O-MMT-2,3'-O-anhydrothymidine (**4**) (Eleuteri, A.; Reese, C.B.; Song, Q., *J. Chem. Soc. Perkin Trans. 1* **1996**, 2237 pp.) with sodium thioacetate to give $^{5'}\text{-MMT}_{\text{araF-T}_{3'\text{SAC}}}$ (**5**) (Elzagheid, M.I.; Mattila, K.; Oivanen, M.; Jones, B.C.N.M.; Cosstick, Lönnberg, H. *Eur. J. Org. Chem.* **2000**, 1987–1991). This nucleoside was then converted to its corresponding phosphoramidite derivative, **6**, as described previously ((a) Sun, S.; Yoshida, A.; Piccirilli, J.A. *RNA*, **1997**, 3, 1352–1363; (b) Matulic-Adamic, J.; Beigelman, L. *Helvetica Chimica Acta* **1999**, 82, 2141–2150; (c) Fettes, K.J.; O'Neil, I.; Roberts, S.M.; Cosstick, R. *Nucleosides, Nucleotides and Nucl. Acids* **2001**, 20, 1351–1354).

Key Words: Thionucleosides; Arabinonucleic acids; Antisense.

*Correspondence: Masad J. Damha, McGill University, Otto Maass Chemistry Building 801 Sherbrooke Street W, H3A 2K6 Montreal, QC, Canada; Fax: +514 398 3797; E-mail: masad.damha@mcgill.ca.



Synthetic oligonucleotides are of considerable importance because of their ability to inhibit gene expression through antisense effects.^[4] The most efficacious antisense oligonucleotides (AON) are those that possess stabilities against nucleases, and adequate lipophilicity to ensure efficient cell permeation. Ideally, they should also suppress gene activity with the assistance of intracellular ribonuclease H (RNase H), an enzyme that degrades the RNA strand of the AON/RNA hybrid (see Ref.^[5], and references therein). However, most AONs lack the appropriate structural attributes for RNase H elicitation and typically display less than optimal cellular potencies. Our laboratory has recently demonstrated that arabinonucleic acids





Figure 1. Structure of modified oligonucleotides.



Scheme 1. i) [Bis(2-methoxyethyl)amino]sulfur trifluoride, CH_2Cl_2 , 50 °C, 83%; ii) HBr, CH_2Cl_2 , rt, 24h, 90%; iii) silylated thymine, CCl_4 , 3d, 77 °C, 75%; iv) NH_4OH , MeOH, rt, 24h, 90%; v) MMTTr-Cl, pyridine, 87%; vi) MsCl, pyridine/ Et_3N , EtOH, reflux, 16h, 92%; vii) NaSAc, dioxane, reflux, 12h, 84%; viii) 1M NaOMe, MeOH, 1h; ix) 2-cyanoethyl-*N,N*-diisopropylchlorophosphoramidite, 1-methylimidazole, diisopropyl-ethylamine, CH_2Cl_2 , 80%.

(e.g., 2'-F-ANA), like DNA phosphorothioates (S-DNA), possess the ability to elicit RNase H-mediated hydrolysis of target RNA.^[6]

The antisense activity of phosphorothioate linked 2'-deoxy-2'-fluoro-arabinonucleic acids (S-2'-F-ANA, Fig. 1) has been evaluated,^[5] however, nothing is known about the isomeric analogue in which a sulfur atom replaces one of the two bridging oxygen atoms, i.e., 2',3'-dideoxy-2'-fluoro-3'-thioarabinonucleic acids (2'-F-3'-S-ANA; Fig. 1). Here we report preparation of one of the four nucleoside monomers required for the solid-phase synthesis of 2'-F-3'-S-ANA.

The synthesis is depicted in Sch. 1. Sugar precursor **1** (from Pfanstiehl) was converted into the α -1-bromoarabinose **2** with retention of sugar C-1 configuration.^[7] Compound **2** was then used, without isolation, for S_N2-type glycosylation reaction with silylated thymine base according to literature procedure^[7] which resulted, after removal of 5'- and 3'-O-benzoyl groups, in the formation of nucleoside **3**. The latter nucleoside then 5'-monomethoxytritylated and 3'-mesylated before being converted to 2,3'-anhydronucleoside **4**.^[2] Direct ring opening of the 2,3'-anhydro linkage, in nucleoside **4**, with thioacetate gave nucleoside **5**^a that was then converted to the phosphoroamidothioite derivative **6**^b following literature protocols.^[3] The synthesis of 2'-F-3'-S-ANA oligonucleotides is in progress.

ACKNOWLEDGMENT

We wish to thank CIHR (Canada) for financial support.

REFERENCES

1. Eleuteri, A.; Reese, C.B.; Song, Q. *J. Chem. Soc. Perkin Trans. 1* **1996**, 2237 pp.
2. Elzagheid, M.I.; Mattila, K.; Oivanen, M.; Jones, B.C.N.M.; Cosstick, R.; Lönnberg, H. *Eur. J. Org. Chem.* **2000**, 1987–1991.
3. (a) Sun, S.; Yoshida, A.; Piccirilli, J.A. *RNA*, **1997**, *3*, 1352–1363; (b) Matulic-Adamic, J.; Beigelman, L. *Helvetica Chimica Acta* **1999**, *82*, 2141–2150; (c) Fettes, K.J.; O'Neil, I.; Roberts, S.M.; Cosstick, R. *Nucleosides, Nucleotides and Nucleic Acids* **2001**, *20*, 1351–1354.
4. De Mesmaeker, A.; Häner, R.; Martin, P.; Moser, H.E. *Acc. Chem. Res.* **1995**, *28*, 366–374.

^aNucleoside **5**: TLC (20:1 [v/v] chloroform/methanol)0.49; ¹H-NMR (400 MHz acetone-d₆, tetramethylsilane as internal reference): 10.15 (1H, s, N-H), 7.60–6.90 (16H, m, trityl and H-6), 6.33 (1H, dd, J_{1',F} = 15.6 Hz J_{1',2'} = 4.4 Hz H-1'), 5.3 (1H, dt, J_{2',F} = 54.8 Hz, J_{2',3'} = 5.2 Hz 5.2 Hz H-2') 4.3 (1H, dm or ddd, J_{3',F} = 16.8 Hz J_{3',F} = 3.6 Hz H-3'), 4.1 (1H, m, H-4'), 3.79 (3H, s, CH₃O), 3.36 (2H, dd, H-5',5''), 2.37 (3H, s, CH₃CO-S), 1.64 (3H, CH₃-C5); ¹⁹F-NMR (283 MHz, acetone-d₆, 99% trifluoroacetic acid as external reference): -106.63 (ddd); ESI-MS: 589.1 (M-1), 590.1 (M⁺), 613.1 (M + Na⁺).

^bPhosphoroamidothioite. **6**: ³¹P-NMR (200 MHz, DMSO-d₆, 85% H₃PO₄ as external reference): 162.2, 158.7.



5. Lok, C.-N.; Viazovkina, E.; Min, K.-L.; Nagy, E.; Wilds, C.J.; Damha, M.J.; Parniak, M.A. *Biochemistry* **2002**, *41*, 3457–3467.
6. (a) Damha, M.J.; Wilds, C.J.; Noronha, A.; Brukner, I.; Borkow, G.; Arion, D.; Parniak, M.A., *J. Am. Chem. Soc.* **1998**, *120*, 12,976–12,977; (b) Noronha, A.M.; Wilds, C.J.; Lok, C.-N.; Viazovkina, K.; Arion, D.; Parniak, M.A.; Damha, M. *J. Biochemistry* **2000**, *39*, 7050–7062.
7. Elzagheid, M.I.; Viazovkina, E.; Masad, M.J. *Current Protocols in Nucleic Acid Chemistry* **2002**, 1.7.1–1.7.19.

