

## THE COUPLING OF ORGANIC GROUPS BY THE ELECTROCHEMICAL REDUCTION OF ORGANIC HALIDES: CATALYSIS BY 2,2'-BIPYRIDINENICKEL COMPLEXES

YOLANDE ROLLIN\*, MICHEL TROUPEL, DENNIS G. TUCK \* and JACQUES PERICHON

*Laboratoire d'Electrochimie, Catalyse et Synthèse Organique, U.M. No. 28, C.N.R.S., 2, rue Henri-Dunant, 94320 Thiais (France)*


(Received September 30th, 1985)

### Summary


The electrochemical reduction of a dilute solution of  $\text{NiX}_2\text{bipy}$  (bipy = 2,2'-bipyridine) in *N*-methylpyrrolidone gives the corresponding  $\text{Ni}^0$  complex, which undergoes oxidative addition with an excess of an organic halide  $\text{RX}$  to form  $\text{RNiX}$ . Decomposition of  $\text{RNiX}$  gives the dimer  $\text{R}_2$  in good yield and nickel(II). The nickel(0) species is regenerated to give an electrocatalytic process. The possible mechanism of these reactions is discussed briefly.

### Introduction

There has been increasing interest in recent years in the homocoupling of the organic moieties of organic halides by reactions which are catalysed by nickel(0) complexes [1]:





Semmelhack et al. [2] were the first to demonstrate the usefulness of nickel(0) in this respect by showing that the nickel(0)-cyclooctadiene complex catalyses the reductive dimerisation of aromatic or vinylic halides. In recent work nickel(0) complexes containing trialkyl- or triaryl-phosphine ligands have been widely used [1,3], but with very reactive substrates such as benzyl or vinyl halides, catalytic species can be obtained by the reduction of uncomplexed nickel(II) salts [4]. Similarly, catalytic quantities of nickel may suffice if a reducing agent such as zinc powder is added to regenerate the nickel(0) species [1,3–5], so that the overall reaction becomes:



\* Permanent address: Department of Chemistry, University of Windsor, Windsor, Ontario, N9B 3P4 (Canada).

The preparation and regeneration of the catalyst can also be conveniently carried out electrochemically [6]. In an earlier publication [6b,6c], it was reported that electroreductive coupling of aryl groups can be achieved by the following sequence:



(THF = tetrahydrofuran; HMPA = hexamethylphosphoramide)

Reaction 1 corresponds to the electrochemical preparation of a zerovalent nickel complex, which then yields a stable species by oxidative addition (reaction 2); further reduction in the presence of excess ArX leads to the biaryl with regeneration of the  $\text{Ni}^0$  catalyst (reaction 3).

The significant drawbacks in this system include the need to use a two-compartment cell with separate anode and cathode compartments, and the presence of a large excess of phosphine in order to avoid the deactivation of the catalytic system. The present paper reports a significant improvement to the method involving use of a single-compartment cell with a nickel complex of 2,2'-bipyridine (bipy) as the catalyst. A further advantage is that 2,2'-bipyridine is insensitive both to oxidation, and to side reactions as alkylation.

## Experimental

### General

The solvent, *N*-methylpyrrolidone (NMP) was distilled from alcoholic potassium hydroxide at reduced pressure. Other reagents were used as supplied. The supporting electrolyte was either  $\text{Bu}_4\text{NClO}_4$  or  $\text{Bu}_4\text{NBF}_4$  (0.1 *M*).

The complex  $\text{NiBr}_2 \cdot \text{bipy}$  precipitated out on mixing solutions of  $\text{NiBr}_2 \cdot 2\text{H}_2\text{O}$  and bipy in ethanol. This compound is reduced by a reversible two-electron process at  $-1.2$  V (relative to the saturated aqueous calomel electrode) in a solution in NMP containing at least one molar equivalent of bipy in excess, giving the soluble dark green complex  $[\text{Ni}(\text{bipy})_2]^0$  [7,8].

### Electrochemistry

The electrochemical cell consisted of a single compartment containing three electrodes (see Fig. 1). The anode was a cylindrical rod of magnesium, which was surrounded by a gold gauze cathode. Electrolysis was carried out at room temperature ( $\cong 20^\circ\text{C}$ ) under an atmosphere of argon, with the cathode potential set versus a reference electrode at the potential of the  $\text{Ni}^{\text{II}}/\text{Ni}^0$  transition.

The initial solution consisted of  $\text{NiBr}_2 \cdot \text{bipy}$  (0.02 *M*) and bipy (0.02–0.04 *M*) in NMP, plus supporting electrolyte. When the organic halide (RX, 0.1–0.4 *M*) was added, electrolysis at potentials at which  $\text{Ni}^{\text{II}}$  is reduced yielded stable solutions of  $\text{Ni}^0$  provided that the total charge exceeded approx. 1 mol of electron per mole of RX. Alternatively, the nickel could be first reduced to  $[\text{Ni}(\text{bipy})_2]^0$  before the addition of RX; in this case, approx. 50% of the element is recovered as  $\text{Ni}^{\text{II}}$  in the first stage of the experiment, implying a rapid reaction between RX and all the  $\text{Ni}^0$ .





Fig. 1. Diagram of the electrochemical cell. Key: 1 and 2. Input and output of argon; 3. Tube with sintered glass cap to isolate the aqueous phase reference electrode; 4. Electrical lead connecting to gold gauze cathode; 5. Magnesium electrode; 6. Gold cathode; 7. Reference electrode.

The remaining 50% of the  $\text{Ni}^{\text{II}}$  is recovered more slowly, presumably from the decomposition of an organonickel complex to give  $\text{R}_2$ . The catalytic cycle then involves continuous electrolysis at the potential of the  $\text{Ni}^{\text{II}}/\text{Ni}^0$  couple, with the cathodic process represented as follows:



#### *Product identification*

At the end of the electrolysis an aliquot of the solution was treated with an excess of dilute aqueous hydrochloric acid and the organic species were extracted into diethyl ether for identification by gas-liquid chromatography, and/or mass spectrometry, or NMR spectroscopy.

### **Results and discussion**

#### *Electrosynthesis*

The results of the electrosynthesis experiments are summarised in Table 1. The

TABLE 1  
ELECTROCHEMICAL COUPLING PRODUCTS

RX	% RX consumed	% R <sub>2</sub> formed vs. RX consumed
PhI	> 99	70
PhBr	> 99	75
PhCl	> 99	90
1-Br-naphthalene	> 99	75
4-BrC <sub>6</sub> H <sub>4</sub> COCH <sub>3</sub>	> 99	<sup>a</sup>
4-ClC <sub>6</sub> H <sub>4</sub> COCH <sub>3</sub>	> 99	<sup>a</sup>
3-ClC <sub>6</sub> H <sub>4</sub> CH <sub>3</sub>	90	80
2-ClC <sub>6</sub> H <sub>4</sub> COCH <sub>3</sub>	> 99	<sup>b</sup>
2-ClC <sub>6</sub> H <sub>4</sub> OCH <sub>3</sub>	75	<sup>b</sup>
1-Br,1-butene	> 99	50 <sup>c</sup>
2-Br,2-butene	> 99	50 <sup>c</sup>
PhCH=CHBr	> 99	> 95 <sup>c</sup>
PhCH <sub>2</sub> Cl	> 99	90
3-PhOC <sub>6</sub> H <sub>4</sub> CHClCH <sub>3</sub>	> 99	<sup>a,d</sup>
1-Br-hexane	90	70
2-Cl-thiophene	> 99	<sup>a</sup>
2-Cl-quinoline	> 95	<sup>a</sup>
5-Br-indole	> 99	<sup>a</sup>
4-BrC <sub>6</sub> H <sub>4</sub> OH	> 99	0
4-ClC <sub>6</sub> H <sub>4</sub> NO <sub>2</sub>	> 99	0

Solutions containing RX 0.3 M + NiBr<sub>2</sub>bipy 2 × 10<sup>-2</sup> M + bipy 4 × 10<sup>-2</sup> M were electrolysed at -1.3 V vs. SCE until formation of stable Ni<sup>0</sup> or at most until the charge reached one mol-electron per mole of RX

<sup>a</sup> Yield not calculated, but dimer identified as the main product. <sup>b</sup> Dimer only identified in trace quantities; main product is RH. <sup>c</sup> Starting material was a mixture of *Z*- and *E*-isomers; yield expressed as overall quantity of R<sub>2</sub>, with no distinction between isomers. <sup>d</sup> Starting material was a racemic mixture; enantiomeric and *meso* forms of R<sub>2</sub> obtained in 1/1 ratio.


yields are generally satisfactory, especially in view of the simplicity of the techniques used. In experiments in which RX was completely consumed, the catalyst was almost completely recovered as the Ni<sup>0</sup> complex, and further dimerisation occurred on adding more RX. The electrochemical coupling can obviously be used with a wide variety of organic halides, with the exception of *ortho*-substituted aryl halides, and compounds which have protonic or other easily reduced substituents such as OH, COOH or NO<sub>2</sub>.

#### Role of the magnesium anode

The sacrificial magnesium anode, whose weight loss corresponds to the process Mg → Mg<sup>2+</sup> + 2e based on the quantity of electricity passed through the cell, does not appear to play a critical role in the coupling reaction, but rather allows anodic oxidation to proceed at a potential at which the reduced species, and especially Ni<sup>0</sup>, are not re-oxidised. This is critical in using a single compartment cell, but we found that experiments in which anode and cathode compartments were separated by a glass sinter gave results identical to those with the apparatus described.

The magnesium may also act as a reducing agent in the catalytic cycle of the

nickel-bipy complexes, as has been reported for the synthesis of polyphenylenes from aromatic dibromides [9], since the reaction



is thermodynamically allowed. This may explain why we observed that the Faradaic yield for the conversion of RX was slightly greater than 100% in some cases. Tests carried out without electrolysis but with solution composition, time, etc. identical to those in the electrochemical experiments showed that such chemical reduction by magnesium metal can only account for at most, a 5–10% of the product yield.

The reductive coupling reaction is therefore seen as proceeding via the  $\text{Ni}^{\text{II}}/\text{Ni}^0$  catalytic cycle, and organometallic derivatives of magnesium are apparently not involved in the primary reaction. In keeping with this, magnesium can be satisfactorily replaced by other strongly electropositive metals such as aluminum or zinc.

*The mechanism of the catalytic coupling process*

The detailed mechanism of the electrochemical coupling which yields  $\text{R}_2$  is not clearly understood, but the present results, and the preliminary results of electroanalytical measurements [8], lead to the following tentative model.

The nickel(0) catalyst is obtained by the two-electron reduction:





The addition of excess RX ( $\text{RX} = n\text{-C}_6\text{H}_{13}\text{Br}$ ,  $\text{C}_6\text{H}_5\text{Br}$ ,  $\text{C}_6\text{H}_5\text{CH}_2\text{Cl}$ , etc.) (see Table 1) to an  $\text{Ni}^0$  solution prepared in this way causes total consumption of  $\text{Ni}^0$ , yielding  $\text{Ni}^{\text{II}}$  in quantities corresponding to 50% of the total nickel present. This excludes both the formation of a charge-transfer complex between  $\text{Ni}^0$  species and RX and the formation of a stable reducible  $\sigma$ -organonickel  $\text{RNiX}$  complex, such as was found in the case of the analogous triphenylphosphine complexes (reactions 2 and 3) [6b,c,d]. We suggest that  $\text{RNiX}$  formed in the oxidative addition process



must react quickly by one of the following sequences, which are based on the mechanism proposed by Tsou and Kochi [10] for the synthesis of biaryls from triethylphosphine complexes of nickel.

SCHEME 1



There is ample evidence for each of these reactions, at least in the thermal decomposition of  $\text{ArNiX}(\text{PPh}_3)_2$  [11], but in the present case one cannot exclude the reaction of  $\text{RNiX}$  with RX (Scheme 2) or with  $\text{Ni}^0$  (Scheme 3).

SCHEME 2



## SCHEME 3



In both of these schemes, the decomposition of  $\text{R}_2\text{Ni}$  to  $\text{R}_2$  completes the sequence (eq. 7). Both require that eq. 5 is the rate-controlling process for the consumption of  $[\text{Ni}(\text{bipy})_2]^0$ . We should point out that while Scheme 2 is consistent with the mechanism proposed for the metal-promoted coupling of organic halides [12], it appears to be inconsistent with the fact that stable  $\text{ArNiX} \cdot \text{bipy}$  compounds were isolated in experiments using  $\text{ArX}$  itself as the solvent at room temperature ( $\text{ArX} = \text{C}_6\text{H}_5\text{Cl}, \text{C}_6\text{H}_5\text{Br}, o, m, p\text{-CH}_3\text{C}_6\text{H}_4\text{Cl}$ ) [13].

In any case, the overall rate of the electrocatalytic process is limited by the regeneration of nickel(II) via eq. 7 and subsequent reactions. Equation 7 probably proceeds via an interaction involving  $\text{R}_2\text{Ni}$  and  $\text{RX}$ , followed by the reductive elimination of  $\text{R}_2$  (c.f. [14]) rather than by the first-order thermal decomposition of  $\text{R}_2\text{Ni}$  to  $\text{R}_2$  and  $\text{Ni}^0$ . In agreement with this, we find that while *o*-substituted aryl halides such as *o*-bromoanisole do not yield the appropriate coupled biphenyl, the addition of a very reactive compound such as ethyl iodide results in the formation of 2,2'-dimethoxy-1,1'-biphenyl, with 1-methoxy-2-ethylbenzene as a minor product. The formation of this biaryl can also be brought about by blowing air through the solution of  $\text{R}_2\text{Ni}$ .

*Mixed coupling reactions*

In terms of the usefulness of this system in achieving the coupling of radicals from two organic halides, we find that the reactivity of the respective halides is critically important, at least in the case of aryl compounds. Thus if one halide is much more reactive than the other (e.g.  $\text{RX} = \text{C}_6\text{H}_5\text{I}$ ,  $\text{R}'\text{X} = p\text{-CH}_3\text{C}_6\text{H}_4\text{Br}$ ), the electrolysis of an equimolar mixture gives essentially  $\text{R}_2$  in the early stages of the reaction, then  $\text{R}'_2$  as  $\text{RX}$  is consumed. On the other hand, when  $\text{RX}$  and  $\text{R}'\text{X}$  are of similar reactivity (e.g.,  $\text{C}_6\text{H}_5\text{Br}$  and  $p\text{-CH}_3\text{C}_6\text{H}_4\text{Br}$ ), a mixture of the three products is found at all stages of the reaction in the statistically determined approximate ratio 25%  $\text{R}_2$ , 50%  $\text{RR}'$ , 25%  $\text{R}'_2$ . Unfortunately these results do not help in identifying the reaction mechanism, since all three schemes predict this same product distribution.

**Conclusion**

The method described provides a simple and efficient route to coupling products from a range of organic halides. The reactions are carried out at controlled potentials, and the catalytic precursor  $\text{NiX}_2 \cdot \text{bipy}$  ( $\text{X} = \text{Cl}, \text{Br}$ ) is recovered in almost quantitative yield at the end of the reaction. The mechanism is not clearly identified, and work on this aspect of the problem is proceeding, but this does not detract from the usefulness of the preparative method.

**Acknowledgement**

One of us (D.G. Tuck) thanks the University of Windsor for the award of sabbatical leave.

## References

- 1 P.W. Jolly, in G. Wilkinson (Ed.), *Comprehensive Organometallic Chemistry*, Pergamon Press, Oxford, 1983, Vol. 8.
- 2 M.F. Semmelhack, P.M. Helquist and L.D. Jones, *J. Am. Chem. Soc.*, 93 (1971) 5908; 94 (1972) 9234.
- 3 K. Takagi, N. Hayama and K. Sasaki, *Bull. Chem. Soc. Japan*, 57 (1984) 1887.
- 4 K. Takagi, H. Mimura and S. Inokawa, *Bull. Chem. Soc. Japan*, 57 (1984) 3517, and refs. therein.
- 5 (a) M. Zembayashi, K. Tamao, J. Yoshida and M. Kumada, *Tetrahedron*, 47 (1977) 4089; (b) M. Tiecco, L. Testaferri, M. Tingoli, D. Chianelli and M. Montanucci, *Synthesis*, 9 (1984) 735.
- 6 (a) M. Mori, Y. Hashimoto and Y. Ban, *Tetrahedron Lett.*, (1980) 631; (b) M. Troupel, Y. Rollin, S. Sibille, J.F. Fauvarque and J. Perichon, *J. Chem. Research (S)*, (1980) 26; (c) *idem.*, *J. Organomet. Chem.*, 202 (1980) 435; (d) G. Bontempelli, S. Daniele and M. Fiorani, *J. Electroanal. Chem.*, 160 (1984) 249.
- 7 (a) B.J. Henne and D.E. Bartak, *Inorg. Chem.* 23 (1984) 369.  
(b) G. Bontempelli, S. Daniele and M. Fiorani, *Annal. di Chim.*, 75 (1985) 19.
- 8 Y. Rollin, M. Troupel and J. Perichon, unpublished results.
- 9 T. Yamamoto, Y. Hayashi and A. Yamamoto, *Bull. Chem. Soc. Japan*, 51 (1978) 2091.
- 10 T.T. Tsou and J.K. Kochi, *J. Am. Chem. Soc.*, 101 (1979) 7577.
- 11 A. Nakamura and S. Otsuka, *Tetrahedron Lett.*, 5 (1974) 463.
- 12 H. Zollinger, *Azo and Diazo Chemistry, Aliphatic and Aromatic Compounds*, Interscience, New York, 1961, Chap. 12.
- 13 M. Uchino, K. Asagi, A. Yamamoto and S. Ikeda, *J. Organomet. Chem.*, 84 (1975) 93.
- 14 T. Yamamoto, T. Kohara, K. Osakada and A. Yamamoto, *Bull. Chem. Soc. Japan*, 56 (1983) 2147.