

A Pd-Catalyzed [4 + 2] Annulation Approach to Fluorinated N-Heterocycles

Víctor García-Vázquez,[§] Larry Hoteite,[§] Christopher P. Lakeland,[§] David W. Watson, and Joseph P. A. Harrity*

Cite This: *Org. Lett.* 2021, 23, 2811–2815

Read Online


ACCESS |

Metrics & More

Article Recommendations

Supporting Information

ABSTRACT: 3-Fluoro- and trifluoromethylthio-piperidines represent important building blocks for discovery chemistry. We report a simple and efficient method to access analogs of these compounds that are armed with rich functionality allowing them to be chemoselectively derivatized with high diastereocontrol.



Fluorine-containing molecules exhibit a variety of useful properties in pharmaceuticals, agrochemicals, and materials science.^{1–3} In particular, the introduction of a C–F bond into a bioactive compound can have a dramatic impact on both the physical and chemical properties of the molecule.⁴ Additionally, nitrogen heterocycles are one of the most highly represented motifs within FDA approved small molecule drugs, with the piperidine ring as the most prevalent example of this class.⁵ Among other modifications of this cyclic amine, the selective incorporation of a fluorine atom at the 3-position of the piperidine scaffold has been demonstrated to be an effective strategy to improve the pharmacological properties of a number of biologically active compounds targeting SYK,⁶ CGRP,⁷ and MET kinase⁸ (Figure 1). In these cases, the fluorine atom plays a key role in preventing metabolism, as well

as modulating the basicity of the nitrogen atom. 3-Fluoropiperidines have also been investigated as radiotracers for NR2B NMDA receptor visualization.⁹

In spite of the importance of 3-fluoropiperidine derivatives, only a few general strategies exist to access these compounds. The electrophilic fluorination of piperidone derived enol equivalents has been reported,¹⁰ but this method faces regioselectivity issues when applied to nonsymmetrical scaffolds. The deoxofluorination of alkoxy piperidines has also been reported, but these reactions require extensive prefunctionalization and exhibit poor atom economy.¹¹ A particularly prevalent strategy relies on the intramolecular aminofluorination of olefins using Pd-catalysis¹² or hypervalent iodine reagents.¹³ However, these methods commonly employ strong oxidizing agents or rely on the use of stoichiometric quantities of toxic^{13b} or expensive reagents.^{13c} A recent report on the direct hydrogenation of fluorinated pyridines¹⁴ provides a diastereoselective synthetic pathway for the synthesis of fluoropiperidines, but the high H₂ pressure required reduces operational simplicity. We envisioned that our recently reported [4 + 2] annulation strategy to N-heterocycles¹⁵ could offer a powerful route to 3-fluoropiperidines using readily available α -fluoro- β -ketoester starting materials.¹⁶ Advantages of this method would include its highly modular




Figure 1. Prominent bioactive 3-fluoropiperidines.

Received: March 4, 2021


Published: March 24, 2021




nature, allowing for the rapid construction of the piperidine core. Moreover, the heterocycle products contain orthogonal functionality that would allow their elaboration to new products through multiple vectors (Scheme 1).

Scheme 1. Synthetic Routes to 3-Fluoropiperidines


a) Intramolecular aminofluorination of olefins:



b) Hydrogenation of fluoropiperidines:



c) Pd-catalyzed [4+2] annulation (this work):




We began our studies by investigating the allylation/condensation reaction of readily available α -fluoro- β -ketoester **1a** and cyclic carbamate **2** as shown in Scheme 2. Subjecting this substrate to 5 mol % of Pd(dba)₂ and 15 mol % of ligand **L1** followed by treatment of the intermediate with TFA led smoothly to the desired 3-fluoropiperidine **4a** in high yield (Scheme 2a). Moreover, this product was also obtained in comparable yield employing a one-pot procedure without isolation of intermediate **3**. To our delight, scaling up the reaction to multigram quantities yielded **4a** with similar results. We then examined the applicability of this methodology to a


number of α -fluoro- β -ketoesters. Aryl substituted imines with either electron-withdrawing or electron-donating groups at the *para*-position on the aryl ring gave excellent yields (**4b–f**). Substitution at other points on the aryl ring such as *ortho*-methyl (**4g**) and naphthyl (**4h**) are also well tolerated. The heterocyclic thiophenyl containing substrate **1i** was also converted into the corresponding piperidine imine **4i** in 78% yield. In addition to α -fluoro- β -ketoesters, numerous other fluorinated nucleophiles could be employed in the allylation/condensation sequence including α -fluoro- β -ketonitriles (**4j**), α -fluoro- β -ketosulfones (**4k**), and α -fluoro- β -ketoamides (**4l**). Unfortunately, however, α -fluoroketones bearing alkyl groups were not transformed to the corresponding heterocycles, and a complex mixture of products was instead produced. Furthermore, this sequence could also be applied to alkyl substituted α -fluoro- β -ketoesters in a regioselective manner (Scheme 2b). Simple alkyl groups containing various levels of substitution at the α -position (Me, 1°, 2°, and 3°) afforded the piperidine imines **4n–4r** in high yields. Finally, a derivative of L-proline was evaluated, with **4s** obtained in good yield and with moderate diastereoselectivity. The excellent functional group tolerance of this reaction sequence serves to highlight the mild nature of this procedure.

We next turned our attention to demonstrating that the functionalized 3-fluoropiperidines were versatile intermediates for organic chemistry (Scheme 3). First, a chemoselective reduction of **4a** using NaBH(OAc)₃ in acetic acid solvent produced saturated piperidine **5** with high diastereoselectivity. Subsequent protection of **5** using di-*tert*-butyl dicarbonate then gave compound **6** in 88% yield, with the X-ray structure of **6** (CCDC 2063492) providing the relative configuration of the major diastereoisomer obtained in this process. Interestingly, a chemoselective reduction of the ester moiety was also achieved using LiAlH₄ to give **7** in moderate yield. A hydrolytic

Scheme 2. Reaction Scope^a



^aReaction conditions: **1** (0.3 mmol), **2** (0.2 mmol), Pd(dba)₂ (10 μ mol, 5 mol %), **L1** (30 μ mol, 15 mol %), DCM (0.1 M), rt, 18 h under N₂.


Scheme 3. Chemoselective Functionalization of 3-Fluoropiperidine Imines^a

^aReagents and conditions: (a) NaBH(OAc)₃ (1.5 equiv), AcOH, rt, 18 h (90%); (b) Boc₂O (2.0 equiv), Et₃N (2.0 equiv), THF, rt, 18 h (88%); (c) LiAlH₄ (2.0 equiv), THF, rt, 3.5 h (50%); (d) aq. HCl (15 equiv), 100 °C, 1 h (quant.); (e) NaBH₄ (2.0 equiv), MeOH, 0 °C to rt, 18 h (77%); (f) Hoveyda–Grubbs second Gen. (5 mol %), pent-4-en-1-yl acetate (3 equiv), DCM, 25 °C 18 h then reflux, 3 h (54%).

decarboxylation using aq. HCl and heating afforded fluoropiperidine **8**, which could then be reduced using NaBH₄ to provide saturated piperidine **9** in 77% yield as a single diastereoisomer following column chromatography.¹⁷ Notably, this decarboxylation circumvents the limitation associated with the poor reactivity of α -fluoroketones in the allylation/condensation cascade. Selective functionalization of the exocyclic alkene is also possible; cross-metathesis produced **10** as a mixture of geometric isomers. The ability to selectively functionalize each functional handle in piperidine imines **4** demonstrates their utility as synthetic intermediates.

The suitability of our method for accessing useful fluorinated heterocycles suggested that it might be adapted to allow the incorporation of trifluoromethylthio (SCF₃) groups. In this regard, and to the best of our knowledge, only two examples of 3-SCF₃-substituted piperidines have been reported.¹⁸ Due to its electron-withdrawing nature and high lipophilicity, the SCF₃ moiety can significantly modulate the pharmacological properties of bioactive compounds.¹⁹ Nevertheless, the availability of synthetic methods that deliver saturated *N*-trifluoromethyl-thiolated six-membered heterocycles is scarce, and those that are documented suffer from limited substrate scope.²⁰

Our efforts to employ the [4 + 2] annulation sequence to α -SCF₃-ketones is summarized in Scheme 4. Aryl substituted ketones proved to be excellent substrates for this transformation, generating a range of 3-SCF₃-substituted piperidines under mild conditions. Unfortunately, these products proved to be unstable to chromatography, and so we used a borohydride reduction step prior to isolation. Accordingly, 2-aryl 3-trifluoromethylthio-piperidines **13a–g** were isolated in excellent yields over three steps, and with very high *cis*-stereocontrol. X-ray crystal structure analysis of aryl substituted products **13a** (CCDC 2063487) and **13e** (CCDC 2063489) confirmed the relative stereochemistry of the major diastereomer in these cases, and the stereochemistry of all other aryl-

Scheme 4. Synthesis of 3-SCF₃-Substituted Piperidines^a

^aReagents and conditions: **11** (0.7 mmol), **2** (0.47 mmol), Pd(dba)₂ (23 μ mol, 5 mol %), **L1** (70 μ mol, 15 mol %), CH₂Cl₂ (0.1 M), RT, 18 h under N₂. ^bHeated at 40 °C.

substituted products was assigned by inference. Unfortunately, however, 2-aryl-substituted ketones containing electron-withdrawing groups (4-nitrophenyl and 4-trifluoromethylphenyl) were found to decompose during the TFA-mediated deprotection–condensation step. Finally, α -SCF₃-propiophenone led to a more substituted analog **13h**, while the potential to access 2-alkyl piperidine products was confirmed in one case, albeit in low yield.

In conclusion, we report that 3-fluoropiperidines bearing orthogonal imine, ester, and alkene functionality can be readily prepared and chemoselectively derivatized, providing a powerful approach to these important substructures. Moreover, this method can be extended to provide the first general means to incorporate the 3-trifluoromethylthio-group into piperidines, offering a new and efficient entry into these important scaffolds.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.orglett.1c00752>.

Details of experimental procedures and spectroscopic data. NMR spectral data included (PDF)

Accession Codes

CCDC 2063487, 2063489, and 2063492 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

■ AUTHOR INFORMATION

Corresponding Author

Joseph P. A. Harrity – Department of Chemistry, University of Sheffield, Sheffield S3 7HF, United Kingdom; orcid.org/0000-0001-5038-5699; Email: j.harrity@sheffield.ac.uk

Authors

Victor García-Vázquez – Department of Chemistry, University of Sheffield, Sheffield S3 7HF, United Kingdom
Larry Hoteite – Department of Chemistry, University of Sheffield, Sheffield S3 7HF, United Kingdom
Christopher P. Lakeland – Department of Chemistry, University of Sheffield, Sheffield S3 7HF, United Kingdom
David W. Watson – Medicinal Chemistry, Oncology R&D Research and Early Development, AstraZeneca Cambridge Science Park, Cambridge CB4 0WG, United Kingdom

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acs.orglett.1c00752>

Author Contributions

[§]V.G.V., L.H., and C.P.L. contributed equally to this work.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We are grateful for support from AstraZeneca and the European Funding Horizon 2020-MSCA (ITN-EJD CAT-MEC 14/06-721223).

■ REFERENCES

- (1) (a) Champagne, P. A.; Desroches, J.; Hamel, J.-D.; Vandamme, M.; Paquin, J.-F. Monofluorination of Organic Compounds: 10 Years of Innovation. *Chem. Rev.* **2015**, *115*, 9073–9174. (b) Fustero, S.; Sedgwick, D. M.; Román, R.; Barrio, P. Recent advances in the synthesis of functionalised monofluorinated compounds. *Chem. Commun.* **2018**, *54*, 9706–9725. (c) Neumann, C. N.; Ritter, T. Late Stage Fluorination: Fancy Novelty or Useful Tool. *Angew. Chem., Int. Ed.* **2015**, *54*, 3216–3221.
- (2) (a) Purser, S.; Moore, P. R.; Swallow, S.; Gouverneur, V. Fluorine in medicinal chemistry. *Chem. Soc. Rev.* **2008**, *37*, 320–330. (b) Gillis, E. P.; Eastman, K. J.; Hill, M. D.; Donnelly, D. J.; Meanwell, N. A. Applications of Fluorine in Medicinal Chemistry. *J. Med. Chem.* **2015**, *58*, 8315–8359.
- (3) Berger, R.; Resnati, G.; Metrangolo, P.; Weber, E.; Hulliger, J. Organic fluorine compounds: a great opportunity to enhanced materials properties. *Chem. Soc. Rev.* **2011**, *40*, 3496–3508.
- (4) Müller, K.; Faeh, C.; Diederich, F. Fluorine in Pharmaceuticals: Looking Beyond Intuition. *Science* **2007**, *317*, 1881–1886.
- (5) Vitaku, E.; Smith, D. T.; Njardarson, J. T. Analysis of the Structural Diversity, Substitution Patterns, and Frequency of Nitrogen Heterocycles among U. S. FDA Approved Pharmaceuticals. *J. Med. Chem.* **2014**, *57*, 10257–10274.
- (6) (a) Curtis, N. R.; Davies, S. H.; Gray, M.; Leach, S. G.; McKie, R. A.; Vernon, L. E.; Walkington, A. J. Asymmetric Fluorination Approach to the Scalable Synthesis of a SYK Inhibitor. *Org. Process Res. Dev.* **2015**, *19*, 865–871. (b) Mócsai, A.; Ruland, J.; Tybulewicz, V. L. The SYK tyrosine kinase: a crucial player in diverse biological functions. *Nat. Rev. Immunol.* **2010**, *10*, 387–402. (c) Tan, S.-L.; Liao, C.; Lucas, M. C.; Stevenson, C.; DeMartino, J. A. Targeting the SYK-BTK axis for the treatment of immunological and hematological disorders: recent progress and therapeutic perspectives. *Pharmacol. Ther.* **2013**, *138*, 294–309.
- (7) Molinaro, C.; Phillips, E. M.; Xiang, B.; Milczek, E.; Shevlin, M.; Balsells, J.; Ceglia, S.; Chen, J.; Chen, L.; Chen, Q.; Fei, Z.; Hoerner, S.; Qi, J.; de Lera Ruiz, M.; Tan, L.; Wan, B.; Yin, J. *J. Org. Chem.* **2019**, *84*, 8006–8018.
- (8) Liederer, B. M.; Berezhkovskiy, L. M.; Dean, B. J.; Dinkel, V.; Peng, J.; Merchant, M.; Plise, E. G.; Wong, H.; Liu, X. Precrinal absorption, distribution, metabolism, excretion, and pharmacokinetic-pharmacodynamic modelling of N-(4-(3-((3S,4R)-1-ethyl-3-fluoropiperidine-4-ylamino)-1H-pyrazolo[3,4-b]pyridin-4-yloxy)-3-fluorophenyl)-2-(4-fluorophenyl)-3-oxo-2,3-dihydropyrazine-4-carboxamide, a novel MET Kinase inhibitor. *Xenobiotica* **2011**, *41*, 327–339.
- (9) (a) Koudih, R.; Gilbert, G.; Dhilly, M.; Abbas, A.; Barré, L.; Debryne, D.; Sobrio, F. Synthesis and in vitro characterization of trans and cis-[¹⁸F]-4-methylbenzyl 4-[(pyrimidin-2-ylamino)methyl]-3-fluoropiperidine-1-carboxylates as new potential PET radiotracer candidates for the NR2B subtype N-methyl-D-aspartate receptor. *Eur. J. Med. Chem.* **2012**, *53*, 408–415. (b) Koudih, R.; Gilbert, G.; Dhilly, M.; Abbas, A.; Barré, L.; Debryne, D.; Sobrio, F. Radiolabelling of 1,4-disubstituted 3-[¹⁸F]fluoropiperidines and its application to new radiotracers for NR2B NMDA receptor visualization. *Org. Biomol. Chem.* **2012**, *10*, 8493–8500.
- (10) (a) Sun, A.; Lankin, D. C.; Hardcastle, K.; Snyder, J. P. 3-Fluoropiperidines and N-Methyl-3-fluoropiperidinium salts: The Persistence of an Axial Fluorine. *Chem. - Eur. J.* **2005**, *11*, 1579–1591. (b) Shaw, S. J.; Goff, D. A.; Boralsky, L. A.; Irving, M.; Singh, R. Enantioselective Synthesis of cis-3-Fluoropiperidin-4-ol, a Building Block for Medicinal Chemistry. *J. Org. Chem.* **2013**, *78*, 8892–8897. (c) Walpole, C.; Liu, Z.; Lee, E. E.; Yang, H.; Zhou, F.; Mackintosh, N.; Sjogren, M.; Taylor, D.; Shen, J.; Batey, R. A. Diastereoselective synthesis of fluorinated piperidine quinazoline spirocycles as iNOS selective inhibitors. *Tetrahedron Lett.* **2012**, *53*, 2942–2947.
- (11) (a) Goldberg, N. W.; Shen, X.; Li, J.; Ritter, T. AlkylFluor: Deoxyfluorination of Alcohols. *Org. Lett.* **2016**, *18*, 6102–6104. (b) Li, X.; Russell, R. K.; Spink, J.; Ballentine, S.; Teleha, C.; Branum, S.; Wells, K.; Beauchamp, D.; Patch, R.; Huang, H.; Player, M.; Murray, W. Process for Development for Scale-Up of a Novel 3,5-Substituted Thiazolidine-2,4-dione Compound as a Potent Inhibitor for Estrogen-Related Receptor 1. *Org. Process Res. Dev.* **2014**, *18*, 321–330.
- (12) Wu, T.; Yin, G.; Liu, G. Palladium Catalyzed Intramolecular Aminofluorination of Unactivated Alkenes. *J. Am. Chem. Soc.* **2009**, *131*, 16354–16355.
- (13) (a) Serguchev, Y. A.; Ponomarenko, M. V.; Ignat'ev, N. V. Azafuorocyclization of nitrogen-containing unsaturated compounds. *J. Fluorine Chem.* **2016**, *185*, 1–16. (b) Vardelle, E.; Martin-Mingot, A.; Jouannetaud, M.-P.; Bachmann, C.; Marrot, J.; Thibaudeau, S. Substitution Effect on the Cyclization/Fluorination Reaction of N-Dienes in Superacid. *J. Org. Chem.* **2009**, *74*, 6025–6034. (c) Kong, W.; Feige, P.; de Haro, T.; Nevado, C. Regio- and Enantioselective Aminofluorination of Alkenes. *Angew. Chem., Int. Ed.* **2013**, *52*, 2469–2473.
- (14) Nairoukh, Z.; Wollenburg, M.; Schlepfforst, C.; Bergander, K.; Glorius, F. The Formation of all-cis-(multi)fluorinated piperidines by a dearomatization-hydrogenation process. *Nat. Chem.* **2019**, *11*, 264–270.
- (15) Allen, B. D. W.; Connolly, M. J.; Harrity, J. P. A. A Pd-Catalyzed Synthesis of Functionalized Piperidines. *Chem. - Eur. J.* **2016**, *22*, 13000–13003.
- (16) Tang, L.; Yang, Z.; Jiao, J.; Cui, Y.; Zou, G.; Zhou, Q.; Zhou, Y.; Rao, W.; Ma, X. Chemoselective Mono- and Difluorination of 1,3-Dicarbonyl Compounds. *J. Org. Chem.* **2019**, *84*, 10449–10458.
- (17) The stereochemical assignment of compound **9** was made on the basis of ¹⁹F–¹H coupling constants in the ¹⁹F NMR spectrum. See [Supporting Information](#) for details.
- (18) (a) Luo, J.; Liu, Y.; Zhao, X. Chiral Selenide-Catalyzed Enantioselective Construction of Saturated Trifluoromethylthiolated Azaheterocycles. *Org. Lett.* **2017**, *19*, 3434–3437. (b) Zhang, C.; Wang, Y.; Song, Y.; Gao, H.; Sun, Y.; Sun, X.; Yang, Y.; He, M.; Yang, Z.; Zhan, L.; Yu, Z.-X.; Rao, Y. Synthesis of quaternary carbon-centered Benzoinolizidines via novel photoredox-catalyzed Alkene

aminoarylation: Facile access to Tylophorine and analogues. *CCS Chemistry* **2019**, *1*, 352–364.

(19) (a) Manteau, B.; Pazenok, S.; Vors, J.-P.; Leroux, F. R. New Trends in the chemistry of α -fluorinated ethers, thioethers, amines and phosphines. *J. Fluorine Chem.* **2010**, *131*, 140–158. (b) Landelle, G.; Panossian, A.; Leroux, F. Trifluoromethylethers and – thioethers as tools for medicinal chemistry and drug discovery. *Curr. Top. Med. Chem.* **2014**, *14*, 941–951. (c) Barata-Vallejo, S.; Bonesi, S.; Postigo, A. Late Stage trifluoromethylthiolation strategies for organic compounds. *Org. Biomol. Chem.* **2016**, *14*, 7150–7182. (d) Xu, X.-H.; Matsuzaki, K.; Shibata, N. Synthetic Methods for Compounds having CF₃-units on Carbon by Trifluoromethylation, Trifluoromethylthiolation, Triflylation and Related Reactions. *Chem. Rev.* **2015**, *115*, 731–764.

(20) (a) Hu, F.; Shao, X.; Zhu, D.; Lu, L.; Shen, Q. Silver-catalyzed decarboxylative trifluoromethylthiolation of aliphatic carboxylic acids in aqueous emulsion. *Angew. Chem., Int. Ed.* **2014**, *53*, 6105–6109. (b) Candish, L.; Pitzer, L.; Gómez-Suárez, A.; Glorius, F. Visible Light-Promoted Decarboxylative Di- and Trifluoromethylthiolation of Alkyl Carboxylic Acids. *Chem. - Eur. J.* **2016**, *22*, 4753–4756. (c) Shao, X.; Liu, T.; Lu, L.; Shen, Q. Copper-Catalyzed Trifluoromethylthiolation of Primary and Secondary Alkylboronic Acids. *Org. Lett.* **2014**, *16*, 4738–4741.