

## Enantioselective Formal Aza-Diels–Alder Reactions of Enones with Cyclic Imines Catalyzed by Primary Aminothioureas

Mathieu P Lalonde, Meredith Ann McGowan, Naomi S Rajapaksa, and Eric Niels Jacobsen

*J. Am. Chem. Soc.*, **Just Accepted Manuscript** • DOI: 10.1021/ja310718f • Publication Date (Web): 15 Jan 2013

Downloaded from <http://pubs.acs.org> on January 16, 2013

### Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



# Enantioselective Formal Aza-Diels–Alder Reactions of Enones with Cyclic Imines Catalyzed by Primary Aminothiureas

Mathieu P. Lalonde, Meredith A. McGowan, Naomi S. Rajapaksa, and Eric N. Jacobsen\*

Department of Chemistry and Chemical Biology, Harvard University, Cambridge, Massachusetts 02138

**ABSTRACT:** A highly enantio- and diastereoselective synthesis of indolo- and benzoquinolizidine compounds has been developed through the formal aza-Diels–Alder reaction of enones with cyclic imines. This transformation is catalyzed by a new bifunctional primary aminothiurea that achieves simultaneous activation of both the enone and imine reaction components.

## Introduction

Chiral indolo- and benzoquinolizidine frameworks reside within a wide assortment of biologically active natural products and synthetic pharmaceutical compounds (Figure 1).<sup>1,2</sup> Among laboratory approaches to access structurally and stereochemically complex members of this class of heterocycles, formal aza-Diels–Alder (FADA) reactions between enones and cyclic imines (Scheme 1) are particularly attractive from the perspective of convergency, resulting in the concomitant formation of a C–C and C–N bond and up to four new stereocenters.<sup>3,4</sup> A proline-based protocol for enantioselective FADA reactions involving dihydro- $\beta$ -carboline has been described by Itoh and coworkers,<sup>5</sup> but the scope of this method has proven to be very limited.<sup>6,7</sup> We describe here the discovery of a new primary aminothiurea catalyst with broad scope for the highly enantio- and diastereoselective synthesis of indolo- and benzoquinolizidine derivatives through the formal [4+2] cycloaddition between enones and cyclic imines.





Figure 1. Selected examples of bioactive indolo- and benzoquinolizidine derivatives

## Scheme 1. FADA Reactions of Enones and Cyclic Imines and Approach to Catalysis




Our approach to catalysis of the FADA reaction was premised on the possibility of a cooperative mechanism, with specific acid activation of the imine combined with activation of the enone as the corresponding dienamine (Scheme 1). Ureas and thiureas have demonstrated broad utility as hydrogen bond donor catalysts for additions to imines, inducing electrophile activation by either direct binding or indirectly by means of anion binding.<sup>8,9</sup> In addition, several studies have highlighted the utility of primary amines as enamine precursors in transformations that involve ketone and hindered aldehyde substrates.<sup>10</sup> On the basis of these precedents, we undertook an investigation of primary amine-containing hydrogen bond donors as bifunctional catalysts for the enantioselective FADA reaction of enones and cyclic imines.

## Results and Discussion

A series of thiureas of the general structure **1** was evaluated for catalysis of the FADA reaction between dihydro- $\beta$ -carboline **2a**<sup>11</sup> and commercially available enone **3a** (Table 1). Systematic variation of the amide, amino acid, and diamine components of the catalyst (e.g., entries 1–6)<sup>12</sup> led to the determination that the primary amino functional group was essential for catalysis (entries 5, 6), and that the steric properties of the amide exerted a significant influence on stereoselectivity. Ultimately, tertiary amide derivative **1c** was shown to be optimal for the generation of the exo cycloadduct **4aa** with respect to both diastereo- and enantioselectivity. A crucial role of a weak Brønsted acid additive was also established: in the absence of catalytic AcOH, very low turnover of the thiurea catalyst was observed, although the enantioselectivity re-

mained high (entry 7). A beneficial role of added AcOH has been observed in other primary aminothiourea-catalyzed reactions,<sup>10</sup> and may be ascribed to acceleration of the condensation and/or hydrolysis steps integral to the enamine catalysis cycle. Additionally, thiourea-assisted, specific acid activation of the imine (Scheme 1, X = OAc) represents another potential role of the acid co-catalyst.<sup>13</sup> Consistent with this hypothesis, the identity of the carboxylic acid was found to have a measurable influence on the enantioselectivity of the transformation (AcOH vs. BzOH, entries 3 and 8). Finally, improved product yields were obtained using an excess of enone (Table 1, entries 3 vs. 9), as this served to curtail the effects of product inhibition.<sup>14</sup>

**Table 1. Catalyst Optimization Studies<sup>a</sup>**




entry	catalyst	t (h)	yield <sup>b</sup> (%)	dr <sup>c</sup> (4aa:5aa)	ee <sup>d</sup> (%) (4aa/5aa)
1	<b>1a</b>	72	67	2.1:1	88/95
2	<b>1b</b>	52	75	3.2:1	88/95
3	<b>1c</b>	60	89	9.4:1	99/97
4	<b>1d</b>	48	77	3.7:1	94/98
5	<b>1e</b>	72	0	–	–
6	<b>1f</b>	60	0	–	–
7 <sup>e</sup>	<b>1c</b>	60	6	5.6:1 <sup>f</sup>	94/90
8 <sup>g</sup>	<b>1c</b>	60	92	5.6:1	97/96
9 <sup>h</sup>	<b>1c</b>	72	68	9.2:1	99/99

<sup>a</sup>Unless noted otherwise, reactions were conducted using **2a** (0.1 mmol, 1.0 equiv), **3a** (2.0 equiv), **1a-e** (5.0 mol%), and AcOH (5.0 mol%) at 4 °C in anhydrous toluene ([**2a**]<sub>0</sub>=0.1 M). The notation **4xy** refers to the FADA adduct derived from imine **2x** and enone **3y**. <sup>b</sup>Isolated yield of diastereomerically pure **4aa** following purification by flash column chromatography. <sup>c</sup>Determined by HPLC analysis of the unpurified reaction mixture. <sup>d</sup>Determined by HPLC analysis of pure, isolated product diastereomers using commercial chiral columns. <sup>e</sup>Reaction carried out with 0 equiv of AcOH. <sup>f</sup>Calculated based on the isolated yields of each diastereomer. <sup>g</sup>Reaction carried out with 5.0 mol% BzOH instead of AcOH <sup>h</sup>Reaction carried out with 1.2 equiv of **3a**.

Under the optimized conditions outlined above, primary aminothiourea **1c** displays broad scope in FADA reactions between dihydro-β-carboline derivatives and conjugated enones (Table 2; see Supporting Information for additional substrates). High yields of highly enantioenriched adducts were obtained with enones bearing β-aryl and heteroaryl sub-

stituents (entries 1–9), linear and branched alkyl substituents (entries 10–13), as well as cyclic enones (entry 14). Longer reaction times and/or higher catalyst loadings were required for α-substituted (entries 13, 14) and electron-rich β-aryl substituted enones (entry 4), presumably due to slower imine/enamine formation with the aminothiourea catalyst.

**Table 2. Primary Aminothiourea-Catalyzed FADA Reactions of Enones and Substituted 9-Tosyl-3,4-dihydro-β-carboline Imines<sup>a</sup>**




entry	imine	enone	t (h)	yield of <b>4</b> <sup>b</sup> (%)	dr <sup>c</sup> (4:5)	ee <sup>d</sup> (%) (4/5)
1	<b>2a</b>	<b>3a</b>	60	90	9.4:1	99/97
2	<b>2a</b>	<b>3b</b>	48	95	18:1	99/n.d.
3	<b>2a</b>	<b>3c</b>	72	87	7.6:1	99/99
4	<b>2a</b>	<b>3d</b>	312	57	1.3:1	99/99
5 <sup>e</sup>	<b>2a</b>	<b>3e</b>	48	91	10:1	99/n.d.
6	<b>2a</b>	<b>3f</b>	30	92	13:1	98/n.d.
7	<b>2a</b>	<b>3g</b>	48	96	>19:1	98/n.d.
8	<b>2a</b>	<b>3h</b>	48	84	4.2:1	96/99
9	<b>2a</b>	<b>3i</b>	48	81	4.5:1	99/99
10	<b>2a</b>	<b>3j</b>	55	91	>19:1	98/n.d.
11	<b>2a</b>	<b>3k</b>	60	>99	>19:1	98/n.d.
12	<b>2a</b>	<b>3l</b>	192	90	>19:1	97/n.d.
13 <sup>f</sup>	<b>2a</b>	<b>3m</b>	192	88	7.0:1	95/92
14 <sup>g</sup>	<b>2a</b>	<b>3n</b>	48	87	>19:1	99/n.d.
15 <sup>h</sup>	<b>2a</b>	<b>3o</b>	408	50	–	92/n.d.
16	<b>2b</b>	<b>3j</b>	48	88	>19:1	97/n.d.
17	<b>2c</b>	<b>3j</b>	48	87	>19:1	97/n.d.

<sup>a</sup>Unless noted otherwise, reactions were conducted using **2** (0.3 mmol, 1.0 equiv), **3** (2.0 equiv), **1c** (5.0 mol%), and AcOH (5.0 mol%) at 4 °C in anhydrous toluene ([**2**]<sub>0</sub>=0.1 M). <sup>b</sup>Yields of isolated diastereomerically pure product following flash column chromatography on silica gel. <sup>c</sup>Determined by chiral HPLC analysis of the unpurified reaction mixture. <sup>d</sup>Determined by HPLC analysis of pure, isolated product diastereomers using commercial chiral columns (see Supporting Information). <sup>e</sup>The absolute con-

Figurations of **4ae** and **5ae** were determined via X-ray crystallography. The stereochemistry of all other adducts is assigned by analogy. <sup>f</sup>**1c** (20 mol%), AcOH (20 mol%), 23 °C. <sup>g</sup>**1c** (10 mol%), AcOH (10 mol%), 23 °C. <sup>h</sup>Incomplete conversion after 408 hours.

The scope of the aminothiurea methodology with respect to the imine component includes both electron-deficient and -rich dihydro- $\beta$ -carbolines (Table 2, entries 16, 17). Substituted 3,4-dihydroisoquinolines (**6**) also underwent FADA reactions with enone **3j** in the presence of catalyst **1c** to generate chiral benzoquinolizidine frameworks in high yield, dr, and ee, although higher catalyst loadings were required for this imine substrate class (Table 3).<sup>15</sup>

**Table 3. Primary Aminothiurea-Catalyzed FADA Reactions of Enones and 3,4-Dihydroisoquinolines<sup>a</sup>**




entry	imine	t (h)	yield <sup>b</sup> (%) (7)	dr <sup>c</sup> (7:8)	ee <sup>d</sup> (%) (7)
1 <sup>e</sup>	6a	60	88	>19:1	92
2	6b	48	89	16:1	91
3 <sup>f</sup>	6c	48	>99	17:1	92
4	6d	72	93	>19:1	92
5	6e	48	88	15:1	91

<sup>a</sup>Unless otherwise noted, reactions were conducted using **6** (0.2 mmol, 1.0 equiv), **3j** (1.5 equiv), **1c** (15 mol%), and AcOH (15 mol%) at 4 °C in anhydrous toluene ([**6**]=0.1 M). <sup>b</sup>Yields of isolated diastereomerically pure product following flash column chromatography on silica gel. <sup>c</sup>Determined by chiral HPLC analysis of the unpurified reaction mixture. <sup>d</sup>Determined by chiral HPLC analysis of pure, isolated product diastereomers. <sup>e</sup>**6a** (0.4 mmol scale). <sup>f</sup>**6c** (0.35 mmol scale).


Examination of the data in Table 2 reveals a trend where FADA adducts bearing electron-rich C4 substituents are obtained in relatively low diastereomeric ratios (e.g. entries 4, 8, 9). In order to determine whether diastereoselectivity is under kinetic or thermodynamic control, diastereomerically pure adduct **4ad** was subjected to the reaction conditions in the presence of one equivalent of enone **3d** for 4 d.<sup>16</sup> A product ratio of 6.3:1 (**4ad**:**5ad**) was determined.<sup>17,18</sup> However, a crossover experiment conducted between diastereomerically pure adduct **4aa** and enone **3j** yielded none of the possible crossover product **4aj**, indicating that the overall reaction is not reversible.<sup>19</sup> Instead, the observed decrease in the product diastereomeric ratio suggests that during the course of the reaction, kinetic adducts **4** can undergo epimerization at C4 to thermodynamic adducts **5**,<sup>20</sup> but that the C10 center is formed irreversibly.

Based on these observations, we outline a catalytic cycle for the primary aminothiurea-catalyzed FADA reaction (Scheme 2).<sup>21</sup> In this proposal, activation of the enone is achieved by the catalyst through formation of the corresponding covalently bound dienamine, and simultaneously the imine is activated as a thiourea-bound iminium ion (**A**). Cyclization to intermediate **C** can then proceed by means of an irreversible concerted [4+2] cycloaddition or by a step-wise Mannich-conjugate addition, in which the C10 center is installed irreversibly. Tautomerization of enamine **C** to iminium ion **D**, followed by hydrolysis releases the FADA adduct and regenerates the aminothiurea catalyst. The diminished diastereomeric ratios of FADA adducts containing electron-rich C4 substituents can be ascribed to either or both of the mechanisms outlined in Scheme 3. An off-cycle retro-Mannich reaction of intermediate **D** or a potentially on-cycle  $\beta$ -elimination from tautomeric intermediate **C** would generate iminium ions **E** and **B**, respectively, which would both be stabilized by electron-rich C4 substituents. Ring closure from either of these iminium ion intermediates could result in the formation of thermodynamic FADA adducts via intermediates **C'** or **D'**.

**Scheme 2. Proposed Catalytic Cycle for the Formation of FADA Adducts**



**Scheme 3. Possible C4-Epimerization Pathways**



## Conclusion

In conclusion, we have developed a primary aminothiourea-catalyzed formal aza-Diels–Alder reaction of enones and cyclic imines for the enantioselective synthesis of a variety of stereochemically complex indolo- and benzoquinolizidine building blocks. The hydrogen bond donor and primary amine are essential functional components of the optimal catalyst, allowing for dual activation of the reaction components. The application of this catalyst to the synthesis of alkaloid natural products is the subject of ongoing research and will be published in due course.

## ASSOCIATED CONTENT

**Supporting Information.** Catalyst optimization studies, complete experimental procedures and characterization data for substrates and FADA products, ee and dr determination, and crystallographic data for compounds **4ae** and **5ae**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## AUTHOR INFORMATION

### Corresponding Author

[jacobsen@chemistry.harvard.edu](mailto:jacobsen@chemistry.harvard.edu)

## ACKNOWLEDGMENT

This work was supported by the NIGMS (GM-43214 and P50 GM-69721) and by predoctoral fellowship support to M.P.L. from NSERC and La Fondation Baxter et Alma Ricard, to M.A.M from Eli Lilly, and to N.S.R from Boehringer Ingelheim.

## REFERENCES

- (1) (a) *The Alkaloids: Chemistry and Biology*; Cordell, G. A., Ed.; Academic Press: New York, 1998; Vol 50. (b) Baxter, E. W.; Mariano, P. S. In *Alkaloids: Chemical and Biological Perspectives*; Pelletier, S. W., Ed.; Springer-Verlag: New York, 1992; Vol 8, pp 197–319.
- (2) (a) Openshaw, H. T.; Whittaker, N. *J. Chem. Soc.* **1963**, 1461–1471. (b) Szántay, C.; Töke, L.; Honty, K. *Tetrahedron Lett.* **1965**, 22, 1665–1670. (c) Szántay, C.; Vedres, A.; Toth, G. *Heterocycles* **1977**, 6, 1793–1798. (d) Maryanoff, B. E.; McComsey, D. F.; Taylor, R. J. Jr.; Gardocki, J. F. *J. Med. Chem.* **1981**, 24, 79–88. (e) Caroon, J. M.; Clark, R. D.; Kluge, A. F.; Lee, C.-H.; Strosberg, A. M. *J. Med. Chem.* **1983**, 26, 1426–1433. (f) Blaskó, G.; Knight, H.; Honty, K.; Szántay, C. *Liebigs Ann. Chem.* **1986**, 655–663, and references therein. (g) Din Belle, D.; Jokela, R.; Tolvanen, A.; Haapalinna, A.; Karjalainen, A.; Sallinen, J.; Ratilainen, J. Polycyclic Compounds as Potent Alpha2-Adrenoreceptor Antagonists. PCT/FI2003/000255, September 10, 2003.
- (3) For the first reports describing the addition of enones to cyclic imines, see: (a) Openshaw, H. T.; Whittaker, N. *J. Chem. Soc.* **1963**, 1449–1460. (b) Brossi, A. Preparation of Substituted 2-Oxobenzoquinolizines. U.S. Patent 3,045,021, September 13, 1960. (c) Beke, D.; Szántay, C. *Chem. Ber.* **1962**, 95, 2132–2136.
- (4) For reviews on alternate syntheses of chiral indolo- and benzoquinolizidines, see: (a) Stöckigt, J.; Antonchick, A. P.; Wu, F.; Waldmann, H. *Angew. Chem., Int. Ed.* **2011**, 50, 8558–8564. (b) Chrzanowska, M.; Rowadowska, M. D. *Chem. Rev.* **2004**, 104, 3341–3370. (c) Cox, E. D.; Cook, J. M. *Chem. Rev.* **1995**, 95, 1797–1842.
- (5) (a) Itoh, T.; Yokoya, M.; Miyauchi, K.; Nagata, K.; Ohsawa, A. *Org. Lett.* **2003**, 5, 4301–4304. (b) Itoh, T.; Yokoya, M.; Miyauchi, K.; Nagata, K.; Ohsawa, A. *Org. Lett.* **2006**, 8, 1533–1535. (c) Nagata, K.; Ishikawa, H.; Tanaka, A.; Miyazaki, M.; Kanemitsu, T.; Itoh, T. *Heterocycles* **2010**, 81, 1791–1798.
- (6) Proline catalysis of the reaction of imine **2a** with three different enones has been reported (ref. 5). High catalyst loadings (30–50 mol%) and large excess of enone (30 equiv relative to imine in two of the reported examples) were required to achieve acceptable yields of cyclization product.
- (7) For organocatalytic cycloaddition reactions of enones to acyclic imines, see: (a) Sundén, H.; Ibrahim, I.; Eriksson, L.; Córdova, A. *Angew. Chem., Int. Ed.* **2005**, 44, 4877–4880. (b) Rueping, M.; Azap, C. *Angew. Chem., Int. Ed.* **2006**, 45, 7832–7835. (c) Liu, H.; Cun, L.-F.; Mi, A.-Q.; Jiang, Y.-Z.; Gong, L.-Z. *Org. Lett.* **2006**, 8, 6023–6026. (d) Aznar, F.; Garcia, A.-B.; Quiñones, N.; Cabal, M.-P. *Synthesis* **2008**, 479–484. (e) Yang, H.; Carter, R. G. *J. Org. Chem.* **2009**, 74, 5151–5156. (f) Quiñones, N.; Aznar, F.; Cabal, M.-P. *Heterocycles* **2009**, 2963–2978. Dynamic resolution of benzoquinolizidine derivatives via crystallization has been carried out successfully in four instances. See: (g) Openshaw, H. T.; Whittaker, N. *J. Chem. Soc.* **1963**, 1449–1460. (h) Szántay, C.; Töke, L.; Honty, K.; Kalas, G. *J. Org. Chem.* **1967**, 32, 423–427. (i) Clark, R. D.; Kern, J. R.; Kurz, L. J.; Nelson, J. T. *Heterocycles* **1990**, 31, 353–366.
- (8) (a) Vachal, P.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2002**, 124, 10012–10014. (b) Zuend, S. J.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2007**, 129, 15872–15883. (c) Fang, Y.-Q.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2008**, 130, 5660–5661.
- (9) For reviews of thiourea anion-binding catalysis, see: (a) Zhang, Z.; Schreiner, P. R. *Chem. Soc. Rev.* **2009**, 38, 1187–1198. (b) Phipps, R. J.; Hamilton, G. L.; Toste, F. D. *Nature Chem.* **2012**, 4, 603–614. (c) Brak, K.; Jacobsen, E. N. *Angew. Chem., Int. Ed.* **2013**, 52, 534–561. See also: (d) Raheem, I. T.; Thiara, P. S.; Peterson, E. A.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2007**, 129, 13404–13405. (e) Zuend, S. J.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2009**, 131, 15358–15374. (f) Xu, H.; Zuend, S. J.; Woll, M.; Tao, Y.; Jacobsen, E. N. *Science* **2010**, 327, 986–990.
- (10) For selected examples of enantioselective organocatalysis with primary amines, see: (a) d’Angelo, J.; Desmaële, D.; Dumas, F.; Guingant, A. *Tetrahedron: Asymmetry* **1992**, 3, 459–505. (b) Tsogoeva, S. B.; Wei, S. *Chem. Commun.* **2006**, 1451–1453. (c) Huang, H.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2006**, 128, 7170–7171. (d) Lalonde, M. P.; Chen, Y.; Jacobsen, E. N. *Angew. Chem., Int. Ed.* **2006**, 45, 6366–6370. (e) Ramasastry, S. S. V.; Zhang, H.; Tanaka, F.; Barbas, C. F. III. *J. Am. Chem. Soc.* **2007**, 129, 288–289. (f) Bartoli, G.; Bosco, M.; Carlone, A.; Pescioli, F.; Sambri, L.; Melchiorre, P. *Org. Lett.* **2007**, 9, 1403–1405. (g) Chen, W.; Du, W.; Duan, Y.-Z.; Wu, Y.; Yang, S.-Y.; Chen, Y.-C. *Angew. Chem., Int. Ed.* **2007**, 46, 7667–7670. (h) Singh, R. P.; Bartelson, K.; Wang, Y.; Su, H.; Lu, X.; Deng, L. *J. Am. Chem. Soc.* **2008**, 130, 2422–2423. (i) Erkkilä, A.; Pihko, P. M.; Clarke, M.-R. *Adv. Synth. Catal.* **2007**, 349, 802–806. (j) Wang, X.; Reisinger, C. M.; List, B. *J. Am. Chem. Soc.* **2008**, 130, 6070–6071. (k) Brown, A. R.; Kuo, W.-H.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2010**, 132, 9286–9288. (l) Burns, N. Z.; Witten, M. R.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2011**, 133, 14578–14581.
- (11) **2a** was obtained from tryptamine in 75% overall yield over three steps. See Supporting Information.
- (12) See Supporting Information for an expanded description of the catalyst and reaction optimization studies.
- (13) This type of anion-binding catalysis has been invoked in other thiourea-promoted reactions of imines. See: Klausen, R. S.; Jacobsen, E. N. *Org. Lett.* **2009**, 11, 887–890.
- (14) The excess enone can be recovered quantitatively from the reaction mixture. See Supporting Information (Scheme S8) for a product inhibition study.
- (15) Proline was found to be far inferior as a catalyst for FADA reactions involving dihydroisoquinolines. For example, reaction of imine **6d** with enone **3j** (entry 4 in Table 3) with 30 mol% proline

1  
2 afforded a 1.3:1 mixture of **7dj** and **8dj** in 66% and 46% ee, respec-  
3 tively.

4 (16) These conditions were chosen to mimic conditions toward the  
5 completion of the reaction, where there is one equivalent of unreacted  
6 enone remaining.

7 (17) The ee of the major diastereomer was 99% whereas that of the  
8 minor was 98.5%. Virtually no epimerization occurred when the anal-  
9 ogous experiment was conducted with catalyst **1e**.

10 (18) Virtually no epimerization is observed in the absence of cata-  
11 lyst.

(19) FADA adduct **4aa** was recovered quantitatively, and with a  
slight diminishment of dr (>99:1 to 99:1).

(20) The absolute configuration of **4ae** and **5ae**, determined via X-  
ray crystal structure analysis, confirms that epimerization occurs at  
C4.

(21) Girling, P. R.; Kiyoi, T.; Whiting, A. *Org. Biomol. Chem.*  
**2011**, *9*, 3105–3121.

