

This article was downloaded by: [Florida State University]

On: 15 March 2013, At: 08:38

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Asian Natural Products Research

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/ganp20>

Ethylenediamine: an effective reagent for deacetylation of natural products

Ramasamy Mohankumar ^a, Kaliappan Ilango ^b, Vichangal Pridiuldi Santhanakrishnan ^a, Viswanathan Radhakrishnan ^a & Srinivasan Narasimhan ^a

^a Asthagiri Herbal Research Foundation, 14/1 Second Main Road, Jaya Nagar, Tambaram Sanatorium, Chennai, 600 047, Tamilnadu, India

^b Department of Pharmaceutical Chemistry, SRM College of Pharmacy, Kattankulathur, Kancheepuram, 603 203, Tamilnadu, India

Version of record first published: 28 Sep 2010.

To cite this article: Ramasamy Mohankumar, Kaliappan Ilango, Vichangal Pridiuldi Santhanakrishnan, Viswanathan Radhakrishnan & Srinivasan Narasimhan (2010): Ethylenediamine: an effective reagent for deacetylation of natural products, *Journal of Asian Natural Products Research*, 12:10, 851-858

To link to this article: <http://dx.doi.org/10.1080/10286020.2010.507545>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

ORIGINAL ARTICLE

Ethylenediamine: an effective reagent for deacetylation of natural products

Ramasamy Mohankumar^a, Kaliappan Ilango^b, Vichangal Pridiuldi Santhanakrishnan^a,
Viswanathan Radhakrishnan^a and Srinivasan Narasimhan^{a*}

^aAsthagiri Herbal Research Foundation, 14/1 Second Main Road, Jaya Nagar, Tambaram Sanatorium, Chennai 600 047, Tamilnadu, India; ^bDepartment of Pharmaceutical Chemistry, SRM College of Pharmacy, Kattankulathur, Kancheepuram 603 203, Tamilnadu, India

(Received 7 January 2010; final version received 8 July 2010)

The use of ethylenediamine in methanol is described for the selective cleavage of the acetate group in nimbin (**1**) to 6-deacetyl nimbin (**1a**) under microwave irradiation. This method enables to deacetylate without affecting other functional groups such as α,β -unsaturated ketone, ester, ether, etc. in certain tetranortriterpenoids and other acetate-containing natural compounds.

Keywords: forskolin; ethylenediamine; nimbin; microwave

1. Introduction

Semi-synthetic modification of natural products has gathered momentum in recent times owing to its potential in generating molecules with promising biological activity. Modification of a given structural entity is best achieved by functional group transformation or by introduction of moieties, which can enhance or alter the activity.


2. Results and discussion

During our ongoing investigations to prepare nitrogen analogs of some selected diterpenes and tetranortriterpenoids, we employed ethylenediamine in methanol to produce an imino product with forskolin [1], a compound known to exhibit excellent vasodilating and cardio-stimulating properties.

The NMR spectrum of the product revealed the disappearance of the C=O

resonance at 170 ppm (acetate carbonyl), indicating hydrolysis of a 7 β -acetoxy moiety. The carbonyl resonance at 203 ppm confirmed the presence of C-11 carbonyl in concurrence with the above-mentioned observation. This prompted us to evaluate the use of ethylenediamine to produce deacetyl products from polyfunctional natural products. The available agents such as NaOH and Na₂CO₃ in alcohol resulted in the formation of deesterified products along with the deacetylated product in nimbin. The Lewis acids would cleave the epoxide moiety [2] and the use of the other reagents like hydrazine hydrate may result in the cleavage of α,β -epoxy ketones [3]. Ethylenediamine seems to be a superior reagent for deacetylation in comparison with NH₃ in MeOH [4], exclusively yielding one product irrespective of the quantity of amine used. Hence, an attempt has been made to deacetylate

*Corresponding author. Email: asthagiri.herbal@gmail.com



Scheme 1. Conversion of nimbin (**1**) to deacetyl nimbin (**1a**).

nimbin using ethylenediamine resulting in deacetyl nimbin (Scheme 1) in 75% yield within 7 min under microwave [5,6] irradiation condition without any side products. The compound was then purified by column chromatography.

It has been observed that this reagent is mild and selectively cleaves the acetate group amid other functional groups. We have demonstrated (Table 1) the selective cleavage of acetate in certain tetranortriterpenoids such as nimbin [7,8], nimonol, salannin, and azadiradione [9–11] with multiple functional groups, and therefore this method has advantages over other methods.

The basicity of ethylenediamine is exploited in deacetylation without affecting other functional groups. The ¹H and ¹³C NMR spectra of this compound showed a similar characteristic to that of nimbin, except for the acetate group at C-7. It was clearly evident from the disappearance of the carbonyl group of acetate at 170.3 ppm and the methyl group signal at 20.8 ppm. The formation of the hydroxyl group at C-7 was confirmed by the appearance of the signal at 86.9 ppm, which in turn showed the connectivity to the proton at 3.76 ppm. The assigned structure for the product obtained is 6-deacetyl nimbin, which is in agreement with the mass 498 [M]⁺.

In summary, we have developed a simple, selective, and mild procedure for the cleavage of the acetate group in natural products, which will have a wide scope in

view of the usefulness of ethylenediamine [12,13] in organic synthesis.

3. Experimental

3.1 Procedure

Various tetranortriterpenoids such as nimbin, salannin, azadiradione, nimonol, etc. were obtained in the pure form after being subjected to repeated purification by column chromatography from the oil, fruit coat, and leaves of *Azadirachta indica*. Similarly, forskolin was obtained from *Coleus forskohlii* and cedrelone from the barks of *Toona ciliata*. The β-amyrin and pregnenolone were obtained from SD Fine Chemicals, Worli, Mumbai, India.

NMR spectra were recorded on a Bruker 200 MHz instrument using TMS as the internal reference for both ¹H and ¹³C NMR experiments. CDCl₃ was used as the solvent. Chemical shifts are given in terms of parts per million (δ scale). MS analyses were recorded on a Shimadzu QP 5000 instrument. IR spectra (cm⁻¹) were recorded on a Perkin-Elmer RX1 FT-IR spectrophotometer. Precoated thin layer chromatography plates (E-Merck, Darmstadt, Germany; Kieselgel 60 F254, 0.2 mm thickness, coated on aluminum sheets) were used. Column chromatography was performed using silica gel (60–120 mesh and 230–400 mesh).

3.2 Deacetylation of nimbin

Nimbin (108 mg, 5 mmol) was added to a solution of ethylenediamine (2 ml in 2 ml

Table 1. Deacetylation of natural products with ethylenediamine under modified microwave irradiation.

















Entry	Substrate	Product	Time (min)	Isolated yield (%)
1	 <p>Nimbin (1)</p>	 <p>Deacetyl nimbin (1a)</p>	7	75
2	 <p>Nimonol</p>	 <p>Deacetyl nimonol</p>	6	88
3	 <p>Salannin</p>	 <p>Deacetyl salannin</p>	12	76
4	 <p>Azadiradione</p>	 <p>Deacetyl azadiradione</p>	10	78
5	 <p>Cedrelone acetate</p>	 <p>Cedrelone</p>	5	82

Table 1 – continued

Entry	Substrate	Product	Time (min)	Isolated yield (%)
6	 <p>Forskolin</p>	 <p>Deacetyl forskolin</p>	5	80
7	 <p>Pregnenolone acetate</p>	 <p>Pregnenolone</p>	4	90
8	 <p>Amyrin acetate</p>	 <p>Amyrin</p>	4	94

Note: Yields were calculated after purification by column chromatography.

of methanol) and irradiated under microwave with stirring for 7 min. The reaction was monitored by TLC (70:30; ethyl acetate–hexane). The solvent was removed under reduced pressure. The reaction mixture was poured into ice-cold water and extracted with ethyl acetate to afford a white powder. The yield obtained after purification by column chromatography was 81 mg (75%). A series of deacetyl products are given in Table 1. The selected ^1H and ^{13}C NMR spectral data of both the starting material and the product are shown in Tables 2–6.

3.2.1 Deacetyl nimbin

$\text{C}_{28}\text{H}_{34}\text{O}_8$; mp 208°C; UV λ_{max} (MeOH): 209 nm; IR (KBr) ν_{max} : 3574, 3640, 2940, 2855, 1455, 1390, 1370, 1355, 1275, 1250, 1140, 1112, 1090, 1060, 1028, 878, 783, 744 cm^{-1} ; ^1H NMR (δ , ppm): 7.32 (m, 1H), 7.24 (m, 1H), 6.41 (d, 1H, $J = 10.1$ Hz), 6.33 (m, 1H), 5.84 (d, 1H, $J = 10.1$ Hz), 5.55 (m, 1H), 4.01 (d, 1H, $J = 3.3$ Hz), 3.91 (dd, 1H, $J = 11.4$, 3.3 Hz), 3.70 (s, 3H), 3.65 (s, 3H), 3.60 (m, 1H), 3.40 (d, 1H, $J = 11.7$ Hz), 2.92 (m, 1H), 2.77 (m, 1H), 2.25 (m, 1H), 2.20 (m, 1H), 2.06 (m, 1H), 1.21–1.69

Table 2. The selected ^1H and ^{13}C NMR spectral data of nimbin (**1**) and 6-deacetyl nimbin (**1a**).

Carbon no.	^1H NMR (δ ppm)		^{13}C NMR (δ ppm)	
	1	1a	1	1a
5	3.70 (m, 1H)	3.40 (d, 1H, $J = 11.7$ Hz)	41.5	43.6
6	5.25 (dd, 1H, $J = 12.3, 3.0$ Hz)	3.91 (dd, 1H, $J = 11.7, 3.3$ Hz)	68.6	66.1
7	4.05 (d, 1H, $J = 3.0$ Hz)	4.01 (d, 1H, $J = 3.3$ Hz)	84.5	86.9
8	–	–	47.1	47.3
9	2.85 (dd, 1H, $J = 5.5, 3.0$ Hz)	2.77 (m, 1H)	38.5	39.0
10	–	–	47.9	47.4

Table 3. The selected ^1H and ^{13}C NMR spectral data of nimonol (**2**) and 7-deacetyl nimonol (**2a**).

Carbon no.	^1H NMR (δ ppm)		^{13}C NMR (δ ppm)	
	2	2a	2	2a
5	2.23 (m, 1H)	2.45 (m, 1H)	49.9	47.2
6	4.37 (m, 1H)	4.22 (m, 1H)	68.1	68.4
7	5.37 (m, 1H)	3.49 (m, 1H)	79.1	75.9
8	–	–	45.5	44.4
9	2.44 (m, 1H)	2.87 (m, 1H)	36.9	35.6
10	–	–	43.3	45.6

Table 4. The selected ^1H and ^{13}C NMR spectral data of salannin (**3**) and 3-deacetyl salannin (**3a**).

Carbon no.	^1H NMR (δ ppm)		^{13}C NMR (δ ppm)	
	3	3a	3	3a
1	4.78 (t, 1H, $J = 2.7$ Hz)	5.30 (m, 1H)	71.3	72.8
2	2.04–2.37 (m, 2H)	2.05–2.35 (m, 2H)	27.5	30.5
3	4.96 (t, 1H, $J = 2.8$ Hz)	3.87 (m, 1H)	71.3	70.8
4	–	–	42.6	44.2
5	2.81 (d, 1H, $J = 12.5$ Hz)	2.63 (d, 1H, $J = 12.3$ Hz)	40.5	38.8
6	4.01 (dd, 1H, $J = 12.5, 3.3$ Hz)	4.08 (dd, 1H, $J = 12.3, 3.3$ Hz)	72.5	72.4

Table 5. The selected ^1H and ^{13}C NMR spectral data of azadiradione (**4**) and 7-deacetyl azadiradione (**4a**).

Carbon no.	^1H NMR (δ ppm)		^{13}C NMR (δ ppm)	
	4	4a	4	4a
5	2.20 (dd, 1H, $J = 3.7, 11.6$ Hz)	2.50 (dd, 1H, $J = 12.8$ Hz)	46.1	36.5
6	1.88 (m, 2H)	1.88 (m, 2H)	23.3	27.1
7	5.32 (t, 1H, $J = 2.7$ Hz)	3.99 (m, 1H)	73.8	71.7
8	–	–	44.4	44.2
9	2.50 (m, 1H)	2.55 (m, 1H)	38.1	44.5
10	–	–	39.9	40.1

Table 6. The selected ^1H and ^{13}C NMR spectral data of forskolin (**6**) and 7-deacetyl forskolin (**6a**).

Carbon no.	^1H NMR (δ ppm)		^{13}C NMR (δ ppm)	
	6	6a	6	6a
5	1.05 (m, 1H)	1.07 (m, 1H)	41.8	42.9
6	4.57 (m, 1H)	4.48 (m, 1H)	74.0	70.5
7	5.48 (d, 1H, $J = 4.3$ Hz)	4.62 (m, 1H)	73.2	74.7
8	–	–	68.8	82.2
9	–	–	80.3	82.2
10	–	–	41.7	42.7

(br s, $4 \times \text{CH}_3$); ^{13}C NMR (δ , ppm): 202.9, 175.5, 173.7, 148.1, 146.9, 143.0, 139.0, 134.9, 126.8, 126.4, 110.4, 87.4, 86.9, 66.2, 53.0, 51.6, 49.6, 47.8, 47.5, 43.6, 47.3, 41.4, 39.0, 34.4. MS m/z : 498 $[\text{M}]^+$.

3.2.2 Deacetyl salannin

$\text{C}_{32}\text{H}_{42}\text{O}_8$; mp 213–216°C; UV λ_{max} (MeOH): 210 nm; IR (KBr) cm^{-1} : 3430, 2993, 1698, 1662, 1593, 1460, 1387, 1151, 1030, 752, 602; ^1H NMR (δ , ppm): 7.31 (m, 1H), 7.25 (m, 1H), 6.92 (m, 1H), 6.27 (m, 1H), 5.38 (m, 1H), 5.30 (m, 1H), 4.18 (d, 1H, $J = 3.3$ Hz), 4.08 (dd, 1H, $J = 12.3, 3.3$ Hz), 3.87 (m, 1H), 3.87 (m, 1H), 3.62 (m, 1H), 3.62 (m, 1H), 3.19 (s, 3H), 2.67 (m, 1H), 2.50–2.75 (d, 1H, $J = 4.5$ Hz), 2.05–2.35 (m, 2H), 2.05–2.35 (m, 1H), 2.05–2.35 (m, 1H), 2.05–2.35 (m, 2H), 1.85 (s, 3H), 0.96–1.69 (br s, $4 \times \text{CH}_3$); ^{13}C NMR (δ , ppm): 172.6, 166.3, 146.4, 142.8, 138.7, 138.1, 134.9, 128.6, 127.1, 110.6, 87.8, 85.8, 77.8, 72.8, 72.4, 70.8, 51.4, 49.3, 48.9, 44.2, 41.2, 40.9, 39.4, 38.8, 30.5, 30.5, 19.8, 16.8, 15.1, 14.5, 13.1, 12.1. MS m/z : 554 $[\text{M}]^+$.

3.2.3 Deacetyl azadiradione

$\text{C}_{26}\text{H}_{32}\text{O}_4$; mp 208°C; UV λ_{max} (MeOH) 237 nm; IR (KBr) cm^{-1} : 3409, 2923, 2826, 1728, 1649, 1435, 1387, 1259, 1155, 1037, 983, 786, 732, 599; ^1H NMR (δ , ppm): 7.43 (m, 1H), 7.34 (m, 1H), 7.10 (d, 1H, $J = 10.2$ Hz), 6.10 (s, 1H), 6.06 (m, 1H), 5.89 (d, 1H, $J = 10.2$ Hz), 3.99

(m, 1H), 3.50 (s, 1H), 2.55 (m, 1H), 2.50 (dd, 1H, $J = 12.8$ Hz), 2.08 (m, 1H), 2.04 (m, 1H), 1.88 (m, 2H), 1.86 (m, 1H), 1.84 (m, 1H), 1.42 (s, 3H), 1.27 (s, 3H), 1.12 (s, 3H), 1.12 (s, 3H), 0.96 (s, 3H); ^{13}C NMR (δ , ppm): 204.6, 201.8, 193.7, 157.1, 142.8, 141.7, 125.9, 123.4, 118.4, 111.1, 71.7, 60.9, 48.3, 44.6, 44.2, 40.2, 36.6, 30.5, 27.1, 26.4, 26.0, 25.4, 21.44, 19.1, 15.7. MS m/z : 408 $[\text{M}]^+$.

3.2.4 Deacetyl nimonol

$\text{C}_{26}\text{H}_{34}\text{O}_4$; mp 194–198°C; UV λ_{max} (MeOH): 223 nm; IR ν_{max} (MeOH) cm^{-1} : 3500, 2900, 2875, 1658, 1460, 1382, 1224, 1157, 1103, 1028, 931, 794, 597, 466; ^1H NMR (δ , ppm): 7.38 (m, 1H), 7.25 (m, 1H), 7.07 (d, 1H, $J = 10.2$ Hz), 6.29 (m, 1H), 5.86 (d, 1H, $J = 10.2$ Hz), 5.58 (d, 1H, $J = 10.2$ Hz), 4.22 (m, 1H), 3.49 (m, 1H), 2.87 (m, 1H), 2.45 (m, 1H), 2.45 (m, 1H), 2.16–2.37 (m, 2H), 2.16–2.37 (m, 2H), 1.95 (m, 2H), 1.11–1.42 (s, 3H), 1.11–1.42 (s, 3H), 1.11–1.42 (s, 3H); ^{13}C NMR (δ , ppm): 206.1, 161.0, 157.3, 142.7, 139.8, 126.2, 124.2, 120.0, 111.0, 76.0, 68.4, 51.6, 49.5, 47.2, 45.6, 44.4, 40.8, 35.6, 34.4, 32.3, 29.7, 26.9, 20.6, 20.3, 16.2. MS m/z : 410 $[\text{M}]^+$.

3.2.5 Cedrelone

$\text{C}_{26}\text{H}_{32}\text{O}_4$; mp 208°C; UV λ_{max} (MeOH): 237 nm; IR (KBr) cm^{-1} : 3392, 3118, 3033, 2933, 2866, 1676, 1618, 1448, 1384, 1340, 1251, 1153, 1060, 1031, 981, 947,

918, 877, 813, 779, 752, 715, 669, 634, 596; $^1\text{H NMR}$ (δ , ppm): 7.37 (m, 1H), 7.15 (m, 1H), 6.90 (d, 1H, $J = 9.7$ Hz), 6.44 (bs, D_2O exchangeable), 6.18 (m, 1H), 5.89 (d, 1H, $J = 9.7$ Hz), 3.80 (s, 1H), 2.30 (m, 1H), 2.75 (m, 1H), 2.67 (m, 1H), 2.07 (m, 1H), 1.94 (m, 1H), 1.85 (m, 1H), 1.69 (m, 1H), 0.75–1.57 (br s, $\times \text{CH}_3$); $^{13}\text{C NMR}$ (δ , ppm): 203.7, 197.9, 152.4, 143.0, 141.1, 139.3, 134.8, 127.2, 123.1, 110.6, 69.8, 55.0, 46.8, 48.5, 43.1, 42.1, 42.0, 41.7, 40.2, 35.1, 26.7, 23.9, 23.0, 21.1, 20.2, 19.4. MS m/z : 422 $[\text{M}]^+$.

3.2.6 Deacetyl forskolin

$\text{C}_{20}\text{H}_{32}\text{O}_6$; mp: 163–166°C; UV λ_{max} (MeOH): 210 nm; IR (KBr) cm^{-1} : 3419, 2939, 1716, 1562, 1402, 1247, 1103, 1060; $^1\text{H NMR}$ (δ , ppm): 6.12 (dd, 1H, $J = 17.4$, 10.7 Hz), 5.19 (d, 1H, $J = 17.3$ Hz), 4.98 (d, 1H, $J = 10.8$ Hz), 4.63 (m, 1H), 4.48 (m, 1H), 4.13 (m, 1H), 3.18 (d, 1H, $J = 17.2$ Hz), 2.50 (d, 1H, $J = 17.2$ Hz), 2.35 (m, 1H), 2.20 (m, 1H), 2.09 (m, 1H), 1.75 (m, 1H), 1.65 (s, 3H), 1.43 (s, 3H), 1.41 (s, 3H), 1.27 (s, 3H), 1.06 (s, 3H); $^{13}\text{C NMR}$ (δ , ppm): 205.8, 146.5, 110.5, 82.3, 82.2, 75.2, 74.8, 74.7, 70.5, 48.8, 48.8, 43.0, 42.8, 36.1, 34.3, 33.0, 30.8, 24.2, 23.4, 20.0. MS m/z : 368 $[\text{M}]^+$.

3.2.7 Pregnenolone

$\text{C}_{21}\text{H}_{32}\text{O}_2$; mp 171–173°C; UV λ_{max} (MeOH): 206 nm; IR (KBr) cm^{-1} : 3506, 3352, 3028, 2933, 2835, 1683, 1461, 1433, 1357, 1292, 1224, 1191, 1168, 1134, 1097, 1053, 985, 950, 914, 840, 800, 596, 468; $^1\text{H NMR}$ (δ , ppm): 5.36 (m, 1H), 3.53 (m, 1H), 2.54 (t, 1H, $J = 9.0$ Hz), 2.20–2.35 (m, 2H), 2.20–2.35 (m, 2H), 2.13 (s, 3H), 2.06 (m, 1H), 2.04 (m, 2H), 1.40–1.75 (m, 2H), 1.40–1.75 (m, 2H), 1.40–1.75 (m, 1H), 1.40–1.75 (m, 1H), 1.40–1.75 (m, 2H), 1.40–1.75 (m, 1H), 1.00–1.30 (m, 2H), 1.0–1.30 (m, 2H), 1.01 (s, 3H), 0.63 (s, 3H); $^{13}\text{C NMR}$ (δ , ppm): 209.6, 140.8, 121.3, 71.6, 63.7, 56.9, 50.0, 44.0, 42.2,

38.8, 36.4, 36.5, 31.8, 31.7, 31.6, 31.5, 24.5, 22.8, 21.1, 19.4, 13.2. MS m/z : 316 $[\text{M}]^+$.

3.2.8 β -Amyrin

$\text{C}_{30}\text{H}_{50}\text{O}$; mp 196–198°C; UV λ_{max} (MeOH): 206 nm; IR (KBr) cm^{-1} : 3316, 2854, 1643, 1456, 1360, 1195, 1036, 988, 881, 824, 660; $^1\text{H NMR}$ (δ , ppm): 4.62 (m, 1H), 3.21 (m, 1H), 1.92–2.10 (m, 2H), 1.65–1.85 (4H, m), 1.40–1.55 (m, 12H), 0.90–1.10 (m, 4H), 1.03 (s, 3H), 0.99 (s, 3H), 0.96 (s, 2H), 0.94 (s, 3H), 0.91 (s, 3H), 0.82 (s, 3H), 0.79 (s, 3H), 0.76 (s, 3H); $^{13}\text{C NMR}$ (δ , ppm): 150.1, 124.4, 79.0, 55.2, 48.0, 48.0, 46.9, 41.5, 40.0, 38.7, 38.7, 37.1, 34.2, 33.3, 32.9, 32.6, 31.3, 28.1, 28.4, 28.0, 27.4, 26.1, 25.1, 23.5, 23.5, 18.9, 16.1, 15.7, 15.4. MS m/z : 426 $[\text{M}]^+$.

Acknowledgements

R.M.K. thanks DBT, V.P.S. and V.R.K. thank CSIR for the financial support and Spic Science Foundation for the Analytical Center Facility.

Supporting Information

Supporting information is available online.

References

- [1] S.V. Bhat, B.S. Bajqwa, H. Dornauer, and N.J.D. Scusa, *Tetrahedron Lett.* **18**, 1669 (1977).
- [2] S. Chandrasekhar, C.R. Reddy, B. Nagenra Babu, and G. Chandrashekar, *Tetrahedron Lett.* **43**, 3801 (2002).
- [3] A.R. Jorge, J.L. Salvador, A. Leitao, M.L.S. Melo, and J.R. Hanson, *Tetrahedron Lett.* **46**, 1067 (2005).
- [4] T. Nelson and E.S. Werstiuk, *Can. J. Chem.* **49**, 493 (1971).
- [5] E.J. Corey and W.C. Taylor, *J. Am. Chem. Soc.* **86**, 3881 (1964).
- [6] S. Narasimhan and S. Velmathi, *Synth. Comm.* **32**, 3791 (2002).
- [7] T.R. Govindachari, G. Suresh, and G. Geetha, *J. Liq. Chromatogr.* **18**, 3465 (1995).
- [8] M. Harris, R. Henderson, R. McCrindle, K.H. Overton, and D.W. Turner, *Tetrahedron* **24**, 1517 (1968).
- [9] A. Chatterjee and P.S. Chandra, *Treatise on Indian Medicinal Plants*, 3rd ed.

- (National Institute of Science Communication, New Delhi, India, 1997).
- [10] L.V. Asoklar, K.K. Kakkar, and O.J. Chakre, *Second Supplement to Glossary of Indian Medicinal Plants with Active Principles*, 1st ed. (Publication and Information Directorate, CSIR, New Delhi, India, 1992).
- [11] K.R. Kiritikar and B.D. Basu, *Indian Medicinal Plants*, 1st ed. (Orient Longman Pvt. Ltd., Chennai, India, 1994).
- [12] A.K. Sen, G. Singh, K. Singh, R.K. Noren, R.N. Handa, and S.N. Dubey, *Indian J. Chem.* **36A**, 891 (1997).
- [13] P.J. Bhattacharya, *J. Indian Chem. Soc.* **59**, 505 (1982).