

Article

## Synthesis of Functionalized Indolines and Dihydrobenzofurans by Iron and Copper Catalyzed Aryl C-N and C-O Bond Formation

Martyn C. Henry, Hans Martin Senn, and Andrew Sutherland

*J. Org. Chem.*, **Just Accepted Manuscript** • DOI: 10.1021/acs.joc.8b02888 • Publication Date (Web): 06 Dec 2018

Downloaded from <http://pubs.acs.org> on December 7, 2018

### Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

# Synthesis of Functionalized Indolines and Dihydrobenzofurans by Iron and Copper Catalyzed Aryl C-N and C-O Bond Formation

*Martyn C. Henry, Hans Martin Senn and Andrew Sutherland\**


WestCHEM, School of Chemistry, The Joseph Black Building, University of Glasgow,

Glasgow G12 8QQ, United Kingdom.

Andrew.Sutherland@glasgow.ac.uk

**RECEIVED DATE (to be automatically inserted after your manuscript is accepted if required  
according to the journal that you are submitting your paper to)**

## Table of Contents Graphic:



**Abstract:** A simple and effective one-pot, two-step intramolecular aryl C-N and C-O bond forming process for the preparation of a wide range of benzo-fused heterocyclic scaffolds using iron and copper catalysis is described. Activated aryl rings were subjected to a highly regioselective, iron(III) triflimide-catalyzed iodination, followed by a copper(I)-catalyzed intramolecular *N*- or *O*-arylation step leading to indolines, dihydrobenzofurans and six-membered analogues. The general applicability and functional group tolerance of this method was exemplified by the total synthesis of the neolignan natural product, (+)-obtusafuran. DFT calculations using Fukui functions were also performed, providing a molecular orbital rationale for the highly regioselective arene iodination process.

## INTRODUCTION


Indoline and dihydrobenzofuran scaffolds are privileged structures, represented in a wide range of natural products and pharmaceutically important agents.<sup>1</sup> For this reason, considerable efforts have focused on the discovery of efficient methods for the preparation of these heterocycles.<sup>2</sup> A commonly used approach for the synthesis of indolines and dihydrobenzofurans is the Buchwald-Hartwig or Ullmann-type C-N and C-O bond forming process of pre-functionalized, halogenated phenylethylamines and phenylethylalcohols (Figure 1a).<sup>3-6</sup> As well as 5-membered rings, this strategy is highly effective for the preparation of 6-membered benzo-fused heterocyclic systems and has been used for the preparation of a wide range of natural products such as (+)-isatisine A,<sup>4g</sup> corsifuran A<sup>5b</sup> and a number of quinoline-containing alkaloids.<sup>3b</sup>

An alternative approach for preparing these ring systems has been developed more recently involving transition metal-catalyzed aryl C-H activation and intramolecular cross-coupling with N-H or O-H bonds.<sup>7-8</sup> Pioneering work by the Yu group showed that triflimide-protected 2-phenylethylamines and Pd(II)/Cu(I) catalysis could be used for the one-pot preparation of indolines via a tandem C-H bond iodination-amination sequence.<sup>8a</sup> The palladium-catalyzed intramolecular aryl C(sp<sup>2</sup>)-H amination process was improved using *N*-chelating groups and the use of oxidizing agents such as hypervalent iodonium salts.<sup>8-10</sup> Among the range of *N*-protected amides used to direct the palladium-catalyzed functionalization of aryl C-H bonds, the Zhao group demonstrated the highly effective use of a *N,O*-bidentate oxalyl amide (Figure 1b).<sup>8g</sup>


While this strategy has also been used to prepare dihydrobenzofurans from phenylethylalcohols, the oxidizing conditions can be problematic for substrates bearing primary and secondary alcohols.<sup>11</sup> More recently, Zakarian and co-workers reported a mechanistically distinct approach for the preparation of dihydrobenzofurans (Figure 1c).<sup>12</sup> A one-pot intramolecular aryl C-O bond forming process was achieved by formation of non-symmetrical diaryliodonium salts by oxidation of electron-rich 2-

phenylethylalcohols, followed by a copper-catalyzed C-O bond forming process. A key feature of this method was the room temperature conditions for copper-catalyzed cyclization.


a) Buchwald-Hartwig or Ullmann-type coupling of aryl halides




b) Palladium-catalyzed, auxiliary directed indoline synthesis



c) Dihydrobenzofuran synthesis via arylidonium intermediates



d) This work: synthesis of *N*- and *O*-heterocycles using iron and copper catalysis



**Figure 1.** Synthesis of indolines and dihydrobenzofuran scaffolds.

While these methods provide an attractive entry to these ring systems, many of the approaches have been specifically developed for either C-N or C-O bond formation or for the preparation of a particular ring size. We were interested in developing a new approach for intramolecular C-N or C-O bond formation that would avoid a pre-functionalization step, precious transition metals, strong oxidizing conditions and could be used for the general preparation of both five- and six-membered heterocyclic systems. Herein, we describe a one-pot intramolecular C-N and C-O bond forming process that utilizes a highly


1 regioselective iron-catalyzed iodination for initial arene activation, followed by a copper-catalyzed C-N  
2 and C-O cyclization (Figure 1d). As well as providing an electronic rationale for the high regioselectivity  
3 of the iron-catalyzed halogenation reaction, we show the general application of this process for the  
4 preparation of a wide range of ring systems and as the key step for the total synthesis of the natural  
5 product, (+)-obtusafuran.  
6  
7  
8  
9  
10

## 11 12 13 14 RESULTS AND DISCUSSION

15  
16 Previously, we have shown that the combination of iron(III) chloride and the inexpensive ionic liquid,  
17 [BMIM]NTf<sub>2</sub> results in the formation of iron triflimide, which can be used as a super Lewis acid catalyst<sup>13</sup>  
18 to activate *N*-halosuccinimides for the fast and efficient regioselective halogenation of aromatic  
19 compounds.<sup>14</sup> In this current study, this transformation was investigated for the regioselective iodination  
20 of a new class of substrate, *N*-protected 2-phenylethylamines (Scheme 1). The initial aim was to evaluate  
21 the 3-methoxy substituent as a directing group for selective *para*-iodination and assess if the resulting  
22 activated aryl intermediate could undergo a copper(I)-catalyzed *N*-arylation reaction for the one-pot  
23 synthesis of indolines. Using standard conditions for halogenation with iron(III) chloride (2.5 mol%) and  
24 [BMIM]NTf<sub>2</sub> (7.5 mol%), the iodination of *N*-benzoyl protected **1a** by *N*-iodosuccinimide (NIS) was  
25 complete in 5 hours at 40 °C.<sup>14a,15</sup> Analysis of the crude reaction mixture by <sup>1</sup>H NMR spectroscopy  
26 showed the formation of the *para*-iodinated regioisomer as the sole product. The aryl ring of **1a** can  
27 undergo iodination with NIS, without the iron(III) triflimide catalyst, however, under the same conditions,  
28 full conversion was only achieved after 22 hours giving a 10:1 mixture of *para* and *ortho*-regioisomers.  
29  
30 The regioselective iron(III) triflimide-catalyzed activation of **1a** was then performed in combination with  
31 a Cu(I)-catalyzed *N*-arylation for the one-pot synthesis of indoline **2a**. Using copper iodide (10 mol%)  
32 and DMEDA (20 mol%) during the C-N bond forming step gave *N*-benzoyl-protected indoline **2a** in 79%  
33 overall yield. It should be noted that when the synthesis of **2a** was done by performing each step  
34 separately, the overall yield (59%) was significantly lower than that for the one-pot process. With the  
35 success of the one-pot synthesis of **2a**, a range of *N*-protecting groups were explored to evaluate the most  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

suitable nucleophile for the Cu(I)-catalyzed *N*-arylation. While *N*-Cbz and *N*-Boc carbamate protected indolines **2c** and **2d** were prepared in good yields, the most efficient one-pot processes involved *N*-acetamide or *N*-sulfonamide protected compounds. In particular, one-pot activation and cyclization of *N*-tosyl phenylethylamine **1f** gave indoline **2f** in 93% yield.


### Scheme 1. One-Pot Activation and Cyclization of *N*-Protected 3-Methoxyphenylethylamines **1**<sup>a</sup>



<sup>a</sup>Isolated yields are shown.

Having identified optimized conditions and the most efficient *N*-nucleophile, the scope of the one-pot activation and cyclization process was explored for the preparation of indolines (Scheme 2). Using a range of *N*-tosyl ethylamine substituted anisoles, anilines and acetanilides gave the corresponding indolines **2f**–**2o** as single regioisomers, in 43–93% yields.<sup>16</sup> As expected, substrates with multiple activating groups were converted to the indolines in shorter overall reaction times. Interestingly, phenylethylamine **1o** containing *N*-acetyl and chlorine substituents failed to undergo the iron(III)-catalyzed iodination even at 70 °C. Instead, activation was achieved by bromination using *N*-bromosuccinimide (NBS) at 40 °C. Completion of the one-pot process gave indoline **2o** in 43% yield. Access to other benzannulated heterocycles was also achieved using the one-pot process. Iron(III)-catalyzed iodination and copper(I)-catalyzed cyclization of 3-methoxyphenylacetamide (**1p**) gave 2-oxindole **2p** in 65% yield, while an *N*-tosyl propylamine substituted anisole led to the corresponding tetrahydroquinoline **2q** in 85% yield.<sup>17</sup>

**Scheme 2. One-Pot Activation and Cyclization for the Synthesis of *N*-Heterocycles<sup>a</sup>**




<sup>a</sup>Isolated yields are shown. <sup>b</sup>The one-pot process was performed via the bromide intermediate using NBS for the activation step.

The use of palladium-catalyzed C-H activation in the presence of phenyliodonium diacetate, followed by C-O cyclization for the synthesis of dihydrobenzofurans can be done using tertiary,<sup>11a</sup> or secondary benzylic alcohol nucleophiles.<sup>11b</sup> However, the general use of substrates bearing primary or secondary hydroxy groups for this process are problematic due to competitive oxidation.<sup>12</sup> Mild oxidative conditions such as reported by the Zakarian group are required for general access to dihydrobenzofurans (Figure



1c).<sup>12</sup> Following the successful application of the one-pot iron(III)-catalyzed activation and copper(I)-catalyzed cyclization for the synthesis of *N*-heterocycles, we were interested to discover whether the mild oxidative conditions for this two-step process could also be applied for the preparation of dihydrobenzofurans. The transformation was initially attempted using 3-methoxyphenylethan-2'-ol (**3a**) for the synthesis of 2,3-dihydro-5-methoxybenzofuran (**4a**) (Scheme 3). While the standard iodination conditions could be used, a slightly higher temperature (150 °C) was required for complete conversion to the cyclized dihydrobenzofuran. This gave **4a** in 65% yield. The scope of this process was then explored using a range of substrates with various aryl activating groups and primary, secondary or tertiary alcohol nucleophiles (**3a–3i**). In general, the one-pot processes were performed under the standard conditions developed for the *N*-heterocycles, giving the corresponding dihydrobenzofurans as single regioisomers in 56–72% yields. It should be noted that while other one-pot methods have had problems with over-oxidation and the generation of benzofuran by-products, especially with electron-rich substrates,<sup>11b</sup> analogous dihydrobenzofurans produced in this study (e.g. **4b–4d**), were formed cleanly as single products. The only limitation was found during the synthesis of natural product, corsifuran A (**4g**).<sup>18</sup> Substrate **3g**, which contains two activated aryl rings gave a mixture of products during the iodination step, resulting in the isolation of corsifuran A (**4g**) in only 29% yield.<sup>19</sup> However, using secondary benzylic alcohols with less electron-rich aryl rings (e.g. **3h** and **3i**) allowed selective iodination of the 3-methoxyphenyl moiety resulting in the synthesis of dihydrobenzofurans **4h** and **4i** in 64% and 63% yields, respectively. This approach was also effective for the one-pot synthesis of dihydrobenzopyrans. Application of (3-methoxyphenyl)propan-3'-ol (**3j**) to the one-pot iron(III)-catalyzed activation and copper(I)-catalyzed cyclization gave dihydrobenzopyran **4j** as the sole product in 57% yield. Similar results were also obtained for dihydrobenzopyrans **4k** and **4l**.

**Scheme 3. One-Pot Activation and Cyclization for the Synthesis of *O*-Heterocycles<sup>a</sup>**




<sup>a</sup>Isolated yields are shown.

To further explore the functional group tolerance of the one-pot process and illustrate its application for natural product synthesis, the method was investigated as a key step for the total synthesis of (+)-obtusafuran (**10**). The neolignan, (+)-obtusafuran was first isolated from the heartwood of *Dalbergia*

1 *retusa*<sup>20</sup> and more recently from several other *Dalbergia* species.<sup>21-23</sup> As well as possessing  
2 antiplasmodial activity,<sup>21</sup> (+)-obtusafuran has been shown to have anticarcinogenic activity as a potent  
3 inducer of the carcinogen-detoxifying enzyme, quinone reductase.<sup>24</sup> Racemic obtusafuran has been  
4 prepared by a thermal rearrangement of the neoflavanoid, obtusaquinol,<sup>25</sup> while the only asymmetric  
5 synthesis of (+)-obtusafuran was reported by Chen and Weisel, who used an enantioselective  
6 hydrogenation to produce a chiral alcohol that was then subjected to an S<sub>N</sub>Ar reaction to form the furan  
7 ring.<sup>26</sup> Our strategy involved the synthesis of  $\alpha$ -methyl phenyl ketone **7** (Scheme 4) and the application  
8 of this to a Merck-type enantioselective hydrogenation involving a base-mediated dynamic kinetic  
9 resolution process.<sup>27,28</sup> The resulting secondary benzylic alcohol **8** would then be used in the one-pot  
10 iron(III)-catalyzed iodination and copper(I)-catalyzed cyclization to complete the synthesis of the  
11 dihydrobenzofuran skeleton. Initially, Weinreb amide **6** was prepared in two steps from phenylacetic acid  
12 **5**, by coupling with *N,O*-dimethylhydroxylamine using EDCI and HOBt, followed by TBDMS protection  
13 of the phenol under standard conditions. Reaction of **6** with phenylmagnesium bromide and then  $\alpha$ -  
14 alkylation with LiHMDS and methyl iodide gave key intermediate **7** in good overall yield. This was then  
15 subjected to the Merck enantioselective hydrogenation using the commercially available Noyori-type  
16 chiral catalyst, RuCl<sub>2</sub>[(*S*)-DM-Segphos][(*S*)-DAIPEN].<sup>27-29</sup> On screening various conditions and catalyst  
17 loadings, the best results were achieved by hydrogenation at 10 bar of pressure, using 2 mol% of the  
18 Ru(II)-catalyst. This gave secondary alcohol **8** as a single diastereomer, in 95% enantiomeric excess and  
19 64% yield.<sup>30</sup> Our one-pot process was then investigated for the final key step. Activation of the aryl ring  
20 using the iron(III)-catalyzed iodination required a slightly higher temperature (50 °C) and longer reaction  
21 time (7 h) than the more simple substrates. Following this step, the standard conditions of the copper(I)-  
22 catalyzed cyclization were then used to complete the one-pot process, which gave dihydrobenzofuran **9**  
23 in 63% yield. Despite using a substrate with a highly activated aryl ring and a secondary alcohol, no by-  
24 products from over-iodination or oxidation were observed at either stage of the one-pot process. Finally,  
25 TBAF mediated removal of the silyl protecting group completed the eight-step synthesis of (+)-  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

obtusafuran (**10**) in 16% overall yield. The spectroscopic data and optical rotation of **10** were entirely consistent with literature data.<sup>20b,26</sup>

#### Scheme 4. Total Synthesis of (+)-Obtusafuran (**10**)<sup>a</sup>




<sup>a</sup>Isolated yields are shown.

Iron(III)-catalyzed activation of the *N*-protected 2-phenylethylamines and phenylethan-2'-ols gave the *para*-iodinated isomers as the sole product. As no reaction was observed at the other activated positions, including the most sterically accessible *ortho*-position, DFT calculations were used to explore electronic reasons for this reactivity.<sup>31</sup> The reactivities of different sites towards electrophilic or nucleophilic attack may be assessed using a computed descriptor such as partial (atomic) charge. In this study, the Hirshfeld


partitioning scheme was used.<sup>32</sup> The Hirshfeld charges calculated for the (unsubstituted) aromatic carbons of *N*-mesyl protected 2-phenylethylamine **1e** single out C-5 as the least preferred site for electrophilic attack, but cannot distinguish which of C-2, C-4, or C-6 would be the most preferred site (Table 1, entry 1). A more refined and powerful reactivity descriptor is provided by the Fukui functions.<sup>33,34</sup> The electrophilic Fukui function  $f^+(\mathbf{r})$  has more positive values at points in space where it is energetically favorable to remove electrons (see supporting information for background and derivations); that is,  $f^+(\mathbf{r})$  identifies sites favored for electrophilic attack. If the reactivity is entirely controlled by the frontier orbitals,  $f^+(\mathbf{r})$  is well approximated by the density of the HOMO. From Figure 2, it is evident that the most positive region of  $f^+(\mathbf{r})$ , and hence the most favorable site for electrophilic attack of **1e**, is located around C-6. More specifically, the  $p_z$  atomic orbital on C-6 makes the largest contribution to the HOMO. By contracting the continuous Fukui functions to distinct sites (e.g. atoms), “condensed” Fukui reactivity indices are obtained; a large (positive) electrophilicity index  $f^+$  indicates a favored site for electrophilic attack. Using frontier-orbital terminology,  $f^+$  for a particular atom can be identified with the contribution of that atom to the HOMO.  $f^+$  values for the aromatic carbons in **1e** are presented in Table 1 (entries 2 and 3). The quantitative reactivity analysis using atomic Fukui indices thus clearly identifies C-6 as the most preferred site for electrophilic attack in this case, in agreement with experiment. C-2 and C-4 have significantly diminished, nearly equal reactivity; C-5 is predicted to be least reactive. Further analysis can be performed using a “dual descriptor”  $\Delta f$ , which combines the separate electrophilic and nucleophilic Fukui functions into one descriptor.<sup>35</sup> More positive values of  $\Delta f$  indicates sites for nucleophilic attack; more negative values sites for electrophilic attack. As can be seen from Table 1 (entries 4 and 5), the analysis based on  $\Delta f$  values fully confirms the regioselectivity observed for activation of the substrates in this study.

**Table 1. Reactivity Descriptors Computed for the Aromatic Carbons of Methoxyphenylethylamine 1e.<sup>a</sup>**



entry	reactivity descriptors	C-2	C-4	C-5	C-6
1	$q_{\text{H}} / e$	-0.065	-0.071	-0.042	-0.065
2	$f(q_{\text{H}})$	0.082	0.078	0.065	0.140
3	$f(\text{HOMO})$	12	10	8	22
4	$\Delta f(q_{\text{H}})$	0.045	0.019	0.068	-0.088
5	$\Delta f(\text{HOMO})$	7	4	12	-15

$q_{\text{H}}$  is the Hirshfeld charge (in units of the elementary charge,  $e$ );  $f(q_{\text{H}})$  and  $\Delta f(q_{\text{H}})$  are the condensed electrophilic Fukui function and the condensed “dual descriptor”, respectively, calculated from Hirshfeld charges;  $f(\text{HOMO})$  and  $\Delta f(\text{HOMO})$  are the same descriptors but calculated from Hirshfeld-atomic contributions to the HOMO (expressed in %). <sup>a</sup>M06-2X/def2-TZVP/PCM(toluene).



**Figure 2.** Isosurface plots of: (A) the HOMO and (B) the electrophilic Fukui function  $f(r)$  for **1e**. The aminoalkyl part is omitted as its contribution to either function is negligible. Positive values are colored in orange, negative values in purple. Isosurface values: 0.006 a.u. for  $f(r)$ , 0.1 a.u. for the HOMO.

## CONCLUSIONS

In summary, a one-pot, two-step method involving iron(III)-catalyzed aryl ring activation and copper(I)-catalyzed C-N or C-O bond forming cyclization has been developed for the general synthesis of valuable *N*- and *O*-heterocyclic scaffolds. Following DFT calculations, that showed the molecular orbital basis for the highly regioselective halogenation step, the novel, one-pot method was applied to the efficient synthesis of indolines and dihydrobenzofurans, as well as six-membered analogues. This one-pot approach does not require pre-functionalization of the substrate as with the traditional Buchwald-Hartwig and Ullmann-type intramolecular couplings and unlike the established palladium-catalyzed dehydrogenative processes, this method has no issues with over-iodination or oxidation and could be applied to substrates with highly activated aryl ring systems and with primary and secondary alcohols. This was exemplified by the use of this one-pot process as the key step for the total synthesis of the neolignan natural product, (+)-obtusafuran. We expect this simple and effective approach to find utilization in the preparation of other heterocyclic scaffolds and for application in the synthesis of natural products and medicinal chemistry targets. Investigation of further applications of the one-pot process is currently underway.

## EXPERIMENTAL SECTION

All reagents and starting materials were obtained from commercial sources and used as received unless otherwise stated. Dry solvents were purified using a solvent purification system. Brine refers to a saturated solution of sodium chloride. All reactions were performed in oven-dried glassware under an atmosphere of argon unless otherwise stated. Flash column chromatography was carried out using silica gel (40–63  $\mu\text{m}$ ) and neutral aluminium oxide (50–200  $\mu\text{m}$ ). Aluminium-backed plates pre-coated with silica gel 60 (UV<sub>254</sub>) were used for thin layer chromatography and were visualized under ultraviolet light and by staining with KMnO<sub>4</sub> or ninhydrin. <sup>1</sup>H NMR spectra were recorded on an NMR spectrometer at 400 or 500 MHz and data are reported as follows: chemical shift in ppm relative to tetramethylsilane or the

solvent as the internal standard ( $\text{CDCl}_3$ ,  $\delta$  7.26 ppm), multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet or overlap of nonequivalent resonances, integration).  $^{13}\text{C}\{^1\text{H}\}$  NMR spectra were recorded on an NMR spectrometer at 101 or 126 MHz and data are reported as follows: chemical shift in ppm relative to tetramethylsilane or the solvent as internal standard ( $\text{CDCl}_3$ ,  $\delta$  77.0 ppm), multiplicity with respect to hydrogen (deduced from DEPT experiments, C, CH,  $\text{CH}_2$  or  $\text{CH}_3$ ). IR spectra were recorded on a FTIR spectrometer; wavenumbers are indicated in  $\text{cm}^{-1}$ . Mass spectra were recorded using electron impact or electrospray techniques. HRMS spectra were recorded using a dual-focusing magnetic analyzer mass spectrometer. Melting points are uncorrected. Optical rotations were determined as solutions irradiating with the sodium D line ( $\lambda = 589 \text{ nm}$ ) using a polarimeter.  $[\alpha]_{\text{D}}$  values are given in units  $10^{-1} \text{ deg cm}^2 \text{ g}^{-1}$ . Chiral HPLC methods were calibrated with the corresponding racemic mixtures.

**3-Methoxy-1-[(E)-2'-nitrovinyl]benzene.**<sup>36</sup> To a solution of *m*-anisaldehyde (0.890 mL, 7.30 mmol) in toluene (30 mL) was added nitromethane (2.00 mL, 37.0 mmol) and ammonium acetate (0.560 g, 7.30 mmol). The resulting solution was heated under reflux for 18 h. The reaction mixture was washed with water ( $2 \times 30 \text{ mL}$ ), followed by brine ( $2 \times 30 \text{ mL}$ ). Purification by flash column chromatography (dichloromethane) gave 3-methoxy-1-[(E)-2'-nitrovinyl]benzene (1.21 g, 97%) as a yellow solid. Mp 91–92 °C (lit.<sup>36</sup> 89–91 °C);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  3.85 (s, 3H), 7.02–7.08 (m, 2H), 7.14 (d,  $J = 7.6 \text{ Hz}$ , 1H), 7.37 (t,  $J = 7.6 \text{ Hz}$ , 1H), 7.57 (d,  $J = 13.6 \text{ Hz}$ , 1H), 7.97 (d,  $J = 13.6 \text{ Hz}$ , 1H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  55.4 ( $\text{CH}_3$ ), 114.0 (CH), 118.0 (CH), 121.7 (CH), 130.4 (CH), 131.4 (C), 137.4 (CH), 139.0 (CH), 160.2 (C); MS (EI)  $m/z$  179 ( $\text{M}^+$ , 100), 136 (82), 135 (72), 84 (86), 77 (50).

**3,4-Dimethoxy-1-[(E)-2'-nitrovinyl]benzene.**<sup>37</sup> 3,4-Dimethoxy-1-[(E)-2'-nitrovinyl]benzene was synthesized as described for 3-methoxy-1-[(E)-2'-nitrovinyl]benzene using 3,4-dimethoxybenzaldehyde (1.00 g, 6.00 mmol). Purification by flash column chromatography (dichloromethane) gave 3,4-dimethoxy-1-[(E)-2'-nitrovinyl]benzene (1.04 g, 83%) as a yellow solid. Mp 134–136 °C (lit.<sup>37</sup> 135–137 °C);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  3.93 (s, 3H), 3.95 (s, 3H), 6.91 (d,  $J = 8.3 \text{ Hz}$ , 1H), 7.01 (d,  $J = 1.7 \text{ Hz}$ , 1H), 7.18 (dd,  $J = 8.3, 1.7 \text{ Hz}$ , 1H), 7.53 (d,  $J = 13.6 \text{ Hz}$ , 1H), 7.96 (d,  $J = 13.6 \text{ Hz}$ , 1H);  $^{13}\text{C}$  NMR



(101 MHz, CDCl<sub>3</sub>)  $\delta$  56.0 (CH<sub>3</sub>), 56.1 (CH<sub>3</sub>), 110.3 (CH), 111.4 (CH), 122.8 (C), 124.6 (CH), 135.2 (CH), 139.3 (CH), 149.6 (C), 152.9 (C); MS (ESI)  $m/z$  232 (M+Na<sup>+</sup>. 100).

**3,5-Dimethoxy-1-[(E)-2'-nitrovinyl]benzene.**<sup>38</sup> 3,5-Dimethoxy-1-[(E)-2'-nitrovinyl]benzene was synthesized as described for 3-methoxy-1-[(E)-2'-nitrovinyl]benzene using 3,5-dimethoxybenzaldehyde (0.350 g, 2.10 mmol). The residue was recrystallized from diethyl ether which gave 3,5-dimethoxy-1-[(E)-2'-nitrovinyl]benzene (0.357 g, 81%) as a yellow solid. Mp 81–83 °C (lit.<sup>38</sup> 78 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  3.83 (s, 6H), 6.59 (t,  $J$  = 2.2 Hz, 1H), 6.66 (d,  $J$  = 2.2 Hz, 2H), 7.54 (d,  $J$  = 13.6 Hz, 1H), 7.92 (d,  $J$  = 13.6 Hz, 1H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$  55.5 (CH<sub>3</sub>), 55.6 (CH<sub>3</sub>), 102.8 (CH), 104.2 (CH), 107.0 (CH), 107.6 (CH), 133.4 (CH), 139.2 (C), 160.7 (C), 161.3 (C); MS (EI)  $m/z$  209 (M<sup>+</sup>. 100), 189 (48), 165 (90), 135 (34), 84 (98).

**1-[(E)-2'-Nitrovinyl]-3,4,5-trimethoxybenzene.**<sup>37</sup> 1-[(E)-2'-Nitrovinyl]-3,4,5-trimethoxybenzene was synthesized as described for 3-methoxy-1-[(E)-2'-nitrovinyl]benzene using 3,4,5-trimethoxybenzaldehyde (1.00 g, 6.00 mmol). The residue was recrystallized from hexane, which gave 1-[(E)-2'-nitrovinyl]-3,4,5-trimethoxybenzene (1.14 g, 94%) as a yellow solid. Mp 119–121 °C (lit.<sup>37</sup> 122–124 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  3.91 (s, 6H), 3.92 (s, 3H), 6.76 (s, 2H), 7.52 (d,  $J$  = 13.6 Hz, 1H), 7.94 (d,  $J$  = 13.6 Hz, 1H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$  56.3 (2 × CH<sub>3</sub>), 61.1 (CH<sub>3</sub>), 106.5 (2 × CH), 125.3 (C), 136.4 (CH), 139.3 (CH), 141.8 (C), 153.7 (2 × C); MS (ESI)  $m/z$  262 (M+Na<sup>+</sup>. 100).

**3-Methoxy-4-methylbenzaldehyde.**<sup>39</sup> To a stirred solution of lithium aluminium hydride (0.425 g, 11.1 mmol) in dry tetrahydrofuran (20 mL) was added a solution of methyl 3-methoxy-4-methylbenzoate (1.00 g, 5.55 mmol) in dry tetrahydrofuran (10 mL) dropwise at 0 °C. The resulting suspension was warmed to room temperature and stirred for 4 h after which, the solution was cooled to 0 °C and diluted with tetrahydrofuran (20 mL). Water (0.5 mL) was added slowly followed by 15% aqueous sodium hydroxide solution (0.5 mL) and water (1.5 mL). The resulting solution was warmed to room temperature, magnesium sulfate (0.50 g) was added and stirred for 0.5 h. The suspension was filtered and the filtrate was concentrated *in vacuo* to give 3-methoxy-4-methylbenzyl alcohol (0.767 g, 91%) as a colorless oil

1 which was used without further purification. To a stirred solution of 3-methoxy-4-methylbenzyl alcohol  
2 (0.737 g, 4.85 mmol) in chloroform (25 mL) was added manganese dioxide (4.22 g, 48.5 mmol). The  
3 resulting suspension was stirred at room temperature for 18 h. The crude reaction mixture was filtered  
4 through Celite® and concentrated *in vacuo* to give 3-methoxy-4-methylbenzaldehyde (0.621 g, 75%) as  
5 a white solid. Mp 34–36 °C (lit.<sup>39</sup> 39–40 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.29 (s, 3H), 3.90 (s, 3H),  
6 7.29 (d, *J* = 7.5 Hz, 1H), 7.34 (d, *J* = 1.4 Hz, 1H), 7.36 (dd, *J* = 7.5, 1.4 Hz, 1H), 9.93 (s, 1H); <sup>13</sup>C NMR  
7 (101 MHz, CDCl<sub>3</sub>) δ 16.9 (CH<sub>3</sub>), 55.5 (CH<sub>3</sub>), 107.9 (CH), 124.5 (CH), 130.9 (C), 134.9 (CH), 135.9 (C),  
8 158.3 (C), 192.0 (CH); MS (EI) *m/z* 150 (M<sup>+</sup>, 100), 121 (19), 91 (28), 84 (21).

### 19 **3-Methoxy-4-methyl-1-[(*E*)-2'-nitrovinyl]benzene.**

### 20 3-Methoxy-4-methyl-1-[(*E*)-2'-

21 nitrovinyl]benzene was synthesized as described for 3-methoxy-1-[(*E*)-2'-nitrovinyl]benzene using 3-  
22 methoxy-4-methylbenzaldehyde (0.624 g, 4.16 mmol). The residue was recrystallized from hexane which  
23 gave 3-methoxy-4-methyl-1-[(*E*)-2'-nitrovinyl]benzene (0.585 g, 73%) as a yellow solid. Mp 142–143  
24 °C; IR (neat) 3119, 2945, 1629, 1602, 1573, 1494, 1414, 1343, 1328, 1248, 1159, 1035, 976, 814 cm<sup>-1</sup>;  
25 <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.26 (s, 3H), 3.87 (s, 3H), 6.92 (s, 1H), 7.06 (d, *J* = 7.6 Hz, 1H), 7.19 (d, *J*  
26 = 7.6 Hz, 1H), 7.57 (d, *J* = 13.6 Hz, 1H), 7.97 (d, *J* = 13.6 Hz, 1H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 16.6  
27 (CH<sub>3</sub>), 55.4 (CH<sub>3</sub>), 109.3 (CH), 122.2 (CH), 128.8 (C), 131.4 (CH), 132.4 (C), 136.3 (CH), 139.6 (CH),  
28 158.3 (C); MS (ESI) *m/z* 216 (M+Na<sup>+</sup>, 100); HRMS (ESI) calcd for C<sub>10</sub>H<sub>11</sub>NNaO<sub>3</sub> (M+Na<sup>+</sup>) 216.0631,  
29 found 216.0626.

### 43 **3,4-Methylenedioxy-1-[(*E*)-2'-nitrovinyl]benzene.<sup>40</sup>**

### 44 3,4-Methylenedioxy-1-[(*E*)-2'-

45 nitrovinyl]benzene was synthesized as described for 3-methoxy-1-[(*E*)-2'-nitrovinyl]benzene using  
46 piperonal (1.00 g, 6.00 mmol). The residue was recrystallized from hexane which gave 3,4-  
47 methylenedioxy-1-[(*E*)-2'-nitrovinyl]benzene (1.03 g, 80%) as a yellow solid. Mp 142–143 °C (lit.<sup>40</sup> 148  
48 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 6.06 (s, 2H), 6.87 (d, *J* = 8.0 Hz, 1H), 7.00 (d, *J* = 1.8 Hz, 1H), 7.08  
49 (dd, *J* = 8.0, 1.8 Hz, 1H), 7.47 (d, *J* = 13.6 Hz, 1H), 7.92 (d, *J* = 13.6 Hz, 1H); <sup>13</sup>C NMR (101 MHz,  
50 51 52 53 54 55 56 57 58 59 60

CDCl<sub>3</sub>)  $\delta$  102.1 (CH<sub>2</sub>), 107.0 (CH), 109.1 (CH), 124.2 (CH), 126.6 (CH), 135.5 (C), 139.1 (CH), 148.8 (C), 151.4 (C); MS (EI)  $m/z$  193 (M<sup>+</sup>. 100), 146 (98), 89 (65), 84 (51), 63 (44).

**3-Nitro-1-[(E)-2'-nitrovinyl]benzene.**<sup>41</sup> 3-Nitro-1-[(E)-2'-nitrovinyl]benzene was synthesized as described for 3-methoxy-1-[(E)-2'-nitrovinyl]benzene using 3-nitrobenzaldehyde (1.00 g, 6.00 mmol). Purification by flash column chromatography (petroleum ether/ethyl acetate, 9:1) gave 3-nitro-1-[(E)-2'-nitrovinyl]benzene (0.420 g, 31%) as a yellow solid. Mp 119–121 °C (lit.<sup>41</sup> 125–126 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.68 (d,  $J$  = 13.7 Hz, 1H), 7.69 (t,  $J$  = 8.0 Hz, 1H), 7.88 (dt,  $J$  = 8.0, 1.6 Hz, 1H), 8.05 (d,  $J$  = 13.7 Hz, 1H), 8.35 (dt,  $J$  = 8.0, 1.6 Hz, 1H), 8.43 (t,  $J$  = 1.6 Hz, 1H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$  123.5 (CH), 126.2 (CH), 130.6 (CH), 131.9 (C), 134.4 (CH), 136.2 (CH), 139.3 (CH), 148.9 (C); MS (EI)  $m/z$  194 (M<sup>+</sup>. 100), 147 (48), 118 (38), 102 (100), 84 (82), 76 (36).

**4-Chloro-3-nitro-1-[(E)-2'-nitrovinyl]benzene.** 4-Chloro-3-nitro-1-[(E)-2'-nitrovinyl]benzene was synthesized as described for 3-methoxy-1-[(E)-2'-nitrovinyl]benzene using 4-chloro-3-nitrobenzaldehyde (1.00 g, 6.00 mmol). Purification by flash column chromatography (petroleum ether/ethyl acetate, 7:3) gave 4-chloro-3-nitro-1-[(E)-2'-nitrovinyl]benzene (0.620 g, 26%) as a yellow solid. Mp 142–143 °C; IR (neat) 3109, 2945, 2361, 1605, 1540, 1342, 1049, 833 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.61 (d,  $J$  = 13.7 Hz, 1H), 7.66–7.72 (m, 2H), 7.96 (d,  $J$  = 13.7 Hz, 1H), 8.06 (d,  $J$  = 1.2 Hz, 1H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$  125.6 (CH), 130.1 (C), 130.4 (C), 132.6 (CH), 133.2 (CH), 133.8 (C), 135.1 (CH), 139.4 (CH); MS (EI)  $m/z$  228 (M<sup>+</sup>. 40), 181 (100), 152 (38), 136 (95), 115 (38), 101 (48), 89 (41), 75 (54); HRMS (EI) calcd for C<sub>8</sub>H<sub>5</sub><sup>35</sup>ClN<sub>2</sub>O<sub>4</sub> (M<sup>+</sup>) 227.9938, found 227.9929.

**1'-(3-Methoxyphenyl)ethyl-2'-amine.**<sup>42</sup> To a suspension of sodium borohydride (0.180 g, 4.80 mmol) in dry tetrahydrofuran (10 mL) was added boron trifluoride diethyl etherate (0.750 mL, 6.00 mmol) dropwise at 0 °C and the contents were stirred at room temperature for 0.25 h. A solution of 3-methoxy-1-[(E)-2'-nitrovinyl]benzene in tetrahydrofuran (3.0 mL) was added dropwise into the reaction mixture which was then heated under reflux for 6.5 h. After cooling to room temperature, the reaction was quenched by the slow addition of ice water (12 mL). The reaction mixture was acidified with 1 M aqueous

hydrochloric acid (12 mL) and heated to 85 °C for 2 h. The reaction mixture was cooled to room temperature, washed with dichloromethane (2 × 10 mL), then 1 M aqueous sodium hydroxide was added until basic (ca. pH 12). The aqueous layer was extracted with dichloromethane (3 × 20 mL), dried (MgSO<sub>4</sub>) and concentrated *in vacuo* to give 1'-(3-methoxyphenyl)ethyl-2'-amine (0.140 g, 92%) as a yellow oil which was used without further purification. Spectroscopic data were consistent with the literature.<sup>42</sup> <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 1.99 (br s, 2H), 2.72 (t, *J* = 6.8 Hz, 2H), 2.96 (t, *J* = 6.8 Hz, 2H), 3.78 (s, 3H), 6.73–6.80 (m, 3H), 7.21 (t, *J* = 7.6 Hz, 1H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 39.0 (CH<sub>2</sub>), 43.0 (CH<sub>2</sub>), 55.2 (CH<sub>3</sub>), 111.6 (CH), 114.6 (CH), 121.2 (CH), 129.5 (CH), 140.9 (C), 159.8 (C); MS (ESI) *m/z* 152 (M+H<sup>+</sup>. 100).

***N*-[(3-Methoxyphenyl)ethyl]benzamide (1a).**<sup>43</sup> 1'-(3-Methoxyphenyl)ethyl-2'-amine (0.050 g, 0.33 mmol) was dissolved in dry dichloromethane (5 mL) and triethylamine (0.070 mL, 0.50 mmol) was added. The reaction mixture was cooled to 0 °C and benzoyl chloride (0.039 mL, 0.33 mmol) was added dropwise. The reaction mixture was stirred at 0 °C for 0.5 h, warmed to room temperature and stirred for 20 h. The reaction mixture was diluted with dichloromethane (10 mL), washed with 1 M aqueous hydrochloric acid (10 mL) then brine (10 mL), dried (MgSO<sub>4</sub>) and concentrated *in vacuo*. Purification by flash column chromatography (petroleum ether/ethyl acetate, 7:3) gave *N*-[(3-methoxyphenyl)ethyl]benzamide (**1a**) (0.077 g, 91% yield) as a white solid. Mp 64–66 °C (lit.<sup>43</sup> 67 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.91 (t, *J* = 6.9 Hz, 2H), 3.71 (q, *J* = 6.9 Hz, 2H), 3.78 (s, 3H), 6.24 (br s, 1H), 6.76–6.84 (m, 3H), 7.20–7.26 (m, 1H), 7.36–7.42 (m, 2H), 7.44–7.50 (m, 1H), 7.67–7.72 (m, 2H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 35.8 (CH<sub>2</sub>), 41.1 (CH<sub>2</sub>), 55.2 (CH<sub>3</sub>), 112.1 (CH), 114.4 (CH), 121.1 (CH), 126.8 (2 × CH), 128.6 (2 × CH), 129.7 (CH), 131.4 (CH), 134.7 (C), 140.5 (C), 159.9 (C), 167.5 (C); MS (EI) *m/z* 255 (M<sup>+</sup>. 25), 134 (100), 105 (62), 77 (25).

***N*-[(3-Methoxyphenyl)ethyl]acetamide (1b).**<sup>44</sup> 1'-(3-Methoxyphenyl)ethyl-2'-amine (0.050 g, 0.33 mmol) was dissolved in dry dichloromethane (10 mL) and acetic anhydride (0.038 mL, 0.40 mmol) was added while stirring. The reaction mixture was stirred at room temperature for 20 h. The reaction mixture

1 was diluted with dichloromethane (15 mL), washed with 1 M aqueous sodium carbonate (15 mL), then  
2 brine (15 mL), dried (MgSO<sub>4</sub>) and concentrated *in vacuo* to give *N*-[(3-methoxyphenyl)ethyl]acetamide  
3 (**1b**) (0.060 g, 94%) as a yellow oil which was used without further purification. Spectroscopic data were  
4  
5 consistent with the literature.<sup>44</sup> <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 1.94 (s, 3H), 2.79 (t, *J* = 6.8 Hz, 2H), 3.51  
6  
7 (q, *J* = 6.8 Hz, 2H), 3.80 (s, 3H), 5.44 (br s, 1H), 6.72–6.82 (m, 3H), 7.23 (t, *J* = 7.9 Hz, 1H); <sup>13</sup>C NMR  
8  
9 (101 MHz, CDCl<sub>3</sub>) δ 23.4 (CH<sub>3</sub>), 35.7 (CH<sub>2</sub>), 40.5 (CH<sub>2</sub>), 50.2 (CH<sub>3</sub>), 111.9 (CH), 114.5 (CH), 121.1  
10  
11 (CH), 129.7 (CH), 140.5 (C), 159.9 (C), 170.0 (C); MS (ESI) *m/z* 216 (M+Na<sup>+</sup>. 100).

12  
13  
14  
15  
16  
17 **Benzyl *N*-[(3-methoxyphenyl)ethyl]carbamate (1c).**<sup>45</sup> 1'-(3-Methoxyphenyl)ethyl-2'-amine (0.100  
18 g, 0.660 mmol) was dissolved in dry dichloromethane (10 mL) and triethylamine (0.142 mL, 1.00 mmol)  
19 was added while stirring. The reaction mixture was cooled to 0 °C and benzyl chloroformate (0.114 mL,  
20 0.800 mmol) was added dropwise. The reaction mixture was stirred at 0 °C for 0.5 h, warmed to room  
21 temperature and stirred for 5 h. The reaction mixture was diluted with dichloromethane (15 mL), washed  
22 with 1 M aqueous hydrochloric acid (15 mL), then brine (15 mL), dried (MgSO<sub>4</sub>) and concentrated *in*  
23 *vacuo*. Purification by flash column chromatography (petroleum ether/ethyl acetate, 4:1) gave benzyl *N*-  
24 [(3-methoxyphenyl)ethyl]carbamate (**1c**) (0.134 g, 72%) as a colorless oil. Spectroscopic data were  
25 consistent with the literature.<sup>45</sup> <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.79 (t, *J* = 6.6 Hz, 2H), 3.45 (q, *J* = 6.6  
26 Hz, 2H), 3.78 (s, 3H), 4.76 (br s, 1H), 5.09 (s, 2H), 6.69–6.81 (m, 3H), 7.21 (t, *J* = 7.9 Hz, 1H), 7.28–7.38  
27 (m, 5H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 36.1 (CH<sub>2</sub>), 42.1 (CH<sub>2</sub>), 55.2 (CH<sub>3</sub>), 66.7 (CH<sub>2</sub>), 111.9 (CH),  
28 114.5 (CH), 121.1 (CH), 128.1 (2 × CH), 128.5 (2 × CH), 128.6 (CH), 129.6 (CH), 136.6 (C), 140.3 (C),  
29 156.3 (C), 159.8 (C); MS (ESI) *m/z* 308 (M+Na<sup>+</sup>. 100).

30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48 ***tert*-Butyl-*N*-[(3-methoxyphenyl)ethyl]carbamate (1d).**<sup>46</sup> 1'-(3-Methoxyphenyl)ethyl-2'-amine  
49 (0.200 g, 1.32 mmol) was dissolved in dry dichloromethane (10 mL) and triethylamine (0.370 mL, 2.64  
50 mmol) was added with stirring. Di-*tert*-butyl dicarbonate (0.870 g, 3.97 mmol) was added and the  
51 resulting solution was stirred at room temperature for 16 h. The reaction mixture was concentrated *in*  
52 *vacuo* and the residue was purified by column chromatography (petroleum ether/ethyl acetate, 4:1) to give  
53  
54  
55  
56  
57  
58  
59  
60

*tert*-butyl-*N*-[(3-methoxyphenyl)ethyl]carbamate (**1d**) (0.253 g, 76%) as a yellow oil. Spectroscopic data were consistent with the literature.<sup>46</sup> <sup>1</sup>H NMR (400 MHz, CHCl<sub>3</sub>) δ 1.44 (s, 9H), 2.77 (t, *J* = 6.9 Hz, 2H), 3.38 (q, *J* = 6.9 Hz, 2H), 3.80 (s, 3H), 4.55 (br s, 1H), 6.72–6.89 (m, 3H), 7.22 (t, *J* = 7.9 Hz, 1H); <sup>13</sup>C NMR (101 MHz, CHCl<sub>3</sub>) δ 28.4 (3 × CH<sub>3</sub>), 36.3 (CH<sub>2</sub>), 41.7 (CH<sub>2</sub>), 55.2 (CH<sub>3</sub>), 79.2 (C), 111.8 (CH), 114.5 (CH), 121.1 (CH), 129.6 (CH), 140.6 (C), 155.9 (C), 159.8 (C); MS (EI) *m/z* 251 (M<sup>+</sup>. 12), 195 (32), 134 (100), 121 (48), 91 (24).

***N*-[(3-Methoxyphenyl)ethyl]methanesulfonamide (1e)**. 1'-(3-Methoxyphenyl)ethyl-2'-amine (0.198 g, 1.32 mmol) was dissolved in dry dichloromethane (10 mL) and triethylamine (0.370 mL, 2.64 mmol) was added with stirring. The reaction mixture was cooled to 0 °C and methanesulfonyl chloride (0.120 mL, 1.59 mmol) was added. The reaction mixture was stirred at 0 °C for 0.5 h, warmed to room temperature and stirred for 4 h. The reaction mixture was diluted with dichloromethane (15 mL), washed with 1 M aqueous hydrochloric acid (15 mL), brine (15 mL), dried (MgSO<sub>4</sub>) and concentrated *in vacuo*. Purification by flash column chromatography (petroleum ether/ethyl acetate, 1:1) gave *N*-[(3-methoxyphenyl)ethyl]methanesulfonamide (**1e**) (0.184 g, 60%) as a yellow oil. IR (neat) 3287, 2936, 1586, 1489, 1312, 1258, 1146, 783 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 2.83–2.88 (m, 5H), 3.41 (q, *J* = 6.4 Hz, 2H), 3.81 (s, 3H), 4.24 (br s, 1H), 6.74–6.82 (m, 3H), 7.25 (t, *J* = 7.9 Hz, 1H); <sup>13</sup>C NMR (126 MHz, CDCl<sub>3</sub>) δ 36.5 (CH<sub>2</sub>), 40.4 (CH<sub>3</sub>), 44.3 (CH<sub>2</sub>), 55.2 (CH<sub>3</sub>), 112.2 (CH), 114.7 (CH), 121.1 (CH), 129.9 (CH), 139.3 (C), 160.0 (C); MS (EI) *m/z* 229 (M<sup>+</sup>. 35), 134 (100), 122 (76), 108 (62), 91 (25); HRMS (EI) calcd for C<sub>10</sub>H<sub>15</sub>NO<sub>3</sub>S (M<sup>+</sup>) 229.0773, found 229.0780.

***N*-[(3-Methoxyphenyl)ethyl]-4''-methylbenzenesulfonamide (1f)**.<sup>47</sup> 1'-(3-Methoxyphenyl)ethyl-2'-amine (0.200 g, 1.32 mmol) was dissolved in dry dichloromethane (10 mL) and triethylamine (0.280 mL, 1.98 mmol) was added with stirring. The reaction mixture was cooled to 0 °C and *p*-toluenesulfonyl chloride (0.302 g, 1.58 mmol) was added. The reaction mixture was stirred at 0 °C for 0.5 h, warmed to room temperature and stirred for 6 h. The reaction mixture was diluted with dichloromethane (15 mL), washed with 1 M aqueous hydrochloric acid (15 mL), brine (15 mL), dried (MgSO<sub>4</sub>) and concentrated *in*

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
*vacuo*. Purification by flash column chromatography (petroleum ether/ethyl acetate, 4:1) gave *N*-[(1-methoxyphenyl)ethyl]-4''-methylbenzenesulfonamide (**1f**) (0.327 g, 81%) as a yellow oil. Spectroscopic data were consistent with the literature.<sup>47</sup> <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.45 (s, 3H), 2.75 (t, *J* = 6.9 Hz, 2H), 3.23 (q, *J* = 6.9 Hz, 2H), 3.79 (s, 3H), 4.54 (br s, 1H), 6.63 (br s, 1H), 6.69 (br d, *J* = 7.9 Hz, 1H), 6.78 (dd, *J* = 7.9, 2.5 Hz, 1H), 7.20 (t, *J* = 7.9 Hz, 1H), 7.31 (d, *J* = 8.2 Hz, 2H), 7.71 (d, *J* = 8.2 Hz, 2H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 21.5 (CH<sub>3</sub>), 35.8 (CH<sub>2</sub>), 44.1 (CH<sub>2</sub>), 55.1 (CH<sub>3</sub>), 112.1 (CH), 114.4 (CH), 121.0 (CH), 127.1 (2 × CH), 129.7 (2 × CH), 129.8 (CH), 136.9 (C), 139.2 (C), 143.4 (C), 159.9 (C); MS (ESI) *m/z* 328 (M+Na<sup>+</sup>. 100).

19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
*N*-[(3,4-Dimethoxyphenyl)ethyl]-4''-methylbenzenesulfonamide (**1g**).<sup>48</sup> To a suspension of sodium borohydride (0.817 g, 21.6 mmol) in dry tetrahydrofuran (50 mL) was added boron trifluoride diethyl etherate (3.37 mL, 27.3 mmol) dropwise at 0 °C and the mixture was stirred at room temperature for 0.25 h. A solution of 3,4-dimethoxy-1-[(*E*)-2'-nitrovinyl]benzene (0.950 g, 4.55 mmol) in tetrahydrofuran (15 mL) was added dropwise into the reaction mixture which was then heated under reflux for 6.5 h. After cooling to room temperature, the reaction was quenched by the slow addition of ice water (30 mL). The reaction mixture was acidified with 1 M aqueous hydrochloric acid (30 mL) and heated to 85 °C for 2 h. The reaction mixture was cooled to room temperature, washed with dichloromethane (2 × 40 mL), then 1 M aqueous sodium hydroxide was added until basic (ca. pH 12). The aqueous layer was extracted with dichloromethane (3 × 40 mL), dried (MgSO<sub>4</sub>) and concentrated *in vacuo* to give 1'-(3,4-dimethoxyphenyl)ethyl-2'-amine (0.582 g, 71%) as a yellow oil which was used without further purification. 1'-(3,4-Dimethoxyphenyl)ethyl-2'-amine (0.509 g, 2.81 mmol) was dissolved in dry dichloromethane (20 mL) and triethylamine (0.588 mL, 4.22 mmol) was added while stirring. The reaction mixture was cooled to 0 °C and *p*-toluenesulfonyl chloride (0.643 g, 3.37 mmol) was added. The reaction mixture was stirred at 0 °C for 0.5 h, warmed to room temperature and stirred for 6 h. The reaction mixture was diluted with dichloromethane (30 mL), washed with 1 M aqueous hydrochloric acid (30 mL), brine (30 mL), dried (MgSO<sub>4</sub>) and concentrated *in vacuo*. Purification by flash column chromatography

(petroleum ether/ethyl acetate, 7:3) gave *N*-[(3,4-dimethoxyphenyl)ethyl]-4''-methylbenzenesulfonamide (**1g**) (0.716 g, 76%) as a yellow oil. Spectroscopic data were consistent with the literature.<sup>48</sup> <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.42 (s, 3H), 2.71 (t, *J* = 6.8 Hz, 2H), 3.19 (q, *J* = 6.8 Hz, 2H), 3.81 (s, 3H), 3.86 (s, 3H), 4.31 (t, *J* = 6.8 Hz, 1H), 6.55 (d, *J* = 2.0 Hz, 1H), 6.62 (dd, *J* = 8.1, 2.0 Hz, 1H), 6.76 (d, *J* = 8.1 Hz, 1H), 7.28 (d, *J* = 8.2 Hz, 2H), 7.67 (d, *J* = 8.2 Hz, 2H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 21.5 (CH<sub>3</sub>), 35.3 (CH<sub>2</sub>), 44.3 (CH<sub>2</sub>), 55.8 (CH<sub>3</sub>), 56.0 (CH<sub>3</sub>), 111.5 (CH), 111.8 (CH), 120.8 (CH), 127.1 (2 × CH), 129.7 (2 × CH), 130.1 (C), 136.9 (C), 143.4 (C), 148.0 (C), 149.2 (C); MS (EI) *m/z* 335 (M<sup>+</sup>. 60), 184 (17), 164 (38), 151 (100), 107 (17), 91 (48).

***N*-[(3,5-Dimethoxyphenyl)ethyl]-4''-methylbenzenesulfonamide (1h).**<sup>49</sup> *N*-[(3,5-Dimethoxyphenyl)ethyl]-4''-methylbenzenesulfonamide (**1h**) was synthesized as described for *N*-[(3,4-dimethoxyphenyl)ethyl]-4''-methylbenzenesulfonamide (**1g**) using 3,5-dimethoxy-1-[(*E*)-2'-nitrovinyl]benzene. Reduction of 3,5-dimethoxy-1-[(*E*)-2'-nitrovinyl]benzene (0.330 g, 1.58 mmol) using sodium borohydride (0.285 g, 7.51 mmol) and boron trifluoride diethyl etherate (1.17 mL, 9.48 mmol) gave 1'-(3,5-dimethoxyphenyl)ethyl-2'-amine (0.165 g, 58%) which was used without further purification. The *N*-protection step was carried out at room temperature for 18 h. Purification by flash column chromatography (petroleum ether/ethyl acetate, 4:1) gave *N*-[(3,5-dimethoxyphenyl)ethyl]-4''-methylbenzenesulfonamide (**1h**) (0.150 g, 50%) as a colorless oil. Spectroscopic data were consistent with the literature.<sup>49</sup> <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.42 (s, 3H), 2.69 (t, *J* = 6.9 Hz, 2H), 3.19 (q, *J* = 6.9 Hz, 2H), 3.74 (s, 6H), 4.48 (t, *J* = 6.9 Hz, 1H), 6.21 (d, *J* = 2.2 Hz, 2H), 6.31 (t, *J* = 2.2 Hz, 1H), 7.27 (d, *J* = 8.2 Hz, 2H), 7.68 (d, *J* = 8.2 Hz, 2H); <sup>13</sup>C NMR (126 MHz, CDCl<sub>3</sub>) δ 21.5 (CH<sub>3</sub>), 36.0 (CH<sub>2</sub>), 44.0 (CH<sub>2</sub>), 55.3 (2 × CH<sub>3</sub>), 98.7 (CH), 106.7 (2 × CH), 127.1 (2 × CH), 129.7 (2 × CH), 136.9 (C), 140.0 (C), 143.4 (C), 161.1 (2 × C); MS (ESI) *m/z* 358 (M+Na<sup>+</sup>. 100).

**4''-Methyl-*N*-[(3,4,5-trimethoxyphenyl)ethyl]benzenesulfonamide (1i).**<sup>50</sup> 4''-Methyl-*N*-[(3,4,5-trimethoxyphenyl)ethyl]benzenesulfonamide (**1i**) was synthesized as described for *N*-[(3,4-dimethoxyphenyl)ethyl]-4''-methylbenzenesulfonamide (**1g**) using 1-[(*E*)-2'-nitrovinyl]-3,4,5-



1 trimethoxybenzene. Reduction of 1-[(*E*)-2'-nitrovinyl]-3,4,5-trimethoxybenzene (1.09 g, 4.56 mmol)  
2 using sodium borohydride (0.821 g, 21.7 mmol) and boron trifluoride diethyl etherate (3.90 mL, 27.4  
3 mmol) gave 1'-(3,4,5-trimethoxyphenyl)ethyl-2'-amine (0.748 g, 78%) which was used without further  
4 purification. The *N*-protection step was carried out at room temperature for 6 h. Purification by flash  
5 column chromatography (petroleum ether/ethyl acetate, 3:2) gave 4''-methyl-*N*-[(3,4,5-  
6 trimethoxyphenyl)ethyl]benzenesulfonamide (**1i**) (0.903 g, 71%) as a yellow oil. Spectroscopic data were  
7 consistent with the literature.<sup>50</sup> <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.42 (s, 3H), 2.71 (t, *J* = 6.8 Hz, 2H), 3.21  
8 (q, *J* = 6.8 Hz, 2H), 3.80 (s, 6H), 3.81 (s, 3H), 4.48 (t, *J* = 6.8 Hz, 1H), 6.28 (s, 2H), 7.28 (d, *J* = 8.3 Hz,  
9 2H), 7.69 (d, *J* = 8.3 Hz, 2H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 21.5 (CH<sub>3</sub>), 36.1 (CH<sub>2</sub>), 44.2 (CH<sub>2</sub>), 56.1  
10 (2 × CH<sub>3</sub>), 60.8 (CH<sub>3</sub>), 105.7 (2 × CH), 127.1 (2 × CH), 129.7 (2 × CH), 133.3 (C), 136.9 (C), 137.0 (C),  
11 143.5 (C), 153.4 (2 × C); MS (ESI) *m/z* 388 (M+Na<sup>+</sup>. 100).

12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26 ***N*-[(3-Methoxy-4-methylphenyl)ethyl]-4''-methylbenzenesulfonamide (1j).** *N*-[(3-Methoxy-4-  
27 methylphenyl)ethyl]-4''-methylbenzenesulfonamide (**1j**) was synthesized as described for *N*-[(3,4-  
28 dimethoxyphenyl)ethyl]-4''-methylbenzenesulfonamide (**1g**) using 3-methoxy-4-methyl-1-[(*E*)-2'-  
29 nitrovinyl]benzene. Reduction of 3-methoxy-4-methyl-1-[(*E*)-2'-nitrovinyl]benzene (0.585 g, 3.03  
30 mmol) using sodium borohydride (0.757 g, 20.5 mmol) and boron trifluoride diethyl etherate (3.42 mL,  
31 26.5 mmol) gave 1'-(3-methoxy-4-methylphenyl)ethyl-2'-amine (0.356 g, 71%) which was used without  
32 further purification. The *N*-protection step was carried out at room temperature for 24 h. Purification by  
33 flash column chromatography (petroleum ether/ethyl acetate, 4:1) gave *N*-[(3-methoxy-4-  
34 methylphenyl)ethyl]-4''-methylbenzenesulfonamide (**1j**) (0.278 g, 45%) as a yellow oil. IR (neat) 3264,  
35 2924, 1586, 1512, 1464, 1414, 1323, 1256, 1155, 1094, 814 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.17 (s,  
36 3H), 2.42 (s, 3H), 2.72 (t, *J* = 6.8 Hz, 2H), 3.20 (q, *J* = 6.8 Hz, 2H), 3.76 (s, 3H), 4.34 (t, *J* = 6.8 Hz, 1H),  
37 6.51 (d, *J* = 1.4 Hz, 1H), 6.51 (dd, *J* = 7.5, 1.4 Hz, 1H), 7.01 (d, *J* = 7.5 Hz, 1H), 7.27 (d, *J* = 8.2 Hz, 2H),  
38 7.67 (d, *J* = 8.2 Hz, 2H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 15.8 (CH<sub>3</sub>), 21.5 (CH<sub>3</sub>), 35.7 (CH<sub>2</sub>), 44.2 (CH<sub>2</sub>),  
39 55.2 (CH<sub>3</sub>), 110.4 (CH), 120.4 (CH), 125.2 (C), 127.1 (2 × CH), 129.7 (2 × CH), 130.8 (CH), 136.3 (C),  
40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60

137.0 (C), 143.4 (C), 158.0 (C); MS (ESI)  $m/z$  342 (M+Na<sup>+</sup>. 100); HRMS (ESI) calcd for C<sub>17</sub>H<sub>21</sub>NNaO<sub>3</sub>S (M+Na<sup>+</sup>) 342.1134, found 342.1125.

**4''-Methyl-*N*-[(3,4-methylenedioxyphenyl)ethyl]benzenesulfonamide (1k).**<sup>51</sup> 4''-Methyl-*N*-[(3,4-methylenedioxyphenyl)ethyl]benzenesulfonamide (**1k**) was synthesized as described for *N*-[(3,4-dimethoxyphenyl)ethyl]-4''-methylbenzenesulfonamide (**1g**) using 3,4-methylenedioxy-1-[(*E*)-2'-nitrovinyl]benzene. Reduction of 3,4-methylenedioxy-1-[(*E*)-2'-nitrovinyl]benzene (1.03 g, 5.32 mmol) using sodium borohydride (0.957 g, 25.3 mmol) and boron trifluoride diethyl etherate (3.90 mL, 31.9 mmol) gave 1'-(3,4-methylenedioxyphenyl)ethyl-2'-amine (0.587 g, 66%) which was used without further purification. The *N*-protection step was carried out at room temperature for 6 h. Purification by flash column chromatography (petroleum ether/ethyl acetate, 4:1) gave 4''-methyl-*N*-[(3,4-methylenedioxyphenyl)ethyl]benzenesulfonamide (**1k**) (0.679 g, 76%) as a white solid. Mp 86–88 °C (lit.<sup>51</sup> 89–90 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.43 (s, 3H), 2.67 (t, *J* = 6.8 Hz, 2H), 3.16 (q, *J* = 6.8 Hz, 2H), 4.34 (br s, 1H), 5.92 (s, 2H), 6.49–6.54 (m, 2H), 6.70 (dd, *J* = 7.1, 1.3 Hz, 1H), 7.29 (d, *J* = 8.4 Hz, 2H), 7.69 (d, *J* = 8.4 Hz, 2H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 21.5 (CH<sub>3</sub>), 35.5 (CH<sub>2</sub>), 44.3 (CH<sub>2</sub>), 101.0 (CH<sub>2</sub>), 108.5 (CH), 109.0 (CH), 121.8 (CH), 127.1 (2 × CH), 129.7 (2 × CH), 131.3 (C), 137.0 (C), 143.5 (C), 146.5 (C), 147.9 (C); MS (ESI)  $m/z$  342 (M+Na<sup>+</sup>. 100).

**4''-Methyl-*N*-[(3-nitrophenyl)ethyl]benzenesulfonamide.** 4''-Methyl-*N*-[(3-nitrophenyl)ethyl]benzenesulfonamide was synthesized as described for *N*-[(3,4-dimethoxyphenyl)ethyl]-4''-methylbenzenesulfonamide (**1g**) using 3-nitro-1-[(*E*)-2'-nitrovinyl]benzene. Reduction of 3-nitro-1-[(*E*)-2'-nitrovinyl]benzene (0.281 g, 1.45 mmol) using sodium borohydride (0.363 g, 9.78 mmol) and boron trifluoride diethyl etherate (1.59 mL, 12.3 mmol) gave 1'-(3-nitrophenyl)ethyl-2'-amine (0.186 g, 77%) which was used without further purification. The *N*-protection step was carried out at room temperature for 6 h. Purification by flash column chromatography (petroleum ether/ethyl acetate, 1:1) gave 4''-methyl-*N*-[(3-nitrophenyl)ethyl]benzenesulfonamide (0.142 g, 82%) as a yellow oil. IR (neat) 3285, 1597, 1526, 1348, 1325, 1155, 1094, 814 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.42 (s,

3H), 2.88 (t,  $J = 7.0$  Hz, 2H), 3.25 (q,  $J = 7.0$  Hz, 2H), 5.03 (br s, 1H), 7.27 (d,  $J = 7.9$  Hz, 2H), 7.42 (t,  $J = 7.8$  Hz, 1H), 7.47 (dt,  $J = 7.8, 1.5$  Hz, 1H), 7.68 (d,  $J = 7.9$  Hz, 2H), 7.90 (t,  $J = 1.5$  Hz, 1H), 8.03 (dt,  $J = 7.8, 1.5$  Hz, 1H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  21.5 (CH<sub>3</sub>), 35.5 (CH<sub>2</sub>), 43.8 (CH<sub>2</sub>), 121.8 (CH), 123.6 (CH), 127.0 (2  $\times$  CH), 129.5 (CH), 129.8 (2  $\times$  CH), 135.2 (CH), 136.7 (C), 140.0 (C), 143.7 (C), 148.3 (C); MS (ESI)  $m/z$  343 (M+Na<sup>+</sup>. 100); HRMS (ESI) calcd for C<sub>15</sub>H<sub>16</sub>N<sub>2</sub>NaO<sub>4</sub>S (M+Na<sup>+</sup>) 343.0723, found 343.0712.

***N*-[(4-Chloro-3-nitrophenyl)ethyl]-4''-methylbenzenesulfonamide.** *N*-[(4-Chloro-3-nitrophenyl)ethyl]-4''-methylbenzenesulfonamide was synthesized as described for *N*-[(3,4-dimethoxyphenyl)ethyl]-4''-methylbenzenesulfonamide (**1g**) using 4-chloro-3-nitro-1-[(*E*)-2'-nitrovinyl]benzene. Reduction of 4-chloro-3-nitro-1-[(*E*)-2'-nitrovinyl]benzene (0.255 g, 1.12 mmol) using sodium borohydride (0.280 g, 7.55 mmol) and boron trifluoride diethyl etherate (1.23 mL, 9.52 mmol) gave 1'-(4-chloro-3-nitrophenyl)ethyl-2'-amine (0.174 g, 78%) which was used without further purification. The *N*-protection step was carried out at room temperature for 6 h. Purification by flash column chromatography (petroleum ether/ethyl acetate, 7:3) gave *N*-[(4-chloro-3-nitrophenyl)ethyl]-4''-methylbenzenesulfonamide (0.238 g, 79%) as a yellow solid. Mp 84–86 °C; IR (neat) 3271, 2922, 2361, 1532, 1327, 1157, 1088, 810 cm<sup>-1</sup>;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  2.43 (s, 3H), 2.82 (t,  $J = 6.8$  Hz, 2H), 3.22 (q,  $J = 6.8$  Hz, 2H), 5.13 (t,  $J = 6.8$  Hz, 1H), 7.24–7.30 (m, 3H, 6-H), 7.38 (d,  $J = 8.2$  Hz, 1H), 7.55 (d,  $J = 2.0$  Hz, 1H), 7.65 (d,  $J = 8.2$  Hz, 2H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  21.6 (CH<sub>3</sub>), 34.9 (CH<sub>2</sub>), 43.6 (CH<sub>2</sub>), 125.2 (C), 125.8 (CH), 127.0 (2  $\times$  CH), 129.8 (2  $\times$  CH), 131.9 (CH), 133.8 (CH), 136.5 (C), 138.6 (C), 143.9 (C), 147.7 (C); MS (ESI)  $m/z$  377 (M+Na<sup>+</sup>. 100); HRMS (ESI) calcd for C<sub>15</sub>H<sub>15</sub><sup>35</sup>ClN<sub>2</sub>NaO<sub>4</sub>S (M+Na<sup>+</sup>) 377.0333, found 377.0326.

***N*-[(3-Aminophenyl)ethyl]-4''-methylbenzenesulfonamide (**1l**).** To a stirred solution of 4''-methyl-*N*-[(3-nitrophenyl)ethyl]benzenesulfonamide (0.142 g, 0.40 mmol) in ethanol (20 mL) was added tin(II) dichloride dihydrate (0.758 g, 3.36 mmol) and the resulting solution was heated under reflux for 18 h. After cooling to room temperature, the reaction mixture was diluted with saturated aqueous sodium

hydrogen carbonate solution (20 mL) and extracted with dichloromethane (4 × 50 mL). The combined extracts were washed with brine (2 × 200 mL), dried (MgSO<sub>4</sub>) and concentrated *in vacuo*. Purification by flash column chromatography (petroleum ether/ethyl acetate, 1:1) gave *N*-[(3-aminophenyl)ethyl]-4''-methylbenzenesulfonamide (**1l**) (0.102 g, 79%) as a white solid. Mp 80–82 °C; IR (neat) 3268, 2922, 1601, 1495, 1460, 1319, 1153, 1093, 814 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.42 (s, 3H), 2.66 (t, *J* = 6.8 Hz, 2H), 3.17 (q, *J* = 6.8 Hz, 2H), 3.62 (br s, 2H), 4.38 (br s, 1H), 6.39 (t, *J* = 1.7 Hz, 1H), 6.42–6.48 (m, 1H), 6.53 (dd, *J* = 7.6, 1.7 Hz, 1H), 7.04 (t, *J* = 7.6 Hz, 1H), 7.29 (d, *J* = 7.9 Hz, 2H), 7.69 (d, *J* = 7.9 Hz, 2H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 21.5 (CH<sub>3</sub>), 35.7 (CH<sub>2</sub>), 44.1 (CH<sub>2</sub>), 113.6 (CH), 115.3 (CH), 118.8 (CH), 127.1 (2 × CH), 129.7 (2 × CH), 129.8 (CH), 137.0 (C), 138.8 (C), 143.4 (C), 146.8 (C); MS (ESI) *m/z* 313 (M+Na<sup>+</sup>. 100); HRMS (ESI) calcd for C<sub>15</sub>H<sub>18</sub>N<sub>2</sub>NaO<sub>2</sub>S (M+Na<sup>+</sup>) 313.0981, found 313.0984.

***N*-[(3-Amino-4-chlorophenyl)ethyl]-4''-methylbenzenesulfonamide (1m).** *N*-[(3-Amino-4-chlorophenyl)ethyl]-4''-methylbenzenesulfonamide (**1m**) was synthesized as described for *N*-[(3-aminophenyl)ethyl]-4''-methylbenzenesulfonamide (**1l**) using 4''-methyl-*N*-[(3-nitro-4-chlorophenyl)ethyl]benzenesulfonamide (0.278 g, 0.790 mmol). Purification by flash column chromatography (petroleum ether/ethyl acetate, 7:3) gave *N*-[(3-amino-4-chlorophenyl)ethyl]-4''-methylbenzenesulfonamide (**1m**) (0.226 g, 89%) as a colorless oil. IR (neat) 3372, 3279, 2924, 2361, 1620, 1497, 1435, 1319, 1156, 1088, 810 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.41 (s, 3H), 2.62 (t, *J* = 6.8 Hz, 2H), 3.13 (q, *J* = 6.8 Hz, 2H), 4.00 (br s, 2H), 4.88 (t, *J* = 6.8 Hz, 1H), 6.36 (dd, *J* = 8.1, 2.0 Hz, 1H), 6.49 (d, *J* = 2.0 Hz, 1H), 7.06 (d, *J* = 8.1 Hz, 1H), 7.26 (d, *J* = 8.3 Hz, 2H), 7.67 (d, *J* = 8.3 Hz, 2H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 21.6 (CH<sub>3</sub>), 35.3 (CH<sub>2</sub>), 44.0 (CH<sub>2</sub>), 116.1 (CH), 117.7 (C), 119.2 (CH), 127.1 (2 × CH), 129.6 (CH), 129.7 (2 × CH), 136.8 (C), 137.5 (C), 143.0 (C), 143.5 (C); MS (ESI) *m/z* 347 (M+Na<sup>+</sup>. 100); HRMS (ESI) calcd for C<sub>15</sub>H<sub>17</sub><sup>35</sup>ClN<sub>2</sub>NaO<sub>2</sub>S (M+Na<sup>+</sup>) 347.0591, found 347.0582.

***N*-[(3-Acetamidophenyl)ethyl]-4''-methylbenzenesulfonamide (1n).** Acetic anhydride (0.110 mL, 1.17 mmol) was added to a stirred solution of *N*-[(3-aminophenyl)ethyl]-4''-methylbenzenesulfonamide

(**1l**) (0.225 g, 0.780 mmol) in dry dichloromethane (10 mL) and stirred at room temperature for 16 h. The reaction mixture was washed with saturated sodium carbonate (15 mL) and brine (15 mL). The organic layer was dried (MgSO<sub>4</sub>) and concentrated *in vacuo*. Purification by flash column chromatography (petroleum ether/ethyl acetate, 3:7) gave *N*-[(3-acetamidophenyl)ethyl]-4''-methylbenzenesulfonamide (**1n**) (0.245 g, 95%) as a colorless oil. IR (neat) 3282, 2921, 1669, 1613, 1595, 1549, 1489, 1440, 1319, 1153, 1094, 814 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.14 (s, 3H), 2.42 (s, 3H), 2.72 (t, *J* = 6.8 Hz, 2H), 3.18 (q, *J* = 6.8 Hz, 2H), 4.69 (t, *J* = 6.8 Hz, 1H), 6.81 (d, *J* = 7.5 Hz, 1H), 7.20 (t, *J* = 7.5 Hz, 1H), 7.25–7.30 (m, 3H), 7.39 (d, *J* = 7.5 Hz, 1H), 7.46 (br s, 1H), 7.69 (d, *J* = 7.9 Hz, 2H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 21.5 (CH<sub>3</sub>), 24.6 (CH<sub>3</sub>), 35.8 (CH<sub>2</sub>), 44.1 (CH<sub>2</sub>), 118.3 (CH), 120.2 (CH), 124.5 (CH), 127.1 (2 × CH), 129.3 (CH), 129.8 (2 × CH), 136.8 (C), 138.3 (C), 138.8 (C), 143.5 (C), 168.6 (C); MS (ESI) *m/z* 355 (M+Na<sup>+</sup>. 100); HRMS (ESI) calcd for C<sub>17</sub>H<sub>20</sub>N<sub>2</sub>NaO<sub>3</sub>S (M+Na<sup>+</sup>) 355.1087, found 355.1080.

***N*-[(3-Acetamido-4-chlorophenyl)ethyl]-4''-methylbenzenesulfonamide (1o).** *N*-[(3-Acetamido-4-chlorophenyl)ethyl]-4''-methylbenzenesulfonamide (**1o**) was synthesized as described for *N*-[(3-acetamidophenyl)ethyl]-4''-methylbenzenesulfonamide (**1n**) using *N*-[(3-amino-4-chlorophenyl)ethyl]-4''-methylbenzenesulfonamide (**1m**) (0.109 g, 0.340 mmol). Purification by flash column chromatography (petroleum ether/ethyl acetate, 1:1) gave *N*-[(3-acetamido-4-chlorophenyl)ethyl]-4''-methylbenzenesulfonamide (**1o**) (0.121 g, 99%) as a colorless oil. IR (neat) 3279, 2932, 2361, 1674, 1582, 1528, 1427, 1319, 1157, 1096 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.22 (s, 3H), 2.42 (s, 3H), 2.72 (t, *J* = 6.9 Hz, 2H), 3.19 (q, *J* = 6.9 Hz, 2H), 4.73 (t, *J* = 6.9 Hz, 1H), 6.78 (dd, *J* = 8.2, 1.7 Hz, 1H), 7.23 (d, *J* = 8.2 Hz, 1H), 7.27 (d, *J* = 8.2 Hz, 2H), 7.59 (br s, 1H), 7.69 (d, *J* = 8.2 Hz, 2H), 8.11 (br s, 1H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 21.5 (CH<sub>3</sub>), 24.9 (CH<sub>3</sub>), 35.6 (CH<sub>2</sub>), 44.0 (CH<sub>2</sub>), 121.0 (C), 121.8 (CH), 125.1 (CH), 127.1 (2 × CH), 129.0 (CH), 129.7 (2 × CH), 134.6 (C), 136.9 (C), 137.9 (C), 143.4 (C), 168.4 (C); MS (ESI) *m/z* 389 (M+Na<sup>+</sup>. 100); HRMS (ESI) calcd for C<sub>17</sub>H<sub>19</sub><sup>35</sup>ClN<sub>2</sub>NaO<sub>3</sub>S (M+Na<sup>+</sup>) 389.0697, found 389.0685.

**3-Methoxyphenylacetamide (1p).**<sup>52</sup> To a stirred solution of 3-methoxyphenylacetic acid (0.500 g, 3.00 mmol) in dry dichloromethane (15 mL) was added thionyl chloride (2.63 mL, 63.0 mmol) at 0 °C. The reaction mixture was heated under reflux for 2.5 h after which the solvent was removed *in vacuo*. The residue was dissolved in tetrahydrofuran (20 mL) and 25% aqueous ammonium hydroxide (4 mL) was added slowly at 0 °C. The reaction mixture was then stirred at room temperature for 16 h. The mixture was concentrated *in vacuo* and water (15 mL) was added. The solution was heated for 0.5 h. The suspension was cooled to 0 °C and the resulting white powder was collected by vacuum filtration and washed with ice-water to give 3-methoxyphenylacetamide (**1p**) (0.209 g, 42%) as a white solid. Mp 139–141 °C (lit.<sup>52</sup> 137–139 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 3.56 (s, 2H), 3.81 (s, 3H), 5.41 (br s, 1H), 5.52 (br s, 1H), 6.80–6.88 (m, 3H), 7.28 (t, *J* = 7.8 Hz, 1H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 43.4 (CH<sub>2</sub>), 55.2 (CH<sub>3</sub>), 113.0 (CH), 115.0 (CH), 121.6 (CH), 130.1 (CH), 136.3 (C), 160.1 (C), 173.2 (C); MS (ESI) *m/z* 186 (M+Na<sup>+</sup>. 100).

**1'-(3-Methoxyphenyl)acrylonitrile.**<sup>53</sup> To a solution of *m*-anisaldehyde (1.79 mL, 14.7 mmol) in dry dichloromethane (25 mL) was added cyanomethylene triphenylphosphorane (4.88 g, 16.2 mmol) and the resulting mixture was stirred at room temperature for 16 h. After this time, the reaction mixture was concentrated *in vacuo* and the residue was purified by flash column chromatography (petroleum ether/ethyl acetate, 9:1) to give 1'-(3-methoxyphenyl)acrylonitrile (3:1 ratio of *E* to *Z* isomers) (1.81 g, 79%) as a colorless oil. Spectroscopic data is reported for the major *E* isomer. Spectroscopic data were consistent with the literature.<sup>53</sup> <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 3.84 (s, 3H), 5.87 (t, *J* = 16.6 Hz, 1H), 6.93–7.05 (m, 3H), 7.30–7.43 (m, 2H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 55.4 (CH<sub>3</sub>), 96.7 (CH), 112.5 (CH), 116.9 (CH), 117.1 (C), 120.0 (CH), 130.2 (CH), 134.8 (C), 150.5 (CH), 160.0 (C); MS (EI) *m/z* 159 (M<sup>+</sup>. 100), 116 (20), 89 (20).

***N*-[(1-Methoxyphenyl)propyl]-4''-methylbenzenesulfonamide (1q).**<sup>54</sup> To a solution of 1'-(3-methoxyphenyl)acrylonitrile (1.00 mL, 6.29 mmol) in ethanol (25 mL) was added 37% aqueous hydrochloric acid (3 mL) and 10% palladium on charcoal (0.073 g). The reaction mixture was

hydrogenated at 2.5 bar for 72 h. The reaction mixture was filtered through Celite® and concentrated *in vacuo*. The crude hydrochloride salt was dissolved in dichloromethane (25 mL) and triethylamine (0.700 mL, 5.00 mmol) was added. *p*-Toluenesulfonyl chloride (0.572 g, 3.00 mmol) was added at 0 °C and the resulting solution was stirred at room temperature for 16 h. The reaction mixture was diluted with dichloromethane (20 mL) and washed with 1 M aqueous hydrochloric acid (30 mL) and brine (30 mL), dried (MgSO<sub>4</sub>) and concentrated *in vacuo*. Purification by flash column chromatography (petroleum ether/ethyl acetate, 7:3) gave *N*-[(1-methoxyphenyl)propyl]-4''-methylbenzenesulfonamide (**1q**) (0.499 g, 63%) as a colorless oil. Spectroscopic data were consistent with the literature.<sup>54</sup> <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 1.70–1.80 (m, 2H), 2.40 (s, 3H), 2.56 (t, *J* = 7.7 Hz, 2H), 2.94 (q, *J* = 6.7 Hz, 2H), 3.75 (s, 3H), 5.04 (t, *J* = 6.7 Hz, 1H), 6.62–6.68 (m, 2H), 6.70 (ddd, *J* = 8.0, 2.5, 0.8 Hz, 1H), 7.14 (t, *J* = 8.0 Hz, 1H), 7.28 (d, *J* = 8.3 Hz, 2H), 7.74 (d, *J* = 8.3 Hz, 2H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 21.5 (CH<sub>3</sub>), 31.0 (CH<sub>2</sub>), 32.8 (CH<sub>2</sub>), 42.6 (CH<sub>2</sub>), 55.2 (CH<sub>3</sub>), 111.5 (CH), 114.1 (CH), 120.8 (CH), 127.1 (2 × CH), 129.4 (CH), 129.8 (2 × CH), 137.0 (C), 142.7 (C), 143.4 (C), 159.7 (C); MS (ESI) *m/z* 342 (M+Na<sup>+</sup>. 100).

**1-Benzoyl-5-methoxyindoline (2a).** Iron(III) chloride (0.50 mg, 3.0 μmol) was dissolved in 1-butyl-3-methylimidazolium bis(trifluoromethanesulfonyl)imide (3.0 μL, 10 μmol) and stirred for 0.5 h at room temperature and then added to a solution of *N*-iodosuccinimide (0.031 g, 0.14 mmol) in toluene (1.0 mL). *N*-[(3-Methoxyphenyl)ethyl]benzamide (**1a**) (0.035 g, 0.14 mmol) was then added and the mixture was stirred at 40 °C for 5 h. Upon completion of the iodination step, the reaction mixture was cooled to room temperature and copper(I) iodide (3.0 mg, 14 μmol), cesium carbonate (0.088 g, 0.27 mmol), *N,N'*-dimethylethylenediamine (3.0 μL, 28 μmol) and water (0.5 mL) were added. The reaction mixture was degassed under argon for 0.1 h and then heated to 130 °C for 24 h. The reaction mixture was then cooled to room temperature, diluted with ethyl acetate (10 mL) and washed with a 1 M aqueous sodium thiosulfate solution (10 mL). The aqueous layer was extracted with ethyl acetate (3 × 10 mL) and the combined organic layers were washed with brine (10 mL). The organic phase was dried (MgSO<sub>4</sub>), filtered and concentrated *in vacuo*. Purification by flash column chromatography (petroleum ether/ethyl acetate,

4:1) gave 1-benzoyl-5-methoxyindoline (**2a**) (0.027 g, 79%) as a brown solid. Mp 102–104 °C; IR (neat) 2922, 1624, 1595, 1487, 1400, 1294, 1140, 1026, 833 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>, 100 °C) δ 3.08 (t, *J* = 8.2 Hz, 2H), 3.76 (s, 3H), 4.00 (t, *J* = 8.2 Hz, 2H), 6.71 (dd, *J* = 8.7, 2.4 Hz, 1H), 6.88 (d, *J* = 2.4 Hz, 1H), 7.46–7.62 (m, 6H); <sup>13</sup>C NMR (126 MHz, DMSO-*d*<sub>6</sub>, 100 °C) δ 28.4 (CH<sub>2</sub>), 50.8 (CH<sub>2</sub>), 56.1 (CH<sub>3</sub>), 111.6 (CH), 112.6 (CH), 117.4 (CH), 127.3 (2 × CH), 128.8 (2 × CH), 130.2 (CH), 134.7 (C), 137.0 (C), 137.9 (C), 156.8 (C), 167.9 (C); MS (EI) *m/z* 253 (M<sup>+</sup>. 43), 148 (12), 105 (100), 77 (29); HRMS (EI) calcd for C<sub>16</sub>H<sub>15</sub>NO<sub>2</sub> (M<sup>+</sup>) 253.1103, found 253.1114.

**1-Acetyl-5-methoxyindoline (2b).**<sup>55</sup> 1-Acetyl-5-methoxyindoline (**2b**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using *N*-[(3-methoxyphenyl)ethyl]acetamide (**1b**) (0.049 g, 0.25 mmol). Purification by flash column chromatography (petroleum ether/ethyl acetate, 1:1) gave 1-acetyl-5-methoxyindoline (**2b**) (0.042 g, 87%) as a colorless oil. Spectroscopic data were consistent with the literature.<sup>55</sup> NMR spectra showed a 5:1 mixture of rotamers. Only signals for the major rotamer are recorded. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.20 (s, 3H), 3.17 (t, *J* = 8.4 Hz, 2H), 3.78 (s, 3H), 4.04 (t, *J* = 8.4 Hz, 2H), 6.68–6.76 (m, 2H), 8.12 (d, *J* = 8.6 Hz, 1H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 24.0 (CH<sub>3</sub>), 28.2 (CH<sub>2</sub>), 48.9 (CH<sub>2</sub>), 55.6 (CH<sub>3</sub>), 110.9 (CH), 111.9 (CH), 117.5 (CH), 132.7 (C), 136.7 (C), 156.2 (C), 167.9 (C); MS (EI) *m/z* 191 (M<sup>+</sup>. 80), 149 (60), 134 (100).

**1-(Benzyloxycarbonyl)-5-methoxyindoline (2c).** 1-(Benzyloxycarbonyl)-5-methoxyindoline (**2c**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using benzyl *N*-[(3-methoxyphenyl)ethyl]carbamate (**1c**) (0.048 g, 0.17 mmol). Purification by flash column chromatography (petroleum ether/ethyl acetate, 9:1) gave 1-(benzyloxycarbonyl)-5-methoxyindoline (**2c**) (0.030 g, 63%) as a colorless oil. IR (neat) 2953, 1701, 1493, 1406, 1325, 1263, 1132, 1024, 756 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>, 100 °C) δ 3.09 (t, *J* = 8.6 Hz, 2H), 3.74 (s, 3H), 4.01 (t, *J* = 8.6 Hz, 2H), 5.25 (s, 2H), 6.72 (dd, *J* = 8.7, 2.6 Hz, 1H), 6.84 (d, *J* = 2.6 Hz, 1H), 7.31–7.45 (m, 5H), 7.56 (d, *J* = 8.7 Hz, 1H); <sup>13</sup>C NMR (126 MHz, DMSO-*d*<sub>6</sub>, 100 °C) δ 27.7 (CH<sub>2</sub>), 48.0 (CH<sub>2</sub>), 56.2 (CH<sub>3</sub>), 66.8 (CH<sub>2</sub>), 112.0 (CH), 112.9 (CH), 115.2 (CH), 128.1 (2 × CH), 128.3 (CH), 128.8 (2 × CH), 133.4 (C), 136.3 (C), 137.3 (C), 152.9



(C), 156.2 (C); MS (ESI)  $m/z$  306 ( $M+Na^+$ , 100); HRMS (ESI) calcd for  $C_{17}H_{17}NNaO_3$  ( $M+Na^+$ ) 306.1101, found 306.1095.

**1-(*tert*-Butoxycarbonyl)-5-methoxyindoline (2d).**<sup>56</sup> 1-(*tert*-Butoxycarbonyl)-5-methoxyindoline (**2d**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using *tert*-butyl-*N*-[(3-methoxyphenyl)ethyl]carbamate (**1d**) (0.063 g, 0.25 mmol). Purification by flash column chromatography (petroleum ether/diethyl ether, 9:1) gave 1-(*tert*-butoxycarbonyl)-5-methoxyindoline (**2d**) (0.035 g, 56%) as a white solid. Mp 84–86 °C (lit.<sup>56</sup> Mp 87–88 °C); <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>, 100 °C)  $\delta$  1.53 (s, 9H), 3.04 (t,  $J$  = 8.6 Hz, 2H), 3.73 (s, 3H), 3.91 (t,  $J$  = 8.6 Hz, 2H), 6.71 (dd,  $J$  = 8.7, 2.5 Hz, 1H), 6.81 (d,  $J$  = 2.5 Hz, 1H), 7.50 (d,  $J$  = 8.7 Hz, 1H); <sup>13</sup>C NMR (126 MHz, DMSO-*d*<sub>6</sub>, 100 °C)  $\delta$  27.5 (CH<sub>2</sub>), 28.7 (3 × CH<sub>3</sub>), 48.1 (CH<sub>2</sub>), 56.2 (CH<sub>3</sub>), 80.4 (C), 111.9 (CH), 112.8 (CH), 115.1 (CH), 133.3 (C), 136.6 (C), 152.3 (C), 155.8 (C); MS (EI)  $m/z$  249 ( $M^+$ , 15), 193 (100), 149 (28), 134 (62), 84 (30).

**1-(Methanesulfonyl)-5-methoxyindoline (2e).** 1-(Methanesulfonyl)-5-methoxyindoline (**2e**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using *N*-[(3-methoxyphenyl)ethyl]methanesulfonamide (**1e**) (0.057 g, 0.25 mmol). Purification by flash column chromatography (petroleum ether/ethyl acetate, 1:1) gave 1-(methanesulfonyl)-5-methoxyindoline (**2e**) (0.041 g, 73%) as a colorless oil. IR (neat) 2932, 1489, 1343, 1157, 1030, 818 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  2.81 (s, 3H), 3.12 (t,  $J$  = 8.5 Hz, 2H), 3.78 (s, 3H), 3.97 (t,  $J$  = 8.5 Hz, 2H), 6.72 (dd,  $J$  = 8.8, 2.5 Hz, 1H), 6.79 (d,  $J$  = 2.5 Hz, 1H), 7.32 (d,  $J$  = 8.8 Hz, 1H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$  28.4 (CH<sub>2</sub>), 33.9 (CH<sub>3</sub>), 50.7 (CH<sub>2</sub>), 55.7 (CH<sub>3</sub>), 111.6 (CH), 112.8 (CH), 115.1 (CH), 133.1 (C), 135.4 (C), 156.8 (C); MS (EI)  $m/z$  227 ( $M^+$ , 45), 148 (100%), 133 (69), 117 (38), 77 (33); HRMS (EI) calcd for  $C_{10}H_{13}NO_3S$  ( $M^+$ ) 227.0616, found 227.0617.

**5-Methoxy-1-(4'-methylbenzenesulfonyl)indoline (2f).** 5-Methoxy-1-(4'-methylbenzenesulfonyl)indoline (**2f**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using *N*-[(3-methoxyphenyl)ethyl]-4'-methylbenzenesulfonamide (**1f**) (0.077 g, 0.25 mmol). Purification by flash column chromatography (petroleum ether/ethyl acetate, 4:1) gave 5-methoxy-1-(4'-

methylbenzenesulfonyl)indoline (**2f**) (0.070 g, 93%) as a yellow oil. IR (neat) 2943, 1597, 1485, 1350, 1163, 1032, 814  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  2.39 (s, 3H), 2.77 (t,  $J = 8.2$  Hz, 2H), 3.77 (s, 3H), 3.92 (t,  $J = 8.2$  Hz, 2H), 6.64 (d,  $J = 2.5$  Hz, 1H), 6.75 (dd,  $J = 8.8, 2.5$  Hz, 1H), 7.22 (d,  $J = 8.2$  Hz, 2H), 7.58 (d,  $J = 8.8$  Hz, 1H), 7.63 (d,  $J = 8.2$  Hz, 2H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  21.5 ( $\text{CH}_3$ ), 28.3 ( $\text{CH}_2$ ), 50.4 ( $\text{CH}_2$ ), 55.6 ( $\text{CH}_3$ ), 111.0 (CH), 112.6 (CH), 116.7 (CH), 127.4 ( $2 \times \text{CH}$ ), 129.6 ( $2 \times \text{CH}$ ), 133.9 (C), 134.0 (C), 135.5 (C), 143.9 (C), 156.9 (C); MS (ESI)  $m/z$  326 ( $\text{M}+\text{Na}^+$ , 100); HRMS (ESI) calcd for  $\text{C}_{16}\text{H}_{17}\text{NNaO}_3\text{S}$  ( $\text{M}+\text{Na}^+$ ) 326.0821, found 326.0810.

**5,6-Dimethoxy-1-(4'-methylbenzenesulfonyl)indoline (2g).** 5,6-Dimethoxy-1-(4'-methylbenzenesulfonyl)indoline (**2g**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using *N*-[(3,4-dimethoxyphenyl)ethyl]-4''-methylbenzenesulfonamide (**1g**) (0.084 g, 0.25 mmol). The *N*-arylation step was carried out at 130 °C for 21 h. Purification by flash column chromatography (petroleum ether/ethyl acetate, 4:1) gave 5,6-dimethoxy-1-(4'-methylbenzenesulfonyl)indoline (**2g**) (0.065 g, 78%) as a white solid. Mp 116–118 °C; IR (neat) 2955, 1597, 1505, 1456, 1348, 1211, 1159, 1089, 814  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  2.37 (s, 3H), 2.70 (t,  $J = 8.2$  Hz, 2H), 3.80 (s, 3H), 3.90 (t,  $J = 8.2$  Hz, 2H), 3.94 (s, 3H), 6.60 (s, 1H), 7.20 (d,  $J = 8.7$  Hz, 2H), 7.32 (s, 1H), 7.59 (d,  $J = 8.7$  Hz, 2H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  21.5 ( $\text{CH}_3$ ), 28.1 ( $\text{CH}_2$ ), 50.7 ( $\text{CH}_2$ ), 56.3 ( $\text{CH}_3$ ), 56.3 ( $\text{CH}_3$ ), 101.2 (CH), 108.2 (CH), 123.6 (C), 127.4 ( $2 \times \text{CH}$ ), 129.6 ( $2 \times \text{CH}$ ), 133.9 (C), 135.4 (C), 143.9 (C), 146.4 (C), 148.7 (C); MS (ESI)  $m/z$  356 ( $\text{M}+\text{Na}^+$ , 100); HRMS (ESI) calcd for  $\text{C}_{17}\text{H}_{19}\text{NNaO}_4\text{S}$  ( $\text{M}+\text{Na}^+$ ) 356.0927, found 356.0918.

**5,7-Dimethoxy-1-(4'-methylbenzenesulfonyl)indoline (2h).** 5,7-Dimethoxy-1-(4'-methylbenzenesulfonyl)indoline (**2h**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using *N*-[(3,5-dimethoxyphenyl)ethyl]-4''-methylbenzenesulfonamide (**1h**) (0.082 g, 0.25 mmol). The iodination step was carried out at 40 °C for 4 h and the *N*-arylation step at 130 °C for 21 h. Purification by flash column chromatography (dichloromethane/diethyl ether, 19:1) gave 5,7-dimethoxy-1-(4'-methylbenzenesulfonyl)indoline (**2h**) (0.061 g, 75%) as a colorless oil; IR (neat) 2361, 1558, 1350, 1165, 813  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  2.32 (t,  $J = 7.4$  Hz, 2H), 2.40 (s, 3H), 3.77 (s, 3H), 3.85 (s, 3H),

4.02 (t,  $J = 7.4$  Hz, 2H), 6.25 (d,  $J = 2.2$  Hz, 1H), 6.38 (d,  $J = 2.2$  Hz, 1H), 7.19 (d,  $J = 8.6$  Hz, 2H), 7.54 (d,  $J = 8.6$  Hz, 2H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  21.6 ( $\text{CH}_3$ ), 29.9 ( $\text{CH}_2$ ), 53.1 ( $\text{CH}_2$ ), 55.6 ( $\text{CH}_3$ ), 56.2 ( $\text{CH}_3$ ), 99.1 (CH), 101.7 (CH), 124.5 (C), 127.7 ( $2 \times \text{CH}$ ), 129.3 ( $2 \times \text{CH}$ ), 135.6 (C), 139.6 (C), 143.6 (C), 152.9 (C), 152.6 (C); MS (ESI)  $m/z$  356 ( $\text{M}+\text{Na}^+$ , 100); HRMS (ESI) calcd for  $\text{C}_{17}\text{H}_{19}\text{NNaO}_4\text{S}$  ( $\text{M}+\text{Na}^+$ ) 356.0927, found 356.0917.

**1-(4'-Methylbenzenesulfonyl)-5,6,7-trimethoxyindoline (2i).** 1-(4'-Methylbenzenesulfonyl)-5,6,7-trimethoxyindoline (**2i**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using 4'-methyl-*N*-[(3,4,5-trimethoxyphenyl)ethyl]benzenesulfonamide (**1i**) (0.092 g, 0.25 mmol). The iodination step was carried out at 40 °C for 4 h and the *N*-arylation step at 130 °C for 22 h. Purification by flash column chromatography (petroleum ether/ethyl acetate, 7:3) gave 1-(4'-methylbenzenesulfonyl)-5,6,7-trimethoxyindoline (**2i**) (0.068 g, 74%) as a white solid. Mp 126–128 °C; IR (neat) 2940, 1597, 1470, 1418, 1350, 1236, 1163, 1125, 1067, 816  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  2.30 (t,  $J = 7.6$  Hz, 2H), 2.40 (s, 3H), 3.80 (s, 3H), 3.87 (s, 3H), 4.00 (s, 3H), 4.02 (t,  $J = 7.6$  Hz, 2H), 6.38 (s, 1H), 7.18 (d,  $J = 8.3$  Hz, 2H), 7.53 (d,  $J = 8.3$  Hz, 2H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  21.6 ( $\text{CH}_3$ ), 29.6 ( $\text{CH}_2$ ), 53.1 ( $\text{CH}_2$ ), 56.3 ( $\text{CH}_3$ ), 60.3 ( $\text{CH}_3$ ), 61.2 ( $\text{CH}_3$ ), 102.9 (CH), 127.6 (C), 127.7 ( $2 \times \text{CH}$ ), 129.3 ( $2 \times \text{CH}$ ), 132.5 (C), 135.5 (C), 141.5 (C), 143.7 (C), 146.9 (C), 152.4 (C); MS (ESI)  $m/z$  386 ( $\text{M}+\text{Na}^+$ , 100); HRMS (ESI) calcd for  $\text{C}_{18}\text{H}_{21}\text{NNaO}_5\text{S}$  ( $\text{M}+\text{Na}^+$ ) 386.1033, found 386.1022.

**5-Methoxy-6-methyl-1-(4'-methylbenzenesulfonyl)indoline (2j).** 5-Methoxy-6-methyl-1-(4'-methylbenzenesulfonyl)indoline (**2j**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using *N*-[(3-methoxy-4-methylphenyl)ethyl]-4'-methylbenzenesulfonamide (**1j**) (0.058 g, 0.18 mmol). Purification by flash column chromatography (hexane/ethyl acetate, 7:3) gave 5-methoxy-6-methyl-1-(4'-methylbenzenesulfonyl)indoline (**2j**) (0.042 g, 74%) as a colorless oil; IR (neat) 2947, 1597, 1497, 1350, 1157, 1088, 1026, 810  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  2.23 (s, 3H), 2.36 (s, 3H), 2.71 (t,  $J = 8.2$  Hz, 2H), 3.75 (s, 3H), 3.88 (t,  $J = 8.2$  Hz, 2H), 6.55 (s, 1H), 7.19 (d,  $J = 8.1$  Hz, 2H), 7.46 (s, 1H), 7.59 (d,  $J = 8.1$  Hz, 2H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  16.7 ( $\text{CH}_3$ ), 21.5 ( $\text{CH}_3$ ), 28.3 ( $\text{CH}_2$ ), 50.4 ( $\text{CH}_2$ ), 55.6

(CH<sub>3</sub>), 107.0 (CH), 118.3 (CH), 126.0 (C), 127.4 (2 × CH), 129.6 (2 × CH), 130.6 (C), 134.0 (C), 134.8 (C), 143.8 (C), 155.0 (C); MS (ESI) *m/z* 340 (M+Na<sup>+</sup>. 100); HRMS (ESI) calcd for C<sub>17</sub>H<sub>19</sub>NNaO<sub>3</sub>S (M+Na<sup>+</sup>) 340.0978, found 340.0970.

**1-(4'-Methylbenzenesulfonyl)-(5,6-methylenedioxy)indoline (2k).** 5,6-Methylenedioxy-1-(4'-methylbenzenesulfonyl)indoline (**2k**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using 4''-methyl-*N*-[(3,4-methylenedioxyphenyl)ethyl]benzenesulfonamide (**1k**) (0.078 g, 0.25 mmol). The iodination step was carried out at 40 °C for 4 h and the *N*-arylation step at 130 °C for 21 h. Purification by flash column chromatography (petroleum ether/ethyl acetate, 9:1) gave 1-(4'-methylbenzenesulfonyl)-(5,6-methylenedioxy)indoline (**2k**) (0.056 g, 73%) as a white solid. Mp 139–141 °C; IR (neat) 2955, 1597, 1476, 1454, 1352, 1306, 1163, 1038, 937 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.38 (s, 3H), 2.66 (t, *J* = 8.2 Hz, 2H), 3.90 (t, *J* = 8.2 Hz, 2H), 5.94 (s, 2H), 6.51 (s, 1H), 7.22 (d, *J* = 8.3 Hz, 2H), 7.24 (s, 1H), 7.62 (d, *J* = 8.3 Hz, 2H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 21.5 (CH<sub>3</sub>), 28.0 (CH<sub>2</sub>), 50.9 (CH<sub>2</sub>), 99.0 (CH), 101.4 (CH<sub>2</sub>), 105.2 (CH), 124.7 (C), 127.4 (2 × CH), 129.6 (2 × CH), 134.0 (C), 136.0 (C), 144.0 (C), 144.6 (C), 147.2 (C); MS (ESI) *m/z* 340 (M+Na<sup>+</sup>. 100); HRMS (ESI) calcd for C<sub>16</sub>H<sub>15</sub>NNaO<sub>4</sub>S (M+Na<sup>+</sup>) 340.0614, found 340.0603.

**5-Amino-1-(4'-methylbenzenesulfonyl)indoline (2l).** 5-Amino-1-(4'-methylbenzenesulfonyl)indoline (**2l**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using *N*-[(3-aminophenyl)ethyl]-4''-methylbenzenesulfonamide (**1l**) (0.080 g, 0.28 mmol). The iodination step was carried out at 40 °C for 4 h and the *N*-arylation step at 130 °C for 21 h. Purification by flash column chromatography (hexane/ethyl acetate, 1:1) gave 5-amino-1-(4'-methylbenzenesulfonyl)indoline (**2l**) (0.055 g, 70%) as a colorless oil. IR (neat) 3475, 3365, 1624, 1597, 1488, 1343, 1161, 1091, 814 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.36 (s, 3H), 2.64 (t, *J* = 8.2 Hz, 2H), 3.53 (br s, 2H), 3.86 (t, *J* = 8.2 Hz, 2H), 6.41 (d, *J* = 2.3 Hz, 1H), 6.53 (dd, *J* = 8.5, 2.3 Hz, 1H), 7.19 (d, *J* = 8.1 Hz, 2H), 7.45 (d, *J* = 8.5 Hz, 1H), 7.58 (d, *J* = 8.1 Hz, 2H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 21.5 (CH<sub>3</sub>), 28.3 (CH<sub>2</sub>), 50.3 (CH<sub>2</sub>), 111.9 (CH), 114.3 (CH), 117.3 (CH), 127.4 (2 × CH), 129.5 (2 × CH),

133.9 (C), 134.0 (2 × C), 143.4 (C), 143.7 (C); MS (ESI)  $m/z$  311 (M+Na<sup>+</sup>. 100); HRMS (ESI) calcd for C<sub>15</sub>H<sub>16</sub>N<sub>2</sub>NaO<sub>2</sub>S (M+Na<sup>+</sup>) 311.0825, found 311.0826.

**5-Amino-6-chloro-1-(4'-methylbenzenesulfonyl)indoline (2m).** 5-Amino-6-chloro-1-(4'-methylbenzenesulfonyl)indoline (**2m**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using *N*-[(3-amino-4-chlorophenyl)ethyl]-4''-methylbenzenesulfonamide (**1m**) (0.059 g, 0.18 mmol). The iodination step was carried out at 40 °C for 4 h. Purification by flash column chromatography (hexane/ethyl acetate, 1:1) gave 5-amino-6-chloro-1-(4'-methylbenzenesulfonyl)indoline (**2m**) (0.032 g, 55%) as a colorless oil. IR (neat) 3472, 3372, 2924, 2361, 1620, 1597, 1481, 1342, 1159, 1088, 810 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.38 (s, 3H), 2.66 (t, *J* = 8.2 Hz, 2H), 3.71–3.96 (m, 4H), 6.49 (s, 1H), 7.22 (d, *J* = 8.2 Hz, 2H), 7.58 (s, 1H), 7.61 (d, *J* = 8.2 Hz, 2H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 21.5 (CH<sub>3</sub>), 28.0 (CH<sub>2</sub>), 50.3 (CH<sub>2</sub>), 112.1 (CH), 117.0 (CH), 118.1 (C), 127.4 (2 × CH), 129.7 (2 × CH), 132.4 (C), 133.7 (C), 134.2 (C), 139.7 (C), 144.0 (C); MS (ESI)  $m/z$  345 (M+Na<sup>+</sup>. 100); HRMS (ESI) calcd for C<sub>15</sub>H<sub>15</sub><sup>35</sup>ClN<sub>2</sub>NaO<sub>2</sub>S (M+Na<sup>+</sup>) 345.0435, found 345.0422.

**5-Acetamido-1-(4'-methylbenzenesulfonyl)indoline (2n).** 5-Acetamido-1-(4'-methylbenzenesulfonyl)indoline (**2n**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using *N*-[(3-acetamidophenyl)ethyl]-4''-methylbenzenesulfonamide (**1n**) (0.148 g, 0.45 mmol). The iodination step was carried out at 40 °C for 4 h and the *N*-arylation step at 130 °C for 21 h. Purification by flash column chromatography (hexane/ethyl acetate, 7:3) gave 5-acetamido-1-(4'-methylbenzenesulfonyl)indoline (**2n**) (0.094 g, 64%) as a white solid. Mp 168–170 °C; IR (neat) 3320, 2924, 1675, 1546, 1487, 1351, 1163, 1091, 815 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 2.13 (s, 3H), 2.37 (s, 3H), 2.83 (t, *J* = 8.1 Hz, 2H), 3.89 (t, *J* = 8.1 Hz, 2H), 7.05 (dd, *J* = 8.3, 1.6 Hz, 1H), 7.21 (d, *J* = 7.7 Hz, 2H), 7.24 (br s, 1H), 7.49 (br s, 1H), 7.54 (d, *J* = 8.3 Hz, 1H), 7.62 (d, *J* = 7.7 Hz, 2H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 21.5 (CH<sub>3</sub>), 24.4 (CH<sub>3</sub>), 28.1 (CH<sub>2</sub>), 50.2 (CH<sub>2</sub>), 115.4 (CH), 117.6 (CH), 119.3 (CH), 127.3 (2 × CH), 129.7 (2 × CH), 133.0 (C), 133.7 (C), 134.2 (C), 138.4 (C), 144.2 (C), 168.3 (C); MS

(ESI)  $m/z$  353 ( $M+Na^+$ . 100); HRMS (ESI) calcd for  $C_{17}H_{18}N_2NaO_3S$  ( $M+Na^+$ ) 353.0930, found 353.0925.

**5-Acetamido-6-chloro-1-(4'-methylbenzenesulfonyl)indoline (2o).** 5-Acetamido-6-chloro-1-(4'-methylbenzenesulfonyl)indoline (**2o**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using *N*-[(3-acetamido-4-chlorophenyl)ethyl]-4''-methylbenzenesulfonamide (**1o**) (0.042 g, 0.11 mmol). The bromination step was carried out at 40 °C for 4 h and the *N*-arylation step at 130 °C for 21 h. Purification by flash column chromatography (hexane/ethyl acetate, 2:3) gave 5-acetamido-6-chloro-1-(4'-methylbenzenesulfonyl)indoline (**2o**) (0.017 g, 43%) as a white solid. Mp 138–140 °C; IR (neat) 3350, 2925, 1653, 1356, 1162  $cm^{-1}$ ;  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  2.21 (s, 3H), 2.39 (s, 3H), 2.85 (t,  $J = 8.4$  Hz, 2H), 3.90 (t,  $J = 8.4$  Hz, 2H), 7.25 (d,  $J = 8.5$  Hz, 2H), 7.49 (s, 1H), 7.65 (d,  $J = 8.5$  Hz, 2H), 7.68 (s, 1H), 8.05 (br s, 1H);  $^{13}C$  NMR (101 MHz,  $CDCl_3$ )  $\delta$  21.5 ( $CH_3$ ), 24.7 ( $CH_3$ ), 27.9 ( $CH_2$ ), 50.3 ( $CH_2$ ), 115.4 (CH), 118.4 (CH), 121.9 (C), 127.3 (2  $\times$  CH), 129.8 (2  $\times$  CH), 130.5 (C), 131.6 (C), 133.6 (C), 138.6 (C), 144.4 (C), 168.2 (C); MS (ESI)  $m/z$  387 ( $M+Na^+$ . 100); HRMS (ESI) calcd for  $C_{17}H_{17}^{35}ClN_2NaO_3S$  ( $M+Na^+$ ) 387.0541, found 387.0527.

**5-Methoxyindolin-2-one (2p).**<sup>57</sup> 5-Methoxyindolin-2-one (**2p**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using 3-methoxyphenylacetamide (**1p**) (0.042 g, 0.25 mmol). The *N*-arylation step was carried out at 130 °C for 21 h. Purification by flash column chromatography (hexane/ethyl acetate, 1:1) gave 5-methoxyindolin-2-one (**2p**) (0.027 g, 65%) as a white solid. Mp 128–130 °C (lit.<sup>57</sup> 132–134 °C);  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  3.53 (s, 2H), 3.78 (s, 3H), 6.75 (dd,  $J = 8.5$ , 2.4 Hz, 1H), 6.79 (d,  $J = 8.5$  Hz, 1H), 6.85 (br s, 1H), 8.53 (br s, 1H);  $^{13}C$  NMR (101 MHz,  $CDCl_3$ )  $\delta$  36.7 ( $CH_2$ ), 55.8 ( $CH_3$ ), 110.0 (CH), 111.8 (CH), 112.5 (CH), 126.7 (C), 135.9 (C), 155.7 (C), 177.5 (C); MS (ESI)  $m/z$  186 ( $M+Na^+$ . 100).

**6-Methoxy-1-(4'-methylbenzenesulfonyl)-1,2,3,4-tetrahydroquinoline (2q).** 6-Methoxy-1-(4'-methylbenzenesulfonyl)-2,3,4-tetrahydroquinoline (**2q**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using *N*-[(1-methoxyphenyl)propyl]-4''-methylbenzenesulfonamide (**1q**) (0.083 g,

0.50 mmol). Purification by flash column chromatography (hexane/ethyl acetate, 4:1) gave 6-methoxy-1-(4'-methylbenzenesulfonyl)-1,2,3,4-tetrahydroquinoline (**2q**) (0.071 g, 85%) as a colorless oil. IR (neat) 2943 (CH), 1609, 1597 (C=C), 1493, 1339, 1162, 1090, 812  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  1.51–1.59 (m, 2H), 2.33 (t,  $J = 6.8$  Hz, 2H), 2.38 (s, 3H), 3.73–3.77 (m, 2H), 3.78 (s, 3H), 6.52 (d,  $J = 2.8$  Hz, 1H), 6.75 (dd,  $J = 9.0, 2.8$  Hz, 1H), 7.18 (d,  $J = 8.3$  Hz, 2H), 7.42 (d,  $J = 8.3$  Hz, 2H), 7.70 (d,  $J = 9.0$  Hz, 1H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  21.3 ( $\text{CH}_2$ ), 21.5 ( $\text{CH}_3$ ), 26.6 ( $\text{CH}_2$ ), 46.4 ( $\text{CH}_2$ ), 55.4 ( $\text{CH}_3$ ), 112.1 (CH), 113.7 (CH), 127.0 (CH), 127.2 ( $2 \times \text{CH}$ ), 129.5 ( $2 \times \text{CH}$ ), 129.9 (C), 132.7 (C), 136.7 (C), 143.4 (C), 157.0 (C); MS (ESI)  $m/z$  340 ( $\text{M}+\text{Na}^+$ , 100); HRMS (ESI) calcd for  $\text{C}_{17}\text{H}_{19}\text{NNaO}_3\text{S}$  ( $\text{M}+\text{Na}^+$ ) 340.0978, found 340.0965.

**(3,4-Methylenedioxy)phenethan-2'-ol (3c).**<sup>58</sup> To a stirred suspension of lithium aluminium hydride (0.211 g, 5.55 mmol) in dry tetrahydrofuran (15 mL) was added 3,4-(methylenedioxy)phenylacetic acid (0.500 g, 2.78 mmol) in tetrahydrofuran (5 mL) dropwise under a constant stream of argon at 0 °C. The suspension was stirred at room temperature for 5 h, cooled to 0 °C and quenched with water (0.20 mL). To this solution was added 15% aqueous sodium hydroxide (0.20 mL), followed by water (0.60 mL). Magnesium sulfate was added and the suspension was stirred for 0.5 h, filtered, then concentrated *in vacuo*. Purification by flash column chromatography (petroleum ether/ethyl acetate, 1:1) gave (3,4-methylenedioxy)phenethan-2'-ol (**3c**) (0.308 g, 67%) as a colorless oil. Spectroscopic data were consistent with the literature.<sup>58</sup>  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  1.50 (br s, 1H), 2.78 (t,  $J = 6.6$  Hz, 2H), 3.79 (br s, 2H), 5.93 (s, 2H), 6.67 (dd,  $J = 7.9, 1.6$  Hz, 1H), 6.72 (d,  $J = 1.6$  Hz, 1H), 6.75 (d,  $J = 7.9$  Hz, 1H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  38.9 ( $\text{CH}_2$ ), 63.7 ( $\text{CH}_2$ ), 100.9 ( $\text{CH}_2$ ), 108.3 (CH), 109.3 (CH), 121.9 (CH), 132.2 (C), 146.2 (C), 147.8 (C); MS (EI)  $m/z$  166 ( $\text{M}^+$ , 30), 135 (100).

**1'-(Dihydro-3,4-benzodioxinyl)ethan-2'-ol (3d).**<sup>59</sup> 1'-(Dihydro-3,4-benzodioxinyl)ethan-2'-ol (**3d**) was synthesized as described for (3,4-methylenedioxy)phenethan-2'-ol (**3c**) using 1,4-benzodioxane-6-acetic acid (0.469 g, 2.51 mmol). Purification by flash column chromatography (petroleum ether/ethyl acetate, 1:1) gave 1'-(dihydro-3,4-benzodioxinyl)ethan-2'-ol (**3d**) (0.425 g, 94%) as a colorless oil.

Spectroscopic data were consistent with the literature.<sup>59</sup> <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 1.78 (br s, 1H), 2.74 (t, *J* = 6.5 Hz, 2H), 3.78 (br s, 2H), 4.22 (s, 4H), 6.68 (dd, *J* = 8.2, 2.0 Hz, 1H), 6.73 (d, *J* = 2.0 Hz, 1H), 6.79 (d, *J* = 8.2 Hz, 1H); δ<sub>C</sub> (101 MHz, CDCl<sub>3</sub>) δ 38.4 (CH<sub>2</sub>), 63.6 (CH<sub>2</sub>), 64.3 (CH<sub>2</sub>), 64.4 (CH<sub>2</sub>), 117.3 (CH), 117.6 (CH), 121.9 (CH), 131.7 (C), 142.1 (C), 143.4 (C); MS (ESI) *m/z* 203 (M+Na<sup>+</sup>. 100).

**1'-(3-Aminophenyl)ethan-2'-ol.**<sup>60</sup> To a stirred solution of 1'-(3-nitrophenyl)ethan-2'-ol (0.350 g, 2.09 mmol) in ethanol (25 mL) was added tin(II) dichloride dihydrate (2.40 g, 10.5 mmol) and the resulting solution was heated under reflux for 18 h. After cooling to room temperature, the reaction mixture was diluted with saturated aqueous sodium hydrogen carbonate solution (20 mL) and extracted with dichloromethane (5 × 50 mL). The combined extracts were washed with brine (2 × 200 mL), dried (MgSO<sub>4</sub>) and concentrated *in vacuo*. Purification by flash column chromatography (petroleum ether/ethyl acetate, 3:7) gave 1'-(3-aminophenyl)ethan-2'-ol (0.090 g, 32%) as a colorless oil. Spectroscopic data were consistent with the literature.<sup>60</sup> <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 1.54 (br s, 1H), 2.78 (t, *J* = 6.5 Hz, 2H), 3.64 (br s, 2H), 3.83 (t, *J* = 6.5 Hz, 2H), 6.54–6.58 (m, 2H), 6.60–6.65 (m, 1H), 7.07–7.13 (m, 1H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 39.2 (CH<sub>2</sub>), 63.6 (CH<sub>2</sub>), 113.3 (CH), 115.7 (CH), 119.2 (CH), 129.6 (CH), 139.7 (C), 146.6 (C); MS (EI) *m/z* 137 (M<sup>+</sup>. 70), 106 (100), 84 (38), 78 (37), 63 (42).

**1'-(3-Acetamidophenyl)ethan-2'-ol (3e).** Acetic anhydride (0.085 mL, 0.900 mmol) was added to a stirred solution of 1'-(3-aminophenyl)ethan-2'-ol (0.082 g, 0.600 mmol) in dichloromethane (10 mL) and stirred for 24 h at room temperature. The reaction mixture was washed with aqueous saturated sodium carbonate (15 mL) and brine (15 mL). The organic layer was dried (MgSO<sub>4</sub>) and concentrated *in vacuo*. Recrystallization from diethyl ether gave 1'-(3-acetamido)phenylethan-2'-ol (**3e**) (0.042 g, 42%) as a white solid. Mp 102–104 °C; IR (neat) 3294, 2924, 1667, 1612, 1551, 1489, 1435, 1319, 1041, 787 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 1.67 (br s, 1H), 2.16 (s, 3H), 2.84 (t, *J* = 6.6 Hz, 2H), 3.85 (t, *J* = 6.6 Hz, 2H), 6.95–7.00 (m, 1H), 7.23–7.45 (m, 4H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 24.6 (CH<sub>3</sub>), 39.1 (CH<sub>2</sub>), 63.5 (CH<sub>2</sub>), 118.1 (CH), 120.5 (CH), 125.0 (CH), 129.2 (CH), 138.1 (C), 139.6 (C), 168.4 (C); MS (ESI) *m/z* 202 (M+Na<sup>+</sup>. 100); HRMS (ESI) calcd for C<sub>10</sub>H<sub>13</sub>NNaO<sub>2</sub> (M+Na<sup>+</sup>) 202.0838, found 202.0838.



**Methyl (3-methoxyphenyl)acetate.**<sup>61</sup> To a stirred solution of 3-methoxyphenylacetic acid (2.00 g, 12.0 mmol) in methanol (20 mL) was added a few drops of concentrated sulfuric acid. The resulting mixture was heated under reflux for 16 h. The methanol was removed *in vacuo* and the residue was diluted with dichloromethane (50 mL). The solution was washed with water (4 × 50 mL) and brine (50 mL), dried (MgSO<sub>4</sub>) and concentrated to give methyl (3-methoxyphenyl)acetate (2.12 g, 99%) as a colorless oil. Spectroscopic data were consistent with the literature.<sup>61</sup> <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 3.59 (s, 2H), 3.67 (s, 3H), 3.78 (s, 3H), 6.76–6.88 (m, 3H), 7.21 (t, *J* = 7.8 Hz, 1H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 41.2 (CH<sub>2</sub>), 52.0 (CH<sub>3</sub>), 55.2 (CH<sub>3</sub>), 112.6 (CH), 114.9 (CH), 121.6 (CH), 129.6 (CH), 135.4 (C), 159.8 (C), 171.9 (C); MS (ESI) *m/z* 203 (M+Na<sup>+</sup>, 100).

**2',2'-Dimethyl-1'-(3-methoxyphenyl)ethan-2'-ol (3f).** Methylmagnesium bromide (3.0 M in diethyl ether, 2.20 mL, 6.50 mmol) was added dropwise to a 0 °C solution of methyl (3-methoxyphenyl)acetate (0.390 g, 2.17 mmol) in dry tetrahydrofuran (20 mL). The yellow solution was warmed to room temperature and stirred for 5 h. The reaction mixture was quenched with saturated aqueous ammonium chloride (30 mL) and diluted with diethyl ether (30 mL). The layers were separated and the aqueous layer was extracted with diethyl ether (3 × 30 mL). The combined ethereal extracts were washed with brine (30 mL), dried (MgSO<sub>4</sub>) and concentrated. Purification by flash column chromatography (petroleum ether/ethyl acetate, 4:1) gave 2',2'-dimethyl-1'-(3-methoxyphenyl)ethan-2'-ol (**3f**) (0.301 g, 77%) as a colorless oil. IR (neat) 3426, 2969, 1601, 1489, 1261, 1153, 1047 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 1.23 (s, 6H), 2.74 (s, 2H), 3.79 (s, 3H), 6.75–6.81 (m, 3H), 7.21 (t, *J* = 7.9 Hz, 1H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 29.2 (2 × CH<sub>3</sub>), 49.8 (CH<sub>2</sub>), 55.2 (CH<sub>3</sub>), 70.7 (C), 111.8 (CH), 116.3 (CH), 122.9 (CH), 129.2 (CH), 139.4 (C), 159.5 (C); MS (ESI) *m/z* 203 (M+Na<sup>+</sup>, 100); HRMS (ESI) calcd for C<sub>11</sub>H<sub>16</sub>NaO<sub>2</sub> (M+Na<sup>+</sup>) 203.1043, found 203.1044.

***N*-Methoxy-1'-(3-methoxyphenyl)-*N*-methylacetamide.**<sup>62</sup> To a solution of 3-methoxyphenylacetic acid (1.50 g, 9.03 mmol) in dichloromethane (50 mL) was added 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (1.70 g, 9.03 mmol), 1-hydroxybenzotriazole hydrate

(1.35 g, 9.94 mmol), *N,O*-dimethylhydroxylamine hydrochloride (0.88 g, 9.03 mmol) and *N,N*-diisopropylethylamine (6.3 mL, 36.1 mmol). The mixture was stirred at room temperature for 24 h. Water (30 mL) and sodium hydrogen carbonate (30 mL) were added and the mixture extracted with dichloromethane ( $4 \times 100$  mL). The combined extracts were washed with brine ( $2 \times 200$  mL), dried ( $\text{MgSO}_4$ ) and concentrated *in vacuo*. Purification by flash column chromatography (petroleum ether/ethyl acetate, 1:1) gave *N*-methoxy-1'-(3-methoxyphenyl)-*N*-methylacetamide (1.40 g, 74%) as a yellow oil. Spectroscopic data were consistent with the literature.<sup>62</sup>  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  3.19 (s, 3H), 3.60 (s, 3H), 3.74 (s, 2H), 3.79 (s, 3H), 6.79 (dd,  $J = 8.0, 2.6$  Hz, 1H), 6.83–6.92 (m, 2H), 7.23 (t,  $J = 8.0$  Hz, 1H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  32.2 ( $\text{CH}_3$ ), 39.4 ( $\text{CH}_2$ ), 55.2 ( $\text{CH}_3$ ), 61.3 ( $\text{CH}_3$ ), 112.4 (CH), 114.8 (CH), 121.7 (CH), 129.4 (CH), 136.4 (C), 159.7 (C), 172.3 (C); MS (ESI)  $m/z$  232 ( $\text{M}+\text{Na}^+$ , 100).

**1'-(3-Methoxyphenyl)-2'-(4''-methoxyphenyl)ethan-2'-one.**<sup>5b</sup> An oven-dried three-neck flask was flushed with argon and charged with magnesium turnings (0.070 g, 2.3 mmol), a crystal of iodine and dry tetrahydrofuran (12 mL). 4-Bromoanisole (0.29 mL, 2.3 mmol) was added and the solution was heated under reflux for 1 h. This solution was then transferred via cannula to a solution of *N*-methoxy-1'-(3-methoxyphenyl)-*N*-methylacetamide (0.40 g, 1.9 mmol) in dry tetrahydrofuran (15 mL). The resulting suspension was stirred at room temperature for 16 h. The reaction mixture was quenched with saturated ammonium chloride solution (30 mL) and extracted with ethyl acetate ( $2 \times 30$  mL). The combined organic layers were washed with brine (30 mL), dried ( $\text{MgSO}_4$ ) and concentrated *in vacuo*. Purification by flash column chromatography (petroleum ether/ethyl acetate, 4:1) gave 1'-(3-methoxyphenyl)-2'-(4''-methoxyphenyl)ethan-2-one (0.19 g, 45%) as a colorless oil. Spectroscopic data were consistent with the literature.<sup>5b</sup>  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  3.77 (s, 3H), 3.85 (s, 3H), 4.19 (s, 2H), 6.78 (dd,  $J = 8.0, 2.4$  Hz, 1H), 6.80–6.83 (m, 1H), 6.85–6.88 (m, 1H), 6.91 (d,  $J = 9.0$  Hz, 2H), 7.22 (t,  $J = 8.0$  Hz, 1H), 7.99 (d,  $J = 9.0$  Hz, 2H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  45.4 ( $\text{CH}_2$ ), 55.2 ( $\text{CH}_3$ ), 55.5 ( $\text{CH}_3$ ), 112.3 (CH), 113.8 ( $2 \times \text{CH}$ ), 115.0 (CH), 121.8 (CH), 129.6 (CH), 131.0 ( $2 \times \text{CH}$ ), 136.5 ( $2 \times \text{C}$ ), 159.8 (C), 163.5 (C), 196.1 (C); MS (ESI)  $m/z$  279 ( $\text{M}+\text{Na}^+$ , 100).

**1'-(3-Methoxyphenyl)-2'-phenylethan-2'-one.**<sup>63</sup> An oven-dried three-neck flask was flushed with argon and charged with *N*-methoxy-1'-(3-methoxyphenyl)-*N*-methylacetamide (0.349 g, 1.67 mmol) in dry tetrahydrofuran (15 mL). Phenylmagnesium bromide (1.84 mL, 1.84 mmol; 1.0 M in tetrahydrofuran) was added dropwise at 0 °C and the solution was warmed to room temperature and stirred for 2.5 h. The reaction mixture was quenched with saturated aqueous ammonium chloride solution (15 mL) and extracted with diethyl ether (4 × 30 mL), dried (MgSO<sub>4</sub>) and concentrated *in vacuo*. Purification by flash column chromatography (petroleum ether/ethyl acetate, 19:1) gave 1'-(3-methoxyphenyl)-2'-phenylethan-2'-one (0.210 g, 56%) as a colorless oil. Spectroscopic data were consistent with the literature.<sup>63</sup> <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 3.78 (s, 3H), 4.25 (s, 2H), 6.78–6.83 (m, 2H), 6.86 (br d, *J* = 7.6 Hz, 1H), 7.24 (t, *J* = 7.6 Hz, 1H), 7.45 (t, *J* = 7.8 Hz, 2H), 7.55 (t, *J* = 7.8 Hz, 1H), 8.01 (d, *J* = 7.8 Hz, 2H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 45.6 (CH<sub>2</sub>), 55.2 (CH<sub>3</sub>), 112.4 (CH), 115.1 (CH), 121.8 (CH), 128.6 (4 × CH), 129.7 (CH), 133.2 (CH), 136.0 (C), 136.6 (C), 159.8 (C), 197.5 (C); MS (ESI) *m/z* 249 (M+Na<sup>+</sup>. 100).

**2'-(4''-Chlorophenyl)-1'-(3-methoxyphenyl)ethan-2'-one.** The reaction was carried out as described for 1'-(3-methoxyphenyl)-2'-phenylethan-2'-one using *N*-methoxy-1'-(3-methoxyphenyl)-*N*-methylacetamide (0.387 g, 1.85 mmol). This gave 2'-(4''-chlorophenyl)-1'-(3-methoxyphenyl)ethan-2'-one (0.339 g, 70%) as a colorless oil. IR (neat) 2940, 2361, 1682, 1589, 1489, 1265, 1157, 1088, 1049, 772 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 3.78 (s, 3H), 4.22 (s, 2H), 6.78–6.85 (m, 3H), 7.24 (t, *J* = 7.6 Hz, 1H), 7.42 (d, *J* = 8.6 Hz, 2H), 7.94 (d, *J* = 8.6 Hz, 2H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 45.6 (CH<sub>2</sub>), 55.2 (CH<sub>3</sub>), 112.5 (CH), 115.1 (CH), 121.7 (CH), 129.0 (2 × CH), 129.8 (CH), 130.1 (2 × CH), 134.8 (C), 135.6 (C), 139.6 (C), 159.9 (C), 196.3 (C); MS (ESI) *m/z* 283 (M+Na<sup>+</sup>. 100); HRMS (ESI) calcd for C<sub>15</sub>H<sub>13</sub><sup>35</sup>ClNaO<sub>2</sub> (M+Na<sup>+</sup>) 283.0496, found 283.0500.

**1'-(3-Methoxyphenyl)-2'-(4''-methoxyphenyl)ethan-2'-ol (3g).**<sup>5b</sup> To a stirred solution of 1'-(3-methoxyphenyl)-2'-(4''-methoxyphenyl)ethan-2-one (0.530 g, 2.07 mmol) in methanol (15 mL) was added sodium borohydride (0.196 g, 5.17 mmol). The resulting suspension was stirred at room

temperature for 4 h after which time the reaction was quenched with water (15 mL) and extracted with dichloromethane (3 × 30 mL). The combined organic layers were washed with water (2 × 60 mL), brine (60 mL), dried (MgSO<sub>4</sub>) and concentrated *in vacuo* to give 1'-(3-methoxyphenyl)-2'-(4''-methoxyphenyl)ethan-2'-ol (**3g**) (0.508 g, 95%) as a colorless oil. Spectroscopic data were consistent with the literature.<sup>5b</sup> <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 1.97 (br s, 1H), 2.94 (dd, *J* = 13.5, 8.0 Hz, 1H), 2.99 (dd, *J* = 13.5, 5.6 Hz, 1H), 3.76 (s, 3H), 3.80 (s, 3H), 4.84 (ddd, *J* = 8.0, 5.6, 2.7 Hz, 1H), 6.70–6.73 (m, 1H), 6.75–6.80 (m, 2H), 6.87 (d, *J* = 8.7 Hz, 2H), 7.21 (t, *J* = 7.9 Hz, 1H), 7.27 (d, *J* = 8.7 Hz, 2H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 46.1 (CH<sub>2</sub>), 55.2 (CH<sub>3</sub>), 55.3 (CH<sub>3</sub>), 74.9 (CH), 112.1 (CH), 113.8 (2 × CH), 115.1 (CH), 121.9 (CH), 127.2 (2 × CH), 129.5 (CH), 136.0 (C), 139.8 (C), 159.1 (C), 159.7 (C); MS (ESI) *m/z* 281 (M+Na<sup>+</sup>. 100).

**1'-(3-Methoxyphenyl)-2'-phenylethan-2'-ol (3h).**<sup>64</sup> The reaction was carried out as described for 1'-(3-methoxyphenyl)-2'-(4''-methoxyphenyl)ethan-2'-ol (**3g**) using 1'-(3-methoxyphenyl)-2'-phenylethan-2-one (0.188 g, 0.830 mmol), except that the reaction was stirred at room temperature for 1 h. This gave 1'-(3-methoxyphenyl)-2'-phenylethan-2'-ol (**3h**) (0.169 g, 89%) as a colorless oil. Spectroscopic data were consistent with the literature.<sup>64</sup> <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 1.97 (d, *J* = 3.0 Hz, 1H), 2.96 (dd, *J* = 13.7, 8.4 Hz, 1H), 3.03 (dd, *J* = 13.7, 5.0 Hz, 1H), 3.77 (s, 3H), 4.89 (ddd, *J* = 8.4, 5.0, 3.0 Hz, 1H), 6.71–6.74 (m, 1H), 6.76–6.83 (m, 2H), 7.22 (t, *J* = 7.9 Hz, 1H), 7.26–7.40 (m, 5H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 46.2 (CH<sub>2</sub>), 55.2 (CH<sub>3</sub>), 75.2 (CH), 112.2 (CH), 115.1 (CH), 121.8 (CH), 125.9 (2 × CH), 127.6 (CH), 128.4 (2 × CH), 129.5 (CH), 139.6 (C), 143.8 (C), 159.7 (C); MS (ESI) *m/z* 251 (M+Na<sup>+</sup>. 100).

**2'-(4''-Chlorophenyl)-1'-(3-methoxyphenyl)ethan-2'-ol (3i).** The reaction was carried out as described for 1'-(3-methoxyphenyl)-2'-(4''-methoxyphenyl)ethan-2'-ol (**3g**) using 2'-(4''-chlorophenyl)-1'-(3-methoxyphenyl)ethan-2'-one (0.243 g, 0.930 mmol). This gave 2'-(4''-chlorophenyl)-1'-(3-methoxyphenyl)ethan-2'-ol (**3i**) (0.210 g, 86%) as a colorless oil. IR (neat) 3402, 2940, 2361, 1597, 1489, 1258, 1157, 1088, 1049, 833, 779 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 1.98 (d,

$J = 2.9$  Hz, 1H), 2.91 (dd,  $J = 13.7, 8.5$  Hz, 1H), 2.99 (dd,  $J = 13.7, 5.0$  Hz, 1H), 3.78 (s, 3H), 4.88 (ddd,  $J = 8.5, 5.0, 2.9$  Hz, 1H), 6.71 (t,  $J = 2.0$  Hz, 1H), 6.74–6.82 (m, 2H), 7.22 (t,  $J = 8.0$  Hz, 1H), 7.27 (d,  $J = 8.8$  Hz, 2H), 7.32 (d,  $J = 8.8$  Hz, 2H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  46.2 ( $\text{CH}_2$ ), 55.2 ( $\text{CH}_3$ ), 74.5 (CH), 112.2 (CH), 115.1 (CH), 121.8 (CH), 127.3 ( $2 \times \text{CH}$ ), 128.5 ( $2 \times \text{CH}$ ), 129.6 (CH), 133.2 (C), 139.1 (C), 142.2 (C), 159.8 (C); MS (ESI)  $m/z$  285 ( $\text{M}+\text{Na}^+$ , 100); HRMS (ESI) calcd for  $\text{C}_{15}\text{H}_{15}^{35}\text{ClNaO}_2$  ( $\text{M}+\text{Na}^+$ ) 285.0653, found 285.0646.

**Ethyl (*E*)-1'-(3-methoxyphenyl)acrylate.**<sup>65</sup> A solution of lithium chloride (0.310 g, 7.30 mmol), triethyl phosphonoacetate (1.45 mL, 7.30 mmol) and 1,8-diazabicyclo[5,4,0]undec-7-ene (1.09 mL, 7.30 mmol) in dry acetonitrile (30 mL) was stirred for 0.5 h. *m*-Anisaldehyde (0.890 mL, 7.30 mmol) was added and the solution was stirred at room temperature for 18 h. The reaction mixture was quenched with brine (30 mL), concentrated and the residue was extracted with diethyl ether ( $5 \times 50$  mL). The combined organic layers were dried ( $\text{MgSO}_4$ ) and concentrated. Purification by flash column chromatography (petroleum ether/ethyl acetate, 9:1) gave ethyl (*E*)-1'-(3'-methoxyphenyl)acrylate (1.36 g, 91%) as a colorless oil. Spectroscopic data were consistent with the literature.<sup>65</sup>  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  1.33 (t,  $J = 7.2$  Hz, 3H), 3.82 (s, 3H), 4.26 (q,  $J = 7.2$  Hz, 2H), 6.42 (d,  $J = 15.9$  Hz, 1H), 6.91 (ddd,  $J = 8.1, 2.5, 0.8$  Hz, 1H), 7.04 (t,  $J = 2.5$  Hz, 1H), 7.10 (br d,  $J = 8.1$  Hz, 1H), 7.29 (t,  $J = 8.1$  Hz, 1H), 7.64 (d,  $J = 15.9$  Hz, 1H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  14.3 ( $\text{CH}_3$ ), 55.3 ( $\text{CH}_3$ ), 60.5 ( $\text{CH}_2$ ), 112.9 (CH), 116.1 (CH), 118.6 (CH), 120.7 (CH), 129.9 (CH), 135.8 (C), 144.5 (CH), 159.9 (C), 166.9 (C); MS (ESI)  $m/z$  207 ( $\text{M}+\text{H}^+$ , 100).

**Ethyl (*E*)-1'-(3,4,5-trimethoxyphenyl)acrylate.**<sup>66</sup> The reaction was conducted as described for the synthesis of ethyl (*E*)-1'-(3'-methoxyphenyl)acrylate using trimethoxybenzaldehyde (1.00 g, 5.10 mmol). The resulting off-white solid was recrystallized from hot hexane which gave ethyl (*E*)-1'-(3,4,5-trimethoxyphenyl)acrylate (0.713 g, 53%) as a white solid. Mp 50–52 °C (lit.<sup>66</sup> 53–55 °C);  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  1.34 (t,  $J = 7.2$  Hz, 3H), 3.88 (s, 3H), 3.89 (s, 6H), 4.26 (q,  $J = 7.2$  Hz, 2H), 6.35 (d,  $J = 15.9$  Hz, 1H), 6.76 (s, 2H), 7.60 (d,  $J = 15.9$  Hz, 1H);  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  14.3 ( $\text{CH}_3$ ), 56.1 (2

1 × CH<sub>3</sub>), 60.5 (CH<sub>2</sub>), 61.0 (CH<sub>3</sub>), 105.2 (2 × CH), 117.5 (CH), 130.0 (C), 140.1 (C), 144.5 (CH), 153.4 (2  
2 × C), 166.9 (C); MS (ESI) *m/z* 289 (M+H<sup>+</sup>. 100).

3  
4  
5  
6 **1'-(3-Methoxyphenyl)propan-3'-ol (3j).**<sup>67</sup> To a stirred suspension of lithium aluminium hydride  
7 (0.150 g, 3.94 mmol) in dry tetrahydrofuran (10 mL) was added ethyl (*E*)-1'-(3-methoxyphenyl)acrylate  
8 (0.325 g, 1.58 mmol) in tetrahydrofuran (10 mL) dropwise under a constant stream of argon at 0 °C. The  
9 suspension was stirred at room temperature for 5 h, then cooled to 0 °C and quenched with a saturated  
10 aqueous solution of potassium sodium tartrate (20 mL) and stirred overnight. The suspension was  
11 extracted with diethyl ether (5 × 50 mL) and the combined organic extracts were dried (MgSO<sub>4</sub>) and  
12 concentrated. Purification by flash column chromatography (petroleum ether/ethyl acetate, 7:3) gave 1'-  
13 (3-methoxyphenyl)propan-3'-ol (**3j**) (0.179 g, 69%) as a colorless oil. Spectroscopic data were consistent  
14 with the literature.<sup>67</sup> <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 1.85–1.94 (m, 2H), 2.69 (t, *J* = 7.9 Hz, 2H), 3.68 (t, *J*  
15 = 6.5 Hz, 2H), 3.80 (s, 3H), 6.72–6.77 (m, 2H), 6.78–6.82 (m, 1H), 7.21 (t, *J* = 7.6 Hz, 1H); <sup>13</sup>C NMR  
16 (101 MHz, CDCl<sub>3</sub>) δ 32.1 (CH<sub>2</sub>), 34.1 (CH<sub>2</sub>), 55.1 (CH<sub>3</sub>), 62.3 (CH<sub>2</sub>), 111.1 (CH), 114.2 (CH), 120.8  
17 (CH), 129.4 (CH), 143.5 (C), 159.7 (C); MS (ESI) *m/z* 189 (M+Na<sup>+</sup>. 100).

18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34 **1'-(3,4,5-Trimethoxyphenyl)propan-3'-ol (3l).**<sup>68</sup> The reaction was conducted as described for the  
35 synthesis of 1'-(3-methoxyphenyl)propan-3'-ol (**3j**) using ethyl (*E*)-1'-(3,4,5-trimethoxyphenyl)acrylate  
36 (0.596 g, 2.24 mmol). The suspension was stirred at room temperature for 5 h, cooled to 0 °C and  
37 quenched with water (0.25 mL). To this solution was added 15% aqueous sodium hydroxide (0.25 mL),  
38 followed by water (0.75 mL). Magnesium sulfate was added and the suspension was stirred for 0.5 h,  
39 filtered and concentrated *in vacuo*. Purification by flash column chromatography (ethyl acetate/petroleum  
40 ether, 3:2) gave 1'-(3,4,5-trimethoxyphenyl)propan-3'-ol (**3l**) (0.228 g, 45%) as a colorless oil.  
41 Spectroscopic data were consistent with the literature.<sup>68</sup> <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 1.42 (br s, 1H),  
42 1.85–1.95 (m, 2H), 2.63–2.69 (m, 2H), 3.68 (t, *J* = 6.5 Hz, 2H), 3.83 (s, 3H), 3.85 (s, 6H), 6.42 (s, 2H);  
43 <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 32.6 (CH<sub>2</sub>), 34.3 (CH<sub>2</sub>), 56.1 (2 × CH<sub>3</sub>), 60.9 (CH<sub>3</sub>), 62.3 (CH<sub>2</sub>), 105.3 (2  
44 × CH), 136.1 (C), 137.7 (C), 153.2 (2 × C); MS (ESI) *m/z* 249 (M+Na<sup>+</sup>. 100).

**2,3-Dihydro-5-methoxybenzofuran (4a).**<sup>12</sup> 2,3-Dihydro-5-methoxybenzofuran (**4a**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using 3-methoxyphenylethan-2'-ol (**3a**) (0.071 mL, 0.50 mmol). The *O*-arylation step was carried out at 150 °C for 21 h. Purification by flash column chromatography (petroleum ether/ethyl acetate, 19:1) gave 2,3-dihydro-5-methoxybenzofuran (**4a**) (0.046 g, 65%) as a colorless oil. Spectroscopic data were consistent with the literature.<sup>12</sup> <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 3.17 (t, *J* = 8.6 Hz, 2H), 3.75 (s, 3H), 4.53 (t, *J* = 8.6 Hz, 2H), 6.64 (dd, *J* = 8.6, 2.6 Hz, 1H), 6.68 (d, *J* = 8.6 Hz, 1H), 6.78 (d, *J* = 2.6 Hz, 1H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 30.3 (CH<sub>2</sub>), 56.1 (CH<sub>3</sub>), 71.3 (CH<sub>2</sub>), 109.1 (CH), 111.4 (CH), 112.8 (CH), 128.0 (C), 154.1 (C), 154.2 (C); MS (ESI) *m/z* 173 (M+Na<sup>+</sup>. 100).

**2,3-Dihydro-5,6-dimethoxybenzofuran (4b).** 2,3-Dihydro-5,6-dimethoxybenzofuran (**4b**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using 3,4-dimethoxyphenylethan-2'-ol (**3b**) (0.091 g, 0.50 mmol). Purification by flash column chromatography (petroleum ether/ethyl acetate, 9:1) gave 2,3-dihydro-5,6-dimethoxybenzofuran (**4b**) (0.065 g, 72%) as a white solid. Mp 58–60 °C; IR (neat) 2940, 1614, 1502, 1446, 1304, 1208, 1186, 1169, 1095, 1003, 828 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 3.15 (t, *J* = 8.7 Hz, 2H), 3.82 (s, 3H), 3.83 (s, 3H), 4.54 (t, *J* = 8.7 Hz, 2H), 6.45 (s, 1H), 6.78 (s, 1H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 30.0 (CH<sub>2</sub>), 56.1 (CH<sub>3</sub>), 57.0 (CH<sub>3</sub>), 71.7 (CH<sub>2</sub>), 94.9 (CH), 109.3 (CH), 116.6 (C), 143.2 (C), 149.2 (C), 154.3 (C); MS (ESI) *m/z* 203 (M+Na<sup>+</sup>. 100); HRMS (ESI) calcd for C<sub>10</sub>H<sub>12</sub>NaO<sub>3</sub> (M+Na<sup>+</sup>) 203.0679, found 203.0679.

**2,3-Dihydro-5,6-methylenedioxybenzofuran (4c).** 2,3-Dihydro-5,6-methylenedioxybenzofuran (**4c**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using (3,4-methylenedioxy)phenylethan-2'-ol (**3c**) (0.083 g, 0.50 mmol). Purification by flash column chromatography (petroleum ether/ethyl acetate, 9:1) gave 2,3-dihydro-5,6-methylenedioxybenzofuran (**4c**) (0.053 g, 64%) as a white solid. Mp 54–56 °C; IR (neat) 2893, 2361, 1620, 1474, 1296, 1142, 1034, 941 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 3.10 (t, *J* = 8.7 Hz, 2H), 4.54 (t, *J* = 8.7 Hz, 2H), 5.87 (s, 2H), 6.37 (s, 1H), 6.65 (s, 1H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 30.0 (CH<sub>2</sub>), 72.0 (CH<sub>2</sub>), 93.0 (CH), 101.1 (CH<sub>2</sub>), 105.0 (CH), 117.7

(C), 141.4 (C), 147.1 (C), 154.6 (C); MS (EI)  $m/z$  164 ( $M^+$ , 100), 133 (18), 84 (48), 78 (20); HRMS (EI) calcd for  $C_9H_8O_3$  ( $M^+$ ) 164.0473, found 164.0469.

**2,3-Dihydro-5,6-ethylenedioxybenzofuran (4d).** 2,3-Dihydro-5,6-ethylenedioxybenzofuran (**4d**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using 1'-(dihydro-3,4-benzodioxinyl)ethan-2'-ol (**3d**) (0.106 g, 0.590 mmol). The iodination step was carried out at 40 °C for 4 h. Purification by flash column chromatography (petroleum ether/ethyl acetate, 9:1) gave 2,3-dihydro-5,6-ethylenedioxybenzofuran (**4d**) (0.062 g, 60%) as a white solid. Mp 64–66 °C; IR (neat) 2992, 1607, 1486, 1327, 1185, 1060, 931  $cm^{-1}$ ;  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  3.10 (t,  $J$  = 8.3 Hz, 2H), 4.15–4.23 (m, 4H), 4.50 (t,  $J$  = 8.3 Hz, 2H), 6.34 (s, 1H), 6.70 (s, 1H);  $^{13}C$  NMR (101 MHz,  $CDCl_3$ )  $\delta$  29.6 ( $CH_2$ ), 64.1 ( $CH_2$ ), 64.5 ( $CH_2$ ), 71.6 ( $CH_2$ ), 98.4 (CH), 113.2 (CH), 119.2 (C), 137.3 (C), 142.8 (C), 154.3 (C); MS (EI)  $m/z$  178 ( $M^+$ , 100), 122 (92), 69 (25); HRMS (EI) calcd for  $C_{10}H_{10}O_3$  ( $M^+$ ) 178.0630, found 178.0625.

**5-Acetamido-2,3-dihydrobenzofuran (4e).**<sup>69</sup> 5-Acetamido-2,3-dihydrobenzofuran (**4e**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using 1'-(3-acetamidophenyl)ethan-2'-ol (**3e**) (0.027 g, 0.15 mmol). The iodination step was carried out at 40 °C for 4 h. Purification by flash column chromatography (petroleum ether/ethyl acetate, 1:1) gave 5-acetamido-2,3-dihydrobenzofuran (**4e**) (0.014 g, 56%) as a white solid. Mp 94–96 °C (lit.<sup>69</sup> 93–95 °C);  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  2.12 (s, 3H), 3.17 (t,  $J$  = 8.4 Hz, 2H), 4.54 (t,  $J$  = 8.4 Hz, 2H), 6.69 (d,  $J$  = 8.5 Hz, 1H), 7.00 (dd,  $J$  = 8.5, 2.2 Hz, 1H) 7.44–7.49 (m, 2H);  $^{13}C$  NMR (101 MHz,  $CDCl_3$ )  $\delta$  24.2 ( $CH_3$ ), 29.9 ( $CH_2$ ), 71.5 ( $CH_2$ ), 109.0 (CH), 118.6 (CH), 120.7 (CH), 127.6 (C), 130.8 (C), 157.0 (C), 168.6 (C); MS (ESI)  $m/z$  200 ( $M+Na^+$ , 100).

**2,3-Dihydro-2,2-dimethyl-5-methoxybenzofuran (4f).**<sup>12</sup> 2,3-Dihydro-2,2-dimethyl-5-methoxybenzofuran (**4f**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using 2',2'-dimethyl-1'-(3-methoxyphenyl)ethan-2'-ol (**3f**) (0.120 g, 0.670 mmol). Purification by flash column chromatography (petroleum ether/diethyl ether, 19:1) gave 2,3-dihydro-2,2-dimethyl-5-methoxybenzofuran (**4f**) (0.082 g, 69%) as a colorless oil. Spectroscopic data were consistent with the



literature.<sup>12</sup> <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  1.45 (s, 6H), 2.98 (s, 2H), 3.74 (s, 3H), 6.60–6.67 (m, 2H), 6.71–6.75 (m, 1H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$  28.1 (2  $\times$  CH<sub>3</sub>), 43.3 (CH<sub>2</sub>), 56.0 (CH<sub>3</sub>), 86.5 (C), 109.3 (CH), 111.6 (CH), 112.8 (CH), 128.1 (C), 153.1 (C), 153.8 (C); MS (ESI)  $m/z$  201 (M+Na<sup>+</sup>. 100).

**2,3-Dihydro-2-(4'-methoxyphenyl)-5-methoxybenzofuran (4g).**<sup>5b</sup> 2,3-Dihydro-2-(4'-methoxyphenyl)-5-methoxybenzofuran (**4g**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using 1'-(3-methoxyphenyl)-2'-(4''-methoxyphenyl)ethan-2'-ol (**3g**) (0.050 g, 0.19 mmol). The *O*-arylation step was carried out at 130 °C for 18 h. Purification by flash column chromatography (hexane/dichloromethane, 1:1) gave 2,3-dihydro-2-(4'-methoxyphenyl)-5-methoxybenzofuran (**4g**) (0.013 g, 29%) as a colorless oil. Spectroscopic data were consistent with the literature.<sup>5b</sup> <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  3.19 (dd,  $J$  = 15.7, 8.3 Hz, 1H), 3.55 (dd,  $J$  = 15.7, 9.0 Hz, 1H), 3.77 (s, 3H), 3.81 (s, 3H), 5.68 (dd,  $J$  = 9.0, 8.3 Hz, 1H), 6.65–6.84 (m, 3H), 6.89 (d,  $J$  = 8.7 Hz, 2H), 7.33 (d,  $J$  = 8.7 Hz, 2H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$  38.7 (CH<sub>2</sub>), 55.3 (CH<sub>3</sub>), 56.1 (CH<sub>3</sub>), 84.2 (CH), 109.2 (CH), 111.2 (CH), 113.0 (CH), 114.0 (2  $\times$  CH), 127.3 (2  $\times$  CH), 127.7 (C), 133.9 (C), 153.7 (C), 154.2 (C), 159.5 (C); MS (ESI)  $m/z$  279 (M+Na<sup>+</sup>. 100).

**2,3-Dihydro-5-methoxy-2-phenylbenzofuran (4h).**<sup>12</sup> 2,3-Dihydro-5-methoxy-2-phenylbenzofuran (**4h**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using 1'-(3-methoxyphenyl)-2'-phenylethan-2'-ol (**3h**) (0.084 g, 0.37 mmol). The iodination step was carried out at 40 °C for 4 h and the *O*-arylation step at 130 °C for 22 h. Purification by flash column chromatography (petroleum ether/ethyl acetate, 19:1) gave 2,3-dihydro-5-methoxy-2-phenylbenzofuran (**4h**) (0.053 g, 64%) as a colorless oil. Spectroscopic data were consistent with the literature.<sup>12</sup> <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  3.18 (dd,  $J$  = 15.7, 8.2 Hz, 1H), 3.59 (dd,  $J$  = 15.7, 9.4 Hz, 1H), 3.76 (s, 3H), 5.72 (dd,  $J$  = 9.4, 8.2 Hz, 1H), 6.69 (dd,  $J$  = 8.7, 2.6 Hz, 1H), 6.74–6.80 (m, 2H), 7.26–7.43 (m, 5H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$  38.9 (CH<sub>2</sub>), 56.1 (CH<sub>3</sub>), 84.3 (CH), 109.2 (CH), 111.2 (CH), 113.1 (CH), 125.8 (2  $\times$  CH), 127.5 (C), 128.0 (CH), 128.7 (2  $\times$  CH), 142.1 (C), 153.8 (C), 154.3 (C); MS (ESI)  $m/z$  249 (M+Na<sup>+</sup>. 100).

**2-(4'-Chlorophenyl)-2,3-dihydro-5-methoxybenzofuran (4i).**<sup>70</sup> 2-(4'-Chlorophenyl)-2,3-dihydro-5-methoxybenzofuran (**4i**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using 2'-(4''-chlorophenyl)-1'-(3-methoxyphenyl)ethan-2'-ol (**3i**) (0.154 g, 0.590 mmol). The iodination step was carried out at 40 °C for 4 h and the *O*-arylation step at 130 °C for 22 h. Purification by flash column chromatography (petroleum ether/ethyl acetate, 19:1) gave 2-(4'-chlorophenyl)-2,3-dihydro-5-methoxybenzofuran (**4i**) (0.096 g, 63%) as a white solid. Mp 58–60 °C (lit.<sup>70</sup> 60–61 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 3.13 (dd, *J* = 15.7, 8.0 Hz, 1H), 3.60 (dd, *J* = 15.7, 9.4 Hz, 1H), 3.76 (s, 3H), 5.70 (dd, *J* = 9.4, 8.0 Hz, 1H), 6.70 (dd, *J* = 8.7, 2.6 Hz, 1H), 6.74–6.80 (m, 2H), 7.33 (br s, 4H); δ<sub>C</sub> (101 MHz, CDCl<sub>3</sub>) δ 38.9 (CH<sub>2</sub>), 56.0 (CH<sub>3</sub>), 83.4 (CH), 109.3 (CH), 111.2 (CH), 113.1 (CH), 127.1 (2 × CH), 127.1 (C), 128.8 (2 × CH), 133.7 (C), 140.6 (C), 153.6 (C), 154.4 (C); MS (ESI) *m/z* 283 (M+Na<sup>+</sup>. 100).

**2,3-Dihydro-6-methoxy-1-benzopyran (4j).**<sup>12</sup> 2,3-Dihydro-6-methoxybenzopyran (**4j**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using 1'-(3-methoxyphenyl)propan-3'-ol (**3j**) (0.083 g, 0.50 mmol). The *O*-arylation step was carried out at 150 °C for 24 h. Purification by flash column chromatography (hexane/dichloromethane, 1:1) gave 2,3-dihydro-6-methoxy-1-benzopyran (**4j**) (0.046 g, 57%) as a colorless oil. Spectroscopic data were consistent with the literature.<sup>12</sup> <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 1.94–2.02 (m, 2H), 2.77 (t, *J* = 6.5 Hz, 2H), 3.74 (s, 3H), 4.13 (t, *J* = 5.2 Hz, 2H), 6.58 (d, *J* = 2.9 Hz, 1H), 6.66 (dd, *J* = 8.8, 2.9 Hz, 1H), 6.71 (d, *J* = 8.8 Hz, 1H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ 22.5 (CH<sub>2</sub>), 25.2 (CH<sub>2</sub>), 55.7 (CH<sub>3</sub>), 66.3 (CH<sub>2</sub>), 113.3 (CH), 114.4 (CH), 117.2 (CH), 122.7 (C), 149.0 (C), 153.2 (C); MS (EI) *m/z* 164 (M<sup>+</sup>. 100), 149 (42), 136 (22), 108 (14), 84 (28), 77 (11).

**2,3-Dihydro-6,7-dimethoxy-1-benzopyran (4k).**<sup>71</sup> 2,3-Dihydro-6,7-dimethoxybenzopyran (**4k**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using 1'-(3,4-dimethoxyphenyl)propan-3'-ol (**3k**) (0.049 g, 0.25 mmol). The *O*-arylation step was carried out at 150 °C for 24 h. Purification by flash column chromatography (hexane/ethyl acetate, 4:1) gave 2,3-dihydro-6,7-dimethoxy-1-benzopyran (**4k**) (0.027 g, 56%) as a colorless oil. Spectroscopic data were consistent with the literature.<sup>71</sup> <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 1.95–2.02 (m, 2H), 2.70 (t, *J* = 6.6 Hz, 2H), 3.81 (s, 3H), 3.81 (s, 3H), 4.10–4.14

(m, 2H), 6.38 (s, 1H), 6.53 (s, 1H);  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  22.6 ( $\text{CH}_2$ ), 24.3 ( $\text{CH}_2$ ), 55.9 ( $\text{CH}_3$ ), 56.5 ( $\text{CH}_3$ ), 66.3 ( $\text{CH}_2$ ), 100.9 (CH), 112.6 (C), 112.7 (CH), 143.0 (C), 148.3 (C), 148.7 (C); MS (EI)  $m/z$  194 ( $\text{M}^+$ , 100), 179 (86), 149 (25), 123 (15), 57 (25).

**2,3-Dihydro-6,7,8-trimethoxy-1-benzopyran (4I).** 2,3-Dihydro-6,7-trimethoxy-1-benzopyran (**4I**) was synthesised as described for 1-benzoyl-5-methoxyindoline (**2a**) using 1'-(3,4,5-trimethoxyphenyl)propan-3'-ol (**3I**) (0.057 g, 0.25 mmol). The *O*-arylation step was carried out at 150 °C for 22 h. Purification by flash column chromatography (petroleum ether/ethyl acetate, 4:1) gave 2,3-dihydro-6,7,8-trimethoxy-1-benzopyran (**4I**) (0.029 g, 51%) as a colorless oil. IR (neat) 2932, 1489, 1462, 1420, 1277, 1219, 1126, 1099, 1072, 1011  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  1.95–2.03 (m, 2H), 2.73 (t,  $J = 6.5$  Hz, 2H), 3.79 (s, 3H), 3.87 (s, 3H), 3.89 (s, 3H), 4.16–4.21 (m, 2H), 6.34 (s, 1H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  22.4 ( $\text{CH}_2$ ), 24.8 ( $\text{CH}_2$ ), 56.4 ( $\text{CH}_3$ ), 61.1 ( $\text{CH}_3$ ), 61.3 ( $\text{CH}_3$ ), 66.4 ( $\text{CH}_2$ ), 107.5 (CH), 117.1 (C), 141.4 (C), 142.3 (C), 142.6 (C), 146.6 (C); MS (ESI)  $m/z$  247 ( $\text{M}+\text{Na}^+$ , 100); HRMS (ESI) calcd for  $\text{C}_{12}\text{H}_{16}\text{NaO}_4$  ( $\text{M}+\text{Na}^+$ ) 247.0941, found 247.0932.

**1'-(3-Hydroxy-4-methoxyphenyl)-*N*-methoxy-*N*-methylacetamide.** The reaction was carried out as described for *N*-methoxy-1'-(3-methoxyphenyl)-*N*-methylacetamide using 3-hydroxy-4-methoxyphenylacetic acid (**5**) (1.00 g, 5.49 mmol), except that the solvent used was acetonitrile (30 mL). Purification by flash column chromatography (petroleum ether/ethyl acetate, 3:7) gave 1'-(3-hydroxy-4-methoxyphenyl)-*N*-methoxy-*N*-methylacetamide (0.831 g, 68%) as a white solid. Mp 62–64 °C; IR (neat) 3323, 2939, 1642, 1590, 1511, 1440, 1271, 1131, 1006, 761  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  3.18 (s, 3H), 3.61 (s, 3H), 3.67 (s, 2H), 3.85 (s, 3H), 5.85 (s, 1H), 6.73–6.82 (m, 2H), 6.87 (d,  $J = 1.3$  Hz, 1H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  32.3 ( $\text{CH}_3$ ), 38.7 ( $\text{CH}_2$ ), 56.0 ( $\text{CH}_3$ ), 61.3 ( $\text{CH}_3$ ), 110.8 (CH and C), 115.7 (CH), 120.7 (CH), 128.1 (C), 145.6 (C), 172.7 (C); MS (ESI)  $m/z$  248 ( $\text{M}+\text{Na}^+$ , 100); HRMS (ESI) calcd for  $\text{C}_{11}\text{H}_{15}\text{NNaO}_4$  ( $\text{M}+\text{Na}^+$ ) 248.0893, found 248.0889.

**1'-(3-*tert*-Butyldimethylsilyloxy-4-methoxyphenyl)-*N*-methoxy-*N*-methylacetamide (6).** To a solution of 1'-(3-hydroxy-4-methoxyphenyl)-*N*-methoxy-*N*-methylacetamide (1.23 g, 5.46 mmol) and

imidazole (0.740 g, 10.9 mmol) in dry dichloromethane (40 mL) was added *tert*-butyldimethylsilyl chloride (0.99 g, 6.56 mmol) portionwise. 4-Dimethylaminopyridine (0.070 g, 0.55 mmol) was added and the resulting suspension was stirred at room temperature for 16 h. The reaction was quenched with water (30 mL) and the mixture was extracted with dichloromethane (4 × 50 mL). The combined organic layers were washed with aqueous sodium hydrogen carbonate (100 mL) and brine (100 mL), dried (MgSO<sub>4</sub>) and concentrated *in vacuo*. Purification by flash column chromatography (petroleum ether/ethyl acetate, 4:1) gave 1'-(3-*tert*-butyldimethylsilyloxy-4-methoxyphenyl)-*N*-methoxy-*N*-methylacetamide (**6**) (1.87 g, 100%) as a colorless oil. IR (neat) 2932, 1663, 1512, 1271, 1136, 988, 839 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 0.15 (s, 6H), 0.99 (s, 9H), 3.18 (s, 3H), 3.58 (s, 3H), 3.65 (s, 2H), 3.78 (s, 3H), 6.78 (d, *J* = 8.2 Hz, 1H), 6.80 (d, *J* = 2.0 Hz, 1H), 6.83 (dd, *J* = 8.2, 2.0 Hz, 1H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ -4.6 (2 × CH<sub>3</sub>), 18.4 (C), 25.8 (3 × CH<sub>3</sub>), 32.8 (CH<sub>3</sub>), 38.8 (CH<sub>2</sub>), 55.6 (CH<sub>3</sub>), 61.3 (CH<sub>3</sub>), 112.2 (CH), 122.1 (CH), 122.3 (CH), 127.5 (C), 144.9 (C), 149.9 (C), 172.7 (C); MS (ESI) *m/z* 362 (M+Na<sup>+</sup>. 100); HRMS (ESI) calcd for C<sub>17</sub>H<sub>29</sub>NNaO<sub>4</sub>Si (M+Na<sup>+</sup>) 362.1758, found 362.1745.

**1'-(3-*tert*-Butyldimethylsilyloxy-4-methoxyphenyl)-2'-phenylethan-2'-one.** The reaction was carried out as described for 1'-(3-methoxyphenyl)-2'-phenylethan-2'-one using 1'-(3-*tert*-butyldimethylsilyloxy-4-methoxyphenyl)-*N*-methoxy-*N*-methylacetamide (**6**) (1.71 g, 5.03 mmol). Purification by flash column chromatography (petroleum ether/ethyl acetate, 4:1) gave 1'-(3-*tert*-butyldimethylsilyloxy-4-methoxyphenyl)-2'-phenylethan-2'-one (1.32 g, 74%) as a white solid. Mp 60–62 °C; IR (neat) 2930, 1680, 1510, 1271, 1136, 984, 839 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 0.12 (s, 6H), 0.97 (s, 9H), 3.77 (s, 3H), 4.16 (s, 2H), 6.76 (d, *J* = 1.4 Hz, 1H), 6.77–6.82 (m, 2H), 7.40–7.46 (m, 2H), 7.50–7.55 (m, 1H), 7.94–8.01 (m, 2H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ -4.6 (2 × CH<sub>3</sub>), 18.4 (C), 25.7 (3 × CH<sub>3</sub>), 45.0 (CH<sub>2</sub>), 55.5 (CH<sub>3</sub>), 112.4 (CH), 122.2 (CH), 122.5 (CH), 127.0 (C), 128.6 (2 × CH), 128.7 (2 × CH), 133.0 (CH), 136.7 (C), 145.1 (C), 149.9 (C), 197.9 (C); MS (ESI) *m/z* 379 (M+Na<sup>+</sup>. 100); HRMS (ESI) calcd for C<sub>21</sub>H<sub>28</sub>NaO<sub>3</sub>Si (M+Na<sup>+</sup>) 379.1700, found 379.1686.

**1'-(3-*tert*-Butyldimethylsilyloxy-4-methoxyphenyl)-1'-methyl-2'-phenylethan-2'-one (7).** An oven dried three-neck flask was flushed with argon and charged with 1'-(3-*tert*-butyldimethylsilyloxy-4-methoxyphenyl)-2'-phenylethan-2'-one (1.32 g, 3.69 mmol) in dry tetrahydrofuran (30 mL). To this solution was added lithium bis(trimethylsilyl)amide (4.06 mL, 4.06 mmol, 1.0 M in tetrahydrofuran) dropwise at  $-78\text{ }^{\circ}\text{C}$ . The reaction mixture was stirred at  $-78\text{ }^{\circ}\text{C}$  for 0.5 h before methyl iodide (0.690 mL, 11.1 mmol) was added dropwise. The resulting solution was stirred for 1 h, at  $-78\text{ }^{\circ}\text{C}$ , then slowly warmed to  $0\text{ }^{\circ}\text{C}$  and stirred for a further 1 h. A saturated solution of ammonium chloride (30 mL) was added at  $0\text{ }^{\circ}\text{C}$  and the solution was extracted with diethyl ether ( $3 \times 50\text{ mL}$ ). The combined ethereal extracts were washed with brine (100 mL), dried ( $\text{MgSO}_4$ ) and concentrated *in vacuo*. Purification by flash column chromatography (petroleum ether/ethyl acetate, 19:1) gave 1'-(3-*tert*-butyldimethylsilyloxy-4-methoxyphenyl)-1'-methyl-2'-phenylethan-2'-one (7) (1.19 g, 87%) as a colorless oil. IR (neat) 2930, 1684, 1506, 1275, 1138, 970, 837  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  0.09 (s, 6H), 0.96 (s, 9H), 1.48 (d,  $J = 6.8\text{ Hz}$ , 3H), 3.73 (s, 3H), 4.53 (q,  $J = 6.8\text{ Hz}$ , 1H), 6.74 (d,  $J = 8.5\text{ Hz}$ , 1H), 6.75 (d,  $J = 2.0\text{ Hz}$ , 1H), 6.79 (dd,  $J = 8.5, 2.0\text{ Hz}$ , 1H), 7.32–7.39 (m, 2H), 7.42–7.48 (m, 1H), 7.89–7.94 (m, 2H);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  -4.6 ( $2 \times \text{CH}_3$ ), 18.5 (C), 19.3 ( $\text{CH}_3$ ), 25.7 ( $3 \times \text{CH}_3$ ), 47.3 (CH), 55.5 ( $\text{CH}_3$ ), 112.6 (CH), 120.7 (CH), 120.8 (CH), 128.4 ( $2 \times \text{CH}$ ), 128.8 ( $2 \times \text{CH}$ ), 132.6 (CH), 134.1 (C), 136.6 (C), 145.3 (C), 149.9 (C), 200.4 (C); MS (ESI)  $m/z$  393 ( $\text{M}+\text{Na}^+$ , 100); HRMS (ESI) calcd for  $\text{C}_{22}\text{H}_{30}\text{NaO}_3\text{Si}$  ( $\text{M}+\text{Na}^+$ ) 393.1856, found 393.1848.

**(1'*R*,2'*R*)-1'-(3-*tert*-Butyldimethylsilyloxy-4-methoxyphenyl)-1'-methyl-2'-phenylethan-2'-ol (8).**

To an oven-dried conical flask was added potassium *tert*-butoxide (0.023 g, 0.16 mmol) and dichloro[(*S*)-(-)-5,5'-bis[di-(3,5-xylyl)phosphino]-4,4'-bi-1,3-benzodioxole][(2*S*-(+)-1,1-bis(4-methoxyphenyl)-3-methyl-1,2-butanediamine]ruthenium(II) (0.025 g, 0.012 mmol). Distilled 2-propanol (1 mL) was added and the resulting yellow solution was stirred at room temperature for 2 h under a constant stream of argon. 1'-(3-*tert*-Butyldimethylsilyloxy-4-methoxyphenyl)-1'-methyl-2'-phenylethan-2'-one (7) (0.270 g, 0.730 mmol) in 2-propanol (2 mL) was added to the conical flask containing the catalyst solution and

hydrogenated at 10 bar for 48 h. The reaction mixture was treated with activated carbon and stirred for 1 h. The mixture was filtered through Celite® and concentrated *in vacuo*. Purification by flash column chromatography (hexane/diethyl ether, 7:3) gave (1'*R*,2'*R*)-1'-(3-*tert*-butyldimethylsilyloxy-4-methoxyphenyl)-1'-methyl-2'-phenylethan-2'-ol (**8**) (0.192 g, 64%) as a colorless oil. IR (neat) 3454, 2929, 1508, 1275, 1139, 962, 836 cm<sup>-1</sup>; [ $\alpha$ ]<sub>D</sub><sup>23</sup> +34.4 (*c* 1.0, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  0.15 (s, 6H), 1.00 (s, 9H), 1.03 (d, *J* = 7.2 Hz, 3H), 1.90 (d, *J* = 1.4 Hz, 1H), 2.89 (dq, *J* = 8.6, 7.2 Hz, 1H), 3.80 (s, 3H), 4.55 (dd, *J* = 8.6, 1.4 Hz, 1H), 6.73–6.77 (m, 1H), 6.80–6.84 (m, 2H), 7.24–7.36 (m, 5H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$  -4.5 (2 × CH<sub>3</sub>), 18.3 (CH<sub>3</sub>), 18.5 (C), 25.8 (3 × CH<sub>3</sub>), 47.5 (CH), 55.6 (CH<sub>3</sub>), 79.7 (CH), 112.2 (CH), 120.6 (CH), 121.2 (CH), 127.0 (2 × CH), 127.7 (CH), 128.2 (2 × CH), 135.6 (C), 142.5 (C), 145.2 (C), 149.9 (C); MS (ESI) *m/z* 395 (M+Na<sup>+</sup>. 100); HRMS (ESI) calcd for C<sub>22</sub>H<sub>32</sub>NaO<sub>3</sub>Si (M+Na<sup>+</sup>) 395.2013, found 395.2023. Enantiomeric excess was determined by HPLC analysis, using a chiralcel® AD-H column (hexane:*i*-propanol 98:2, flow rate 1.0 mL min<sup>-1</sup>); *t*<sub>minor</sub> = 10.65 min, *t*<sub>major</sub> = 14.55 min, er = 2.5:97.5.

**(2*R*,3*R*)-5-(*tert*-Butyldimethylsilyloxy)-2,3-dihydro-6-methoxy-3-methyl-2-phenylbenzofuran (9).**

(2*R*,3*R*)-5-(*tert*-Butyldimethylsilyloxy)-2,3-dihydro-6-methoxy-3-methyl-2-phenylbenzofuran (**9**) was synthesized as described for 1-benzoyl-5-methoxyindoline (**2a**) using (1'*R*,2'*R*)-1'-(3-*tert*-butyldimethylsilyloxy-4-methoxyphenyl)-1'-methyl-2'-phenylethan-2'-ol (**8**) (0.060 g, 0.16 mmol). The iodination step was carried out at 50 °C for 7 h. Purification by flash column chromatography (hexane/dichloromethane, 3:2) gave (2*R*,3*R*)-5-(*tert*-butyldimethylsilyloxy)-2,3-dihydro-6-methoxy-3-methyl-2-phenylbenzofuran (**9**) (0.038 g, 63%) as a colorless oil. IR (neat) 2929, 1493, 1449, 1215, 1188, 1169, 904, 837 cm<sup>-1</sup>; [ $\alpha$ ]<sub>D</sub><sup>20</sup> +23.0 (*c* 1.0, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  0.14 (s, 3H), 0.14 (s, 3H), 1.00 (s, 9H), 1.37 (d, *J* = 6.7 Hz, 3H), 3.36 (dq, *J* = 8.9, 6.7 Hz, 1H), 3.77 (s, 3H), 5.11 (d, *J* = 8.9, 1H), 6.46 (s, 1H), 6.61 (s, 1H), 7.27–7.45 (m, 5H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$  -4.7 (CH<sub>3</sub>), -4.6 (CH<sub>3</sub>), 18.4 (CH<sub>3</sub>), 18.5 (C), 25.8 (3 × CH<sub>3</sub>), 45.6 (CH), 55.7 (CH<sub>3</sub>), 92.9 (CH), 95.0 (CH), 115.7 (CH), 122.4 (C), 126.1 (2 × CH), 128.1 (CH), 128.6 (2 × CH), 139.0 (C), 141.1 (C), 150.8 (C), 153.6 (C); MS

(EI)  $m/z$  370 ( $M^+$ . 61), 313 (47), 298 (100), 257 (33), 146 (32), 91 (37), 73 (40); HRMS (EI) calcd for  $C_{22}H_{30}O_3Si$  ( $M^+$ ) 370.1964, found 370.1955.

**(2*R*,3*R*)-2,3-Dihydro-5-hydroxyl-6-methoxy-3-methyl-2-phenylbenzofuran (10), [(+)-*Obtusafuran*].**<sup>26</sup> To a stirred solution of (2*R*,3*R*)-5-(*tert*-butyldimethylsilyloxy)-2,3-dihydro-6-methoxy-3-methyl-2-phenylbenzofuran (**9**) (0.038 g, 0.10 mmol) in dry tetrahydrofuran (10 mL) was added tetrabutylammonium fluoride solution (0.15 mL, 0.15 mmol; 1.0 M in tetrahydrofuran) at 0 °C. The resulting solution was stirred at room temperature for 2 h. The reaction mixture was diluted with diethyl ether (10 mL), washed with water (2 × 10 mL), dried ( $MgSO_4$ ) and concentrated *in vacuo*. Purification by flash column chromatography (hexane/diethyl ether, 7:3) gave (2*R*,3*R*)-2,3-dihydro-5-hydroxyl-6-methoxy-3-methyl-2-phenylbenzofuran (**10**) (0.023 g, 88%) as a white solid. Mp 108–110 °C (lit.<sup>26</sup> 111–113 °C);  $[\alpha]_D^{23} +48.2$  ( $c$  0.5, MeOH), lit.<sup>26</sup>  $[\alpha]_D^{25} +50.0$  ( $c$  0.33, MeOH);  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  1.37 (d,  $J = 6.9$  Hz, 3H), 3.38 (dq,  $J = 8.6, 6.9$  Hz, 1H), 3.87 (s, 3H), 5.11 (d,  $J = 8.6$  Hz, 1H), 5.24 (s, 1H), 6.50 (s, 1H), 6.72 (s, 1H), 7.29–7.44 (m, 5H);  $^{13}C$  NMR (101 MHz,  $CDCl_3$ )  $\delta$  18.4 ( $CH_3$ ), 45.7 (CH), 56.2 ( $CH_3$ ), 92.8 (CH), 94.2 (CH), 109.5 (CH), 122.9 (C), 126.0 (2 × CH), 128.1 (CH), 128.6 (2 × CH), 139.9 (C), 141.0 (C), 146.2 (C), 152.4 (C); MS (EI)  $m/z$  256 ( $M^+$ . 100), 239 (11), 165 (12), 91 (10).

**SUPPORTING INFORMATION AVAILABLE.** Computational reactivity analyses, synthetic routes to all one-pot substrates, HPLC traces for **8** and  $^1H$  and  $^{13}C$  NMR spectra for all compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

**ACKNOWLEDGEMENT.** Financial support from the EPSRC (studentship to M.C.H., EP/M508056/1) and the University of Glasgow is gratefully acknowledged.

**REFERENCES:**

- 1  
2  
3 (1) (a) Shen, T.; Wang, X.-N.; Lou, H.-X. Natural Stillbenes: An Overview. *Nat. Prod. Rep.* **2009**, *26*,  
4 916–935. (b) Xu, W.; Gavia, D. J.; Tang, Y. Biosynthesis of Fungal Indole Alkaloids. *Nat. Prod.*  
5  
6 *Rep.* **2014**, *31*, 1474–1487.  
7  
8  
9  
10  
11 (2) (a) Horton, D. A.; Bourne, G. T.; Smythe, M. L. The Combinatorial Synthesis of Bicyclic Privileged  
12 Structures or Privileged Substructures. *Chem. Rev.* **2003**, *103*, 893–930. (b) Anas, S.; Kagan, H. B.  
13 Routes Toward Enantiopure 2-Substituted Indolines: An Overview. *Tetrahedron: Asymmetry* **2009**,  
14 *20*, 2193–2199. (c) Zhang, M. Construction of Heterocycle Scaffolds via Transition Metal Catalyzed  
15  $sp^2$  C-H Functionalization. *Adv. Synth. Catal.* **2009**, *351*, 2243–2270. (d) Song, J. J.; Reeves, J. T.;  
16 Fandrick, D. R.; Tan, Z.; Yee, N. K.; Senanayake, C. H. Construction of the Indole Nucleus Through  
17 C-H Functionalization Reactions. *Arkivoc* **2010**, 390–449. (e) Yu, J.-T.; Pan, C. Radical C-H  
18 Functionalization to Construct Heterocyclic Compounds. *Chem. Commun.* **2016**, *52*, 2220–2236.  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31 (3) For representative indoline synthesis using Buchwald-Hartwig-type reactions, see: (a) Guram, A.  
32 S.; Rennels, R. A.; Buchwald, S. L. A Simple Catalytic Method for the Conversion of Aryl Bromides  
33 to Arylamines. *Angew. Chem. Int. Ed.* **1995**, *34*, 1348–1350. (b) Peat, A. J.; Buchwald, S. L. Novel  
34 Syntheses of Tetrahydropyrroloquinolines: Applications to Alkaloid Synthesis. *J. Am. Chem. Soc.*  
35 **1996**, *118*, 1028–1030. (c) Deboves, H. J. C.; Hunter, C.; Jackson, R. F. W. Synthesis of 2-  
36 Substituted Indolines Using Sequential Pd-Catalyzed Processes. *J. Chem. Soc., Perkin Trans. 1*  
37 **2002**, 733–736. (d) Omar-Amrani, R.; Schneider, R.; Fort, Y. Novel Synthetic Strategy of N-  
38 Arylated Heterocycles via Sequential Palladium-Catalyzed Intra- and Inter-Arylation  
39 Reactions. *Synthesis* **2004**, 2527–2534. (e) Anderson, J. C.; Noble, A.; Tocher, D. A. Reductive  
40 Nitro-Mannich Route for the Synthesis of 1,2-Diamine Containing Indolines and  
41 Tetrahydroquinolines. *J. Org. Chem.* **2012**, *77*, 6703–6727.  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- (4) For representative indoline synthesis using Ullmann-type reactions, see: (a) Kametani, T.; Ohsawa, T.; Ihara, M. A Novel Synthesis of Indole Derivatives. *Heterocycles* **1980**, *14*, 277–280. (b) Klapars, A.; Huang, X.; Buchwald, S. L. A General and Efficient Copper Catalyst for the Amidation of Aryl Halides. *J. Am. Chem. Soc.* **2002**, *124*, 7421–7428. (c) Yamada, K.; Kubo, T.; Tokuyama, H.; Fukuyama, T. A Mild Copper-Mediated Intramolecular Amination of Aryl Halides. *Synlett* **2002**, 231–234. (d) Minatti, A.; Buchwald, S. L. Synthesis of Indolines via a Domino Cu-Catalyzed Amidation/Cyclization Reaction. *Org. Lett.* **2008**, *10*, 2721–2724. (e) Kubo, T.; Katoh, C.; Yamada, K.; Okano, K.; Tokuyama, H.; Fukuyama, T. A Mild Inter- and Intramolecular Amination of Aryl Halides with a Combination of CuI and CsOAc. *Tetrahedron* **2008**, *64*, 11230–11236. (f) Liu, J.-Q.; Qian, C.; Chen, X.-Z. A Facile Chiral Pool Synthesis of (*S*)-6-Nitroindoline-2-carboxylic Acids from L-Phenylalanine. *Synthesis* **2010**, 403–406. (g) Lee, J.; Panek, J. S. Total Synthesis of (+)-Isatisine A. *Org. Lett.* **2011**, *13*, 502–505. (h) Cleghorn, L. A. T.; Albrecht, S.; Stojanovski, L.; Simeons, F. R. J.; Norval, S.; Kime, R.; Collie, I. T.; De Rycker, M.; Campbell, L.; Hallyburton, I.; Frearson, J. A.; Wyatt, P. G.; Read, K. D. Discovery of Indoline-2-carboxamide Derivatives as a New Class of Brain-Penetrant Inhibitors of *Trypanosoma brucei*. *J. Med. Chem.* **2015**, *58*, 7695–7706.
- (5) For representative dihydrobenzofuran synthesis using Ullmann-type reactions, see: (a) Zhu, J.; Price, B. A.; Zhao, S. X.; Skonezny, P. M. Copper(I)-Catalyzed Intramolecular Cyclization Reaction of 2-(2'-Chlorophenyl)ethanol to give 2,3-Dihydrobenzofuran. *Tetrahedron Lett.* **2000**, *41*, 4011–4014. (b) Adams, H.; Gilmore, N. J.; Jones, S.; Muldowney, M. P.; von Reuss, S. H.; Vemula, R. Asymmetric Synthesis of Corsifuran A by an Enantioselective Oxazaborolidine Reduction. *Org. Lett.* **2008**, *10*, 1457–1460. (c) Liu, J.; Fitzgerald, A. E.; Mani, N. S. Facile Assembly of Fused Benzo[4,5]furo Heterocycles. *J. Org. Chem.* **2008**, *73*, 2951–2954.

- (6) For other *N*- and *O*-heterocycle synthesis using Ullmann-type reactions, see: (a) Joyeau, R.; Yadav, L. D. S.; Wakselman, M. Synthesis of Benzocarbacephem and Benzocarbapenam Derivatives by Copper-Promoted Intramolecular Aromatic Substitution. *J. Chem. Soc., Perkin Trans. I* **1987**, 1899–1907. (b) Masse, C. E.; Ng, P. Y.; Fukase, Y.; Sanchez-Rosello, M.; Shaw, J. T. Divergent Structural Complexity from a Linear Reaction Sequence: Synthesis of Fused and Spirobicyclic  $\gamma$ -Lactams from Common Synthetic Precursors. *J. Comb. Chem.* **2006**, *8*, 293–296. (c) Yadav, L. D. S.; Yadav, B. S.; Rai, V. K. Active-Copper-Promoted Expeditious *N*-Arylations in Aqueous Media under Microwave Irradiation. *Synthesis* **2006**, 1868–1872. (d) Boonya-udtayan, S.; Yotapan, N.; Woo, C.; Bruns, C. J.; Ruchirawat, S.; Thasana, N. Synthesis and Biological Activities of Azalamellarins. *Chem. Asian. J.* **2010**, *5*, 2113–2123. (e) Sahn, J. J.; Martin, S. F. Facile Syntheses of Substituted, Conformationally-Constrained Benzoxazocines and Benzazocines via Sequential Multicomponent Assembly and Cyclization. *Tetrahedron Lett.* **2011**, *52*, 6855–6858.
- (7) For reviews, see: (a) Thansandote, P.; Lautens, M. Construction of Nitrogen-Containing Heterocycles by C-H Bond Functionalization. *Chem. Eur. J.* **2009**, *15*, 5874–5883. (b) Henry, M. C.; Mostafa, M. A. B.; Sutherland, A. Recent Advances in Transition Metal-Catalyzed, Directed Aryl C-H/N-H Cross-Coupling Reactions. *Synthesis* **2017**, *49*, 4586–4598. (c) Park, Y.; Kim, Y.; Chang, S. Transition Metal-Catalyzed C-H Amination: Scope, Mechanism and Applications. *Chem. Rev.* **2017**, *117*, 9247–9301.
- (8) (a) Li, J.-J.; Mei, T.-S.; Yu, J.-Q. Synthesis of Indolines and Tetrahydroisoquinolines from Arylethylamines by Pd<sup>II</sup>-Catalyzed C-H Activation Reactions. *Angew. Chem. Int. Ed.* **2008**, *47*, 6452–6455. (b) He, G.; Zhao, Y.; Zhang, S.; Lu, C.; Chen, G. Highly Efficient Syntheses of Azetidines, Pyrrolidines and Indolines via Palladium Catalyzed Intramolecular Amination of C(sp<sup>3</sup>)-H and C(sp<sup>2</sup>)-H at  $\gamma$  and  $\delta$  Positions. *J. Am. Chem. Soc.* **2012**, *134*, 3–6. (c) Nadres, E. T.; Daugulis, O. Heterocycle Synthesis via Direct C-H/N-H Coupling. *J. Am. Chem. Soc.* **2012**, *134*, 7–

10. (d) He, G.; Lu, C.; Zhao, Y.; Nack, W. A.; Chen, G. Improved Protocol for Indoline Synthesis via Palladium-Catalyzed Intramolecular C(sp<sup>2</sup>)-H Amination. *Org. Lett.* **2012**, *14*, 2944–2947. (e) Mei, T.-S.; Leow, D.; Xiao, H.; Laforteza, B. N.; Yu, J.-Q. Synthesis of Indolines via Pd(II)-Catalyzed Amination of C-H Bonds Using PhI(OAc)<sub>2</sub> as the Bystanding Oxidant. *Org. Lett.* **2013**, *15*, 3058–3061. (f) Ye, X.; He, Z.; Ahmed, T.; Weise, K.; Akhmedov, N. G.; Petersen, J. L.; Shi, X. 1,2,3-Triazoles as Versatile Directing Group for Selective sp<sup>2</sup> and sp<sup>3</sup> C-H Activation: Cyclization vs Substitution. *Chem. Sci.* **2013**, *4*, 3712–3716. (g) Wang, C.; Chen, C.; Zhang, J.; Han, J.; Wang, Q.; Guo, K.; Liu, P.; Guan, M.; Yao, Y.; Zhao, Y. Easily Accessible Auxiliary for Palladium-Catalyzed Intramolecular Amination of C(sp<sup>2</sup>)-H and C(sp<sup>3</sup>)-H Bonds at δ- and ε-Positions. *Angew. Chem. Int. Ed.* **2014**, *53*, 9884–9888. (h) He, Y.-P.; Zhang, C.; Fan, M.; Wu, Z.; Ma, D. Assembly of Indoline-2-Carboxylate-Embodied Dipeptides via Pd-Catalyzed C(sp<sup>2</sup>)-H Bond Direct Functionalization. *Org. Lett.* **2015**, *17*, 496–499. (i) Zheng, Y.; Song, W.; Zhu, Y.; Wei, B.; Xuan, L. Pd-Catalyzed Intramolecular C(sp<sup>2</sup>)-H Amination of Phenylalanine Moieties in Dipeptides: Synthesis of Indoline-2-Carboxylate-Containing Dipeptides. *Org. Biomol. Chem.* **2018**, *16*, 2402–2405.
- (9) A similar strategy using copper catalysis has also been reported: Takamatsu, K.; Hirano, K.; Satoh, T.; Miura, M. Synthesis of Indolines by Copper-Mediated Intramolecular Aromatic C-H Amination. *J. Org. Chem.* **2015**, *80*, 3242–3249.
- (10) A mechanistically distinct, metal-free method involving oxidation and activation of the aryl ring before cyclization has also been reported for the synthesis of indolines: Pouységu, L.; Avellan, A.-V.; Quideau, S. Iodine(III)-Mediated Generation of Nitrogen Tethered Orthoquinol Acetates for the Construction of Oxygenated Indole, Quinoline and Phenanthridine Alkaloid Motifs. *J. Org. Chem.* **2002**, *67*, 3425–3436.

- 1 (11) (a) Wang, X.; Liu, Y.; Dai, H. X.; Yu, J.-Q. Pd(II)-Catalyzed Hydroxyl-Directed C-H  
2 Activation/C-O Cyclization: Expedient Construction of Dihydrobenzofurans. *J. Am. Chem. Soc.*  
3 **2010**, *132*, 12203–12205. (b) Wang, H.; Li, G.; Engle, K. M.; Yu, J.-Q.; Davies, H. W. Sequential  
4 C-H Functionalization Reactions for the Enantioselective Synthesis of Highly Functionalized 2,3-  
5 Dihydrobenzofurans. *J. Am. Chem. Soc.* **2013**, *135*, 6774–6777.  
6  
7  
8  
9  
10  
11  
12  
13 (12) Alvarado, J.; Fournier, J.; Zakarian, A. Synthesis of Functionalized Dihydrobenzofurans by Direct  
14 Aryl C-O Bond Formation Under Mild Conditions. *Angew. Chem. Int. Ed.* **2016**, *55*, 11625–11628.  
15  
16  
17  
18  
19 (13) Antoniotti, S.; Dalla, V.; Duñach, E. Metal Triflimidates: Better Than Metal Triflates as Catalysts  
20 in Organic Synthesis- The Effect of a Highly Delocalized Counterion. *Angew. Chem. Int. Ed.* **2010**,  
21 *49*, 7860–7888.  
22  
23  
24  
25  
26 (14) (a) Racys, D. T.; Warrilow, C. E.; Pimlott, S. L.; Sutherland, A. Highly Regioselective Iodination  
27 of Arenes via Iron(III)-Catalyzed Activation of *N*-Iodosuccinimide. *Org. Lett.* **2015**, *17*, 4782–4785.  
28  
29 (b) Racys, D. T.; Sharif, S. A. I.; Pimlott, S. L.; Sutherland, A. Silver(I)-Catalyzed Iodination of  
30 Arenes: Tuning the Lewis Acidity of *N*-Iodosuccinimide Activation. *J. Org. Chem.* **2016**, *81*, 772–  
31 780. (c) Mostafa, M. A. B.; Bowley, R. M.; Racys, D. T.; Henry, M. C.; Sutherland, A. Iron(III)-  
32 Catalyzed Chlorination of Activated Arenes. *J. Org. Chem.* **2017**, *82*, 7529–7537.  
33  
34  
35  
36  
37  
38  
39  
40  
41 (15) Mostafa, M. A. B.; Calder, E. D. D.; Racys, D. T.; Sutherland, A. Intermolecular Aryl C-H  
42 Amination Through Sequential Iron and Copper Catalysis. *Chem. Eur. J.* **2017**, *23*, 1044–1047.  
43  
44  
45  
46 (16) The scope of iron-catalyzed iodination is limited by the electronics of the aryl ring system. As  
47 previously reported by us (see reference 14a), aryl compounds bearing only electron-withdrawing  
48 groups are not activated enough to undergo iodination. At least one electron-rich, activating group  
49 is required for the reaction to proceed. Conversely, some phenols were found to be too activated and  
50 led to poly-iodinated products. For this substrate class, less active Lewis acid catalysts are required  
51 (see reference 14b).  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- (17) For other examples of tetrahydroquinoline synthesis via arene C-H reactions, see: (a) Togo, H.; Hoshina, Y.; Muraki, T.; Nakayama, H.; Yokoyama, M. Study on Radical Amidation onto Aromatic Rings with (Diacloxyiodo)arenes. *J. Org. Chem.* **1998**, *63*, 5193–5200. (b) Yang, M.; Su, B.; Wang, Y.; Chen, K.; Jiang, X.; Zhang, Y.-F.; Zhang, X.-S.; Chen, G.; Cheng, Y.; Cao, Z.; Guo, Q.-Y.; Wang, L.; Shi, Z.-J. Silver-Catalyzed Direct Amination of Unactivated C-H Bonds of Functionalized Molecules. *Nat. Commun.* **2014**, *5*, 4707. (c) Cosgrove, S. C.; Plane, J. M. C.; Marsden, S. P. Radical-Mediated Direct C-H Amination of Arenes with Secondary Amines. *Chem. Sci.* **2018**, *9*, 6647–6652.
- (18) von Reuss, S. H.; König, W. A. Corsifurans A-C, 2-Arylbenzofurans of Presumed Stilbenoid Origin from *Corsinia coriandrina* (Hepaticae). *Phytochemistry* **2004**, *65*, 3113–3118.
- (19) While *p*-iodination of the 3-methoxyphenyl ring was the main product, *o*-iodination of the side-chain 4-methoxyphenyl group was also observed by NMR spectroscopy.
- (20) (a) Gregson, M.; Ollis, W. D.; Redman, B. T.; Sutherland, I. O.; Dietrichs, H. H. The Constitution and Stereochemistry of Obtusafuran. *Chem. Commun.* **1968**, 1394–1395. (b) Gregson, M.; Ollis, W. D.; Redman, B. T.; Sutherland, I. O.; Dietrichs, H. H.; Gottlieb, O. R. Obtusastylene and Obtustylene, Cinnamylphenols from *Dalbergia retusa*. *Phytochemistry* **1978**, *17*, 1395–1400.
- (21) Beldjoudi, N.; Mambu, L.; Labaïed, M.; Grellier, P.; Ramanitrahasimbola, D.; Rasoanaivo, P.; Martin, M. T.; Frappier, F. Flavonoids from *Dalbergia louvelii* and Their Antiplasmodial Activity. *J. Nat. Prod.* **2003**, *66*, 1447–1450.
- (22) An, R.-B.; Jeong, G.-S.; Kim, Y.-C. Flavonoids from the Heartwood of *Dalbergia odorifera* and Their Protective Effect on Glutamate-Induced Oxidative Injury in HT22 Cells. *Chem. Pharm. Bull.* **2008**, *56*, 1722–1724.

- (23) Liu, R.-H.; Mei, D.-Y.; Lin, S.; Wang, D.-Q.; Shao, F.; Chen, L.-Y.; Guo, S.-L. A New Benzofuran from the Heartwood of *Dalbergia latifolia*. *Nat. Prod. Res.* **2018**, DOI:10.1080/14786419.2018.1431639.
- (24) Yin, H.-Q.; Lee, B.-W.; Kim, Y.-C.; Sohn, D.-H.; Lee, B.-H. Induction of the Anticarcinogenic Marker Enzyme, Quinone Reductase, by *Dalbergiae Lignum*. *Arch. Pharm. Res.* **2004**, *27*, 919–922.
- (25) (a) Jurd, L.; Manners, G.; Stevens, K. Isolation and Synthesis of (±)-Obtusafuran. *Chem. Commun.* **1972**, 992–993. (b) Jurd, L.; Stevens, K.; Manners, G. Acid-Catalyzed and Thermal Rearrangements of Obtusaquinol and Related 3,3-Diarylpropenes. *Tetrahedron* **1973**, *29*, 2347–2353.
- (26) Chen, C.-y.; Weisel, M. Concise Asymmetric Synthesis of (+)-Conocarpan and Obtusafuran. *Synlett* **2013**, *24*, 189–192.
- (27) Chen, C.-y.; Frey, L. F.; Shultz, S.; Wallace, D. J.; Marcantonio, K.; Payack, J. F.; Vazquez, E.; Springfield, S. A.; Zhou, G.; Liu, P.; Kieczkowki, G. R.; Chen, A. M.; Phenix, B. D.; Singh, U.; Strine, J.; Izzo, B.; Krska, S. W. Catalytic, Enantioselective Synthesis of Taranabant, a Novel, Acyclic Cannabinoid-1 Receptor Inverse Agonist for the Treatment of Obesity. *Org. Process Res. Dev.* **2007**, *11*, 616–623.
- (28) Chung, J. Y. L.; Mancheno, D.; Dormer, P. G.; Variankaval, N.; Ball, R. G.; Tsou, N. N. Diastereoselective Friedel-Crafts Alkylation of Indoles with Chiral  $\alpha$ -Phenyl Benzylic Cations. Asymmetric Synthesis of *Anti*-1,1,2-Triarylalkanes. *Org. Lett.* **2008**, *10*, 3037–3040.
- (29) Ohkuma, T.; Ooka, H.; Yamakawa, M.; Ikariya, T.; Noyori, R. Stereoselective Hydrogenation of Simple Ketones Catalyzed by Ruthenium(II)-Complexes. *J. Org. Chem.* **1996**, *61*, 4872–4873.
- (30) For chiral HPLC of **8**, see supporting information.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- (31) Leblanc, Y.; Boudreault, N. Para-Directed Amination of Electron-Rich Arenes with Bis(2,2,2-trichloroethyl) Azodicarboxylate. *J. Org. Chem.* **1995**, *60*, 4268–4271.
- (32) Hirshfeld, F. L. Bonded-Atom Fragments for Describing Molecular Charge Densities. *Theor. Chim. Acta* **1977**, *44*, 129–138.
- (33) Parr, R. G.; Yang, W. *Density-Functional Theory of Atoms and Molecules*, Oxford University Press, New York, **1989**, Sect. 5.4.
- (34) Parr, R. G.; Yang, W. Density Functional Approach to the Frontier-Electron Theory of Chemical Reactivity. *J. Am. Chem. Soc.* **1984**, *106*, 4049–4050.
- (35) Morell, C.; Grand, A.; Toro-Labbé, A. New Dual Descriptor for Chemical Reactivity. *J. Phys. Chem. A* **2005**, *109*, 205–212.
- (36) Jakubec, P.; Cockfield, D. M.; Hynes, P. S.; Cleator, E.; Dixon, D. J. Enantio- and Diastereoselective Michael Additions of C-Succinimidyl Esters to Nitro Olefins Using Cinchonine-Derived Bifunctional Organocatalysts. *Tetrahedron: Asymmetry* **2011**, *22*, 1147–1155.
- (37) Sathish, M.; Chetna, J.; Hari Krishna, N.; Shankaraiah, N.; Alarifi, A.; Kamal, A. Iron-Mediated One-Pot Synthesis of 3,5-Diarylpyridines from  $\beta$ -Nitrostyrenes. *J. Org. Chem.* **2016**, *81*, 2159–2165.
- (38) Limaye, R. A.; Joseph, A. R.; Natu, A. D.; Paradkar, M. V. Convenient Syntheses of 3-Aryl-3,4-Dihydroisocoumarins. *J. Chem. Res.* **2015**, *39*, 191–194.
- (39) Irie, H.; Matsumoto, R.; Nishimura, M.; Zhang, Y. Synthesis of ( $\pm$ )-Heritol, a Sesquiterpene Lactone Belonging to the Aromatic Cadinane Group. *Chem. Pharm. Bull.* **1990**, *38*, 1852–1856.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- (40) Jalal, S.; Sarkar, S.; Bera, K.; Maiti, S.; Jana, U. Synthesis of Nitroalkenes Involving a Cooperative Catalytic Action of Iron(III) and Piperidine: A One-Pot Synthetic Strategy to 3-Alkylindoles, 2H-Chromenes and N-Arylpyrrole. *Eur. J. Org. Chem.* **2013**, 4823–4828.
- (41) Maity, S.; Manna, S.; Rana, S.; Naveen, T.; Mallick, A.; Maiti, D. Efficient and Stereoselective Nitration of Mono- and Disubstituted Olefins with AgNO<sub>2</sub> and TEMPO. *J. Am. Chem. Soc.* **2013**, *135*, 3355–3358.
- (42) Yang, S. H.; Song, C.-H.; Van, H. T. M.; Park, E.; Khadka, D. B.; Gong, E.-Y.; Lee, K.; Cho, W.-J. SAR Based Design of Nicotinamides as a Novel Class of Androgen Receptor Antagonists. *J. Med. Chem.* **2013**, *56*, 3414–3418.
- (43) Minor, D. L.; Wyrick, S. D.; Charifson, P. S.; Watts, V. J.; Nichols, D. E.; Mailman, R. B. Synthesis and Molecular Modeling of 1-Phenyl-1,2,3,4-Tetrahydroisoquinolines and Related 5,6,8,9-Tetrahydro-13bH-dibenzo[a,h]quinolizines as D1 Dopamine Antagonists. *J. Med. Chem.* **1994**, *37*, 4317–4328.
- (44) Han, J.; Liu, P.; Wang, C.; Wang, Q.; Zhang, J.; Zhao, Y.; Shi, D.; Huang, Z.; Zhao, Y. Oxalyl Amide Assisted Palladium-Catalyzed Arylation of C(sp<sup>2</sup>)-H Bond at the  $\delta$  Position. *Org. Lett.* **2014**, *16*, 5682–5685.
- (45) Molander, G. A.; Jean-Gérard, L. Scope of the Suzuki-Miyaura Aminoethylation Reaction Using Organotrifluoroborates. *J. Org. Chem.* **2007**, *72*, 8422–8426.
- (46) In, J.; Hwang, S.; Kim, C.; Seo, J. H.; Kim, S. Synthesis of 3,4-Dihydroisoquinolin-1-ones from N-Boc-( $\beta$ -Arylethyl)carbamates via Isocyanate Intermediates. *Eur. J. Org. Chem.* **2013**, 965–971.



- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- (47) Kinderman, S. S.; Wekking, M. M. T.; van Maarseveen, J. H.; Schoemaker, H. E.; Hiemstra, H.; Rutjes, F. P. J. T. Catalytic N-Sulfonyliminium Ion-Mediated Cyclizations to  $\alpha$ -Vinyl-Substituted Isoquinolines and  $\beta$ -Carbolines and Applications in Metathesis. *J. Org. Chem.* **2005**, *70*, 5519–5527.
- (48) Kiruthika, S. E.; Perumal, P. T. CuI-Catalyzed Coupling of *gem*-Dibromovinylanilides and Sulfonamides: An Efficient Method for the Synthesis of 2-Amidoindoles and Indolo[1,2-*a*]quinazolines. *Org. Lett.* **2014**, *16*, 484–487.
- (49) Kargbo, R. B.; Sajjadi-Hashemi, Z.; Roy, S.; Jin, X.; Herr, R. J. Synthesis of 3-Benzazepines and Azepino[4,5-*b*]heterocyclic Ring Systems via Intramolecular Friedel-Crafts Cyclization. *Tetrahedron. Lett.* **2013**, *54*, 2018–2021.
- (50) Silveira, C. C.; Bernardi, C. R.; Braga, A. L.; Kaufman, T. S. Pictet-Spengler Condensation of *N*-Sulfonyl- $\beta$ -Phenethylamines with  $\alpha$ -Chloro- $\alpha$ -phenylselenoesters. New Synthesis of 1,2,3,4-Tetrahydroisoquinoline-1-carboxylates. *Tetrahedron. Lett.* **1999**, *40*, 4969–4972.
- (51) McKillop, A.; Davies, H. M. L.; Taylor, E. C. Thallium in Organic Synthesis. 67 Intramolecular Capture of Aromatic Radical Cations by An *N*-Tosyl Group. *Synth. Commun.* **1986**, *16*, 267–281.
- (52) Bonne, D.; Dekhane, M.; Zhu, J. Mild Oxidative One-Carbon Homologation of Aldehyde to Amide. *J. Am. Chem. Soc.* **2005**, *127*, 6926–6927.
- (53) Franzmann, P.; Trosien, S.; Schubert, M.; Waldvogel, S. R. Modular Approach to 9-Monosubstituted Fluorene Derivatives Using Mo<sup>V</sup> Reagents. *Org. Lett.* **2016**, *18*, 1182–1185.
- (54) Nishikata, T.; Nagashima, H. N Alkylation of Tosylamides Using Esters as Primary and Tertiary Alkyl Sources: Mediated by Hydrosilanes Activated by a Ruthenium Catalyst. *Angew. Chem. Int. Ed.* **2012**, *51*, 5363–5366.

- 1 (55) Jin, N.; Pan, C.; Zhang, H.; Xu, P.; Cheng, Y.; Zhu, C. Rhodium-Catalyzed Direct C7  
2 Alkynylation of Indolines. *Adv. Synth. Catal.* **2015**, *357*, 1149–1153.  
3  
4  
5  
6 (56) Iwao, M.; Kuraishi, T. Directed Lithiation of 1-(*tert*-Butoxycarbonyl)indolines. A Convenient  
7 Route to 7-Substituted Indolines. *Heterocycles* **1992**, *34*, 1031–1038.  
8  
9  
10  
11 (57) Jhan, Y.-H.; Kang, T.-W.; Hsieh, J.-C. Efficient Copper-Catalysed Intramolecular *N*-Arylation for  
12 the Synthesis of Oxindoles. *Tetrahedron. Lett.* **2013**, *54*, 1155–1159.  
13  
14  
15  
16  
17 (58) Shahane, S.; Louafi, F.; Moreau, J.; Hurvois, J.-P.; Renaud, J.-L.; van de Weghe, P.; Roisnel, T.  
18 Synthesis of Alkaloids of *Galipea officinalis* by Alkylation of an  $\alpha$ -Amino Nitrile. *Eur. J. Org.*  
19 *Chem.* **2008**, 4622–4631.  
20  
21  
22  
23  
24  
25 (59) Yoshinaga, H.; Masumoto, S.; Koyama, K.; Kinomura, N.; Matsumoto, Y.; Kato, T.; Baba, S.;  
26 Matsumoto, K.; Horisawa, T.; Oki, H.; Yabuuchi, K.; Kodo, T. Discovery of SMP-304, a Novel  
27 Benzylpiperidine Derivative with Serotonin Transporter Inhibitory Activity and 5-HT1A Weak  
28 Partial Agonistic Activity Showing the Antidepressant-Like Effect. *Bioorg. Med. Chem.* **2017**, *25*,  
29 293–304.  
30  
31  
32  
33  
34  
35  
36  
37 (60) Moreau, E.; Fortin, S.; Desjardins, M.; Rousseau, J. L. C.; Petitclerc, E.; Gaudreault, R. C.  
38 Optimized *N*-Phenyl-*N'*-(2-Chloroethyl)ureas as Potential Antineoplastic Agents: Synthesis and  
39 Growth Inhibition Activity. *Bioorg. Med. Chem.* **2005**, *13*, 6703–6712.  
40  
41  
42  
43  
44  
45 (61) Pflueger, J. J.; Morrill, L. C.; deGruyter, J. N.; Perea, M. A.; Sarpong, R. Magnesiato  
46 Addition/Ring-Expansion Strategy To Access the 6-7-6 Tricyclic Core of Hetsine-Type C<sub>20</sub>-  
47 Diterpenoid Alkaloids. *Org. Lett.* **2017**, *19*, 4632–4635.  
48  
49  
50  
51  
52  
53 (62) Wang, H.; Denton, J. R.; Davies, H. M. L. Sequential Rhodium-, Silver-, and Gold-Catalyzed  
54 Synthesis of Fused Dihydrofurans. *Org. Lett.* **2011**, *13*, 4316–4319.  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- (63) Fu, W. C.; So, C. M.; Yuen, O. Y.; Lee, I. T. C.; Kwong, F. Y. Exploiting Aryl Mesylates and Tosylates in Catalytic Mono- $\alpha$ -arylation of Aryl- and Heteroarylketones. *Org. Lett.* **2016**, *18*, 1872–1875.
- (64) Suh, Y.; Lee, J.-s.; Kim, S.-H.; Rieke, R. D. Direct Preparation of Benzylic Manganese Reagents from Benzylic Halides, Sulfonates and Phosphonates and their Reactions: Applications in Organic Synthesis. *J. Organomet. Chem.* **2003**, *684*, 20–36.
- (65) Spoehrle, S. S. M.; West, T. H.; Taylor, J. E.; Slawin, A. M. Z.; Smith, A. D. Tandem Palladium and Isothiourea Relay Catalysis: Enantioselective Synthesis of  $\alpha$ -Amino Acid Derivatives via Allylic Amination and [2,3]-Sigmatropic Rearrangement. *J. Am. Chem. Soc.* **2017**, *139*, 11895–11902.
- (66) Das, M.; Manvar, A.; Fox, I.; Roberts, D. J.; O'Shea, D. F.  $\text{Bu}_4\text{N}^+$ -Controlled Addition and Olefination with Ethyl 2-(Trimethylsilyl)acetate via Silicon Activation. *Synlett* **2017**, *18*, 2401–2406.
- (67) Paudyal, M. P.; Adebessin, A. M.; Burt, S. R.; Ess, D. H.; Ma, Z.; Kürti, L.; Falck, J. R. Dirhodium-Catalyzed C-H Arene Amination using Hydroxylamines. *Science* **2016**, *353*, 1144–1147.
- (68) Laurent, M. Y.; Stocker, V.; Temgoua, V. M.; Dujardin, G.; Dhal, R. New Two-Step Sequence Involving a Hetero-Diels-Alder and a Nonphenolic Oxidative Coupling Reaction: A Convergent Access to Analogs of Steganacin. *Tetrahedron Lett.* **2011**, *52*, 1608–1611.
- (69) Hartz, R. A.; Ahuja, V. T.; Rafalski, M.; Schmitz, W. D.; Brenner, A. B.; Denhart, D. J.; Ditta, J. L.; Deskus, J. A.; Yue, E. W.; Arvanitis, A. G.; Lelas, S.; Li, Y.-W.; Molski, T. F.; Wong, H.; Grace, J. E.; Lentz, K. A.; Li, J.; Lodge, N. J.; Zaczek, R.; Combs, A. P.; Olson, R. E.; Mattson, R. J.; Bronson, J. J.; Macor, J. E. In Vitro Intrinsic Clearance-Based Optimization of  $N^3$ -

1 Phenylpyrazinones as Corticotropin-Releasing Factor-1 (CRF<sub>1</sub>) Receptor Antagonists, *J. Med.*  
2 *Chem.* **2009**, *52*, 4161–4172.  
3  
4

5  
6 (70) Hu, Y.; Kamitanaka, T.; Mishima, Y.; Dohi, T.; Kita, Y. Brønsted Acid-Controlled [3 + 2]  
7  
8 Coupling Reactions of Quinone Monoacetals with Alkene Nucleophiles: A Catalytic System of  
9  
10 Perfluorinated Acids and Hydrogen Bond Donor for the Construction of Benzofurans. *J. Org. Chem.*  
11  
12 **2013**, *78*, 5530–5543.  
13  
14

15  
16 (71) Hata, K.; Hamamoto, H.; Shiozaki, Y.; Cämmerer, S. B.; Kita, Y. Nucleophilic Attack of  
17  
18 Intramolecular Hydroxyl Groups on Electron-Rich Aromatics Using Hypervalent Iodine(III)  
19  
20 Oxidation. *Tetrahedron* **2007**, *63*, 4052–4060.  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60