

## STANNOBOROXANES

### II\*. SYNTHESIS AND REACTIONS OF TRIS(TRIALKYL TIN) BORATES

S K MEHROTRA, G. SRIVASTAVA and R C. MEHROTRA

*The Chemical Laboratories, University of Rajasthan Jaipur 302004 (India)*

(Received June 15th, 1973)

#### Summary

Tris(trialkyltin) borates  $B(OSnR_3)_3$  ( $R = Me, Et, Pr, Bu, i-Bu$  and  $Ph$ ) have been prepared by esterification of boric acid with trialkyltin hydroxide or bis(trialkyltin) oxide. These reacted with boric anhydride to give the corresponding trialkyltin metaborates  $(R_3SnOBO)_3$ . Reactions of tris(tributyltin) borate with hydrogen chloride, acetyl chloride, acetic acid, acetic anhydride, phenol, butanethiol, thiophenol and isopropyl acetate have been studied. The infrared spectra of these compounds are discussed


---

#### Introduction

In the first part of this series [1], we described the synthesis of 2-(tributylstannoxy)-1,3,2-dioxaborolanes, boroles and borinanes by various routes. The present paper deals with some new tris(trialkyltin) borates

#### Results and discussion

The new compounds were prepared in quantitative yields by the following general route:



Water formed during the reactions was fractionated out azeotropically with benzene. The completion of the reactions was confirmed by the complete disappearance of Sn—O—Sn or OH peak and also by the formation of a new broad band due to B—O—Sn in the range 1270-1300  $cm^{-1}$  range [1]. The

---

\* For part I see ref 1

TABLE 1  
PREPARATION OF TRIS (TRIALKYL(TIN) BORATES

Organotin hydroxide or oxide (g)/(mmoles)	Boric acid (g) (mmoles)	Product	Yield (%)	B p [°C(mm)]	$\nu_{\text{as}}(\text{BO-Sn})$	Analysis (%)		Calcd			
						Found	Found	B	C	H	Sn
4.1 <sup>c</sup> (22.69)	0.47 (7.58)	B(OSnMe <sub>3</sub> ) <sub>3</sub>	89	150-132(1.4)	1300-1285	2.1	20.8	5.2	1.96	19.6	5.0
3.4 <sup>d</sup> (15.26)	0.34 (5.48)	B(OSnEt <sub>3</sub> ) <sub>3</sub>	70	140-145(0.5)	1300-1286	1.4			1.6		
3.2 (9.01)	0.25 (4.03)	B(OSnPr <sub>3</sub> ) <sub>3</sub>	98	152-153(0.1)	1290-75	1.4	40.5	7.9	1.4	40.4	7.9
17.2 (28.89)	1.22 (19.68)	B(OSnBu <sub>3</sub> ) <sub>3</sub>	91	195-200(0.1)	1290-75	1.2			1.2	38.0	38.1
3.3 (5.64)	0.22 (3.55)	B(OSnBu-t <sub>3</sub> ) <sub>3</sub>	99	180-185(0.05)	1290-70	1.1	47.4	8.7	1.2	46.6	8.8
2.8 <sup>c</sup> (7.64)	0.16 (2.58)	B(OSnPh <sub>3</sub> ) <sub>3</sub>	98	128-130 <sup>b</sup>	1270	0.98	57.8	4.0	0.98	58.5	4.1

<sup>a</sup> In all these cases respective hydroxides are taken <sup>b</sup> The compound does not distill, so m p is reported


TABLE 2  
PREPARATION OF TRIALKYLTIN METABORATES

B(OSnR <sub>3</sub> ) (g) (mmoles)	B <sub>2</sub> O <sub>3</sub> (g) (mmoles)	Temp (°C) (h)	Product	ν <sub>as</sub> (BOSn) (cm <sup>-1</sup> )	Metaborate peak (cm <sup>-1</sup> )	Boron found (calcd.) (%)
0.84 (1.26)	0.09 (1.29)	100 (1)	(OBOSnEt <sub>3</sub> ) <sub>3</sub>	1295	732m 742m	4.2 (4.4)
0.92 (1.15)	0.09 (1.29)	140 (3)	(OBOSnPr <sub>3</sub> ) <sub>3</sub>	1295	732m 742m	3.7 (3.8)
2.68 (2.88)	0.20 (2.87)	240 (3)	(OBOSnBu <sub>3</sub> ) <sub>3</sub>	1296	730m 742m	3.3 (3.3)
0.67 (0.72)	0.05 (0.72)	180 (4)	(OBOSnBu-1 <sub>3</sub> ) <sub>3</sub>	1290	732vs. 744s	3.1 (3.3)


reactions are complete within 4-5 hours. On distillation under reduced pressure, these compounds undergo slight disproportionation into bis(trialkyltin) oxide and boric anhydride, as is demonstrated by the appearance of a weak Sn—O—Sn peak in the range 740-84 cm<sup>-1</sup>.

Data for the tris(trialkyltin) borates are presented in Table 1.



On heating tris(trialkyltin) borates with boric anhydride, the latter slowly dissolves with the formation of trialkyltin metaborate (Table 2), and the temperature of dissolution increases from 120-240° from ethyl to butyl derivatives. The ease of such group exchange, leading to metaborates, appears to increase in the following order.




Like tris(trialkyltin) borates, trialkyltin metaborates may also be prepared directly from bis(trialkyltin) oxide and boric anhydride. These are colourless, viscous liquids, miscible with common organic solvents. Like other metaborate esters they have high thermal stability, but disproportionate into orthoborates on attempted distillation under reduced pressure:



Reactions of tris(tributyltin) borate with hydrogen chloride and acetyl chloride are exothermic, and give tributyltin chloride.



Reactions of tris(tributyltin) borate with acetic acid, phenol, butanethiol and thiofenol also proceed with the cleavage of tin-oxygen bond and the negative group attaches to the organotin moiety.




A = CH<sub>3</sub>COO, SBu, SPh, OPh

It is advantageous to carry out these reactions in benzene medium, as one of the products, boric acid, is precipitated out, and can be easily separated by filtration. All these reactions are quite fast, as shown by their exothermic nature, but to ensure completion, the reaction mixtures were kept overnight at room temperature.

Tris(tributyltin) borate also reacts exothermically with acetic anhydride with the formation of tributyltin acetate and boron triacetate (or diboron tetraacetate oxide), but it is rather difficult to separate the products, fractional crystallization from benzene gave tributyltin acetate in 50% yield.

Analogous reactions of trialkyltin alkoxides,  $R_3SnOR$  with  $AcCl$  [5],  $RSH$  [6], acetic acid [7], have been previously reported. Qualitatively, there does not appear any appreciable difference in the reactivities of  $Sn-O-C$  and  $Sn-O-B$  bonds towards these reagents. However, there is a notable difference towards alkyl acetates. In the case of former, transesterification is facile, but there appears to be no reaction with the latter. Because of ( $p \rightarrow p$ ) $\pi$  bonding between boron and oxygen, the nucleophilicity of boroxy-oxygen may be somewhat reduced, and the following intermediate necessary for transesterification is less likely to be formed:



It should be mentioned that trialkyl borates likewise fail to react with esters under normal conditions [8].

#### IR and NMR spectra

Like tris(tributyltin) borate [1], other tris(trialkyltin) borates and trialkyltin metaborates show the strongest peak in their infrared spectra in the range  $1270-1300\text{ cm}^{-1}$ . This may be assigned to  $\nu_{as}(B-O-Sn)$ . The exact positions are given in Tables 1 and 2. The assignment is supported by the fact that on cleavage of the  $B-O-Sn$  bond, this peak disappears completely.

In metaborates, a strong absorption appears near  $1360-1365\text{ cm}^{-1}$ , which may be due to  $\nu(B-O)$  in the ring. Exact positions are as follows.  $(OBOSnEt_3)_3$   $1360\text{ cm}^{-1}$ ,  $(OBOSnPr_3)_3$   $1365\text{ cm}^{-1}$ ,  $(OBOSnBu_3)_3$   $1365\text{ cm}^{-1}$ ,  $(OBOSnBu_{-1}_3)_3$   $1365\text{ cm}^{-1}$ .

Strong absorptions near  $720$  and  $735\text{ cm}^{-1}$  are characteristic of metaborates, and probably arise from out-of-plane vibrations of the  $B_3O_6$  skeleton. These peaks appear at  $732$  and  $742\text{ cm}^{-1}$  in trialkyltin metaborates, and are absent in  $(R_3SnO)_3B$ .

The purity of  $(Me_3SnO)_3B$  [singlet at  $\tau$  9.80 ppm;  $J(^{119}Sn-C-H)$  59 Hz;  $J(^{117}Sn-C-H)$  57 Hz] and  $(Ph_3SnO)_3B$  (complex multiplet at  $\tau$  2.52 ppm) was also checked by PMR spectra.

## Experimental

All the trialkyltin hydroxides or bis(trialkyltin) oxides were used as supplied. Benzene (B.D.H., L.R.) was kept overnight on sodium wire and distilled, and was finally dried azeotropically with ethanol.

Methods of analysing boron and tin have been described earlier [1]. Carbon and hydrogen analyses were carried out by M/s Australian Microanalytical Service CSIRO, Melbourne, Australia.

Infrared spectra were recorded on Perkin-Elmer 337 grating spectrophotometer. PMR spectra were recorded on an Mz 60 D machine.

### (1) Preparation of tris(trimethyltin) borate

Azeotropic removal of water from a mixture of trimethyltin hydroxide (4.1 g, 22.69 mmoles) and boric acid (0.47 g, 7.58 mmoles) gave a colourless liquid (4.0 g, 89% yield), b.p. 130-132°/14 mm (Found: B, 21, C, 20.8, H, 5.2.  $C_9H_{27}O_3BSn_3$  calcd.: B, 2.0; C, 19.6, H, 5.0%)

Tris(trialkyltin) borates prepared by this method are listed in Table 1.

### (2) Preparation of tributyltin metaborate

(a) On heating a mixture of tris(tributyltin) borate (2.68 g, 2.89 mmoles) and boric anhydride (0.202 g, 2.902 mmoles) at 240° ca 3 h, the latter dissolved to give clear colourless liquid, miscible with benzene

(b). Heating a mixture of bis(tributyltin) oxide (4.16 g, 6.99 mmoles) and boric anhydride (0.49 g, 7.04 mmoles) for ca 2 h at 200° yielded tributyltin metaborate (characterized by IR spectrum)

### (3) Reaction between tris(tributyltin) borate and hydrogen chloride

On passing dry HCl gas through a solution of tris(tributyltin) borate (2.56 g) in benzene ( $\approx$  70 ml) for ca 3 h, an exothermic reaction took place with precipitation of boric acid (0.16 g, 95% yield) (Found: B, 17.0.  $H_3BO_3$  calcd.: B, 17.4%). Drying and distillation of the filtrate gave tributyltin chloride (2.3 g, 85% yield) (authentic IR spectrum).

### (4) Reaction between tris(tributyltin) borate and acetyl chloride

Acetyl chloride (0.54 g, 6.88 mmoles) and tris(tributyltin) borate (1.68 g, 1.81 mmoles) reacted exothermically in benzene ( $\approx$  10 ml). A white gelatinous solid separated and was removed by decantation. The liquid on distillation gave tributyltin chloride, b.p. 89°/0.1 mm (1.4 g, 79% yield) (authentic IR spectrum)

### (5) Reaction between tris(tributyltin) borate and acetic acid

On adding acetic acid (0.35 g, 5.84 mmoles) dropwise to a solution of tris(tributyltin) borate (1.80 g, 1.94 mmoles) in benzene ( $\approx$  10 ml), an exothermic reaction occurred with immediate separation of boric acid (0.106 g, 91% yield) (Found: B, 16.7.  $H_3BO_3$  calcd.: B, 17.4%). The filtrate after removal of solvent gave tributyltin acetate, m.p. 84°; 2.0 g, 98% yield (Found: OAc, 17.0.  $C_{14}H_{30}O_2Sn$  calcd.: OAc, 17.3%) (authentic IR spectrum).

**(6). Reaction between tris(tributyltin) borate and phenol**

A solution of phenol (0.41 g, 4.36 mmoles) in benzene ( $\approx 7$  ml) was added to tris(tributyltin) borate (1.41 g, 1.52 mmoles) in benzene ( $\approx 10$  ml); boric acid precipitated out (0.057 g, 77% yield) (Found: B, 17.0.  $H_3BO_3$  calcd. B, 17.4%). Distillation of the filtrate gave tributyltin phenoxide (1.3 g, 74% yield) at  $140^\circ/0.4$  mm (lit. b.p.  $152^\circ/1$  mm.) (Found: Sn, 30.8.  $C_{18}H_{32}OSn$  calcd.: Sn, 31%.) (authentic IR spectrum).

**(7). Reaction between tris(tributyltin) borate and butanethiol**

On adding butanethiol (0.52 g, 5.78 mmoles) dropwise to tris(tributyltin) borate (1.78 g, 1.92 mmoles) in benzene ( $\approx 10$  ml), an exothermic reaction took place with immediate precipitation of boric acid (0.08 g, 67% yield) (Found: B, 17.5.  $H_3BO_3$  calcd.: B, 17.4%). Tributyltin butanethiolate (2.1 g, 95% yield) ( $115^\circ/0.1$  mm; lit.  $122-23^\circ/0.2$  mm) was obtained from the filtrate (Found: Sn, 30.1.  $C_{16}H_{36}SSn$  calcd.: Sn, 30.6%) (authentic IR spectrum).

**(8). Reaction between tris(tributyltin) borate and thiophenol**

Similarly, an exothermic reaction between thiophenol (1.4 g, 12.71 mmoles) and tris(tributyltin) borate (3.1 g, 3.34 mmoles) in benzene ( $\approx 10$  ml) gave boric acid (0.098 g, 58% yield) (Found: B, 16.7.  $H_3BO_3$  calcd. B, 17.4%) and tributyltin benzenethiolate (b.p.  $148^\circ/0.2$  mm lit.  $147^\circ/0.2$  mm) (3.7 g, 87% yield) (Found: Sn, 29.8.  $C_{18}H_{32}SSn$  calcd.: Sn, 29.7%) (authentic IR spectrum).

**(9). Reaction between tris(tributyltin) borate and acetic anhydride**

Some heat was evolved on adding acetic anhydride (0.7 g, 6.86 mmoles) dropwise to a solution of tris(tributyltin) borate (1.9 g, 2.05 mmoles) in benzene ( $\approx 10$  ml). On keeping the mixture overnight, white crystals of tributyltin acetate (m.p.  $83^\circ$ ) (Found: OAc, 17.5.  $C_{14}H_{30}O_2Sn$  calcd.: OAc, 17.3%) were obtained (authentic IR spectrum).

**(10). Attempted reaction between tris(tributyltin) borate and isopropyl acetate**

A mixture of isopropyl acetate (0.61 g, 5.98 mmoles) and tris(tributyltin) borate (1.65 g, 1.78 mmoles) in benzene was refluxed for ca 5 h. Distillation gave unchanged reactants.

**Acknowledgements**

One of the authors (S.K.M.) is grateful to the Council of Scientific and Industrial Research, New Delhi, for J.R.F. Thanks are also due to Nitto Kasei and Co. Japan, for a gift of bis(tributyltin) oxide.

**References**

- 1 S.K. Mehrotra, G. Srivastava and R.C. Mehrotra, *J. Organometal. Chem.* 47 (1973) 39
- 2 E.W. Abel and A. Singh, *J. Chem. Soc.*, (1959) 690.
- 3 M.F. Luggert, *J. Chem. Soc.*, (1958) 3790.
- 4 S.K. Mehrotra, G. Srivastava and R.C. Mehrotra, unpublished results.
- 5 J.C. Pommier and J. Valade, *C.R. Acad. Sci. Paris, Ser. C*, 260 (1965) 4549
- 6 D. Sukhani, Ph.D. Thesis, University of Rajasthan India, 1967
- 7 J.C. Pommier and J. Valade, *Bull. Soc. Chim. Fr.*, (1965) 2697
- 8 H. Steinberg, *Organoboron Chemistry*, Vol. I, Wiley, New York, p. 64