


A NEW SYNTHETIC METHOD FOR PELLITORINE

Tadakatsu MANDAI, Jiso GOTOH, Junzo OTERA, and Mikiyo KAWADA
 Okayama University of Science, Ridai-cho, Okayama 700

A stereoselective synthesis of N-isobutyl-E,E-2,4-decadienamamide, so called pellitorine is described in which the elimination reaction of acetic acid from 2-acetoxy-E-3-decenenitrile providing 2,4-decadienenitrile in a high yield was used as a key reaction.

We have reported the palladium catalyzed elimination reaction of acetic acid or phenol affording a terminal conjugated diene from a variety of allylic acetates and allylic phenyl ethers,¹⁾ and the reactions have been successfully applied to the efficient syntheses of several natural products.^{2,3)} In this communication, we wish to report a new synthetic method for pellitorine by applying the conjugated diene formation reaction to the preparation of 2,4-decadienenitrile, a good precursor for pellitorine. Pellitorine is an insecticidal compound isolated from *Anacyclus Pyrethrum* roots, and its structure was determined as N-isobutyl-E,E-2,4-decadienamamide.⁴⁾ A number of syntheses of pellitorine have proceeded via E,E-2,4-decadienoic acid or its esters as the key intermediates.⁵⁻⁹⁾ Our synthetic route is depicted in the following scheme.

Scheme



The starting E-2-nonenal (**2**) was prepared as the following procedures. The dianion of propargyl alcohol¹⁰⁾ was alkylated with 1-bromohexane in liquid NH_3 to give 2-nonyl-1-ol in 40% yield, which was reduced to E-2-nonenol in 93% yield with $LiAlH_4$ ¹¹⁾ in refluxing THF. Oxidation of the allylic alcohol by CrO_3 gave E-2-nonenal (**2**) in 73% yield. E-2-nonenal (**2**) (5.60 g, 40 mmol) was treated with Me_3SiCN ¹²⁾ (4.75 g, 48 mmol) and a catalytic amount of ZnI_2 in methylene chloride at 20° for 1 hr and the mixture was then hydrolyzed to a cyanohydrine with 1N HCl. The crude cyanohydrine was converted to 2-acetoxy-E-3-decenenitrile (**3**) with Ac_2O-Py in 90% yield from **2** after distillation (150°/8 Torr): NMR (CCl_4) δ 0.70-1.10 (t, 3H, CH_3), 1.10-1.70 (m, 8H, CH_2), 1.90-2.40 (m, 2H, $CH_2C=C$), 2.05 (s, 3H, $OCOCH_3$), 5.20-6.40 (m, 3H, $CH=CH$); IR (film) 1755, 1642, 965, 920 cm^{-1} . The acetate **3** (2.09 g, 10 mmol) thus obtained was heated at 140° (oil bath) in dry diglyme or xylene under nitrogen atmosphere in the presence of $Pd(OAc)_2$ (22.4 mg, 0.1 mmol) and PPh_3 (262 mg, 1 mmol) for 1 hr to give the dienenitriles **4** in 86% yield after column chromatographic purification (silica

gel, hexane-ether 30:1). Unfortunately, the dienenitriles **4** were found to be a mixture of four geometrical isomers by GLC analysis (PEG 20M Celite 545, area ratio 26:5:9:60); NMR (CCl_4) δ 0.70-1.10 (t, 3H, CH_3), 1.10-1.70 (m, 6H, CH_2), 1.90-2.50 (m, 2H, $\text{CH}_2\text{C}=\text{C}$), 4.90-5.40 (m, 1H, olefinic), 5.70-7.30 (m, 3H, olefinic); IR (film) 2200, 1640, 995 cm^{-1} .

The dienenitriles **4** (1.49 g, 10 mmol) were treated with $^i\text{Bu}_2\text{AlH}$ (1.2 eq.) in dry ether at -50° for 30 min, at 20° for 30 min, and at 45° for 30 min. The mixture was hydrolyzed with 1N HCl to give dienals **5a** in 80% yield, which were a mixture of two geometrical isomers by GLC analysis (PEG 20M Celite 545, area ratio 10:90); NMR (CCl_4) δ 0.70-1.10 (t, 3H, CH_3), 1.10-1.70 (m, 6H, CH_2), 2.00-2.50 (m, 2H, $\text{CH}_2\text{C}=\text{C}$), 5.75-6.40 (m, 3H, olefinic), 6.70-7.30 (m, 1H, olefinic), 9.45 (d, 1H, $J=8$ Hz, CHO); IR (film) 1685, 1640, 990 cm^{-1} . It is known that cis- α,β -unsaturated aldehyde readily changes to its trans isomer under acidic condition. In our results, isomerization of 2-olefin accompanied by that of 4-olefin probably occurred during the hydrolysis of aldimines to give E,E-isomer stereoselectively. The isomeric dienals **5a** were oxidized by AgNO_3 -NaOH mixture¹³⁾ to the corresponding dienoic acids **5b** in 52% yield. These acids **5b** were converted to their methyl esters **5c** with CH_2N_2 and they were analyzed by GLC (area ratio 10:90); NMR (CCl_4) δ 0.70-1.10 (t, 3H; CH_3), 1.10-1.80 (m, 6H, CH_2), 1.80-2.50 (m, 2H, $\text{CH}_2\text{C}=\text{C}$), 3.65 (s, 3H, OCH_3), 5.50-6.20 (m, 3H, olefinic), 6.90-7.40 (m, 1H, olefinic); IR (film) 1720, 1640, 1620 cm^{-1} ; MS (m/e) 182 (M^+). The acid chlorides **5d** were treated with $^i\text{BuNH}_2$ in ether to give a crude oil of pellitorine (**1**) which was purified by recrystallization from hexane to afford white crystals: NMR (CDCl_3) δ 0.70-1.10 (t, 3H, CH_3 , d, 6H, CH_3CCH_3), 1.10-1.80 (m, 7H, CH_2 , CH), 1.80-2.40 (m, 2H, $\text{CH}_2\text{C}=\text{C}$), 3.10 (t, 2H, NCH_2), 5.60-6.20 (m, 3H, olefinic), 6.90-7.40 (m, 1H, olefinic); IR (nujol) 3300, 1660, 1630 cm^{-1} ; MS (m/e) 223 (M^+), 151 ($\text{M}^+ - ^i\text{BuNH}_2$); mp 91° - 92° (lit.⁴⁾ 90°).

References

- 1) J. Tsuji, T. Yamakawa, M. Kaito, and T. Mandai, *Tetrahedron Lett.*, 2075 (1978).
- 2) T. Mandai, H. Yasuda, M. Kaito, J. Tsuji, R. Yamaoka, and H. Fukami, *Tetrahedron*, **35**, 309 (1979).
- 3) J. Tsuji, T. Yamakawa, and T. Mandai, *Tetrahedron Lett.*, 3741 (1979).
- 4) L. Crombie, *J. Chem. Soc.*, 999 (1955).
- 5) L. Crombie, *J. Chem. Soc.*, 1007 (1955).
- 6) M. Jacobson, *J. Am. Chem. Soc.*, **75**, 2584 (1953).
- 7) G. Ohloff and M. Pawlak, *Helv. Chim. Acta*, **56**, 1176 (1973).
- 8) F. Naf and R. Decorzant, *Helv. Chim. Acta*, **57**, 1309 (1974).
- 9) J. Tsuji, H. Nagashima, T. Takahashi, and K. Masaoka, *Tetrahedron Lett.*, 1917 (1977).
- 10) G. Rickards and L. Weiler, *J. Org. Chem.*, **43**, 3607 (1978).
- 11) E. J. Corey, J. A. Katzenellenbogen, and G. H. Posner, *J. Am. Chem. Soc.*, **89**, 4245 (1967).
- 12) D. A. Evans, G. L. Carroll, and L. K. Truesdale, *J. Org. Chem.*, **39**, 914 (1974).
- 13) J. C. Sheehan, C. A. Robinson, *J. Am. Chem. Soc.*, **73**, 1207 (1951).

(Received January 5, 1980)