


This article was downloaded by: [Lulea University of Technology]

On: 26 July 2013, At: 10:30

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

Easy and Rapid Method for Disulfide Syntheses Using Nanophase-Manganese (VII) Oxide Coated Clay

Sudershan R. Gondi ^a, David Y. Son ^b, Edward R. Biehl ^b & Rajan K. Vempati ^{a b}

^a ChK Group, Inc., Plano, Texas, USA

^b Southern Methodist University, Dallas, Texas, USA

Published online: 28 Dec 2009.

To cite this article: Sudershan R. Gondi, David Y. Son, Edward R. Biehl & Rajan K. Vempati (2009) Easy and Rapid Method for Disulfide Syntheses Using Nanophase-Manganese (VII) Oxide Coated Clay, Phosphorus, Sulfur, and Silicon and the Related Elements, 185:1, 34-39

To link to this article: <http://dx.doi.org/10.1080/10426500903147175>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

EASY AND RAPID METHOD FOR DISULFIDE SYNTHESSES USING NANOPHASE-MANGANESE (VII) OXIDE COATED CLAY

Sudershan R. Gondi,¹ David Y. Son,² Edward R. Biehl,²
and Rajan K. Vempati^{1,2}

¹ChK Group, Inc., Plano, Texas, USA

²Southern Methodist University, Dallas, Texas, USA

Disulfide bond formation by the oxidation of several thiol compounds using nanophase manganese (VII) oxide coated clay (NM7O coated clay) in chlorinated, nonchlorinated (toluene), and polar solvents (water) is described. During the process, nitro and methoxy groups remain unaffected. The NM7O coated clay is easily prepared, stable, and inexpensive to manufacture. Furthermore, this and other studies prove that NM7O coated clay properties, i.e., mineralogy, chemical, and reactivity, are different from KMnO₄. This rapid and facile synthesis coupled with the use of the recyclable NM7O coated clay catalyst will save energy due to low temperature and rapid reaction times, as well as minimal disposal problems, thus decreasing production costs.

Keywords Catalyst; coupling; easy and rapid synthesis energy; stable

INTRODUCTION

Disulfide bond formation is important in peptide synthesis,¹ protein stabilization,² and in bioactive molecules.³ Thiols are the main precursors to disulfides, but they are highly susceptible to air oxidation. Therefore, in multistep syntheses, the protection of thiols as acetates or esters and their subsequent transformation into -SS-linkages is often necessary. There are a few methods for the cleavage of thiol acetates under abnormal conditions, such as longer reaction time (12 days), argon atmosphere, and photolysis.⁴ Recently, two improved methods have been reported: one requires activation by stannates,⁵ while the other uses a metallic reagent⁶ with a longer reaction time. The last decade has witnessed two major changes in chemical manufacturing: 1) a large interest in “green” chemistry syntheses in which toxic reagents, catalysts, and solvents have been replaced by nontoxic counterparts, and 2) energy efficient processes to decrease CO₂ footprint. This has been achieved by the use of environmentally friendly solid-supported catalysts⁷ and/or in combination with microwave irradiation.⁸ This modification not only increases the reaction rate, but affords

Received 18 May 2009; accepted 26 June 2009.

S.R.G. would like to acknowledge Prodev, Inc., for partial funding on this project. Furthermore, the authors would like to thank Mr. Xiang Hua and Mrs. Jyothi Lekla for their assistance on the project.

Address correspondence to Dr. Edward R. Biehl, Chemistry, Southern Methodist University, 3215 Daniel Ave., Dallas, TX 75275, USA. E-mail: ebiehl@smu.edu



R = a) C₆H₅-, b) HOCH₂CH₂-, c) 2-NH₂-C₆H₅- d) 4-HO-C₆H₅-, e) 4-NO₂-C₆H₅-, f) 4-CH₃-C₆H₅-CH₂- g) CH₂=CH-CH₂-, h) (CH₃)₂CH-, i) (CH₃)₃C-, j) (C₆H₅)₃C-, k) HSCH₂CH₂CH₂-

Scheme 1

a simple workup and ease of manipulation of the reaction. Herein, we report a rapid and convenient method for the oxidative coupling of thiols using a nanophase manganese (VII) oxide coated clay (NM70 coated clay)⁹ catalyst as shown in Scheme 1. The NM70 coated clay properties, i.e., mineralogical, chemical, and reactivity, are distinct and unique from KMnO₄.¹⁰ This method affords a rapid, low temperature method for the synthesis of disulfides that avoids the use of activating reagents or excess chlorinated solvent.

RESULTS AND DISCUSSION

To determine the optimum reaction conditions and solvent for the above thiol-coupling reactions shown in Scheme 1, we studied the conversion of benzenethiol (**1a**) to diphenyl disulfide (**2a**) in various solvents, temperatures, and reaction times. The results, shown in Table I, show that nearly quantitative yields (~98%) of **2a** were obtained when the reactions were carried out in chloroform (entry 3) at room temperature for 2 h and in toluene (entry 1) at 40°C. The other solvents listed, acetonitrile (entry 2) and dichloromethane (entry 4), required longer reaction times (3 h and 6 h, respectively) and gave significantly lower yields of **2a** (66% and 87%, respectively). Thus we decided to carry out subsequent reactions in toluene at 40°C and chloroform at room temperature.

We next used these conditions to the conversion of a variety of thiols (**1a–k**) to the corresponding disulfides (**2a–k**), and the results are listed in Table II. All products were characterized by MS, NMR, melting point, and by comparison of their R_f values to those of authentic samples.












As shown in Table II, most of the listed disulfides (**2a**, **2c**, **2f**, **2g**, **2h**, **2i**, and **2k**) were obtained in essentially quantitative yields within 1–3 h when the reactions were run in toluene or chloroform. Additionally, the reaction of the water-soluble butane-1-thiol (**1a**) carried out in water at room temperature gave the corresponding disulfide **2b** in 98% yield. It is noteworthy that diallylic disulfide (**2g**) (entry 7) was obtained in 98% yield when the reaction was run in toluene; however, in chloroform the yield of **2g** was lowered to 65%.

Table I Thiophenol reaction conditions using NM70 coated clay

Entry	Substrate	Solvent	Time	Temperature	% Yield
1	Thiophenol	Toluene	2 h	40°C	98
2	Thiophenol	Acetonitrile	3 h	rt	66
3	Thiophenol	Chloroform	2 h	rt	98
4	Thiophenol	Dichloromethane	6 h	rt	87

rt = room temperature.

Table II Syntheses of disulfide compounds using NM70 coated clay

Entry No.	Thiol	Solvent	Time	Reaction Conditions	Product	Yield (%)	Ref. [†]
1	 1a	Toluene CHCl ₃	2h 2h	40°C rt	2a	97 98	11
2	 1b	Water	2 h	rt	2b	98	12
3	 1c	Toluene CHCl ₃	1 h	40°C rt	2c	97 99	13
4	 1d	Toluene	2h	40°C	2d	98	14
5	 1e	Toluene	2 h	40°C	2e	94	15
6	 1f	Toluene CHCl ₃	2 h	40°C rt	2f	98 97	16
7	 1g	Toluene CHCl ₃	3 h	40 °C rt	2g	99 65	17
8	 1h	Toluene CHCl ₃	2 h	40°C rt	2h	98 97	18
9	 1i	Toluene CH ₃ Cl	2 h	40°C rt	2i	99 98	19
10	 1j	Toluene	3 h [‡]	40°C	2j	60	20
11	 1k	Toluene CHCl ₃	2 h	40°C rt	2k	99	21

[†]References.[‡]Reaction incomplete even with additional NM70 (1:3) and longer reaction time (18 h).

rt = room temperature.

Interestingly, both the electron-withdrawing 4-nitro group and electron-donating 4-hydroxy group on the respective phenylthiols (entries 5 and 4) gave respective coupling products **2e** and **2d** in 94% and 98% yields. However, despite the utility and simplicity of these reactions, limitations were encountered. For example, ditrityl disulfide **2j** (entry 10) was obtained in 60% when the reaction was run for 3 h in toluene at 40°C; attempts to increase the yield of **2j** by prolonged heating for 18 h failed. However, even though the yield of ditrityl disulfide was low (60%), to the best of our knowledge, it is the highest yield reported to date.

The kinetics and mechanism of the oxidation of a thiol to a disulfide by the one-equivalent oxidants neptunium (VI) and cobalt (III) have been reported.²² The major features of that proposed mechanism involve an initial 1-equiv oxidation of a thiol to a thiol radical in the rate determining step (RDS). The resulting radical then is thought to react with another thiol to give the stable disulfide radical, which presumably is converted to the disulfide. It is possible that the nanophase Mn(VII)O used in our study behaves similarly. We, however, have insufficient evidence for the role of Mn(VII)O in the process. Studies are underway, and the results will be submitted in due course.

CONCLUSION

In summary, we have developed a convenient and practical alternative to disulfide synthesis by coupling various types of thiol groups using NM70 coated clay. The lower temperature and rapid syntheses of disulfides coupled with the use of recyclable NM70 coated clay catalyst will save energy and decrease production costs. Furthermore, the reaction appears to be surface-based and catalytic, since the NM70 coated clay can be reused multiple times for making additional disulfide compounds. We are carrying out mechanistic studies and will report the conclusions in due course.

EXPERIMENTAL

Thiophenol and trityl thiol were obtained from Acros and Aldrich, respectively. The natural montmorillonite clay, Bentonite L-10, was purchased from Southern Clay Products, Gonzalez, TX, USA. All other materials were reagent grade unless otherwise specified. ¹H and ¹³C NMR spectra were obtained on a 400-MHz Bruker Advance NMR spectrometer. Melting points were collected on a TA Instruments DSC 2010 Differential Scanning Calorimeter using a heating rate of 10 °C min⁻¹ and nitrogen as a purge gas. Thin layer chromatography was done on precoated silica gel 60f 254 (0.5 mm) glass plates. The NM70 coated clay (Bentonite-L-10) was prepared by adding MnCl₂ 4H₂O (720 g) to water (500 mL), and stirring for 5 min to dissolve the salt to yield a solution (termed Solution A). Then montmorillonite clay (2000 g) was added to water (8 L) and stirred for 30 min, or until a uniform suspension was achieved (termed Suspension B). Solution A and Suspension B were added and mixed in another vessel to make a beige-colored suspension of Mn(OH)₂. After 30 min of stirring, 1.5 L, 5.2 M NaOH (1.5 L) was added, and the resultant pH of the suspension was 13. Subsequently, 1,4-phenylenediamine (PDA, 40 g) was added to the suspension and stirred for 30 min. The PDA-added suspension was lowered to pH 7.8 by the addition of 6 N HCl, during which time the color of the suspension changed from green to violet within 1 h. The violet color is the indication of NM70 coated clay. The oxidation states were characterized and confirmed by cyclic voltametric techniques and visible spectroscopy.¹⁰ The suspension contained 22% by weight NM70 coated clay

with a Mn content of 7%. Unless otherwise indicated, the compounds were obtained commercially and used without further purification.

In a typical experimental procedure, the thiol compounds were treated with excess NM7O coated clay, i.e., 1:2 molar ratio of thiol compounds and Mn (based on its content in NM7O coated clay) in chloroform at room temperature or in toluene at 40°C in the presence of air for 2 h. After completion of the reaction, the NM7O coated clay was filtered, and the filtrate was concentrated using a rotary evaporator under reduced pressure. Thin layer chromatography was done on precoated silica gel 60f 254 (0.5 mm) glass plates. Column chromatography was used to yield the pure disulfide.

REFERENCES

1. M. Bodanszky, *Principles of Peptide Synthesis* (Springer-Verlag, Berlin, 1984), Chapter 4.
2. K. Fuwa, W. E. C. Wacker, R. Druyan, A. F. Bartholomay, and B. L. Vallee, *Proc. Natl. Acad. Sci. (USA)*, **46**, 1298 (1960).
3. (a) J. R. Johnson, W. F. Bruce, and J. D. Dutcher, *J. Am. Chem. Soc.*, **65**, 2005 (1943); (b) P. A. Miller, K. P. Milstrey, and P. W. Trown, *Science*, **159**, 431 (1968).
4. (a) A. P. Breen, J. A. Murphy, C. W. Patterson, and N. F. Woosfer, *Tetrahedron*, **49**, 10643 (1993); (b) P. Dubs and R. Stussi, *Helv. Chim. Acta*, **61**, 2351 (1978); (c) J. F. Grunwell, D. L. Foerst, and M. J. Sanders, *J. Org. Chem.*, **42**, 1142 (1977).
5. T. Sato, J. Otera, and H. Nozaki, *Tetrahedron Lett.*, **31**, 3595 (1990).
6. H. M. Meshram, *Tetrahedron Lett.*, **34**, 2521 (1993).
7. (a) A. McKillop and D. W. Young, *Synthesis*, 401 and 481 (1979); (b) P. Laszlo, *Preparative Chemistry Using Supported Reagent* (Academic Press, San Diego, 1987), pp. 2, 4; (c) H. M. Meshram, G. S. Reddy, and J. S. Yadav, *Tetrahedron Lett.*, **38**, 8891 (1997); (d) H. M. Meshram, G. S. Reddy, D. Srinivas, and J. S. Yadav, *Synth. Commun.*, **28**, 2593 (1998); (e) J. S. Yadav, H. M. Meshram, G. S. Reddy, and G. Sumithra, *Tetrahedron Lett.*, **39**, 3043 (1998).
8. (a) R. S. Varma and H. M. Meshram, *Tetrahedron Lett.*, **38**, 5247 (1997); (b) R. S. Varma, R. K. Saini, and H. M. Meshram, *Tetrahedron Lett.*, **38**, 6525 (1997); (c) R. S. Varma, R. K. Saini, and R. Dahiya, *Tetrahedron Lett.*, **38**, 7823 (1997); (d) R. S. Varma and R. Saini, *Tetrahedron Lett.*, **38**, 2623 (1997); (e) E. E. Reid, *Organic Chemistry of Bivalent Sulphur Compounds* (Chemical Publishing Co., Inc., New York, 1960), vol. III; (f) P. Laszlo and A. Camelis, *Aldrichimica Acta*, **21**, 97 (1988).
9. R. K. Vempati and D. Y. Son, Solid Support Stabilized Mn(III) and Mn(VII) and Method of Preparation, U.S. Patent No. 6,953,763 B2, (2006)
10. R. K. Vempati, G. Bhat, and R. W. Wagner, *Proceedings of the INTC '06*, Houston, TX, USA, Sept. 25–28, 2006.
11. (a) A. Herrera, *J. Org. Chem.*, **69**, 4545 (2004); (b) M. Furukawa, *Synthesis*, 165 (1975).
12. (a) A. R. Hajipour, *J. Sulfur Chem.*, **27**, 441 (2006); (b) A. Christoforou, *Tetrahedron Lett.*, **47**, 9211 (2006).
13. (a) L. V. Saloutina, *Russ. J. Org. Chem.*, **42**, 1307 (2006); (b) W. E. Fristad, *Synth. Commun.*, **15**, 1 (1985).
14. (a) R. S. Sengar, *New J. Chem.*, **27**, 1115 (2003); (b) A. Nakahashi, *J. Poly. Sci. Part A: Poly. Chem.*, **45**, 3580 (2007).
15. (a) D. Suresh Kumar, *J. Org. Chem.*, **72**, 2106 (2007); (b) H. Boerzel, *Inorg. Chem.*, **42**, 1604 (2003).
16. (a) A. K. Misra, *Synth. Commun.*, **34**, 1079 (2004); (b) S. A. Okecha, *Chem. & Ind. (London, United Kingdom)*, **19**, 788 (1980); (c) E. M. Gordon, *J. Med. Chem.*, **31**, 2199 (1988).
17. (a) O. Higuchi, *J. Agri. Food. Chem.*, **51**, 7208 (2003); (b) F. E. Chen, *Synth. Commun.*, **32**, 3487 (2002).

18. (a) N. Taniguchi, *J. Org. Chem.*, **71**, 7874 (2006); (b) S. Chorbadzhiev, *J. Prakt. Chem. (Leipzig)*, **319**, 1036 (1977).
19. (a) F. Freeman, *Magn. Reson. Chem.*, **26**, 813 (1988); (b) G. J. Derbesy, *J. Org. Chem.*, **60**, 1044 (1995); (c) R. Priefer, *J. Am. Chem. Soc.*, **124**, 5626 (2002); (d) F. J. Arnaiz, *Polyhedron*, **21**, 1635 (2002); (e) S. M. Bonesi, *Photochem. & Photobiol. Sci.*, **3**, 489 (2004).
20. (a) J. C. Cochran, S. R. Friedman, and J. P. Frazier, *J. Org. Chem.*, **61**, 1533 (1996); (b) G. Derbesy and D. N. Harpp, *Tetrahedron Lett.*, **35**, 5381 (1994); (c) R. Steudel, S. Foerster, and J. Albertsen, *Chem. Ber.*, **124**, 2357 (1991); (d) M. Ostrowski, J. Jeske, P. G. Jones, and W. W. du Mont, *Chem. Ber.*, **126**, 1355 (1993).
21. (a) D. Camp, *Aus. J. Chem.*, **43**, 161 (1990); (b) A. R. Hajipour, *J. Sulfur Chem.*, **27**, 441 (2006); (c) A. R. Hajipour, *J. Chem. Res.*, 286 (2004); (d) M. C. Caserio, *Phosphorous, Sulfur, & Silicon*, **23**, 169 (1985).
22. M. Woods, J. Karbwang, J. S. Sullivan, and E. Deutsch, *Inorg. Chem.*, **15**, 1678 (1976).