

Re-investigating old pharmacophores: Are 4-aminoquinolines and tetraoxanes potential two-stage antimalarials?

Bogdan A. Šolaja, Natasa Terzic Jovanovic, Jelena Konstantinovic, Miklos Tot, Jovana Burojevic, Olgica Djurkovic Djakovic, Jelena Srblijanovic, Tijana Stajner, Tatjana Z Verbic, Mario Zlatovi#, Marta Monteiro Maia Machado, Inês S. Albuquerque, Miguel Prudencio, Richard J. Sciotti, Stevan Pecic, Sarah D'Alessandro, and Donatella Taramelli

J. Med. Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.jmedchem.5b01374 • Publication Date (Web): 07 Dec 2015

Downloaded from <http://pubs.acs.org> on December 10, 2015

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Re-investigating old pharmacophores: Are 4-aminoquinolines and tetraoxanes potential**
4 **two-stage antimalarials?**
5
6
7
8
9

10
11 Natasa Terzić,[#] Jelena Konstantinović,[¶] Mikloš Tot,[¶] Jovana Burojević,[¶] Olgica Djurković-
12 Djaković,[∇] Jelena Sribljanović,[∇] Tijana Štajner,[∇] Tatjana Verbić,[¶] Mario Zlatović,[¶] Marta
13 Machado,[§] Inês S. Albuquerque,[§] Miguel Prudêncio,^{§,*} Richard J. Sciotti,[†] Stevan Pecic,[∇]
14 Sarah D'Alessandro,[‡] Donatella Taramelli,[‡] Bogdan A. Šolaja^{¶,*}
15
16
17
18
19
20
21
22
23

24 [#]Institute of Chemistry, Technology, and Metallurgy, Belgrade, Serbia
25
26

27 [¶]Faculty of Chemistry, University of Belgrade, Studentski trg 16, P.O. Box 51, 11158,
28 Belgrade, Serbia
29
30
31

32 [∇]Institute for Medical Research, University of Belgrade, Dr.Subotića 4, 11129 Belgrade,
33 Serbia
34
35
36

37 [§]Instituto de Medicina Molecular, Faculdade de Medicina, Universidade de Lisboa, 1649-028
38 Lisboa, Portugal
39
40
41

42 [†] Experimental Therapeutics Branch, Walter Reed Army Institute of Research, Silver Spring,
43 MD20910, USA
44
45
46

47 [∇] Division of Experimental Therapeutics Department of Medicine, Columbia University, New
48 York, NY 10032, USA
49
50
51

52 [‡] Dipartimento di Scienze Farmacologiche e Biomolecolari, Università degli Studi di Milano,
53 20133 Milan, Italy
54
55
56
57
58
59
60

Abstract

The syntheses and antiplasmodial activities of various substituted aminoquinolines coupled to an adamantane carrier are described. The compounds exhibited pronounced in vitro, and in vivo activity against *P. berghei* in the Thompson test. Tethering a fluorine atom to the aminoquinoline C(3) position afforded fluoroaminoquinolines that act as intra-hepatocytic parasite inhibitors, with compound **25** having an $IC_{50} = 0.31 \mu\text{M}$ and reducing the liver load in mice by up to 92% at 80 mg/kg dose. Screening our peroxides as inhibitors of liver stage infection revealed that the tetraoxane pharmacophore itself is also an excellent liver stage *P. berghei* inhibitor (**78**: $IC_{50} = 0.33 \mu\text{M}$). Up to 91% reduction of the parasite liver load in mice was achieved at 100 mg/kg. Examination of tetraoxane **78** against the transgenic 3D7 strain expressing luciferase under a gametocyte-specific promoter revealed its activity against stage IV-V *P. falciparum* gametocytes ($IC_{50} = 1.16 \pm 0.37 \mu\text{M}$). To the best of our knowledge, compounds **25** and **78** are the first examples of either an 4-aminoquinoline or a tetraoxane liver stage inhibitors.

Introduction

Malaria is a widespread parasitic disease affecting mainly the population of Sub-Saharan Africa, South-East Asia and South America. With its heavy burden of ca. 207 million cases in 2013 and a death toll of ca. 627,000, malaria is one of the major public health problems worldwide (WHO Malaria report 2014).¹ In addition, with as much as ca. 10% of the global population being transiently ill and temporarily unable to work, malaria poses an immense economic burden to malaria-endemic regions. Fortunately, over the last 15 years the death toll and the number of cases decreased steadily by the combined action of the widespread use of insecticide-treated bed nets, the implementation of artemisinin-based combination therapies (ACTs)² and appropriate diagnosis.

In particular, ACTs, in which a fast parasitemia clearing peroxide molecule is combined with a slow-acting antimalarial pharmacophore, an aminoquinoline or alike, proved to be a very effective therapeutic approach. Thanks to the public and private contributions and the efforts of Medicine for Malaria Venture (MMV), new combinations (i.e. DHA-piperaquine; pyronaridine-artesunate) or formulations (i.e., artemether-lumefantrine dispersible for children) have been approved and distributed.³

Nevertheless, drug resistance already pushed aside many good and low-toxic anti-malarial drugs, most relevantly chloroquine. A few years ago, the first signs of resistance of *Plasmodium falciparum* (*Pf*) to artemisinin in Western Cambodia were reported as unusual prolonged parasite clearance phenotype following treatment with an ACT.^{4,5} Artemisinin resistance may presently be spreading throughout South-East Asia.⁶ Multiple point mutations in the propeller domains of the *Pf*kelch gene on chromosome 13 (K13), have been associated with the resistance phenotype in vivo and represent a molecular marker for tracking the spread of artemisinin resistance.⁷

1
2
3 In this context, to counter the problems outlined above, new, cheap and effective anti-
4
5 plasmodial drugs are needed. The life cycle of *Plasmodium* parasites includes several stages,
6
7 in both its mammalian and its invertebrate hosts. Malaria infection in mammals starts with the
8
9 bite of an infected female *Anopheles* mosquito, followed by migration of *Plasmodium*
10
11 sporozoites to the liver, where they infect and develop inside hepatocytes. Hepatic infection is
12
13 asymptomatic and malaria symptoms do not arise until hepatic merozoites are released into
14
15 the blood stream and infect erythrocytes. Therefore, *Plasmodium* liver stages appear to be
16
17 promising targets for new antimalarial drugs aiming at prophylactic intervention.⁸ In the
18
19 blood, the asexual cyclic replication of the parasites causes the symptoms of malaria and leads
20
21 to the differentiation of the parasite's sexual forms, the gametocytes, which are then taken up
22
23 by a mosquito, where the sexual phase of the parasite's life cycle takes place. The discovery
24
25 and development of antimalarials that act through several, if not all, stages of the parasite life
26
27 cycle would greatly enhance their effectiveness, thus allowing much cheaper and considerably
28
29 more effective regimens. This is also the most recent indication of MMV for new
30
31 antimalarials.³
32
33
34
35
36

37 The activity of several chemotypes that target multiple stages of the parasite's life
38
39 cycle was reported in the last few years: (+)-SJ733,⁹ imidazopyrazines,¹⁰ quinolone-3-
40
41 diarylethers,¹¹ flavones,¹² N-cinnamoylated chloroquine analogues,¹³ quinolin-4(1H)-imines,¹⁴
42
43 3,5-bis(benzylidene)-4-piperidones,¹⁵ imidazopyrazines,¹⁶ primaquine-chloroquine hybrid,¹⁷
44
45 ketotifen, norketotifen,¹⁸ decoquinatate,¹⁹ thienopyrimidinones.²⁰ All these multiple stage
46
47 inhibitors are active against both liver and blood *Plasmodium* asexual stages. Remarkably,
48
49 imidazopyrazines,^{10,16} quinolone-3-diarylethers,¹¹ primaquine-chloroquine hybrid,¹⁷ and
50
51 decoquinatate¹⁹ are active against blood sexual stages too. In vivo liver-stage (LS) activity was
52
53 confirmed for imidazopyrazines,^{10,16} quinolone-3-diarylethers,¹¹ primaquine-chloroquine
54
55 hybrid,¹⁷ ketotifen, norketotifen,¹⁸ decoquinatate,¹⁹ and fluoroquinoline DDD107498.²¹
56
57
58
59
60

1
2
3 Peroxide-containing hybrid compounds, in which the LS drug primaquine was bound
4 to an artemisinin or tetraoxane moiety, were synthesized as potential new antimalarial drugs,
5 active against both the liver and the erythrocytic asexual stages of the parasite. In fact, both
6 peroxide classes possess excellent intra-erythrocytic activity. Several primaquine-artemisinin
7 (PQ-ART) hybrids,²² tetraoxane–pyrimidine nitrile hybrids,²³ and tetraoxane-, and
8 artemisinin-primaquine hybrids²⁴ were reported. The PQ-ART and tetraoxane-primaquine
9 amalgams showed very interesting LS activity in vitro. However, the observed LS activity
10 could be primarily ascribed to the partner employed in the hybrids molecules (based on the
11 relative LS inactivity of the respective partner peroxide entities –tetraoxanes and ART^{23, 25, 26}).
12 The in vivo LS activity was confirmed for primaquine-ART hybrids.²²
13
14
15
16
17
18
19
20
21
22
23
24
25

26 Recently, it was observed that reducing drug (chloroquine) pressure results in re-
27 emergence of chloroquine-sensitive *P. falciparum* malaria.²⁷ Although it is very unlikely that
28 any aminoquinoline would be used in monotherapy regimens, new aminoquinolines were
29 developed as promising antimalarials, and as possible partners in combination treatments.^{28, 29,}
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

30, 31, 32, 33, 34, 35, 36 7-chloro-4-aminoquinolines (7-ACQ) are considered remarkably active
against asexual blood stages, but not against LS parasites.²⁶ It is therefore not surprising that
only one successful effort towards designing a LS-active 7-ACQ has been reported.¹⁷ Lödige
et al. showed that a PQ-CQ hybrid is efficient against asexual blood stages of CQ-susceptible
and CQ-resistant *P. falciparum* strains 3D7, Dd2, and K1, and, importantly, against in vitro
LS of *P. berghei*.¹⁷ In addition, they showed that the PQ-CQ hybrid molecule is able to
appreciably decrease the parasite liver load when dosed at 3× 90 mg/kg.

Our own efforts in the field of 7-ACQ antimalarials resulted in promising compounds
with adamantane³⁷ and thiophene-based carriers (Figure 1).³⁸





Figure 1. Recently developed aminoquinoline antimalarials.

Here, we report on a new generation of aminoquinolines with adamantane carriers and benzothiophene terminal amine substitutions. We show that compounds of the new series have improved blood stage antiplasmodial activity in comparison to the initial series. In addition, the first LS activity of an 4-aminoquinoline is reported for the first time, followed by first-in-class LS activity of a tetraoxane antimalarial.

Results


Chemistry

Here, we present the synthesis of three types of aminoquinoline antiplasmodials: a) adamantane derivatives, b) benzothiophene derivatives, and c) 3-fluoro-4-aminoquinolines, which include chloroquine and AQ-13 fluoro derivatives (Scheme 1, Scheme 2). The key reaction for the preparation of adamantane and benzothiophene *P. falciparum* inhibitors is the coupling of prepared fragments by reductive, or Buchwald-Hartwig amination as outlined in Scheme 1. Full details on the syntheses of these compounds and their respective precursors are given as part of Supporting information.



Scheme 1. Amination affords antiplasmodials **9-12, 22-24, 27-31, 33-37, 40-44, 46-67**.

The synthesis of the 3-fluoro-substituted aminoquinoline core was initiated with the introduction of fluorine at the C(3) position via corresponding diazonium salt (not isolated, Balz–Schiemann reaction³⁹) in 70% yield, starting with known 3-aminochloroquinolines **13**, **14**, Scheme 2. The introduction of iodine in position C(4) was accomplished using low temperature generation of intermediate lithio species followed by low temperature iodination thereof,⁴⁰ affording the iodine regioisomers (**17** and **18**, R=Cl) in (4:1)-ratio. It is interesting to note that when R=H, the lithiation/iodination sequence afforded only 3-fluoro-4-iodoaminoquinoline (**19**) in 70% yield. The synthesis of the aminoquinoline part was concluded by Buchwald–Hartwig or nucleophilic amination (41-77%, Scheme 2). The yields of amination reactions at C(2) and C(4) varied; the cross-coupling reaction proceeded smoothly except on two occasions when possibly steric congestion of fluorine and a short spacer lowered the reaction yields (compounds **20** and **68** in Supporting information). All compounds screened for antiplasmodial activity were fully characterized and were found to be >95% pure (HPLC).



Scheme 2. Synthesis of 7-chloro-3-fluoro-4-aminoquinolines. R₁: (N-alkyl-2-(adamant-1-yl)alkanamine); 5-(diethylamino)pentan-2-yl, 3-(diethylamino)propan-1-yl.


***Plasmodium* asexual blood stages**













In vitro antiplasmodial activity









Synthesized compounds were tested in vitro for their antiplasmodial activity against three *P. falciparum* strains: D6 (CQ susceptible (CQS) strain), W2 (CQ resistant (CQR) strain), and TM91C235 (Thailand, a multidrug-resistant (MDR) strain), by measuring the inhibition of incorporation of radiolabeled hypoxanthine by drug-treated parasites, as previously reported.⁴¹ CQ, mefloquine (MFQ) and artemisinin (ART) were used as positive controls. The antiplasmodial activity of all compounds tested is given in Table S1 (Supporting Information). Twenty-one of these compounds were more potent than CQ and MFQ (IC₅₀) against the CQS strain D6, and twenty-three were more active against MDR C235 strain than MFQ. Of 49 compounds more active than CQ against the CQR strain W2, 26 were ≥10-fold more active against the given strain. In addition, three compounds, **9**, **33**, and **34**, were more active than ART against all three *P. falciparum* strains tested.

The in vitro activity data of the most potent compounds with the adamantane carrier is selected to Table 1. Detailed analysis revealed several interesting SAR issues within the series differing in linker length:

- In general, a series with shortest linker ($m=1$, Scheme 1) was more active than the ones with $m=2, 3$ against all three *P. falciparum* strains (compounds **9**, **10**, **33**, **34**). All four compounds had excellent selectivity indices against HepG2 and RAW264.7 cells ($SI_{\text{HepG2/D6}} = 4,188$; $SI_{\text{RAW 264.72/D6}} = 1,886-2,901$), indicating that they could constitute good candidates for in vivo evaluation. The introduction of methyl groups in the α -positions to nitrogens (in an attempt to subdue possible metabolic hydroxylation of the respective α -carbons) afforded remarkably more active antiplasmodials, e.g., **9**, **10**, **33**, **34** vs. **1** (for chemistry and activity details of **1** see ref. 37); however, the screening in human and mouse liver microsomal preparations indicated that only **33** and **34** are acceptably metabolically stable (MLM). Compounds **33** and **34** ($m=1, n=2$) appeared to be 2.5-6 times more stable than those with shorter linkers, $m=1, n=1$ (Table 1). In addition, the position of the methyl group in N-C(α)-C(α')-N-C(α'') linker also had a favorable effect on activity: antiplasmodials **9**, **33** ($\text{CH}_3\text{-C}(\alpha)$) vs. **10**, **34** ($\text{CH}_3\text{-C}(\alpha')$, Table 1) and **11** ($\text{CH}_3\text{-C}(\alpha'')$, Table S1).

Table 1. In vitro antiplasmodial activity								
Compound	MW	In vitro antimalarial activity ^{a,b} (<i>P. falciparum</i> , IC ₅₀ , nM)			Toxicity HepG2 ^f (RAW 264.7) ^g IC ₅₀ , nM	MLM ^h (min)	HLM ⁱ (min)	SI HepG2 (RAW) /D6
		D6 ^c	C235 ^d	W2 ^e				
9 	383.97	1	1	3	(2901)	10	11	(2901)
10	383.97	2	9	10	(3771)	10	13	(1886)

								
22 	398.00	4	15	22	(3161)	11	16	790
23 	398.00	10	10	10	3045	60	60	305
24 	363.54	17	135	185	2317	60	60	136
25 	415.99	135	197	238	3967	21 (79 ^j)	13 (78 ^k)	29
26 	381.53	60	259	157				3
29 	412.02	21	23	22	24271	57	49	1156
30 	412.02	7	19	25	3090			441
31 	412.02	51	165	282	1612			32
32 	430.01	198	342	488	3549			18
33 	398.00	1	1	4	(2910)	60	57	(2910)
34 	398.00	6	7	7	25127	60	27	4188

36 	412.02	13	13	13	24271	60	60	1867
37 	377.57	7	102	126	1590			227
38 	430.01	143	171	203	3212			22
39 	395.56	154	546	394				3
42 	426.05	7	14	26	13273	60	60	1896
43 	426.05	9	47	81	2598			289
44 	391.59	12	286	393	1593			133
45 	444.04	288	590	547	2608			9
CQ	319.87	12	139	456				
MFQ	378.31	16	36	5				
ART	282.33	9	13	7				
<p>^a Antiplasmodial IC₅₀ values (nM) ([³H]hypoxanthine incorporation method) for isolates and clones of <i>P. falciparum</i>. ^b All in vitro experiments were performed as technical triplicates with R2 within 0.96–1. ^c CQ susceptible <i>P. falciparum</i> African D6 clone. ^d <i>P. falciparum</i> multidrug resistant C235 strain (Thailand). ^e CQ resistant <i>P. falciparum</i> Indochina W2 clone. ^f Hepatocellular carcinoma. ^g Rat Macrophage Cell Line. ^h Mouse liver microsomes. ⁱ Human liver microsomes. ^j Mouse hepatocytes. ^k Human hepatocytes.</p>								

1
2
3 Based on the above findings we turned our attention to antiplasmodials with methyls
4 at C(α) position in the series with extended linkers:
5

6
7
8 The examination of the series with m=2; n=1, n=2 (Scheme 1, Ad derivatives), revealed that
9 antiplasmodials with a 7-chloro-4-aminoquinoline moiety (**22**, **23**, **24**, **36**, **37**) had acceptable
10 SI (136-1,867), but we found low SI values for fluorine derivatives **25**, **38** (29, 22,
11 respectively), as well as considerable in vivo toxicity issues for compound **39** (*vide infra*). In
12 this series, good activity against CQS D6, CQR W2 and MDR C235 strains was afforded by
13 7-chloro derivatives **22**, **23**, and **36** (4-22 nM, Table 1). In order to directly compare the
14 influence of chlorine at C(7), des-chloroaminoquinolines **24** and **37** were prepared along with
15 their respective fluoro derivatives **26** and **39**. We found that omitting the Cl-C(7) substituent
16 did not affect the in vitro activity of **24** and **37** against the CQS D6 strain (IC₅₀=17 nM, 7 nM,
17 respectively, Table 1; c.f. compounds **35**, **44**, Table S1). However, the activity against the
18 CQR W2 and MDR C235 strains was significantly impaired; all des-chloro compounds were
19 8-86-fold less active than their 7-chloro-4-aminoquinoline analogs against the MDR C235
20 strain, and 10-24-fold less active against the CQR W2 strain (**23** vs. **24**; **36** vs. **37**; **33** vs. **35**;
21 **42** vs. **44**, Table 1 and Table S1). The introduction of a fluorine atom in position C(3)
22 failed to improve the in vitro activity. All F-C(3) derivatives with the adamantane carrier, **25**,
23 **38**, **26**, **39**, **32**, **45** (Table 1) and **20**, **21** (Table S1), were significantly less active than their
24 C(3)-H isosteres. On the nM scale they could be considered virtually inactive. The lack of in
25 vitro activity of C(3) fluorine derivatives was extended to CQ derivative **74**, which is ca. 34-
26 fold less active than CQ against the CQS D6 strain (Table S1).
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

50 The thiophene moiety was investigated on several occasions as an aromatic
51 contributor to the aminoquinoline pharmacophore.^{34, 38} As part of this report we analyzed the
52 effect of the benzothiophene contributor to the aminoquinoline moiety on antiplasmodial
53 activity (Table S1, Table 2). Two series of this class were prepared: C(2)- and C(3)-
54
55
56
57
58
59
60

1
2
3 substituted benzothiophenes. As a general remark, we note that benzothiophenes did not
4
5 improve the in vitro activity of aminoquinolines, and that the whole series was less potent
6
7 than the 7-chloro-4-aminoquinoline-thiophene-benzene combination.³⁸ C(2)-substituted
8
9 benzothiophenes were found less active in vitro than their C(3) isomers, with only **52** being as
10
11 active as CQ and MFQ against *P. falciparum* strains D6 and C235, respectively (Table S1).
12
13 Of twelve C(3) substituted benzothiophenes, five antiplasmodials were more active than CQ
14
15 against the CQS strain D6, and six were more active than MFQ against the MDR strain C235.
16
17 Compounds without a chlorine atom at the quinoline C(7) position (**54**, **62**, **66**), were much
18
19 less active than the whole benzothiophene series against all three *P. falciparum* strains. The
20
21 introduction of fluorine and cyano group at positions C(5) and C(6) (benzothiophene
22
23 numbering) did not appreciably influence the benzothiophene-aminoquinoline hybrids' in
24
25 vitro activity.
26
27
28
29
30
31

32 In vivo blood-stage efficacy studies

34
35 The compounds presented in Table 2 were examined for their in vivo efficacy in a mouse
36
37 model. A modified Thompson test was employed to determine the blood schizonticidal
38
39 efficacy of 14 compounds. In addition, in a separate host toxicity study, groups of five healthy
40
41 mice were dosed with 160 mg/kg/day \times 3 days of **34**, **33**, **23**, **24**, **25**, **42**, **74**, **73**, **75**, **58**, **59**, **63**,
42
43 without any overt clinical manifestations of toxicity. This was monitored by observations of
44
45 individual mouse behavior and appearance two times a day for 31 days. Compound **58** was
46
47 toxic at 160 mg/kg/day, and was therefore tested at 80 mg/kg/day, at which concentration it
48
49 did not exhibit any signs of toxicity.
50
51
52

53
54 Compounds were administered po at given doses on days 3, 4, and 5 after infection
55
56 using the Thompson test in C57Bl6 female mice infected with 1×10^6 *P. berghei* parasites.⁴²
57
58 The compounds with an ethylene linker (**1**, **34**, **33**) cleared the parasites on D6. However,
59
60

recrudescence occurred on days 10-14 and mice succumbed to malaria later on. Only at the higher dose of 160 mg/kg/day did the compound **33** cure 2 mice, while the minimal active dose (MAD) for **1**, and **34** was estimated at 40 and 80 mg/kg/day, respectively. Next, we tested the series of compounds with a propylene linker: **23**, **24**, **25**, and **26**. The 7-chloro-4-aminoquinoline isosteres **23** and **25** were tested in a dose-dependent manner and a minimal curative dose (MCD) of 80 mg/kg/day for both compounds was estimated. Although none of the two compounds showed curative effect at the lower doses employed, a small difference was noticed: aminoquinoline **23** was more potent than **25**, exhibiting MAD even at 10 mg/kg/day, and clearance with recrudescence at 40 and 20 mg/kg/day, while **25** was ineffective at 20 and 10 mg/kg/day (not shown). The analogs of **23** and **25**, isosteres **24** and **26**, were selected for examining the in vivo 4-aminoquinoline activity dependence on the chlorine substituent at C(7), at 80 mg/kg/day. As shown in Table 2, none of the compounds cured infection. However, isostere **24** (H-C(3)) was more active than **26** providing clearance with recrudescence at the administered concentration.

Compound	mg/kg/day	Parasitemia	Mice dead/ day died	Mice alive on day 31/total	Mean survival time (MST, day)
1	40	D6: negative, clearance. Two mice positive at D31	3/15-24	2/5	>21.4
34	80	D3: 3 mice 0.4%-0.5%; D6: 3 mice negative; D10: 2 mice negative, 1 mouse 0.2%; D14: 3 mice 1.6%-3.6%	1/14, 1/15, 1/17	0/3	15.3
33^c	160	D6: mice negative; D24: 3 mice 2.7%-9.6%; D28: 1 mouse 67.4%. D31: 2 mice negative	2/24, 1/28	2/5	27.6

23	160	D3: 5 mice 0.7%-1.2%; D6: 5 mice negative; D17: 5 mice negative; D31: 5 mice negative		5/5	>31
	80	D3: 5 mice 0.4%-0.5%; D6: 5 mice negative; D17: 5 mice negative; D31: 5 mice negative		5/5	>31
	40	D3: 5 mice 0.3%-2.4%; D6: 5 mice negative; D10: 5 mice negative; D14: 5 mice 0.2%-1.2%; D17: 2 mice 2.1%-4.6%	1/16, 2/17, 2/18	0/5	17.2
	20	D3: 4 mice 0.5%-3.5%; D6: 4 mice negative; D10: 4 mice 0.2%-0.4%; D14: 2 mice 1%-3.9%; D17: 1 mouse 3.5%	2/14, 1/15, 1/18	0/4	15.2
	10	D3: 5 mice 0.4%-1.6%; D6: 5 mice 0.18%-0.5%; D10: 5 mice 1.6%-8.9%	1/11, 3/13, 1/15	0/5	13
24	80	D3: 5 mice 0.2%-1%; D6: 5 mice negative; D10: 3 mice negative, 2 mice 0.2%-0.9%; D14: 5 mice 1.4%-2.5%	2/14, 3/16	0/5	15.2
25	160	D3: 5 mice 0.3%-0.5%; D6: 1 mouse ex., 4 mice negative; D17: 4 mice negative; D31: 4 mice negative	1/7	4/5	26.2
	80	D3: 6 mice 0.3%-1%; D6: 6 mice negative; D17: 6 mice negative; D31: 6 mice negative		6/6	>31
	40	D3: 5 mice 0.5%-3%; D6: 5 mice 1%-4.7%; D10: 5 mice 3.1%-16.3%; D14: 3 mice 5.6%-23%; D21: 2 mice 37.5%-62.4%	2/12, 1/21, 1/23, 1/24	0/5	18.4
26	80	D3: 5 mice 0.4%-1.4%; D6: 5 mice 1.3%-26.8%; D10: 3 mice 3.2%-25%; D14→D21: 1 mouse 7.7%→35%	2/8, 1/12, 1/13, 1/23	0/5	12.8
42	160	D6: 5 mice negative; D14: 4 mice negative, 1 mouse 0.3%; D17: 3 mice recrudescence, 1 mouse negative	1/17, 1/19, 1/21, 1/24, 1/26	0/5	21.4
	40	D6-D14: 0.4%-52.3%	1/12, 4/14	0/5	13.6

74	160	D3: 5 mice 0.5%-1.5%; D6: 5 mice negative; D10: 2 mice negative, 3 mice 0.1%-0.4%; D14: 5 mice 0.16%-3.4%; D17: 2 mice 6.9-10%	1/15, 1/16, 1/17, 2/18	0/5	16.8
73	160	D3: 5 mice 0.3%-1.3%; D6: 5 mice negative; D10: 5 mice 0.2% -0.4%; D14: 5 mice 3%-7%	3/14, 1/15, 1/16	0/5	14.6
75	160	D3: 6 mice 0.1%-1.5%; D6: 6 mice 0.2%-2.6%; D10: 5 mice 1.7%-4.9%; D14: 4 mice 9.6%-23.5%; D21: 4 mice 52-58%; D28: 1 mouse 60%.	1/9, 1/12, 2/22, 1/27, 1/29	0/6	20.2
58	80	D3: 4 mice 0.2%-0.7%; D6: 4 mice 0.2%-0.3%; D10: 4 mice 0.4%-0.6%; D14: 4 mice 1.3%-2.7%	4/15	0/4	14
59	160	D3: 5 mice 0.9%-2.3%; D6: 5 mice negative; D10: 5 mice negative; D14: 5 mice negative; D17: 5 mice 0.3%-4.4%; D21: 4 mice 2.3%-3.8%; D31: 3 mice positive	1/18, 1/21	3/5	26.4
63	160	D3: 4 mice 0.4%-0.9%; D6: 4 mice negative; D10: 3 mice negative, 1 mouse 0.2%; D14: 4 mice 0.2%-1.2%; D17: 4 mice 0.5%-1.7%; D21: 4 mice 1%-37%; D31: 3 mice positive	1/27	3/4	30
	80	D3: 5 mice 0.1%-1.7%; 5 mice negative; D10: 5 mice 0.2%-0.4%; D14: 5 mice 0.6%-3.5%; D17: 5 mice 1.2%-6.5%; D21: 5 mice 2.4%-15.2%; D31: 1 mouse positive	1/27, 3/31	1/5	30.2
CQ	160	D3: 5 mice 0.4%-0.9%; D6-D31: 5 mice negative		5/5	>31
	80	D3: 5 mice 0.5%-0.9%; D6: 5 mice negative; D10: 5 mice negative; D14: 3 mice negative, 2 mice 0.2%; D17: 3 mice negative, 1 mouse 4%; D21: 3 mice negative; D31: 3 mice negative.	1/17, 1/18	3/5	25.6
Infected controls	0	All mice died on day 6-8			

^a Groups of five *P. berghei* (ANKA strain) infected mice were treated with aminoquinolines suspended in 0.5% hydroxyethylcellulose-0.1% Tween 80 administered po once per day on days on 3-5 postinfection. Mice alive on day 31 with no parasites in a blood film were considered cured. ^b All compounds were tested in separate experiments (mouse groups) for toxicity. None exerted any toxic effect when dosed 3× 160 mg/kg/day. ^c Body temperatures were recorded to predict death according to IACUC protocols and parasitemia was captured prior to euthanizing the animal. The animal euthanasia point was 35.5 °C.

Next, we examined CQ-, and AQ-13-fluoro derivatives **74** and **73**, respectively, at the non-toxic dose of 160 mg/kg/day. Both compounds led to clearance with recrudescence, with no cure even at such a high dose.

Three benzothiophenes were examined for their efficacy at the selected doses (Table 2), and since 5-fluorobenzothiophene **58** was toxic at the higher dose it was tested at 80 mg/kg/day with no cure or survival. Its F-C(6) isomer **59** was examined at the 160 mg/kg/day dose, and exhibited clearance with recrudescence and survival of 3/5 animals, with MST >26 days (D31: 3 mice positive). The higher homolog of **58**, compound **63**, also provided clearance with recrudescence with an MST of 30 days, and survival of 3/4 animals. However, a dramatic difference emerged between the two homologs at 80 mg/kg/day. While **58** provided no clearance, homolog **63** led to clearances with recrudescence, 1/5 survival and twice the MST of **58**.

Inhibitory activity of *Plasmodium* liver stages

In vitro LS activity

In an attempt to develop multistage antimalarials, the possible LS-activity of our aminoquinoline blood stage antiplasmodials was investigated. The evaluation of the LS activity of our steroidal and non-steroidal tetraoxanes was also performed by measuring the luminescence intensity in human hepatoma cell line (Huh-7) cells infected with a firefly luciferase-expressing *P. berghei* line, as previously described.⁴³




Figure 2. Activity of compounds **23**, **25**, **36**, **38**, and **69** against *P. berghei* liver stages. Anti-*P. berghei* activity (infection scale in Relative Luminescence Units (RLU), bars) and toxicity to Huh7 cells (cell confluency scale in Relative Fluorescence Units (RFU), circles) are shown at several concentrations. The infection load of human hepatocellular carcinoma (Huh-7) cells was determined by bioluminescence measurements of cell lysates 48 h after infection with luciferase-expressing *P. berghei* parasites.

Initially, we evaluated the aminoquinoline blood-stage antiplasmodials with an adamantane carrier **23**, **36**, **25**, **38**, **69** for LS activity using primaquine as a positive control (Table 3). The percentage of Huh7 cell confluency was used as a measure of relative cell toxicity (Figure 2).

Table 3: Liver stage activity of 4-aminoquinoline and tetraoxane antiplasmodials.			
Compound	% LS infection (5 μM) ^a	LS IC ₅₀ (μM) ^b	LS in vivo, (mg/kg/day) ^c
23	0.1	1-2.5	
25	13 (2 μM)	0.31±0.14	16 (50); 8 (80)
69	~2	1-5	
36	8	2.5-5	
38	3	1-2.5	
74	~35	1-5	
76	~40	1-5	
77	31	<1	
78	0.55	0.33 ±0.05	40 (50); 9 (100)

PQ	62 (10 μ M)		
CQ		ca. 9 ²⁵	
ART	>75 ²³	ca. 12 ²⁵	
ATQ		22 nM ²⁵	

^a 10 000 Huh7 cells/well seeded 24 h before infection. 10 000 PbA-LuciGFPcon spz/well. Confluency measured by a fluorimetric assay. Infection was measured by a bioluminescence assay.

^b All dose-responses employed at least 5 dosages.


^c Measurement of liver load by qRT-PCR or by luminescence measurement. Mean of 2 independent experiments. All mice were infected by i.v. injection of 10,000 *P. berghei* sporozoites. Control groups were treated by oral gavage with vehicle (equivalent% of DMSO, in sunflower oil). Experimental groups were treated by oral gavage administration of **25** and **78** at the given doses.

The in vitro results presented in Table 3 indicate that 7-ACQ-Ad antiplasmodials **23**, **25**, **36**, **38**, **69** all exhibit relatively high potency against the liver forms of the parasite. LS inhibitory activity was assessed at seven concentrations (10 μ M – 0.1 μ M) and several compounds showed a dose-dependent behavior (Figure 2). Table 3 and Figure 2 show that at 5 μ M concentration, homologs **23** and **36** appear to be potent inhibitors of LS infection (0.1%, 8%, respectively). However, their activity dropped at lower concentrations yielding IC₅₀ values (1-2.5 μ M, 2.5-5 μ M), comparable to that of AQ-13, CQ.²⁵ We then explored the effect of the F-C(3) aminoquinoline substituent on LS inhibitory activity by assaying another homolog pair of fluoro isosteres **25** (\leftarrow **23**) and **38** (\leftarrow **36**). It was found that fluorine at C(3) significantly enhanced the activity of **25** in comparison to **23** (IC₅₀= 0.31 μ M vs. IC₅₀ = 1-2.5 μ M, respectively), and of **38** in comparison to **36** (IC₅₀=1-2.5 μ M vs. IC₅₀ = 2.5-5 μ M, respectively). Significant in vitro inhibitory activity (38%) of **25** was observed even at 125 nM concentration. To the best of our knowledge, the 3-fluoro compound **25** is the first 4-aminoquinoline that exerts evident and significant in vitro activity against *Plasmodium* LS (Figure 2). The effect of the fluorine on *Plasmodium* LS stage is additionally indicated by enhanced activity of 2-aminoquinoline **69** (a regioisomer of **25**, LS infection of **69** at 5 μ M

1
2
3 ~2%, IC_{50} = 1-5 μ M), and the activities of CQ fluoro derivative **74** and its 2-aminoquinoline
4 analog **76** (both IC_{50} = 1-5 μ M), Table 1, Figure 2. For comparison, LS activity of CQ is ~9
5 μ M.²⁵
6
7

8
9
10 Tetraoxanes were coupled to pyrimidine-nitriles²³ and primaquine,²⁴ and the
11 respective hybrids were found to be active against LS *Plasmodium* infection in vitro. During
12 our work on steroidal and non-steroidal tetraoxane molecules, we exposed their unexpected
13 stability to various reaction conditions (e.g., stability to LAH, pH1.6), and during in vitro
14 metabolism studies we noticed that tetraoxane moiety was resistant to metabolic
15 transformations.^{44, 45, 46} Therefore, we decided to evaluate the activity of these compounds
16 during the LS of parasite development. Tetraoxanes evaluated here have proven to be
17 excellent intra-erythrocytic antimalarials without any observed toxicity (Figure S1, Table
18 S2).^{44, 45, 46, 47, 48, 49}
19
20
21
22
23
24
25
26
27
28
29
30

31 The structures of the two most LS-active steroidal tetraoxanes are given in Figure 3,
32 and their respective activities are presented in Table 3.
33
34



48 **Figure 3.** Liver stage active tetraoxanes.
49

50 The two tetraoxanes differing only in their side chain exhibited pronounced inhibition
51 potencies with *n*-Pr amide **78** being ca. 3-fold more active than the respective methyl ester **77**
52 (IC_{50} = 0.33 μ M vs IC_{50} = <1 μ M, respectively). While methyl ester **77** was as active as
53 artemether, the in vitro potency of *n*-Pr amide **78** was comparable to the only LS-active
54
55
56
57
58
59
60

peroxide, artemisone ($IC_{50} \sim 0.04 \mu M^{25}$). Although the activity of other examined tetraoxanes was lower than that of **77** and **78** (Table S2), the overall results confirm that the tetraoxane moiety itself is probably capable of killing hepatic parasites without the aid of the 8-aminoquinoline moiety, which would lead to primaquine-like in vitro LS inhibition.

Tetraoxane **78** was also found to be moderately active when tested in vitro against stage IV-V *P. falciparum* gametocytes of the 3D7elo1-pfs16-CBG99 transgenic strain.⁵⁰ Compound **78** inhibited gametocytes viability with $IC_{50} = 1.16 \pm 0.37 \mu M$ (mean of three different experiments in duplicate). Control drugs employed were DHA ($IC_{50} = 0.44 \pm 0.29 \mu M$) and epoxomicin ($IC_{50} = 11.8 \pm 4.2 nM$).

In vivo LS-activity of 7-ACQ and tetraoxane molecules **25** and **78**.

To obtain a better insight into LS activity of our most active aminoquinoline **25** and tetraoxane **78**, we examined their in vivo activity against *P. berghei* LS. Thus, two groups of





Figure 4. Luminescence-based measurement of liver parasite loads after oral administration of compound **25** measured 48 h after infection and plotted as percentage of the infection levels in control mice.

1
2
3
4
5
6 five C57BL/6J mice were treated orally with **25** (DMSO+sunflower oil) at 50 mg/kg and 80
7
8 mg/kg doses during 3 consecutive days (D-1, D0, D+1, two independent experiments each).
9
10 Mice were infected with luciferase-expressing *P. berghei* sporozoites on D0, followed by
11
12 assessment of liver parasite load 48 h after infection by luminescence, and of the appearance
13
14 of parasites in the blood, disease symptoms, and survival. Our experiments employing 50
15
16 mg/kg and 80 mg/kg of the compound showed an 84% and 92% decrease of the *P. berghei*
17
18 liver load relative to controls, respectively (Figure 4).
19
20
21

22 In a similar experiment tetraoxane **78** administered at 50 mg/kg and 100 mg/kg
23
24 reduced the *P. berghei* liver load 60% and 91%, respectively (Figure 5, Table 3).
25
26
27



28
29
30
31
32
33
34
35
36
37
38
39
40
41 **Figure 5.** Parasite liver load in C57BL/6 mice measured 48 h after infection by qRT-PCR
42
43 of *P. berghei* 18S rRNA, normalized to hypoxanthine-guanine phosphoribosyltransferase
44
45 (*Hprt*) and plotted as percentage of the infection levels in control mice.
46
47
48

49 Discussion

50
51
52 The effect of numerous substituents at C(5), C(6), and C(7) of aminoquinoline
53
54 antiplasmodials is well explored.⁵¹ However, the antimalarial data on C(3) substituted
55
56 aminoquinolines are scarce,^{52, 53, 54} and the introduction of several alkyl/aryl groups at C(3) as
57
58
59
60

1
2
3 the logical consequence of the slightly improved activity of purified sontochin over that of
4 chloroquine, revealed that 4-aminoquinolines with C(3) alkyl/aryl substituents
5 (“pharmachins”)⁵⁵ are generally more active against CQR strains than CQ in vitro. We sought
6 to investigate the effect of a fluorine substituent at aminoquinoline position C(3) on β -hematin
7 formation, and on in vitro and in vivo blood stage activity; in addition we investigated the
8 activity of our inhibitors against *P. berghei* LS.
9
10
11
12
13
14
15
16

17 As part of our wider investigation of 4-aminoquinoline-adamantane combination as
18 pharmacophore-carrier symbiosis,³⁷ aminoquinoline **42** was first tested in vivo due to its
19 favorable microsomal metabolic stability and excellent SI data (Table 1, Table S1). Its rather
20 high logPow (QPlogPow 5.67,⁵⁶ experimental 5.87) was assisted by favorably predicted
21 apparent MDCK cell permeability QPPMDCK=1.29 \times 10⁻⁵ cm/s, rather low predicted
22 QPlogHERG toxicity IC₅₀= -5.076 (8.40 μ M), and its MTD was estimated at >160 mg/kg (no
23 host toxicity in mice up to 160 mg/kg/day \times 3 days). At 160 mg/kg/day, **42** showed excellent
24 suppression activity on D6-D14 (Table 2). However, the suppression activity was maintained
25 only with 1/5 mice on D17 and recrudescence occurred in 3/5 cases. MST was estimated at
26 21.4 days without any survival. Screening the compounds containing the ethylene bridge
27 (m=1, Fig. 1) with lower logP_{ow} **1**, **33**, **34** (Table 2), showed that only **33** cured 2/5 mice at
28 160 mg/kg/day. Next, we focused on compounds with a propylene linker (m=2, Fig. 1) and
29 with F-C(3) substituents. As shown in Scheme 2 all F-C(3) derivatives with an adamantane
30 carrier, **25**, **38**, **26**, **39**, **32**, **45** (Table 1) and **20**, **21** (Table S1), were significantly less active in
31 vitro than their H-C(3) isosteres. Homologs **23** and **36** possessed excellent potency across all
32 three *P. falciparum* strains, displayed excellent microsomal metabolic stability (Table 1) and
33 were therefore tested in vivo; in addition, their fluorine isosteres **25** and **38** were also chosen
34 for examination in a *P. berghei* challenge assay, despite the lower in vitro activity and low
35 microsomal stability of **25**. The respective des-chloro derivatives **24**, **26**, **37** and **39** were also
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 prepared in order to explore the influence of Cl-C(7) substituent at the aminoquinoline moiety
4
5 in a *P. berghei* challenge test. First, the compounds were tested for toxicity in mice at doses of
6
7 160 mg/kg/day \times 3 days. To our surprise, compounds **36**, **37**, **38**, **39** (all m=2, n=2; Scheme 1)
8
9 were all found toxic regardless of their respective substitution pattern at C(3) and C(7), which
10
11 is in sharp contrast to compounds **42** (m=3, n=2), **23**, **24**, **25**, **26** (m=2, n=1), and other
12
13 compounds listed in Table 2. It is interesting to note that in vivo toxicity results could not be
14
15 predicted based on in vitro toxicity parameters (c.f. RAW 264.7, HepG2 activity, Table 1).
16
17
18

19
20 As indicated above, isosteres **23** and **25** cured all mice from blood-stage malaria in the
21
22 Thompson test when dosed with 160, 80 mg/kg/day. The difference between the two became
23
24 evident at lower doses with isostere **23** providing clearance on D6 even at 20 mg/kg/day.
25
26 Fluoro derivative **25**, which demonstrated cures of blood stage malaria, was found very active
27
28 against LS *P. berghei* infection. The activity difference (and lack of mouse toxicity) between
29
30 the two isosteres warranted further investigation of their MOA. Thus, elementary information
31
32 about possible mechanism of action was obtained by submitting compounds **23** and **25**, **24**
33
34 and **26**, fluoro AQ-13 derivative **73**, F-CQ derivative **74** and its isomer **76**, and CQ as
35
36 standard, to a β -hematin inhibitory activity assay (BHIA).⁵⁷ The assay screens the interference
37
38 of potential antimalarials with the heme detoxification process employed by *P. falciparum*.
39
40 The results obtained indicated that the introduction of fluorine at the C(3) position of 4-
41
42 aminoquinolines greatly enhances the inhibition of β -hematin formation – **23** (1.84) vs **25**
43
44 (0.61); **24** (1.22) vs **26** (0.61); **CQ** (1.23) vs **74** (1.07) all IC₅₀ values, c.f. Table S1. The
45
46 influence of the F-C(3) substituent is consistent: it significantly enhances the BHIA and
47
48 indicates that such compounds would strongly interfere with heme detoxification process
49
50 within FV. However, this observation did not completely match with the compounds'
51
52 antimalarial activity: all tested fluoro derivatives exhibited slightly lower in vivo activity with
53
54 respect to their H isosteres (Table 2). Additional information on the influence of amine
55
56
57
58
59
60

1
2
3 regiochemistry on BHIA is obtained from the activity of **76**, the 2-aminoalkyl isomer of F-CQ
4
5 **74**. The lower activity of **76** as compared to **74** ($IC_{50}=2.43$ vs $IC_{50}=1.07$, respectively)
6
7 indicates that the interference of alkylaminoquinolines with the hematin formation probably
8
9 depends on the position of the alkylamino side chain as well.
10

11 The introduction of the F-C(3) substituent had a profound effect on the respective
12
13 dipoles (Table S1), e.g., **23** (6.04) vs. **25** (3.10), logP, **23** (3.85) vs. **25** (3.25), and pKa values.
14
15 Experimentally determined pKa values given in Table S1 clearly indicate that all examined
16
17 fluoro derivatives are monoprotinated at physiological pH, and to a great extent in FV.
18
19 Therefore, it appears that while 7-ACQ **23** interacts with hematin as diprotonated species, its
20
21 fluoro isostere **25** is expected to be mostly monoprotinated in FV, as well as within the liver
22
23 hepatocytes, where it exerts considerable inhibition of intra-hepatocytic parasites (for
24
25 distribution diagrams see Supporting information). Since the fluoro derivative **25** showed
26
27 remarkable in vivo activity both against LS and in asexual blood stages, we checked this
28
29 compound for hepatocyte metabolic stability followed by metabolite profiling. The intrinsic
30
31 clearance of $8.9 \mu\text{L}/\text{min}/10^6$ cells and $8.7 \mu\text{L}/\text{min}/10^6$ cells (human (h), mouse (m),
32
33 respectively) was estimated, therefore, significant stability in hepatocytes resulted – $t_{1/2} = 78$
34
35 min, 79 min (h, m), contrary to the reported poor metabolic stability when incubated with
36
37 hepatic microsomes (Table 1). The amount of remaining parent compound **25** at 120 min is
38
39 32% (h) and 37% (m). Metabolite identification exposed the monohydroxylation product as
40
41 the main one, and the product of dihydroxylation as minor - after 60 min: **25** : **25+16** : **25+32** =
42
43 35:26:1 (h); **25** : **25+16** : **25+32** = 33:26:1 (m). No Phase II products were detected. Further
44
45 metabolite analysis, isolation of the main metabolite and its testing will be carried out.
46
47
48
49
50

51 Arrhythmia can be induced by blockage of hERG channels by certain antimalarial
52
53 agents such as lumefantrine⁵⁸ and quinidine.⁵⁹ Both, **23** and **25** were submitted to hERG test
54
55 and $IC_{50} > 5 \mu\text{M}$ was estimated for both. At $5 \mu\text{M}$ concentration (highest concentration tested)
56
57 low inhibition was found – **23**, 11.7%; **25**, 8.54% (positive control quinidine $IC_{50} = 2.67 \mu\text{M}$).
58
59
60

1
2
3 The Ames test used to assess the mutagenic potential of **23** afforded positive results:
4
5 antiplasmodial **23** was negative at 500 $\mu\text{g/mL}$ against *S. typhimurium* strains TA98 and
6
7 TA100 (+ S9, - S9).
8

9
10 Current data on isosteres **23** and **25** indicate that both 4-aminoquinolines with
11
12 adamantane carrier are non-toxic and non-mutagenic antiplasmodials, they are relatively
13
14 metabolically stable in microsomal preparations (**23**) and in hepatocyte preparations (**25**).
15
16 Both compounds are efficient in vivo against *P. berghei* asexual RBC stages and a possible
17
18 indication on their MOA was obtained from BHIA experiments, which showed that all fluoro
19
20 7-ACQ-Ad derivatives were ca. 2-fold more active than their hydrogen isosteres.
21
22 Benzothiophene antiplasmodials were also found to be more active than CQ. This might
23
24 suggest that our compounds exert their activity on asexual RBC stages by interfering with the
25
26 *P. falciparum* detoxification process within the FV. Having in mind the pronounced two-stage
27
28 activity of **25** and the lack of host toxicity in mice for >31 days (indicating the non-toxicity of
29
30 monohydroxylation metabolite too), a strategy for the elucidation of LS MOA will be
31
32 developed.
33
34
35

36
37 In vitro LS activities of tetraoxanes with cholic acid carrier appear to be higher than
38
39 the simple cyclohexylidene analogs and their IC_{50} fall within ca. 1-5 μM (Table S2). The
40
41 highest activity was observed for *N*-propyl amide **78** (IC_{50} = 0.33 μM) and the proof of
42
43 concept was obtained after in vivo experiments with **78** reducing the *P. berghei* liver load by
44
45 91% at 100 mg/kg. Their mechanism of action is not clear at present; however, since the only
46
47 pharmacophore present is the tetraoxane moiety, we speculate that peroxide-generated
48
49 radicals could be involved in parasite elimination.
50

51
52 Asexual *P. falciparum* blood stage infection is followed by the development of sexual
53
54 parasite forms, gametocytes, which are transmitted to mosquitoes to continue their
55
56 development. Each round of schizogony is followed by differentiation of merozoites (ca. 1%)
57
58 into the male and female gametocytes through stages I–V. Tetraoxane **78** was found to
59
60

1
2
3 moderately inhibit the viability of *P. falciparum* late gametocytes (stage IV-V) with $IC_{50} =$
4
5 $1.16 \pm 0.37 \mu\text{M}$, which is only 3.5-fold higher than the IC_{50} against LS. Our results indicate
6
7 that tetraoxane **78** constitutes a good starting material for upgrading this pharmacophore
8
9 against LS and late stage gametocytes, in addition to its established asexual erythrocytic stage
10
11 activity.
12
13

14 15 16 **Conclusion**

17
18
19 The syntheses and antiplasmodial activities of variously substituted 4-aminoquinolines
20
21 coupled to adamantane carrier were described. The compounds exhibited pronounced in vitro
22
23 and in vivo activity against *P. berghei* in the Thompson test. The series with the
24
25 benzothiophene carrier showed lower activity against D6, C235, and W2 *P. falciparum* strains
26
27 in comparison to the adamantane series, and did not afford any cure in vivo. Tethering
28
29 fluorine atom to C(3) position of 4-aminoquinoline yielded fluoroaminoquinolines that
30
31 constitute intra-hepatocytic parasite inhibitors, with compound **25** having $IC_{50} = 0.33 \mu\text{M}$.
32
33 Compound **25** afforded up to 92% reduction of the in vivo *P. berghei* liver load at 80 mg/kg
34
35 dose. Testing our peroxide antimalarials as inhibitors of LS infection revealed that the
36
37 tetraoxane pharmacophore itself is also a powerful LS *P. berghei* parasite inhibitor (**78**: $IC_{50} =$
38
39 $0.33 \mu\text{M}$). A 91% reduction of the parasite liver load in mice was achieved at 100 mg/kg. To
40
41 the best of our knowledge, compounds **25** and **78** are the first examples of an 4-
42
43 aminoquinoline and a tetraoxane liver stage inhibitor, respectively. Neither of the compounds
44
45 was toxic in vivo and both possessed good metabolic stability. Their mode of action in the LS
46
47 is currently unknown. However, in the case of **78**, it could be anticipated that it is likely that
48
49 peroxide-generated radicals are involved in parasite elimination since the only pharmacophore
50
51 present is a tetraoxacyclohexane moiety. Compound **25** and its hepatocyte monohydroxylated
52
53 metabolite are not toxic, as confirmed by challenging the host, **25** is not mutagenic, and has a
54
55
56
57
58
59
60

1
2
3 good hERG profile. The results warrant focusing on the MOA of aminoquinoline **25** and its
4
5 monohydroxylated metabolite, as well as of tetraoxane **78** against *Plasmodium* LS.
6
7

8 Finally, the compounds exhibiting their action through two (**25**), or more (**78**) parasite
9
10 stages have good potential to be developed into efficient inhibitors to add to already existing
11
12 antimalarial drugs.³ The results presented here open the door to new LS-active chemotypes.
13
14

15 16 17 18 **Experimental section**

19
20 **Chemistry.** Melting points were determined on a Boetius PMHK apparatus and were not
21
22 corrected. IR spectra were taken on a Thermo-Scientific Nicolet 6700 FT-IR diamond crystal.
23
24 ¹H and ¹³C NMR spectra were recorded on a Varian Gemini-200 spectrometer (at 200 and 50
25
26 MHz, respectively), and a Bruker Ultrashield Advance III spectrometer (at 500 and 125 MHz,
27
28 respectively) in the indicated solvent (*vide infra*) using TMS as the internal standard.
29
30

31
32 Chemical shifts are expressed in ppm (δ) and coupling constants (J) in Hz. ESI-MS (HRMS)
33
34 spectra of the synthesized compounds were acquired on a Agilent Technologies 1200 Series
35
36 instrument equipped with Zorbax Eclipse Plus C18 (100 \times 2.1 mm i.d. 1.8 μ m) column and
37
38 DAD detector (190-450 nm) in combination with a 6210 Time-of-Flight LC/MS instrument in
39
40 positive and negative ion modes. The samples were dissolved in MeOH (HPLC grade). The
41
42 selected values were as follows: capillary voltage 4 kV; gas temperature 350°C; drying gas 12
43
44 L min⁻¹; nebulizer pressure 45 psig; fragmentator voltage: 70 V. Mass spectral analyses were
45
46 performed using electrospray ionization in positive ion mode on a Surveyor separations
47
48 module coupled to a ThermoFinnigan TSQ AM triple quadrupole mass spectrometer.
49
50

51
52 Compounds were analyzed for purity using an Agilent 1200 HPLC system equipped with
53
54 Quat Pump (G1311B), Injector (G1329B) 1260 ALS, TCC 1260 (G1316A) and Detector
55
56 1260 DAD VL+ (G1315C), and Waters 1525 HPLC dual pump system equipped with an
57
58 Alltech Select degasser system, and a dual k 2487 UV-Vis detector. LKB 5060-006 Micro
59
60

1
2
3 Plate Reader (Vienna, Austria) was used for BHIA test. Potentiometric titrations were
4
5 performed using CRISON pH-Burette 24 2S equipped with CRISON 50 29 micro-combined
6
7 pH electrode (CRISON INSTRUMENTS, S.A. Spain). The electrode was calibrated by means
8
9 of a strong acid – strong base titration in 0.1 M NaCl in MeOH:H₂O (1:1, v:v), using GLEE –
10
11 GLass Electrode Evaluation software;⁶⁰ standard potential E⁰=397.9±0.2 mV, slope 59.2±0.1
12
13 mV, and pK_w 13.84±0.01 are obtained as mean values of four titrations. All compounds were
14
15 >95% pure. For details see Supporting information.
16
17
18
19

20
21 N1-(7-Chloroquinolin-4-yl)ethane-1,2-diamine (**AQ2**), N1-(7-chloroquinolin-4-yl)propane-
22
23 1,3-diamine (**AQ3**), N1-(7-chloroquinolin-4-yl)butane-1,4-diamine (**AQ4**), N1-(7-
24
25 chloroquinolin-4-yl)-ethane-1,6-hexane (**AQ6**) and *N*-(quinolin-4-yl)propane-1,3-diamine
26
27 (**AQ7**) and *N*-(quinolin-4-yl)butane-1,4-diamine (**AQ8**) were prepared according to known
28
29 procedures.⁶¹
30
31
32
33

34 **General procedures for amination.**

35 Compounds 9, 27, 33, 40.

36
37 To a solution of an amine in CH₂Cl₂ were added the appropriate aldehyde (1 equiv) and
38
39 NaBH(OAc)₃ (2 equiv). After stirring the reaction mixture at r.t. for 24 h, aqueous NaOH was
40
41 added. The organic layer was separated, and the aqueous layer was washed with CH₂Cl₂. The
42
43 organic layer was dried over anh. Na₂SO₄, and the solvent was evaporated under reduced
44
45 pressure.
46
47
48
49

50 Compounds 23, 29, 36, 42, 46, 49, 52-67.

51
52 Amine (1.25-1.5 equiv) and appropriate aldehyde (1 equiv) were dissolved in MeOH/CH₂Cl₂
53
54 mixture (v:v; 2:1), anhydrous AcOH (1.25-1.5 equiv) was added, and the mixture was stirred
55
56 under Ar at r.t for 3 h. Then, NaBH₄ (6 equiv) was added and stirring was continued for
57
58
59
60

1
2
3 another 18 h. The solvent was removed under reduced pressure, and the residue was dissolved
4
5 in CH₂Cl₂. The organic layer was washed with 2M NH₄OH, water and brine, and dried over
6
7 Na₂SO₄. Finally, the solvent was evaporated under reduced pressure.
8
9

10
11 Compounds 10, 11, 20-22, 28, 30-31, 34-35, 41, 43, 68.

12
13
14 A mixture of diamine linker (1.2 equiv), 4,7-dichloroquinoline/4-chloroquinoline (1 equiv)
15
16 and phenol (15 equiv) were heated at 120-130 °C, with stirring for 24 h or the reaction
17
18 mixture was subjected to MW irradiation using a *Biotage Initiator 2.5 apparatus*. The
19
20 reaction mixture was cooled to r. t. and taken up in CH₂Cl₂. The organic layer was
21
22 successively washed several times with NaOH and finally with brine. The organic layer was
23
24 dried over anhydrous Na₂SO₄ and solvent was removed under reduced pressure to get a final
25
26 product.
27
28
29
30
31

32 General procedure for palladium catalyzed amination of quinolines (12, 24-26, 32, 37-39, 44-
33
34 45, 69-76).⁶²

35
36 Vial was charged with mixture of Pd(OAc)₂ (4 mol%) and DPEphos (8 mol%)/SPhos (8
37
38 mol%) in dioxane and stirred for a few minutes in Ar at r.t. Subsequently, a haloquinoline (1.0
39
40 equiv), amine (1.2 equiv) and K₃PO₄ (2.5 equiv) were added to the reaction mixture.
41
42 Resulting suspension was flushed with argon for several minutes. The vial was quickly
43
44 capped, heated to 85 °C overnight and then cooled down to r.t. The mixture was adsorbed
45
46 onto silica gel and purified.
47
48
49
50
51

52 General procedure for N-methylation of aminoquinolines (47-48, 50-51).⁶³

53
54 To a stirred solution of aminoquinolines (1equiv) in MeOH containing 37% aqueous
55
56 formaldehyde/acetone (2 equiv) was added mixture of ZnCl₂ (2 equiv) and NaHB₃CN (4
57
58 equiv) in MeOH. After the reaction mixture was stirred at r.t. for 4 h, the solution was taken
59
60

up in 0.1M NaOH and most of methanol was evaporated under reduced pressure. Aqueous solution was extracted with CH₂Cl₂, the combined extracts were washed with water and brine, dried over anh. Na₂SO₄ and the solvent was evaporated under reduced pressure.

7-Chloro-3-fluoroquinoline (**15**).

The mixture of amine **13** (500 mg, 2.80 mmol) in 48% tetrafluoroboric acid (730 μL, 5.60 mmol) was stirred on ice bath. Solution of sodium nitrite (193 mg, 2.80 mmol) in water (1 mL) was added dropwise after 15 min. The reaction mixture was stirred as the temperature rose to r.t., then filtered under reduced pressure to separate the yellow precipitate, diazonium tetrafluoroborate, which was washed with cold EtOH and Et₂O and then dried under reduced pressure. Diazonium salt was suspended in 1,2-dichlorobenzene and the reaction mixture was heated to 110 °C for 2 h. 7-Chloro-3-fluoroquinoline (**15**) was obtained after dry-flash chromatography (SiO₂, eluent: Hex/EtOAc = 9/1). **15**: Light-yellow solid softens at 76-80 °C (Hex). Yield: 356 mg (70%). IR (ATR): 3076m, 3053m, 2957w, 2857w, 1615m, 1596s, 1565m, 1485m, 1443m, 1414w, 1359w, 1333s, 1263m, 1184s, 1149w, 1129w, 1094w, 1071m, 988m, 961w, 935w, 913m, 891s, 812m, 776m, 645m, 537m, 512m, 477m, 461m cm⁻¹. ¹H-NMR (500 MHz, CDCl₃, δ): 8.81 (d, *J* = 2.8, H-C(2)), 8.11 (d, *J* = 1.8, H-C(8)), 7.80-7.70 (m, 2H, H-C(4) and H-C(5)), 7.55-7.50 (m, H-C(6)). ¹³C-NMR (125 MHz, CDCl₃, δ): 156.25 (d, *J* = 257.0, C(3)); 145.60 (C(8a)); 142.52 (d, *J* = 27.2, C(2)); 134.41 (C(7)); 128.73 (C(6)); 128.53 (C(5)); 129.20 (C(4a)); 128.35 (d, *J* = 5.1, C(8)); 118.31 (d, *J* = 17.3, C(4)). HRMS: *m/z* 182.01646 corresponds to molecular formula C₉H₅ClFNH⁺ (error in ppm: - 1.49).

3-Fluoro-quinoline (**16**).

Compound **16** was prepared according to the procedure described for compound **15**, starting from 3-aminoquinoline (**14**) (3 g, 20.96 mmol). 3-Fluoroquinoline (**16**) was obtained after chromatography (SiO₂, eluent: Hex/EtOAc = 9/1) as light-yellow solid (4.90 g, 97%). IR

(ATR): 3414w, 3064m, 2927w, 1652w, 1612s, 1560w, 1498s, 1463s, 1426m, 1371w, 1339s, 1213s, 1155s, 984m, 957w, 894m, 859w, 782m, 735m, 710w, 611w, 472w cm^{-1} . $^1\text{H-NMR}$ (500 MHz, CDCl_3 , δ): 8.82 (d, $J = 2.8$, H-C(2)), 8.13 (d, $J = 8.5$, H-C(8)), 7.85-7.75 (m, 2H, H-C(4) and H-C(5)), 7.75-7.65 (m, H-C(7)), 7.65-7.55 (m, H-C(6)). $^{13}\text{C-NMR}$ (125 MHz, CDCl_3 , δ): 156.25 (d, $J = 255.0$, C(3)); 145.40 (C(8a)); 141.54 (d, $J = 27.6$, C(2)); 129.49 (C(8)); 128.54 (C(7)); 128.42 (d, $J = 5.7$, C(4a)); 127.67 (C(6)); 127.27 (d, $J = 4.6$, C(5)); 118.32 (d, $J = 17.2$, C(4)). HRMS: m/z 148.05503 corresponds to molecular formula $\text{C}_9\text{H}_6\text{FNH}^+$ (error in ppm: - 4.52).

7-Chloro-3-fluoro-4-iodoquinoline (17).

Diisopropylamine (1.8 mL, 12.89 mmol) was added to n-butyllithium in hexane (4.9 mL, 12.28 mmol) in THF (6.5 mL) at -78°C . The stirring continued for 30 min, and a solution of **15** (2.23 g, 12.28 mmol) in 5.5 mL of THF was slowly added to formed LDA solution at -85°C . The resulting reaction mixture was stirred for 4 h at -78°C , and iodine (3.40 g, 13.51 mmol) in 6 mL THF was slowly added at -85°C . Stirring was continued for 2 h at -78°C before hydrolysis by 10 mL of THF/ $\text{H}_2\text{O} = 9/1$. Extraction with EtOAc, drying over anhydrous Na_2SO_4 , filtration and solvent removal afforded crude product which was purified by multiple chromatography: dry-flash (SiO_2 , gradient: Hex/EtOAc = 95/5 \rightarrow 9/1) and flash chromatography (Biotage SP1 NH column, eluent: Hex/EtOAc = 9/1). **17**: Yield 1.53 g (41%). IR (ATR): 3080m, 3031w, 2959w, 2925w, 1837w, 1738w, 1602m, 1585s, 1551s, 1481s, 1441s, 1384w, 1357s, 1317s, 1307s, 1254m, 1210m, 1179m, 1139m, 1075m, 1018w, 949w, 925s, 895w, 870m, 851w, 814m, 755m, 642w, 576w, 536m, 509m, 429w cm^{-1} . $^1\text{H-NMR}$ (500 MHz, CDCl_3 , δ): 8.60 (s, H-C(2)), 8.11 (d, $J = 2.0$, H-C(8)), 7.97 (d, $J = 8.9$, H-C(5)), 7.65-7.60 (m, H-C(6)). $^{13}\text{C-NMR}$ (125 MHz, CDCl_3 , δ): 157.11 (d, $J = 255.3$, C(3)); 145.36 (C(8a)); 140.80 (d, $J = 30.0$, C(2)); 135.37 (C(7)); 132.16 (C(5)); 129.93 (C(6));

1
2
3 129.20 (C(4a)); 128.63 (C(8)); 94.90 (d, $J = 23.0$, C(4)). HRMS: m/z 307.91326 corresponds
4
5 to molecular formula $C_9H_4ClFINH^+$ (error in ppm: - 0.36).

6
7 The compound **18** was also isolated as side product (359 mg, 9.5%).

8 9 **7-Chloro-3-fluoro-2-iodoquinoline (18).**

10
11 IR (ATR): 3171w, 3071w, 3037s, 2923s, 2853m, 1729w, 1689w, 1603m, 1481m, 1404m,
12
13 1373w, 1330s, 1297w, 1255w, 1191s, 1135w, 1071w, 1016m, 918m, 866w, 815w cm^{-1} . 1H -
14
15 NMR (500 MHz, $CDCl_3$, δ): 8.08 (d, $J = 1.8$, H-C(8)), 7.72 (d, $J = 8.7$, H-C(5)), 7.64 (d, $J =$
16
17 7.1, H-C(4)), 7.55 (dd, $J_1 = 2.0$, $J_2 = 8.8$, H-C(6)). ^{13}C -NMR (125 MHz, $CDCl_3$, δ): 154.77 (d,
18
19 $J = 256.0$, C(3)); 146.73 (C(8a)); 135.35 (C(7)); 129.27 (C(6)); 128.21 (d, $J = 20.0$, C(5));
20
21 127.94 (C(8)); 126.36 (C(4a)); 117.50 (d, $J = 85.2$, C(4)); 111.95 (d, $J = 120.1$, C(2)). HRMS:
22
23 m/z 307.91325 corresponds to molecular formula $C_9H_4ClFINH^+$ (error in ppm: - 0.40).
24
25
26
27
28

29 **3-Fluoro-4-iodoquinoline (19)**⁶⁴

30
31 Compound **19** was prepared according to the procedure described for compound **17**, starting
32
33 from 3-fluoroquinoline (1 g, 6.80 mmol). The crude product was purified by dry-flash
34
35 chromatography (SiO_2 , gradient: Hex/EtOAc = 95/5 \rightarrow 9/1). **19**: Yield: 1.30 g (70%). IR
36
37 (ATR): 3069m, 2924m, 2854w, 1586s, 1558s, 1491s, 1455s, 1414m, 1382w, 1339s, 1306s,
38
39 1262m, 1213s, 1141s, 1011w, 959w, 920m, 797w, 754s, 723m, 630w, 506w cm^{-1} . 1H -NMR
40
41 (500 MHz, $CDCl_3$, δ): 8.61 (s, H-C(2)), 8.09 (dd, $J_1 = 0.6$, $J_2 = 8.1$, H-C(8)), 8.02 (dd, $J_1 =$
42
43 1.5, $J_2 = 8.4$, H-C(5)), 7.75-7.70 (m, H-C(7)), 7.70-7.62 (m, H-C(6)). ^{13}C -NMR (125 MHz,
44
45 $CDCl_3$, δ): 157.02 (d, $J = 252.7$, C(3)); 145.33 (d, $J = 1.8$, C(8a)); 139.85 (d, $J = 30.7$, C(2));
46
47 130.87 (d, $J = 5.4$, C(5)); 130.62 (C(4a)); 129.86 (C(8)); 129.15 (d, $J = 2.7$, C(7)); 129.07
48
49 (C(6)); 95.19 (d, $J = 21.7$, C(4)). HRMS: m/z 273.95195 corresponds to molecular formula
50
51 $C_9H_5FINH^+$ (error in ppm: - 1.46).
52
53
54
55
56
57
58
59
60

N^2 -[2-(1-Adamantyl)ethyl]- N^1 -(7-chloroquinolin-4-yl)propane-1,2-diamine (34).

Compound **34** was prepared from amine linker **S32** (140 mg, 0.59 mmol) and 4,7-dichloroquinoline (97 mg, 0.49 mmol) in phenol (692 mg, 7.35 mmol) and was isolated after multiple chromatography: dry-flash (SiO₂, eluent: CH₂Cl₂/MeOH(NH₃ sat.) = 95/5) and flash chromatography (Biotage SP1 NH column, eluent: Hex/EtOAc = 1/1) as colorless powder softening at 134-136 °C. Yield: 174 mg (74%). IR (ATR): 3245w, 2895s, 2842m, 1737m, 1670w, 1566s, 1489w, 1449m, 1427m, 1367m, 1331w, 1304w, 1284w, 1238m, 1202w, 1156w, 1136m, 1076m, 1046m, 989w, 966w, 897w, 869w, 843m, 804m, 769m, 704w, 646w, 622w, 594w, 570w, 544w, 524w, 517w, 509w cm⁻¹. ¹H-NMR (500 MHz, CDCl₃, δ): 8.52 (d, J = 5.3, H-C(2)), 7.95 (d, J = 2.1, H-C(8)), 7.70 (d, J = 8.9, H-C(5)), 7.36 (dd, J_1 = 2.2, J_2 = 8.9, H-C(6)), 6.36 (d, J = 5.4, H-C(3)), 6.01 (bs, H-NAr), 3.35-3.25 (m, 1H, ArNHCH₂CH(CH₃)-), 3.15-3.05 (m, 1H, ArNHCH₂CH(CH₃)-), 3.05-2.95 (m, 1H, ArNHCH₂CH(CH₃)-), 2.80-2.70 (m, 1H, -NHCH₂CH₂Ad), 2.65-2.55 (m, 1H, -NHCH₂CH₂Ad), 1.94 (bs, 3H, -Ad), 1.75-1.55 (m, 6H, -Ad), 1.55-1.45 (m, 6H, -Ad), 1.35-1.25 (m, 3H, -NHCH₂CH₂Ad, H-N-), 1.22 (d, 3H, J = 6.0, CH₃). ¹³C-NMR (125 MHz, CDCl₃, δ): 152.12; 150.01; 149.19; 134.75; 128.76; 125.14; 121.25; 117.47; 99.20; 51.74; 47.22; 45.24; 42.77; 41.27; 37.10; 32.00; 28.64; 19.30. HRMS: m/z 398.23395 corresponds to molecular formula C₂₄H₃₂ClN₃H⁺ (error in ppm: - 4.53). Anal. (C₂₄H₃₂ClN₃) Calcd: C, 72.43; H, 8.10; N, 10.56. Found: C, 71.98; H, 8.49; N, 10.59. HPLC purity (λ = 330 nm): method A: RT 8.017 min, area 99.00%; method B: RT 9.696 min, area 98.87%.

 N^1 -[2-(1-Adamantyl)ethyl]- N^2 -(7-chloroquinolin-4-yl)propane-1,2-diamine (33).

Compound **33** was prepared from amine **S13** (180 mg, 0.76 mmol) and 1-adamantylacetaldehyde (136 mg, 0.76 mmol) using NaBH(OAc)₃ (58 mg, 1.52 mmol) and

1
2
3 was isolated after dry-flash chromatography: SiO₂ eluent: CH₂Cl₂/MeOH(NH₃ sat.) = 95/5)
4
5 and flash chromatography (Biotage SP1 RP column, gradient: MeOH/H₂O = 7/3→9/1) as
6
7 colorless solid (227 mg, 75%) softening at 46-47 °C. IR (ATR): 3252m, 3063w, 2896s,
8
9 2843m, 1610w, 1573s, 1537m, 1447m, 1426w, 1376m, 1330m, 1279w, 1241w, 1205w,
10
11 1134m, 1078w, 966w, 901w, 870m, 843w, 805m, 763m, 645w, 623w, 600w, 540w, 513w,
12
13 490w, 421w cm⁻¹. λ_{max}(ε) = 327 (10189), 253 (16143) nm. ¹H-NMR (500 MHz, CDCl₃, δ):
14
15 8.48 (d, *J* = 5.5, H-C(2)), 7.92 (d, *J* = 1.0, H-C(8)), 7.74 (d, *J* = 9.0, H-C(5)), 7.32 (dd, *J*₁ =
16
17 1.8, *J*₂ = 8.8, H-C(6)), 6.40 (d, *J* = 5.5, H-C(3)), 5.90 (br s, H-N), 3.75-3.65 (m, 1H,
18
19 ArNHCH(CH₃-), 2.95-2.85 (m, 2H, ArNHCH(CH₃)CH₂-), 2.70-2.60 (m, 2H, -CH₂CH₂Ad),
20
21 1.92 (br s, 3H, -Ad), 1.75-1.55 (m, 6H, -Ad), 1.48 (br s, 6H, -Ad), 1.35-1.20 (m, 6H, -CH₂Ad,
22
23 CH₃, H-N-). ¹³C-NMR (125 MHz, CDCl₃, δ): 151.82; 149.31; 149.13; 134.75; 128.46;
24
25 125.04; 121.44; 117.55; 99.31; 54.36, 47.41; 44.59; 44.44; 42.62; 37.03; 31.86; 28.58; 18.11.
26
27 HRMS: *m/z* 398.23435 corresponds to molecular formula C₂₄H₃₂ClN₃H⁺ (error in ppm: -
28
29 3.51). Anal. (C₂₄H₃₂ClN₃ × 1/2 H₂O) Calcd: C, 70.83; H, 8.17; N, 10.32. Found: C, 70.44; H,
30
31 7.91; N, 10.27. HPLC purity (λ = 330 nm): method A: RT 7.891 min, area 97.79%; method B:
32
33 RT 9.904 min, area 95.06%.

40 41 ***N*¹-(1-Adamantylmethyl)-*N*³-(7-chloroquinolin-4-yl)butane-1,3-diamine (23).**

42
43 Compound **23** was prepared from amine **S14** (430 mg, 1.72 mmol) and adamantane-1-
44
45 carboxaldehyde (283 mg, 1.72 mmol) using AcOH (141 μL, 2.15 mmol) and NaBH₄ (390 mg,
46
47 10.32 mmol), and was obtained after dry-flash chromatography: SiO₂, eluent:
48
49 CH₂Cl₂/MeOH(NH₃ sat.) = 95/5) as colorless foam (610 mg, 89%) softening at 135-138 °C.
50
51 IR (ATR): 3250m, 3061w, 2962w, 2897s, 2842m, 1611w, 1570s, 1542m, 1490w, 1450m,
52
53 1430m, 1365m, 1345w, 1330m, 1283w, 1253w, 1206w, 1184w, 1136w, 1079w, 928w, 901w,
54
55 870m, 850m, 821w, 801m, 770w, 756m, 637w, 612w, 602w, 502w cm⁻¹. λ_{max}(ε) = 326
56
57 (10000), 254 (16260) nm. ¹H-NMR (500 MHz, CDCl₃, δ): 8.49 (d, *J* = 5.4, H-C(2)), 7.92 (d, *J*
58
59
60

1
2
3 = 2.0, H-C(8)), 7.80 (d, $J = 8.9$, H-C(5)), 7.35-7.20 (m, H-C(6)), 7.08 (d, $J = 5.1$, H-N), 6.40
4 (d, $J = 5.4$, H-C(3)), 3.90-3.80 (m, 1H, ArNHCH(CH₃)-), 2.95-2.85 (m, 1H, -CH₂NHCH₂Ad),
5 2.85-2.70 (m, 1H, -CH₂NHCH₂Ad), 2.30 (s, 2H, -CH₂Ad), 2.01 (s, 3H, -Ad), 1.95-1.85 (m,
6 1H, ArNHCH(CH₃)CH₂-), 1.85-1.65 (m, 7H, ArNHCH(CH₃)CH₂-, -Ad), 1.65-1.50 (m, 7H, -
7 Ad, H-N-), 1.32 (d, 3H, $J = 6.3$, CH₃). ¹³C-NMR (125 MHz, CDCl₃, δ): 152.05; 149.61;
8 149.40; 134.59; 128.60; 124.54; 122.30; 117.67; 98.90; 63.68; 48.32; 47.81; 41.16; 37.17;
9 35.22; 33.36; 28.43; 19.36. HRMS: m/z 398.23673 corresponds to molecular formula
10 C₂₄H₃₂ClN₃H⁺ (error in ppm: + 2.45). Anal. (C₂₄H₃₂ClN₃ × 1/2H₂O) Calcd: C, 70.83; H, 8.17;
11 N, 10.32. Found: C, 70.71; H, 7.91; N, 10.53. HPLC purity ($\lambda = 330$ nm): method A: RT
12 7.989 min, area 99.68%; method B: RT 9.636 min, area 99.31%.

13
14
15
16
17
18
19
20
21
22
23
24
25 **N¹-(1-Adamantylmethyl)-N³-quinolin-4-ylbutane-1,3-diamine (24).**

26
27 Compound **24** was prepared from amine linker **S52** (120 mg, 0.64 mmol) and 4-
28 chloroquinoline (70 mg, 0.43 mmol) using Pd(OAc)₂ (3.79 mg 0.017 mmol), SPhos (13.88
29 mg, 0.036 mmol) and K₃PO₄ (224mg, 1.08 mmol), and was obtained after dry-flash
30 chromatography: SiO₂, gradient: EtOAc/[MeOH/(NH₃ aq) = 9/1] = 9/1 → 7/3) as light yellow
31 foam softening at 160-161 °C. Yield: 114 mg (73%). IR (ATR): 3270m, 3070m, 2965m,
32 2902s, 2846s, 1618s, 1580w, 1539s, 1451s, 1396m, 1395m, 1373w, 1340m, 1264m, 1151m,
33 1126m, 1103w, 809m, 763 m, 738m, 703w cm⁻¹. ¹H-NMR (500 MHz, CDCl₃, δ): 8.51 (d, $J =$
34 5.3, H-C(2)), 7.95 (dd, $J_1 = 0.8$, $J_2 = 8.4$, H-C(8)), 7.88 (dd, $J_1 = 0.8$, $J_2 = 8.4$, H-C(5)), 7.65-
35 7.55 (m, H-C(7)), 7.40-7.30 (m, H-C(6)), 6.87 (br s, H-N), 6.42 (d, $J = 5.6$, H-C(3)), 3.95-
36 3.80 (m, 1H, ArNHCH(CH₃)-), 2.95-2.85 (m, 1H, -CH₂NHCH₂Ad), 2.80-2.70 (m, 1H, -
37 CH₂NHCH₂Ad), 2.29 (s, 2H, -NHCH₂Ad), 2.05-1.95 (m, 3H, -Ad), 1.95-1.85 (m, 1H,
38 ArNHCH(CH₃)CH₂-), 1.84-1.60 (m, 8H, ArNHCH(CH₃)CH₂-, -Ad, H-N-), 1.58 (m, 6H, -
39 Ad), 1.32 (d, 3H, $J = 6.3$, CH₃). ¹³C-NMR (125 MHz, CDCl₃, δ): 150.88; 149.60; 148.43;
40 129.56; 128.87; 124.00; 120.55; 119.14; 98.58; 63.65; 48.12; 47.83; 41.10; 37.17; 35.46;
41 33.44; 28.44; 19.15. HRMS: m/z 364.27556 corresponds to molecular formula C₂₄H₃₃N₃H⁺
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

(error in ppm: + 2.30). Anal. ($C_{24}H_{33}N_3 \times 1/2H_2O$) Calcd: C, 77.37; H, 9.20; N, 11.28. Found: C, 77.37; H, 9.03; N, 11.18. HPLC purity ($\lambda = 330$ nm) method A: RT 0.953 min, area 99.05%; method B: RT 9.386 min, area 97.27%.

***N*¹-(1-Adamantylmethyl)-*N*³-(7-chloro-3-fluoroquinolin-4-yl)butane-1,3-diamine (25).**

Compound **25** was prepared from amine linker **S52** (135 mg, 0.57 mmol) and **17** (159.64 mg, 0.52 mmol) using Pd(OAc)₂ (5.2 mg, 0.023 mmol), SPhos (18.88 mg, 0.046 mmol) and K₃PO₄ (303 mg, 1.43 mmol), and was obtained after multiple chromatography: dry-flash (SiO₂, gradient: CH₂Cl₂/MeOH(NH₃ sat.) = 95/5 → 8/2) and flash chromatography (Biotage SP1 NH column, eluent: Hex/EtOAc = 7/3 and RP column, eluent: MeOH/H₂O = 8/2) as colorless viscous oil (101.7 mg, 47%). IR (ATR): 3280m, 3070w, 2902s, 2846s, 2674w, 1596s, 1572s, 1540m, 1486w, 1450m, 1422m, 1380m, 1346m, 1295w, 1262m, 1191m, 1148m, 1115m, 1078m, 983w, 927m, 902w, 879m, 813m, 761w, 737m, 656w, 539w, 426w cm⁻¹. $\lambda_{\max}(\epsilon) = 250$ (15462), 337 (9685) nm. ¹H-NMR (500 MHz, CDCl₃, δ): 8.48 (d, $J = 5.8$, H-C(2)), 7.92 (d, $J = 1.6$, H-C(8)), 7.86 (d, $J = 9.0$, H-C(5)), 7.35-7.20 (m, H-C(6)), 6.63 (d, $J = 4.6$, H-N), 4.40-4.25 (m, 1H, ArNHCH(CH₃)-), 3.00-2.85 (m, 1H, -CH₂NHCH₂Ad), 2.85-2.70 (m, 1H, -CH₂NHCH₂Ad), 2.30 (s, 2H, -CH₂Ad), 2.05-1.90 (m, 4H, -Ad, ArNHCH(CH₃)CH₂-), 1.90-1.60 (m, 8H, ArNHCH(CH₃)CH₂-, -Ad, H-N), 1.60-1.50 (m, 6H, -Ad), 1.30 (d, 3H, $J = 6.2$, CH₃). ¹³C-NMR (125 MHz, CDCl₃, δ): 146.60 (C(8a)); 143.19 (d, $J = 239.1$, C(3)); 142.46 (d, $J = 27.5$, C(2)); 136.12 (d, $J = 5.4$, C(4)); 133.76 (d, $J = 7.2$, C(7)); 128.63 (C(8)); 125.37 (C(6)); 122.94 (d, $J = 5.0$, C(5)); 119.64 (d, $J = 5.4$, C(4a)); 50.58 (d, $J = 8.6$, ArNHCH(CH₃)-); 47.52; 41.05; 37.08; 35.75; 33.30; 28.36; 21.60. HRMS: m/z 416.22602 corresponds to molecular formula C₂₄H₃₁ClFN₃H⁺ (error in ppm: - 0.75). Anal. (C₂₄H₃₁ClFN₃ × 2/3 H₂O) Calcd: C, 67.35; H, 7.61; N, 9.82. Found: C, 67.17; H, 7.47; N, 9.74. HPLC purity ($\lambda = 330$ nm) method A: RT 8.589 min, area 97.89%; method B: RT 11.009 min, area 97.73%.

***N*¹-(1-Adamantylmethyl)-*N*³-(3-fluoroquinolin-4-yl)butane-1,3-diamine (26).**

Compound **26** was prepared from amine linker **S52** (880 mg, 3.72 mmol) and **19** (1 g, 3.72 mmol) using Pd(OAc)₂ (33.4 mg, 0.15 mmol), DPEphos (160 mg, 0.30 mmol) and K₃PO₄ (1.97 g, 9.14 mmol), and was obtained after multiple chromatography: dry-flash (SiO₂, gradient:CH₂Cl₂/(NH₃ sat.) 95/5 → 7/3) and flash chromatography (Biotage SP1 NH column, eluent: Hex/EtOAc = 7/3 and RP column, gradient: MeOH/H₂O = 8/2 → 9/1) as colorless oil (572 mg, 49%). IR (ATR): 3265m, 2902s, 2845m, 1601s, 1574s, 1540s, 1496w, 1451m, 1398m, 1366w, 1345w, 1270w, 1202w, 1148w, 759m cm⁻¹. ¹H-NMR (500 MHz, CDCl₃, δ): 8.51 (d, *J* = 5.7, H-C(2)), 8.00-7.88 (m, H-C(8) and H-C(5)), 7.60-7.54 (m, H-C(7)), 7.41-7.35 (m, H-C(6)), 6.31 (s, H-N), 4.37-4.27 (m, 1H, ArNHCH(CH₃)-), 2.95-2.85 (m, 1H, -CH₂NHCH₂Ad), 2.85-2.75 (m, 1H, -CH₂NHCH₂Ad), 2.30 (s, 2H, -CH₂Ad), 2.01-1.96 (m, 3H, -Ad), 1.80-1.60 (m, 9H, ArNHCH(CH₃)CH₂-, -Ad, H-N-), 1.60-1.50 (m, 6H, -Ad), 1.30 (dd, *J*₁ = 0.9, *J*₂ = 6.42, -CH₃). ¹³C-NMR (125 MHz, CDCl₃, δ): 146.01 (C(8a)); 143.22 (d, *J* = 238.3, C(3)); 141.60 (d, *J* = 27.1, C(2)); 135.80 (d, *J* = 5.4, C(4)); 129.85 (C(8)); 127.90 (C(7)); 124.83 (C(6)); 124.68 (C(4a)); 121.27 (d, *J* = 5.4, C(5)); 63.70; 50.44 (d, *J* = 8.1, CH); 41.06; 37.15; 36.21; 33.33; 28.42; 21.71 (d, *J* = 2.7, CH₃). HRMS: *m/z* 382.26498 corresponds to molecular formula C₂₄H₃₂FN₃H⁺ (error in ppm: - 0.83). HPLC purity (λ = 330 nm) method A: RT 7.545 min, area 99.28%; method B: RT 8.251 min, area 99.45%.

***N*¹-[2-(1-Adamantyl)ethyl]-*N*⁴-(7-chloroquinolin-4-yl)pentane-1,4-diamine (42).**

Compound **42** was prepared from amine **S15** (320 mg, 1.21 mmol) and adamantane-1-carboxaldehyde (216 mg, 1.21 mmol) using AcOH (86 μL, 1.51 mmol) and NaBH₄ (92 mg, 2.42 mmol), and was obtained after dry-flash chromatography: SiO₂, eluent: CH₂Cl₂/MeOH(NH₃ sat.) = 9/1) as colorless foam (314 mg, 61%) softening at 49-51 °C. IR (ATR): 3422m, 2901s, 2845m, 1611w, 1578s, 1540w, 1450m, 1426w, 1379w, 1332w,

1
2
3 1280w, 1253w, 1201w, 1150w, 1081w, 905w, 877w, 854w, 806w, 768w, 646w, 601w, 401w
4
5 cm^{-1} . $\lambda_{\text{max}}(\epsilon) = 328 (11161), 255 (17040) \text{ nm}$. $^1\text{H-NMR}$ (500 MHz, CDCl_3 , δ): 8.50 (d, $J =$
6
7 5.4, H-C(2)), 7.93 (d, $J = 2.1$, H-C(8)), 7.73 (d, $J = 9.0$, H-C(5)), 7.33 (dd, $J_1 = 2.2, J_2 = 8.9$,
8
9 H-C(6)), 6.39 (d, $J = 5.5$, H-C(3)), 5.52 (br s, H-N), 3.75-3.60 (m, 1H, $\text{ArNHCH}(\text{CH}_3)-$),
10
11 2.75-2.65 (m, 2H, $-\text{CH}_2\text{NHCH}_2\text{CH}_2\text{Ad}$), 2.65-2.55 (m, 2H, $-\text{CH}_2\text{CH}_2\text{Ad}$), 1.93 (br s, 3H, -
12
13 Ad), 1.85-1.75 (m, 1H, $\text{ArNHCH}(\text{CH}_3)\text{CH}_2-$), 1.75-1.50 (m, 10H, $\text{ArNHCH}(\text{CH}_3)\text{CH}_2-$, -
14
15 $\text{CH}_2\text{CH}_2\text{NHCH}_2\text{CH}_2\text{Ad}$, -Ad, H-N-), 1.45 (m, 6H, -Ad), 1.35-1.25 (m, 5H, CH_3 , $-\text{CH}_2\text{Ad}$).
16
17 $^{13}\text{C-NMR}$ (125 MHz, CDCl_3 , δ): 151.95; 149.32; 149.11; 134.72; 128.68; 124.96; 121.35;
18
19 117.35; 99.12; 49.57; 48.30; 44.48; 44.33; 42.60; 37.08; 34.05; 31.82; 28.61; 26.33; 20.16.
20
21 HRMS: m/z 426.26752 corresponds to molecular formula $\text{C}_{26}\text{H}_{36}\text{ClN}_3\text{H}^+$ (error in ppm: +
22
23 1.10). Anal. ($\text{C}_{26}\text{H}_{36}\text{ClN}_3 \times 3/2 \text{ H}_2\text{O}$) Calcd: C, 68.93; H, 8.68; N, 9.27. Found: C, 69.08; H,
24
25 9.04; N, 9.48. HPLC purity ($\lambda = 330 \text{ nm}$): method A: RT 8.000 min, area 95.37%; method B:
26
27 RT 9.821 min, area 95.54%.

31
32
33
34 **N^4 -(7-Chloro-3-fluoroquinolin-4-yl)- N^1, N^1 -diethylpentane-1,4-diamine (74).**

35
36 Compound **74** was prepared from **17** (300 mg, 0.98 mmol) and 2-amino-5-
37
38 diethylaminopentane (378 μL , 1.95 mmol) using $\text{Pd}(\text{OAc})_2$ (8.8 mg, 0.039 mmol), DPEphos
39
40 (42 mg, 0.078 mmol) and K_3PO_4 (518 mg, 2.44 mmol), and was obtained after multiple
41
42 chromatography: dry-flash (SiO_2 , gradient: $\text{CH}_2\text{Cl}_2/\text{MeOH}(\text{NH}_3 \text{ sat.}) = 95/5 \rightarrow 8/2$) and flash
43
44 chromatography (Biotage SP1 NH column, eluent: Hex/EtOAc = 8/2) as colorless foam (329
45
46 mg, 54%) softening at 48-50 $^\circ\text{C}$. IR (ATR): 3304m, 3069w, 2970s, 2936m, 2871w, 2810w,
47
48 1595s, 1574s, 1536m, 1488w, 1452m, 1418m, 1380m, 1351m, 1294w, 1265m, 1221w,
49
50 1197w, 1140m, 1078m, 929w, 906w, 877w, 815m, 764w, 735w, 656w, 540w cm^{-1} . $\lambda_{\text{max}}(\epsilon) =$
51
52 336 (11554), 252 (16869) nm. $^1\text{H-NMR}$ (500 MHz, CDCl_3 , δ): 8.52 (d, $J = 5.5$, H-C(2)), 7.96
53
54 (d, $J = 2.2$, H-C(8)), 7.75 (d, $J = 9.1$, H-C(5)), 7.42-7.36 (m, H-C(6)), 4.73 (d, $J = 7.7$, -NH),
55
56 4.20-4.05 (m, 1H, $\text{ArNHCH}(\text{CH}_3)-$), 2.54 (q, 4H, $J = 7.2$, $-\text{NH}(\text{CH}_2\text{CH}_3)_2$), 2.50-2.43 (m, 2H,
57
58
59
60

1
2
3 $-CH_2NH(CH_2CH_3)_2$, 1.75-1.55 (m, 4H, ArNHCH(CH₃)CH₂-, ArNHCH(CH₃)CH₂CH₂-), 1.29
4
5 (dd, 3H, $J_1 = 0.9$, $J_2 = 6.3$, ArNHCH(CH₃)-), 1.01 (t, 6H, $J = 7.0$, -NH(CH₂CH₃)₂). ¹³C-NMR
6
7 (125 MHz, CDCl₃, δ): 146.62 (C(8a)); 143.42 (d, $J = 240.1$, C(3)); 142.45 (d, $J = 27.3$, C(2));
8
9 135.77 (d, $J = 6.3$, C(4)); 133.97 (C(7)); 128.96 (C(8)); 125.95 (C(6)); 121.89 (d, $J = 5.4$,
10
11 C(5)); 119.45 (d, $J = 4.5$, C(4a)); 52.54; 51.34 (d, $J = 8.1$, ArNHCH(CH₃)-); 46.78; 36.12;
12
13 23.70; 22.30; 11.26. HRMS: m/z 338.18020 corresponds to molecular formula
14
15 C₁₈H₂₅ClFN₃H⁺ (error in ppm: + 2.43. Anal. (C₁₈H₂₅ClFN₃ × 1/2 H₂O) Calcd: C, 62.33; H,
16
17 7.56; N, 12.11. Found: C, 62.00; H, 7.32; N, 11.85. HPLC purity (λ = 330 nm) method A: RT
18
19 7.677 min, area 98.24%; method B: RT 9.229 min, area 98.45%.

20
21
22
23
24
25
26 ***N'*-(7-Chloro-3-fluoroquinolin-4-yl)-*N,N*-diethylpropane-1,3-diamine (73).**

27
28 Compound **73** was prepared from **17** (100 mg, 0.32 mmol) and 3-diethylamino-1-propilamine
29
30 (105 μL, 0.65 mmol) using Pd(OAc)₂ (2.9 mg, 0.013 mmol), SPhos (10.7 mg, 0.026 mmol)
31
32 and K₃PO₄ (178 mg, 0.81 mmol), and was obtained after multiple chromatography: dry-flash
33
34 (SiO₂, gradient: CH₂Cl₂/MeOH(NH₃ sat.) = 95/5 → 8/2) and flash chromatography (Biotage
35
36 SP1 NH column, eluent: Hex/EtOAc = 85/15) as colorless foam (64 mg, 63%) softening at
37
38 47-46 °C. IR (ATR): 3234m, 3065w, 2970s, 2934m, 2872w, 2822m, 1640w, 1597s, 1574s,
39
40 1540m, 1489w, 1470w, 1454w, 1429m, 1380m, 1360m, 1294w, 1268w, 1246w, 1195m,
41
42 1166w, 1138m, 1077m, 934m, 892m, 812m, 762m, 737w, 654w, 622w, 540w cm⁻¹. λ_{max}(ε) =
43
44 337 (9690), 252 (15073) nm. ¹H-NMR (500 MHz, CDCl₃, δ): 8.45 (d, $J = 6.0$, H-C(2)), 7.91
45
46 (d, $J = 2.2$, H-C(8)), 7.75 (d, $J = 9.0$, H-C(5)), 7.32 (dd, $J_1 = 1.6$, $J_2 = 9.0$, H-C(6)), 3.90-3.80
47
48 (m, 2H, ArNHCH₂-), 2.72-2.68 (m, 2H, -CH₂NH(CH₂CH₃)₂), 2.65 (q, 4H, $J = 7.1$, -
49
50 NH(CH₂CH₃)₂), 2.04 (br s, H-N), 1.90-1.80 (m, 2H, ArNHCH₂CH₂-), 1.10 (t, 6H, $J = 7.1$, -
51
52 NH(CH₂CH₃)₂). ¹³C-NMR (125 MHz, CDCl₃): 146.60 (C(8a)); 143.50 (d, $J = 226.3$, C(3));
53
54 142.45 (d, $J = 27.2$, C(2)); 137.28 (d, $J = 5.9$, C(4)); 133.72 (d, $J = 7.2$, C(7)); 128.65 (C(8));
55
56
57
58
59
60

1
2
3 125.23 (C(6)); 122.74 (d, $J = 5.0$, C(5)); 119.28 (d, $J = 5.9$, C(4a)); 53.50; 47.44 (d, $J = 9.9$,
4 ArNHCH₂-); 47.03; 25.31; 11.28. HRMS: m/z 310.14830 corresponds to molecular formula
5 C₁₆H₂₁ClFN₃H⁺ (error in ppm: + 0.71). (+)ESI-HRMS (m/z (%)): 310.14830 ([M+H]⁺, 100);
6
7
8
9
10 calculated 310.14808 (error in ppm: 0.71). Anal. (C₁₆H₂₁ClFN₃ × 1/2 H₂O) Calcd: C, 60.28;
11
12 H, 6.96; N, 13.18. Found: C, 60.65; H, 7.07; N, 12.79. HPLC purity ($\lambda = 330$ nm) method A:
13
14 RT 7.585 min, area 99.71%; method B: RT 8.693 min, area 95.30%.

15
16
17
18
19 ***N'*-(7-Chloro-3-fluoroquinolin-2-yl)-*N,N*-diethylpropane-1,3-diamine (75).**

20
21 Compound **75** was prepared from 7-chloro-3-fluoro-2-iodoquinoline **18** (100mg, 0.32 mmol)
22
23 and 3-diethylamino-1-propilamine (74 μ L, 0.46 mmol) using Pd(OAc)₂ (2.0 mg, 0.009
24
25 mmol), SPhos (7.5 mg, 0.018 mmol) and K₃PO₄ (124 mg, 0.57 mmol), and was obtained after
26
27 multiple chromatography: dry-flash (SiO₂, gradient: CH₂Cl₂/MeOH(NH₃ sat.) = 95/5 → 8/2)
28
29 and flash chromatography (Biotage SP1 RP column, eluent; MeOH/H₂O = 8/2) as a yellow
30
31 viscous oil (51 mg, 72%). IR (ATR): 3450w, 3241m, 2970m, 2931m, 2873w, 2815m, 1639m,
32
33 1610w, 1563w, 1537s, 1497w, 1459m, 1416m, 1378w, 1338m, 1292m, 1256m, 1193m,
34
35 1146m, 1124m, 1098w, 1070m, 1032w, 979w, 921w, 885m, 800w, 761w, 739w, 604w,
36
37 514w, 476w cm⁻¹. $\lambda_{\max}(\epsilon) = 343$ (8682), 332 (8760), 242 (35685) nm. ¹H-NMR (500 MHz,
38
39 CDCl₃, δ): 7.69 (d, $J = 2.0$, H-C(8)), 7.50-7.40 (bs, 1H, *H*-N-), 7.41 (d, $J = 8.5$, H-C(5)), 7.33
40
41 (d, $J = 11.2$, H-C(4)), 7.14 (dd, $J_1 = 2.0$, $J_2 = 8.4$, H-C(6)), 3.75-3.60 (m, 2H, ArNHCH₂-),
42
43 2.64 (t, 2H, $J = 6.0$, -CH₂NH(CH₂CH₃)₂), 2.57 (q, 4H, $J = 7.1$, -NH(CH₂CH₃)₂), 1.90-1.75 (m,
44
45 2H, ArNHCH₂CH₂-), 1.07 (t, 6H, $J = 7.2$, -NH(CH₂CH₃)₂). ¹³C-NMR (125 MHz, CDCl₃, δ):
46
47 149.16 (d, $J = 14.0$, C(8a)); 147.58 (d, $J = 259.0$, C(3)); 145.54 (d, $J = 2.7$, C(2)); 133.77 (d, J
48
49 = 7.2, C(7)); 127.77 (d, $J = 5.0$, C(5)); 125.03 (C(8)); 122.80 (C(6)); 121.26 (d, $J = 4.5$,
50
51 C(4a)); 115.56 (d, $J = 15.4$, C(4)); 53.00; 46.83; 41.57; 25.11; 11.53. HRMS: m/z 310.14742
52
53 corresponds to molecular formula C₁₆H₂₁ClFN₃H⁺ (error in ppm: - 2.14). Anal. (C₁₆H₂₁ClFN₃
54
55
56
57
58
59
60

1
2
3 × 1/4 H₂O) Calcd: C, 61.14; H, 6.89; N, 13.37. Found: C, 61.33; H, 6.96; N, 13.40. HPLC
4
5 purity (λ = 330 nm) method A: RT 7.559 min, area 99.45%; method B: RT 8.647 min, area
6
7 99.28%.
8
9

10
11
12 ***N*-(7-chloroquinolin-4-yl)-*N'*-[(5-fluoro-1-benzothiophen-3-yl)methyl]propane-1,3-**
13
14 **diamine (58).**
15

16
17 Compound **58** was prepared from aldehyde **S75** (181 mg, 1.00 mmol) using amine **AQ3** (353
18
19 mg, 1.50mmol). The product was purified using column chromatography (dry-flash, SiO₂,
20
21 eluent EtOAc/Hex gradient 1/9 → EtOAc, EtOAc/MeOH gradient 95/5 → 4/6, flash, Biotage
22
23 SP1, NH column, eluent EtOAc/Hex gradient 8/2 → EtOAc, EtOAc/MeOH gradient 95/5 →
24
25 MeOH and flash, Biotage SP1, SiO₂ column, eluent EtOAc/MeOH+NH₃(9/1) gradient 95/5
26
27 → 7/3). Final product **58** was obtained as colorless powder (269 mg, 67%). M.p. = 133 – 134
28
29 °C. IR (ATR): 3240m, 3060w, 2953w, 2852w, 1607m, 1579s, 1535m, 1435m, 1360w,
30
31 1329m, 1279w, 1251w, 1230w, 1205w, 1137w, 1104w, 1082w, 913w, 850w, 800w, 761w,
32
33 718w, 665w, 642w cm⁻¹. ¹H NMR (500 MHz, CDCl₃, δ): 8.50-8.47 (m, H-C(2')), 7.90-7.88
34
35 (m, H-C(8')), 7.83 (dd, *J*₁ = 4.8, *J*₂ = 8.9, H-C(7)), 7.46 (dd, *J*₁ = 2.4, *J*₂ = 9.5, H-C(4)), 7.43
36
37 (s, H-C(2)), 7.35 (d, *J* = 9.0, H-C(5')), 7.17-7.09 (m, H-C(6) and H-N exchangeable with
38
39 D₂O), 7.01-6.97 (m, H-C(6')), 6.32 (d, *J* = 5.5, H-C(3')), 4.06 (s, 2H, -
40
41 CH₂NHCH₂CH₂CH₂NH-), 3.45-3.40 (m, 2H, -CH₂NHCH₂CH₂CH₂NH-), 3.06-3.02 (m, 2H, -
42
43 CH₂NHCH₂CH₂CH₂NH-), 2.01-1.94 (m, 2H, -CH₂NHCH₂CH₂CH₂NH-), 1.69 (bs, H-N
44
45 exchangeable with D₂O). ¹³C NMR (125 MHz, CDCl₃, δ): 160.88 (d, *J* = 240.1), 152.04,
46
47 150.24, 149.05, 139.37 (d, *J* = 9.0), 135.98, 134.48 (d, *J* = 22.5), 128.48, 125.89, 124.81,
48
49 124.17 (d, *J* = 9.9), 121.61, 117.36, 113.51 (d, *J* = 25.3), 107.28 (d, *J* = 22.5), 98.39, 49.36,
50
51 47.64, 43.68, 27.60. HRMS: *m/z* 400.10388 corresponds to molecular formula
52
53 C₂₁H₁₉ClFN₃SH⁺ (error in ppm -1.56). Anal. (C₂₁H₁₉ClFN₃S) Calcd: C, 63.07; H, 4.79; N,
54
55
56
57
58
59
60

1
2
3 10.51; S, 8.02. Found: C, 63.09; H, 4.77; N, 10.30; S, 7.72. HPLC purity ($\lambda = 330$ nm):
4
5 method A: RT 8.998 min, area 98.73%; method B: RT 8.544 min, area 99.17%.
6
7
8
9

10 ***N*-(7-chloroquinolin-4-yl)-*N'*-[(6-fluoro-1-benzothiophen-3-yl)methyl]propane-1,3-**
11 **diamine (59).**
12
13

14 Compound **59** was prepared from aldehyde **S76** (161 mg, 0.888 mmol) using amine **AQ3**
15 (314 mg, 1.33 mmol). The product was purified using column chromatography (dry-flash,
16 SiO₂, eluent EtOAc/Hex gradient 1/9 → EtOAc, EtOAc/MeOH gradient 95/5 → 4/6, flash,
17 Biotage SP1, NH column, eluent EtOAc/Hex gradient 8/2 → EtOAc, EtOAc/MeOH gradient
18 95/5 → MeOH and flash, Biotage SP1, SiO₂ column, eluent EtOAc/MeOH+NH₃(9/1)
19 gradient 95/5 → 3/7). Final product **59** was obtained as colorless powder (194 mg, 54%).
20
21 M.p. = 149 – 150 °C. IR (ATR): 3433w, 3294m, 3210m, 3067m, 3015m, 2927m, 2853m,
22 1606m, 1581s, 1539m, 1466m, 1430w, 1369w, 1330w, 1282w, 1251w, 1208w, 1168w,
23 1139w, 1109w, 1080w, 896w, 854w, 807w, 761w, 737w, 683w, 646w cm⁻¹. ¹H NMR (500
24 MHz, CDCl₃, δ): 8.50-8.47 (m, H-C(2')), 7.90 (bs, H-C(8')), 7.74-7.70 (m, H-C(4)), 7.60-7.56
25 (m, H-C(7)), 7.37-7.33 (m, H-C(5')), 7.30 (s, H-C(2)), 7.22 (bs, H-N exchangeable with D₂O),
26 7.07-7.01 (m, H-C(5)), 7.00-6.96 (m, H-C(6')), 6.31 (d, $J = 5.2$, H-C(3')), 4.08 (s, 2H, -
27 CH₂NHCH₂CH₂CH₂NH-), 3.45-3.40 (m, 2H, -CH₂NHCH₂CH₂CH₂NH-), 3.06-3.03 (m, 2H, -
28 CH₂NHCH₂CH₂CH₂NH-), 2.01-1.95 (m, 2H, -CH₂NHCH₂CH₂CH₂NH-), 1.76 (bs, H-N
29 exchangeable with D₂O). ¹³C NMR (125 MHz, CDCl₃, δ): 160.70 (d, $J = 243.7$), 151.97,
30 150.31, 141.61 (d, $J = 9.9$), 134.71, 134.12, 128.39, 124.83, 122.98, 122.46 (d, $J = 10.0$),
31 121.70, 117.33, 113.28 (d, $J = 23.5$), 109.07 (d, $J = 25.3$), 98.36, 49.37, 47.69, 43.75, 27.52.
32
33 HRMS: m/z 400.10427 corresponds to molecular formula C₂₁H₁₉ClFN₃SH⁺ (error in ppm -
34 0.56). HPLC purity ($\lambda = 330$ nm) method A: RT 9.049 min, area 95.99%; ($\lambda = 254$ nm)
35 method B: RT 7.455 min, area 97.07%.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 ***N*-(7-chloroquinolin-4-yl)-*N'*-[(5-fluoro-1-benzothiophen-3-yl)methyl]butane-1,4-**
4
5 **diamine (63).**
6

7 Compound **63** was prepared from aldehyde **S75** (181 mg, 1.00 mmol) using amine **AQ4** (376
8 mg, 1.51 mmol). The product was purified using column chromatography (dry-flash, SiO₂,
9 eluent EtOAc/Hex gradient 1/9 → EtOAc, EtOAc/MeOH gradient 95/5 → 1/1, flash, Biotage
10 SP1, NH column, eluent EtOAc/Hex gradient 8/2 → EtOAc, EtOAc/MeOH gradient 95/5 →
11 MeOH and flash, Biotage SP1, SiO₂ column, eluent EtOAc/MeOH+NH₃(9/1) gradient 95/5
12 → 1/1). Final product **63** was obtained as a white powder (251 mg, 60%). M.p. = 110 – 113
13 °C. IR (ATR): 3228m, 3063w, 2945w, 2855w, 2810w, 1579s, 1543m, 1490w, 1433m, 1366w,
14 1330w, 1279w, 1245w, 1225w, 1196w, 1161w, 1134w, 1078w, 954w, 898w, 854w, 806w,
15 782w, 640w, 619w, 543w, 483w, 452w, 423w cm⁻¹. ¹H NMR (500 MHz, CDCl₃, δ): 8.50 (d,
16 *J* = 5.2, H-C(2')), 7.94-7.91 (m, H-C(8')), 7.77 (dd, *J*₁ = 4.8, *J*₂ = 8.7, H-C(7)), 7.59 (d, *J* =
17 8.9, H-C(5')), 7.51-7.46 (m, H-C(4)), 7.38 (s, H-C(2)), 7.18-7.14 (m, H-C(6')), 7.10 (td, *J*₁ =
18 2.4, *J*₂ = 8.8, H-C(6)), 6.36 (d, *J* = 5.3, H-C(3')), 5.65 (bs, H-N exchangeable with D₂O), 4.00
19 (s, 2H, -CH₂NHCH₂CH₂CH₂CH₂NH-), 3.32-3.27 (m, 2H, -CH₂NHCH₂CH₂CH₂CH₂NH-),
20 2.80 (t, *J* = 6.6, 2H, -CH₂NHCH₂CH₂CH₂CH₂NH-), 1.88 (quin, 2H, *J* = 6.8, -
21 CH₂NHCH₂CH₂CH₂CH₂NH-), 1.75-1.55 (m, 2H, -CH₂NHCH₂CH₂CH₂CH₂NH- and H-N
22 exchangeable with D₂O). ¹³C NMR (125 MHz, CDCl₃, δ): 160.74 (d, *J* = 240.1), 152.01,
23 149.82, 149.08, 139.46 (d, *J* = 9.0), 135.97, 134.81, 134.74 (d, *J* = 9.0), 128.68, 125.43,
24 125.02, 123.98 (d, *J* = 9.0), 121.05, 117.18, 113.29 (d, *J* = 24.4), 107.47 (d, *J* = 22.6), 98.91,
25 48.99, 47.67, 43.14, 27.69, 26.40. HRMS: *m/z* 414.11939 corresponds corresponds to
26 molecular formula C₂₂H₂₁ClFN₃SH⁺ (error in ppm -1.85). HPLC purity (λ = 330 nm) method
27 A: RT 9.136 min, area 99.72%; (λ = 254 nm) method B: RT 7.522 min, area 95.22%.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53

54 ***Plasmodium* asexual blood stages.**

55
56 In vitro antiplasmodial activity
57
58
59
60

1
2
3 The in vitro antimalarial drug susceptibility screen is a modification of the procedures
4
5 previously published by Desjardins et al.⁶⁵ with modifications developed by Milhous et al.⁴¹
6
7 and full details are given in Ref.42. All synthesized aminoquinolines were screened in vitro
8
9 against the following *P. falciparum* strains: CQ and MFQ susceptible strain D6 (clone of
10
11 SierraI/UNC isolate), CQ resistant but MFQ susceptible strain W2 (clone of Indochina I
12
13 isolate), and CQ and MFQ resistant strain TM91C235 (clone of South East Asian isolate).
14
15 Assessment of compound toxicity in a HepG2 (hepatocellular carcinoma) cells exactly
16
17 followed the protocol described in detail.⁶⁶
18
19

20 In vivo efficacy studies

21
22 The *P. berghei* mouse efficacy tests were conducted using a modified version of the
23
24 Thompson test. Briefly, groups of five mice were inoculated intraperitoneally with
25
26 erythrocytes infected with *P. berghei* on day 0. Drugs were suspended in 0.5%
27
28 hydroxyethylcellulose-0.1% Tween 80 and administered orally once a day beginning on day 3
29
30 post infection. Dosings are shown in Table 2. All untreated infected (control) mice showed
31
32 parasitemia on day 3 and high parasite counts on day 6, with levels of parasitemia between
33
34 37.8% and 57.8% (mean 45.96%), succumbed to infection on day 6-8. Accordingly, the test
35
36 was considered valid. Cure was defined as survival (with no parasitemia) until day 31 post-
37
38 treatment. Parasitemia was determined by thin-blood Giemsa-stained smears prepared from
39
40 mice tail blood of each animal on study days 0, 3, 6, 10, 13, 17, 20, 24, 27, and 31
41
42 (postinfection). The slides were examined under a light microscope.
43
44

45
46 In a separate host toxicity study, groups of five healthy mice were administered 160
47
48 mg/kg/day ×3 days of a compound under investigation. Monitoring individual mouse
49
50 behavior and appearance two times a day for 31 days revealed no overt clinical manifestations
51
52 of toxicity.
53
54

55
56 The study followed the International Guiding Principles for biomedical research involving
57
58 animals, and was reviewed by a local Ethics Committee and approved by the Veterinary
59
60

1
2
3 Directorate at the Ministry of Agriculture and Environmental Protection of Serbia (decision
4
5 no. 323-07-02444/2014-05/1).
6
7

8 9 **Antimalarial activity against stage IV-V *P. falciparum* gametocytes.**

10 Gametocytes cultures were conducted as described recently.⁵⁰ The 3D7elo1-pfs16-CBG99
11
12 transgenic strain⁶⁷ expressing the CBG99 luciferase under the pfs16 gametocyte specific
13
14 promoter was used in all the experiments and the luciferase activity was taken as measure of
15
16 gametocytes viability. Epoxomicin and dihydroartemisinin have been used as positive
17
18 controls.
19
20

21
22 The IC₅₀, the dose which induces 50% inhibition of gametocytes viability, was extrapolated
23
24 from the non-linear regression analysis of the dose –response curve (software Gen5 1.10
25
26 provided with the Synergy4 plate reader, Biotek). The percentage of gametocytes viability
27
28 was calculated as $100 \times [(OD \text{ treated sample} - OD \text{ blank}) / (OD \text{ untreated sample} - OD$
29
30 $\text{blank})]$ where “blank” is the sample treated with 100 nM epoxomicin which completely kills
31
32 gametocytes.
33
34
35
36
37

38 **Liver stage of *P. berghei* infection.**

39 In vitro LS activity

40
41 Inhibition of liver-stage infection by test compounds was determined by measuring the
42
43 luminescence intensity in Huh-7 cells infected with a firefly luciferase-expressing *P. berghei*
44
45 line, as previously described.⁴³ Briefly, Huh-7 cells, a human hepatoma cell line, were
46
47 cultured in 1640 RPMI medium supplemented with 10% v/v fetal bovine serum, 1% v/v
48
49 nonessential amino acids, 1% v/v penicillin/streptomycin, 1% v/v glutamine, and 10 mM 4-
50
51 (2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES), pH 7, and maintained at 37 °C
52
53 with 5% CO₂. For infection assays, Huh-7 cells (1.0×10^4 per well) were seeded in 96-well
54
55 plates the day before drug treatment and infection. The medium was replaced by medium
56
57
58
59
60

1
2
3 containing the appropriate concentration of each compound approximately 1 h prior to
4
5 infection with sporozoites freshly obtained through disruption of salivary glands of infected
6
7 female *Anopheles stephensi* mosquitoes. Sporozoite addition was followed by centrifugation
8
9 at 1700 g for 5 min. Parasite infection load was measured 48 h after infection by a
10
11 bioluminescence assay (Biotium). The effect of the compounds on the viability of Huh-7 cells
12
13 was assessed by the AlamarBlue assay (Invitrogen, U.K.) using the manufacturer's protocol.
14
15

16 In vivo liver stage assay

17
18 C57Bl/6J mice were infected by intravenous inoculation of 5×10^3 luciferase-expressing *P.*
19
20 *berghei* ANKA sporozoites freshly dissected from the salivary glands of infected *Anopheles*
21
22 *stephensi* mosquitoes. The compound was administered by intraperitoneal injection 24 h prior
23
24 to infection, at the time of infection, and 24 h after infection. An equivalent amount of drug
25
26 vehicle was injected in control mice.
27
28

29
30 Determination of liver parasite loads *in vivo* was carried out as previously described, either by
31
32 quantitative real-time PCR (qRT-PCR)⁶⁸ or by luminescence measurement.⁴³ For qRT-PCR
33
34 analyses, livers were collected and homogenized in a denaturing solution (4 M guanidine
35
36 thiocyanate; 25 mM sodium citrate [pH 7], 0.5% sarcosyl, and 0.7% β -mercaptoethanol in
37
38 diethyl pyrocarbonate-treated water) 48 h after sporozoite injection. Total RNA was extracted
39
40 using the Qiagen RNeasy Mini kit according to the manufacturer's instructions. RNA for
41
42 infection measurements was converted into cDNA by using the Nzytech kit according to the
43
44 manufacturer's protocol. The quantitative real-time PCRs (qRT-PCRs) used the Applied
45
46 Biosystems Power SYBR green PCR master mix and were performed according to the
47
48 manufacturer's instructions on an ABI Prism 7500 Fast system (Applied Biosystems).
49
50 Amplification reactions were carried out in a total reaction volume of 20 μ L, using 1 μ g
51
52 cDNA and employing *P. berghei* ANKA 18S rRNA gene- or housekeeping gene-specific
53
54 primers. Relative amounts of *P. berghei* ANKA mRNA were calculated against the amount of
55
56 the hypoxanthine guanine phosphoribosyltransferase (HPRT) housekeeping gene. Primer
57
58
59
60

1
2
3 sequences specific to each gene were as follows: for the *P. berghei* ANKA 18S rRNA gene,
4
5 5'-AAG CAT TAA ATA AAG CGA ATA CAT CCT TAC-3' and 5'-GGA GAT TGG TTT
6
7 TGA CGT TTA TGT G-3'; for the mouse HPRT gene, 5'-TGC TCG AGA TGT GAT GAA
8
9 GG-3' and 5'-TCC CCT GTT GAC TGG TCA TT-3'; and for the human HPRT gene, 5'-
10
11 TGC TCG AGA TGT GAT GAA GG-3' and 5'-TCC CCT GTT GAC TGG TCA TT-3'. For
12
13 bioluminescence measurements, mice were infected with firefly luciferase-expressing *P.*
14
15 *berghei* sporozoites and liver parasite loads were measured 46 h later on an IVIS Lumina
16
17 system, following subcutaneous injection of the luciferin substrate.
18
19

20 **Metabolic Stability Studies.**

21 Liver microsomal preparations

22
23 Selected samples were tested in human and mouse liver microsomal preparations exactly
24
25 following the protocol described in detail.³⁸
26
27
28
29

30 Hepatocyte Metabolic Stability (performed at Cyprotex US, LLC, Watertown, MA)

31
32 Compound **25** was incubated in duplicate with cryopreserved hepatocytes at 37 °C. The cells
33
34 were thawed; viable cells counted, and equilibrated. After 30 min equilibration at 37 °C with
35
36 gentle agitation, the test compounds were added into the cells to give the desired final
37
38 concentration of 1 μM. The cell suspension was incubated at 37 °C as described above. At the
39
40 indicated times, samples were removed and mixed with an equal volume of ice-cold stop
41
42 solution (methanol containing internal analytical standards). Stopped reactions were incubated
43
44 at least ten minutes on ice, and an additional volume of water was added. The samples are
45
46 centrifuged to remove precipitated protein, and the supernatants were analyzed by LC-
47
48 MS/MS to monitor the parent depletion using the MRM method and known Phase I and II
49
50 metabolites (oxidation, demethylation, dehydrogenation, glucuronidation, and oxidation +
51
52 glucuronidation) using SRM method. The amount of monitored **25** is converted to %
53
54 remaining by dividing by the time zero concentration value. Data are fit to a first-order decay
55
56
57
58
59
60

1
2
3 model to determine half-life. From a plot of log (ln) peak area against time, the slope of the
4
5 line is determined. Half-life and intrinsic clearance are calculated using the equations below:
6
7 Elimination rate constant (k) = (- slope); Half-life ($T_{1/2}$) min = 0.693/k; Intrinsic Clearance
8
9 (CLint) (mL/min/million cells) = $(V \times 0.693) / T_{1/2}$; where V= incubation volume mL/number of
10
11 cell. Controls: Verapamil - intrinsic clearance 11.9 $\mu\text{L}/\text{min}/10^6$ cells and 29.5 $\mu\text{L}/\text{min}/10^6$
12
13 cells (human (h), mouse (m), respectively). Warfarin - intrinsic clearance $<1.4 \mu\text{L}/\text{min}/10^6$
14
15 cells and $<1.4 \mu\text{L}/\text{min}/10^6$ cells (human (h), mouse (m), respectively).
16
17
18
19
20
21
22

23 **β -Hematin Inhibitory Activity Assay.**

24
25 The inhibition of β -hematin formation is expressed as the molar equivalent of compound,
26
27 relative to hemin, that inhibits β -hematin formation by 50% and determined by slightly
28
29 modified BHIA assay introduced by Parapini et al.⁵⁷ Briefly, 50.0 μL of 16 mM solution of
30
31 hemin in DMSO was distributed in 1.5 mL centrifuge tubes; 80 mM solution of the compound
32
33 in DMSO was added in doses ranging from 0.125 to 5 molar equivalents to hemin (pure
34
35 DMSO was added to control samples). One hundred μL of ultrapure water was added to each
36
37 tube, and β -hematin formation initiated by the addition of 200.0 μL of 8 M acetate buffer (pH
38
39 5.2). The final concentration of DMSO per tube was kept constant at 25%. Tubes were
40
41 incubated at 37.0 ± 0.1 °C for 18 h and then centrifuged. The remaining pellet was resuspended
42
43 in 0.500 mL DMSO to remove unreacted hematin. Tubes were then centrifuged again,
44
45 DMSO-soluble fraction removed and the pellet, consisting of a pure precipitate of β -hematin,
46
47 dissolved in 1.000 mL 0.1 M NaOH. 10.0 μL Aliquots were transferred to 96-well microplate
48
49 and diluted with 190.0 μL 0.1 M NaOH (200.0 μL of 0.1 M NaOH was used as a blank).
50
51 Absorbance was measured at 405 nm, with correction at 670 nm. A calibration curve of hemin
52
53 dissolved in 0.1 M NaOH was made in the range $c_{\text{HE}} = (0.4 - 4.0) \times 10^{-5}$ M.
54
55
56
57
58
59
60

Acidity constant (pK_a) values determination

Acidity constant were potentiometrically determined in MeOH:H₂O (1:1, v:v) at $t=25\pm 1$ °C and at constant ionic strength ($I=0.1$ M (NaCl)). Solutions of NaOH (0.1 M) and HCl (0.1 M) were prepared in MeOH:H₂O (1:1, v:v) and potentiometrically standardized. For working solution, studied compound was dissolved in methanol and diluted with the equivalent volume of aqueous 0.2 M NaCl ($c=(0.5-1.1)\times 10^{-3}$ M). Prior to titration, 100.0 μ L of standard 0.1 M HCl solution was added to 4.00 mL of working compound solution. All probes were titrated with 2.0 μ L increments of standard 0.1 M NaOH solution in 1.8–12.2 pH range. HyperQuad 2008 software⁶⁹ was used to calculate the value of acidity constant values (pK_{a1} and pK_{a2}) from four repeated titrations.

Inhibition potential of antiplasmodials **23** and **25** against the hERG K⁺ channel.

Compounds **23** and **25** were submitted for a hERG K⁺ channel inhibition test to evaluate the proarrhythmic potential that leads to Q–T prolongation cardiac toxicity. Extent of hERG channel inhibitory activity of the test compounds was assessed by Cyprotex US, LLC, Watertown, MA. A detailed protocol is available in the Supporting Information.

AMES mutagenicity screening of antiplasmodial **23**.

Compound **23** was submitted to Cyprotex US, LLC, Watertown, MA for Ames mutagenicity screening and was found to be negative. A detailed protocol is available in the Supporting Information.

ASSOCIATED CONTENT

Supporting information

The additional figures, tables and syntheses of all intermediates is presented. Procedures for the determination of the purity of tested compounds are provided. NMR spectra of all tested

1
2
3 compounds are appended in addition to hERG and AMES details. This material is available
4
5 free of charge via the Internet at <http://pubs.acs.org>.
6
7

8 9 10 **AUTHOR INFORMATION**

11 **Corresponding author**

12
13 *B.Š.: Phone: +381-11-263-86-06, Fax: +381-11-263-60-61, E-mail: bsolaja@chem.bg.ac.rs;

14
15 *M.P.: Phone: +351217999513, Fax: +351217999504, E-mail:

16
17
18 mprudencio@medicina.ulisboa.pt
19
20
21
22

23 24 **NOTES**

25
26
27 The authors declare no competing financial interest.
28
29
30
31

32 33 **ACKNOWLEDGMENTS**

34 This research was supported by: The Ministry of Science and Technological Development of
35 Serbia (grants no. 172008 and 41019), the Serbian Academy of Sciences and Arts, Fundação
36 para a Ciência e Tecnologia, Portugal (grant PTDC/SAU-MIC/117060/2010), the
37 Bill&Melinda Gates Foundation (grant OPP1040394). The contents of this manuscript have
38
39 been reviewed by the Walter Reed Army Institute of Research. There is no objection to its
40
41 presentation or publication.
42
43
44
45
46
47
48

49 50 **ABBREVIATIONS USED**

51 Ad, adamantane; ART, artemisinin; AQ, 4-amino-quinoline; CQ, chloroquine; MFQ,
52 mefloquine; MLM, HLM, mouse and human liver microsomes, respectively; LS, liver stage;
53
54 MOA, method of action; MAD, minimal active dose; MCD, minimal curative dose; MST,
55
56 mean survival time; MTD, maximum tolerated dose; QPlogPow, predicted octanol/water
57
58
59
60

1
2
3 partition coefficient; QPPMDCK, predicted apparent MDCK cell permeability for non-active
4
5 transport; QPlogHERG; predicted IC₅₀ value for blockage of HERG K⁺ channels.
6
7
8
9
10
11
12
13

14 References

-
- 15
16
17 (1) <http://www.who.int/mediacentre/factsheets/fs094/en/>. Accessed 16 October 2014.
18
19 (2) a) WHO, facts on acts (artemisinin-based combination therapies),
20
21 <http://www.malaria.org/ABOUT%20MALARIA/Facts%20on%20ACTs%20WHO.pdf>.
22
23 Accessed 24 September 2014. b) Noedl, H. Combination Therapy in Light of Emerging
24
25 Artemisinin Resistance. In *Treatment and Prevention of Malaria: Antimalarial Drug*
26
27 *Chemistry, Action and Use*; Staines, H. M., Krishna, S., Eds.; Springer:
28
29 Basel, Switzerland, 2012; pp 213–225.
30
31 (3) Wells, T. N. C.; van Huijsduijnen, R. H.; van Voorhis, W. C.. Malaria medicines: a glass
32
33 half full? *Nature Rev. Drug Discov.* **2015**, *14*, 424-442.
34
35 (4) Noedl, H.; Se, Y.; Schaecher, K.; Smith, B. L.; Socheat, D.; Fukuda, M. M. Evidence of
36
37 artemisinin-resistant malaria in western Cambodia. *N. Engl. J. Med.* **2008**, *359*, 2619-2620.
38
39 (5) Dondorp, A.M., Nosten, F., Yi, P., Das, D., Phyto, A.P., Tarning, J., Lwin, K.M., Ariey, F.,
40
41 Hanpithakpong, W., Lee, S.J., Ringwald, P., Silamut, K., Imwong, M., Chotivanich, K., Lim,
42
43 P., Herdman, T., An, S.S., Yeung, S., Singhasivanon, P., Day, N.P., Lindegardh, N., Socheat,
44
45 D., White, N.J. Artemisinin resistance in Plasmodium falciparum malaria. *N. Engl. J. Med.*
46
47 **2009**, *361*, 455-467. [Erratum, *N. Engl. J. Med.* **2009**, *361*, 1714.]
48
49 (6) Ashley, E.A.; Dhorda, M.; Fairhurst, R.M. et al. Spread of artemisinin resistance in
50
51 Plasmodium falciparum malaria. *New Engl. J. Med.* **2014**, *371*, 411–423.
52
53
54
55
56
57
58
59
60

- 1
2
3
4 (7) a) Straimer, J.; Gnadig, N.F.; Witkowski, B.; Amaratunga, C.; Duru, V.; Ramadani, A.P.;
5
6 Dacheux, M.; Khim, N.; Zhang, L.; Lam, S.; Gregory, P.D.; Urnov, F.D.; Mercereau-Puijalon,
7
8 O.; Benoit-Vical, F.; Fairhurst, R.M.; Ménard, D.; Fidock D.A. Drug resistance. K13-
9
10 propeller mutations confer artemisinin resistance in *Plasmodium falciparum* clinical isolates.
11
12 *Science* **2015**; *347*, 428–431. b) Ariey, F.; Witkowski, B.; Amaratunga, C.; Beghain, J.;
13
14 Langlois, A-C.; Khim, N.; Kim, S.; Duru, V.; Bouchier, C.; Ma, L.; Lim, P.; Leang, R.;
15
16 Duong, S.; Sreng, S.; Suon, S.; Chuor, C.M.; Bout, D.M.; Ménard, S.; Rogers, W.O.;
17
18 Genton, B.; Fandeur, T.; Miotto, O.; Ringwald, P.; Le Bras, J.; Berry, A.; Barale, J-C.;
19
20 Fairhurst, R.M.; Benoit-Vical, F.; Mercereau-Puijalon, O.; Ménard, D. A molecular marker
21
22 of artemisinin resistant *Plasmodium falciparum* malaria. *Nature* **2014**, *505*, 50–55.
23
24
25
26 (8) a) Rodrigues, T.; Prudêncio, M.; Moreira, R.; Mota, M.M.; Lopes F. Targeting the liver
27
28 stage of malaria parasites: A yet unmet goal. *J. Med. Chem.* **2012**, *55*, 995-1012. b)
29
30 Rodrigues, C. A. B.; Frade, R. F. M.; Albuquerque, I. S.; Perry, M.J.; Gut, J.; Machado, M.;
31
32 Rosenthal, P.J.; Prudêncio, M; Afonso C.A.M.; Rui Moreira, R. Targeting the Erythrocytic
33
34 and Liver Stages of Malaria Parasites with s-Triazine-Based Hybrids. *ChemMedChem* **2015**,
35
36 *10*, 883 – 890.
37
38
39 (9) Jiménez-Díaz, M.B.; Ebert, D.; Salinas, Y.; Pradhan, A.; Lehane, A.M.; Myrand-Lapierre,
40
41 M-E.; O’Loughlin, K.G.; Shackleford, D.M.; Justino de Almeida, M.; Carrillo, A.K.; Clark,
42
43 J.A.; Dennis, A.S.M.; Diep, J.; Deng, X.; Duffy, J.; Endsley, A.N.; Fedewa, G.; Guiguemde,
44
45 W.A.; Gómez, M.G.; Holbrook, G.; Horst, J.; Kim, C.C.; Liu, J.; Lee, M.C.S.; Matheny, A.;
46
47 Martínez, M.S.; Miller, G.; Rodríguez-Alejandre, A.; Sanz, L.; Sigal, M.; Natalie J. Spillman,
48
49 N.J.; Stein, P.D.; Wang, Z.; Zhu, F.; Waterson, D.; Knapp, S.; Shelat, A.; Avery, V.M.;
50
51 Fidock, D.A.; Gamoa, F.-J.; Charman, S.A.; Mirsalis, J.C.; Ma, H.; Ferrer, S.; Kirk, K.; Iñigo
52
53 Angulo-Barturen, I.; Kyle, D.E.; DeRisi, J.L.; Floyd, D.M.; Guy, R.K. (+)-SJ733, a clinical
54
55
56
57
58
59
60

1
2
3
4 candidate for malaria that acts through ATP4 to induce rapid host-mediated clearance of
5
6 Plasmodium. *Proc Natl Acad Sci USA*, **2014**, *111*, 5455–5462.

7
8 (10) Zou, B.; Nagle, A.; Chatterjee, A. K.; Leong, S. Y.; Tan, L. J.; Sim, W. L. S.; Mishra, P.;
9
10 Guntapalli, P.; Tully, D. C.; Lakshminarayana, S. B.; Lim, C. S.; Tan, Y. C.; Abas, S. N.;
11
12 Bodenreider, C.; Kuhlen, K. L.; Gagaring, K.; Borboa, R.; Chang, J.; Li, C.; Hollenbeck, T.;
13
14 Tuntland, T.; Zeeman, A.-M.; Kocken, C. H. M.; McNamara, C.; Kato, N.; Winzeler, E. A.;
15
16 Yeung, B. K. S.; Diagona, T. T.; Smith, P. W.; Roland, J. Lead Optimization of
17
18 Imidazopyrazines: A New Class of Antimalarial with Activity on Plasmodium Liver Stages.
19
20 *ACS Med. Chem. Lett.* **2014**, *5*, 947-950.
21
22

23
24 (11) Nilsen, A.; LaCrue, A.; White, K.; Forquer, I.; Cross, R.; Marfurt, J.; Mather, M.;
25
26 Delves, M.; Shackelford, D.; Saenz, F.; Morrisey, J.; Steuten, J.; Mutka, T.; Li, Y.; Wirjanata,
27
28 G.; Ryan, E.; Duffy, S.; Kelly, J.; Sebayang, B.; Zeeman, A.-M.; Noviyanti, R.; Sinden, R.;
29
30 Kocken, C.; Price, R.; Avery, V.; Angulo-Barturen, I.; Jiménez-Díaz, M.; Ferrer, S.; Herreros,
31
32 E.; Sanz, L.; Gamo, F.-J.; Bathurst, I.; Burrows, J.; Siegl, P.; Guy, R.; Winter, R.; Vaidya, A.;
33
34 Charman, S.; Kyle, D.; Manetsch, R.; Riscoe, M. Quinolone-3-diarylethers: A new class of
35
36 antimalarial drug. *Science Tran. Med.* **2013**, *5*, 177ra37.
37
38

39
40 (12) Rodrigues, T.; Ressurreição, A.; Cruz, F.; Albuquerque, I.; Gut, J.; Carrasco, M.;
41
42 Gonçalves, D.; Guedes, R.; Santos, D.; Mota, M.; Rosenthal, P.; Moreira, R.; Prudêncio, M.;
43
44 Lopes, F. Flavones as isosteres of 4(1H)-quinolones: Discovery of ligand efficient and dual
45
46 stage antimalarial lead compounds. *Eur. J. Med. Chem.* **2013**, *69*, 872-880.
47

48
49 (13) Pérez, B. C.; Teixeira, C.; Albuquerque, I. S.; Gut, J.; Rosenthal, P. J.; Gomes, J. R. B.;
50
51 Prudêncio, M.; Gomes, P. N-cinnamoylated chloroquine analogues as dual-stage antimalarial
52
53 leads. *J. Med. Chem.* **2013**, *56*, 556–567.
54

55
56 (14) Rodrigues, T.; da Cruz, F. P.; Lafuente-Monasterio, M. J.; Gonçalves, D.; Ressurreição,
57
58 A. S.; Siteo, A.; Bronze, M. R.; Gut, J.; Schneider, G.; Mota, M. M.; Rosenthal, P. J.;
59
60

1
2
3
4 Prudêncio, M.; Gamo, F.-J.; Lopes, F.; Moreira, R. Quinolin-4(1H)-imines are potent
5 antiplasmodial drugs targeting the liver stage of malaria. *J. Med. Chem.* **2013**, *56*, 4811–
6 4815.
7
8

9
10 (15) Das, U.; Singh, R. S.; Alcorn, J.; Hickman, M. R.; Sciotti, R. J.; Leed, S. E.; Lee, P. J.;
11 Roncal, N.; Dimmock, J. R. 3,5-Bis(benzylidene)-4-piperidones and related N-acyl analogs:
12 A novel cluster of antimalarials targeting the liver stage of *Plasmodium falciparum*. *Bioorg.*
13 *Med. Chem.* **2013**, *21*, 7250-7256.
14
15

16
17 (16) McNamara, C. W.; Lee, M. C. S.; Lim, C. S.; Lim, S. H.; Roland, J.; Nagle, A.; Simon,
18 O.; Yeung, B. K. S.; Chatterjee, A. K.; McCormack, S. L.; Manary, M. J.; Zeeman, A.-M.;
19 Dechering, K. J.; Kumar, T. R. S.; Henrich, P. P.; Gagaring, K.; Ibanez, M.; Kato, N.; Kuhlen,
20 K. L.; Fischli, C.; Rottmann, M.; Plouffe, D. M.; Bursulaya, B.; Meister, S.; Rameh, L.;
21 Trappe, J.; Haasen, D.; Timmerman, M.; Sauerwein, R. W.; Suwanarusk, R.; Russell, B.;
22 Renia, L.; Nosten, F.; Tully, D. C.; Kocken, C. H. M.; Glynne, R. J.; Bodenreider, C.; Fidock,
23 D. A.; Diagana, T. T.; Winzeler, E. A. Targeting *Plasmodium* PI(4)K to eliminate malaria.
24 *Nature* **2013**, *504*, 248–253.
25
26

27
28 (17) Lödige, M.; Lewis, M. D.; Paulsen, E. S.; Esch, H. L.; Pradel, G.; Lehmann, L.; Brun, R.;
29 Bringmann, G.; Mueller, A.-K. A primaquine–chloroquine hybrid with dual activity against
30 *Plasmodium* liver and blood stages. *Int. J. Med. Microbiol.* **2013**, *303*, 539-547.
31
32

33
34 (18) Milner, E.; Sousa, J.; Pybus, B.; Auschwitz, J.; Caridha, D.; Gardner, S.; Grauer, K.;
35 Harris, E.; Hickman, M.; Kozar, M. P.; Lee, P.; Leed, S.; Li, Q.; Melendez, V.; Moon, J.;
36 Ngundam, F.; O'Neil, M.; Parriott, S.; Potter, B.; Sciotti, R.; Tangteung, A.; Dow, G. S.
37 Ketotifen is an antimalarial prodrug of norketotifen with blood schizonticidal and liver-stage
38 efficacy. *Eur. J. Drug Metab. Ph.* **2012**, *37*, 17-22.
39
40

41
42 (19) da Cruz, F. P.; Martin, C.; Buchholz, K.; Lafuente-Monasterio, M. J.; Rodrigues, T.;
43 Sönnichsen, B.; Moreira, R.; Gamo, F.-J.; Marti, M.; Mota, M. M.; Hannus, M.; Prudêncio,
44
45
46
47
48
49
50
51
52
53
54

1
2
3
4 M. Drug screen targeted at Plasmodium liver stages identifies a potent multistage antimalarial
5 drug. *J. Infect. Dis.* **2012**, *205*, 1278–1286.

6
7
8 (20) Cohen, A.; Suzanne, P.; Lancelot, J.-C.; Verhaeghe, P.; Lesnard, A.; Basmaciyan, L.;
9
10 Hutter, S.; Laget, M.; Dumètre, A.; Paloque, L.; Deharo, E.; Crozet, M.D.; Rathelot, P.;
11
12 Dallemagne, P.; Lorthiois, A.; Sibley, C.H.; Vanelle, P.; Valentin, A.; Mazier, D.; Rault, S.;
13
14 Azas, N. Discovery of new thienopyrimidinone derivatives displaying antimalarial properties
15
16 toward both erythrocytic and hepatic stages of Plasmodium. *Eur. J. Med. Chem.* **2015**, *95*, 16-
17
18 28.

19
20
21 (21) Baragaña, B.; Hallyburton, I.; Lee, M. C. S.; Norcross, N. R.; Grimaldi, R.; Otto, T. D.;
22
23 Proto, W. R.; Blagborough, A. M.; Meister, S.; Wirjanata, G.; Ruecker, A.; Upton, L.M.;
24
25 Abraham, T. S.; Almeida, M. J.; Pradhan, A.; Porzelle, A.; Martínez, M. S.; Bolscher, J. M.;
26
27 Woodland, A.; Norval, S.; Zuccotto, F.; Thomas, J.; Simeons, F.; Stojanovski, L.; Osuna-
28
29 Cabello, M.; Brock, P. M.; Churcher, T. S.; Sala, K. A.; Zakutansky, S. E.; Jiménez-Díaz, M.
30
31 B.; Sanz, L. M.; Riley, J.; Basak, R.; Campbell, M.; Avery, V. M.; Sauerwein, R. W.;
32
33 Dechering, K. J.; Noviyanti, R.; Campo, B.; Frearson, J. A.; Angulo-Barturen, I.; Ferrer-
34
35 Bazaga, S. F.; Gamo, F. J.; Wyatt, P. G.; Leroy, D.; Siegl, P.; Delves, M. J.; Kyle, D. E.;
36
37 Wittlin, S.; Marfurt, J.; Price, R. N.; Sinden, R. E.; Winzeler, E. A.; Charman, S. A.;
38
39 Bebrevska, L.; Gray, D. W.; Campbell, S.; Fairlamb, A. H.; Willis, P. A.; Rayner, J. C.;
40
41 Fidock, D. A.; Read, K. D.; Ian H. Gilbert, I. H. A novel multiple-stage antimalarial agent
42
43 that inhibits protein synthesis. *Nature*, **2015**, *522*, 315-320.

44
45
46 (22) Capela, R.; Cabal, G. G.; Rosenthal, P. J.; Gut, J.; Mota, M. M.; Moreira, R.; Lopes, F.;
47
48 Prudêncio, M. Design and evaluation of primaquine-artemisinin hybrids as a multistage
49
50 antimalarial strategy. *Antimicrob. Agents Chemother.* **2011**, *55*, 4698–4706.

51
52
53 (23) Oliveira, R.; Guedes, R. C.; Meireles, P.; Albuquerque, I. S.; Gonçalves, L. M.; Pires, E.;
54
55 Bronze, M. R.; Gut, J.; Rosenthal, P. J.; Prudêncio, M.; Moreira, R.; O'Neill, P. M.; Lopes, F.
56
57
58
59
60

1
2
3
4 Tetraoxane–Pyrimidine Nitrile Hybrids as Dual Stage Antimalarials. *J. Med. Chem.* **2014**,
5
6 57, 4916-4923.

7
8 (24) Miranda, D.; Capela, R.; Albuquerque, I. S.; Meireles, P.; Paiva, I.; Nogueira, F.;
9
10 Amewu, R.; Gut, J.; Rosenthal, P. J.; Oliveira, R.; Mota, M. M.; Moreira, R.; Marti, F.;
11
12 Prudêncio, M.; O'Neill, P. M.; Lopes, F. Novel Endoperoxide-Based Transmission-Blocking
13
14 Antimalarials with Liver- and Blood-Schizontocidal Activities. *ACS Med. Chem. Lett.* **2014**,
15
16 5, 108-112.

17
18 (25) Delves, M.; Plouffe, D.; Scheurer, C.; Meister, S.; Wittlin, S.; Winzeler, E. A.; Sinden,
19
20 R. E.; Leroy, D. The activities of current antimalarial drugs on the life cycle stages of
21
22 plasmodium: A comparative study with human and rodent parasites. *PLoS Med.* **2012**, 9,
23
24 e1001169.

25
26 (26) Charman, S. A.; Arbe-Barnes, S.; Bathurst, I. C.; Brun, R.; Campbell, M.; Charman, W.
27
28 N.; Chiu, F. C.; Chollet, J.; Craft, J. C.; Creek, D. J.; Dong, Y.; Matile, H.; Maurer, M.;
29
30 Morizzi, J.; Nguyen, T.; Papastogiannidis, P.; Scheurer, C.; Shackelford, D. M.; Sriraghavan,
31
32 K.; Stingelin, L.; Tang, Y.; Urwyler, H.; Wang, X.; White, K. L.; Wittlin, S.; Zhou, L.;
33
34 Vennerstrom, J. L. Synthetic ozonide drug candidate OZ439 offers new hope for a single-dose
35
36 cure of uncomplicated malaria. *Proc. Natl. Acad. Sci. U.S.A.* **2011**, 108, 4400–4405.

37
38 (27) a) Kublin, J.G.; Cortese, J.F.; Njunju, E.M.; Mukadam, R.A.G.; Wirima, J.J.; Kazembe,
39
40 P.N.; Djimdé, A.A.; Kouriba, B.; Taylor, T.E.; Plowe, C.V. Reemergence of Chloroquine-
41
42 Sensitive Plasmodium falciparum Malaria after Cessation of Chloroquine Use in Malawi. *J.*
43
44 *Infect. Dis.* **2003**, 187, 1870–1875. b) Wang, X.; Mu, J.; Li, G.; Chen, P.; Guo, X.; Fu, L.;
45
46 Chen, L.; Su, X.; Wellems T.E. Decreased prevalence of the Plasmodium falciparum
47
48 chloroquine resistance transporter 76t marker associated with cessation of chloroquine use
49
50 against P. falciparum malaria in Hainan, People's Republic of China. *Am. J. Trop. Med. Hyg.*,
51
52 **2005**, 72, 410–414.
53
54
55
56
57
58
59
60

- 1
2
3
4 (28) a) Sáenz, F. E.; Mutka, T.; Udenze, K.; Oduola, A. M. J.; Kyle, D. E. Novel 4-
5 aminoquinoline analogs highly active against the blood and sexual stages of Plasmodium in
6 vivo and in vitro. *Antimicrob. Agents Chemother.* **2012**, *56*, 4685-4692. b) Egan, T.J.; Kuter,
7 D. Dual-functioning antimalarials that inhibit the chloroquine-resistance transporter. *Future*
8 *Microbiol.* **2013** *8*, 475-489. c) Next-Generation Antimalarial Drugs: Hybrid Molecules
9 as a New Strategy in Drug Design. Muregi, F.W.; Ishih, A. *Drug Devel. Res.* **2010**, *71*, 20–32.
10
11 (29) Casagrande, M.; Barteselli, A.; Basilico, N.; Parapini, S.; Taramelli, D.; Sparatore, A.
12 Synthesis and antiplasmodial activity of new heteroaryl derivatives of 7-chloro-4-
13 aminoquinoline. *Bioorg. Med. Chem.* **2012**, *20*, 5965-5979.
14
15 (30) Gemma, S.; Camodeca, C.; Coccone, S. S.; Joshi, B. P.; Bernetti, M.; Moretti, V.; Brogi,
16 S.; de Marcos, M. C. B.; Savini, L.; Taramelli, D.; Basilico, N.; Parapini, S.; Rottmann, M.;
17 Brun, R.; Lamponi, S.; Caccia, S.; Guiso, G.; Summers, R. L.; Martin, R. E.; Saponara, S.;
18 Gorelli, B.; Novellino, E.; Campiani, G.; Butini, S. Optimization of 4-aminoquinoline/
19 clotrimazole-based hybrid antimalarials: further structure–activity relationships, in vivo
20 studies, and preliminary toxicity profiling. *J. Med. Chem.* **2012**, *55*, 6948-6967.
21
22 (31) Manohar, S.; Rajesh, U. C.; Khan, S. I.; Tekwani, B. L.; Rawat, D. S. Novel 4-
23 aminoquinoline-pyrimidine based hybrids with improved in vitro and in vivo antimalarial
24 activity. *ACS Med. Chem. Lett.* **2012**, *3*, 555-559.
25
26 (32) Tukulula, M.; Njoroge, M.; Abay, E. T.; Mugumbate, G. C.; Wiesner, L.; Taylor, D.;
27 Gibhard, L.; Norman, J.; Swart, K. J.; Gut, J.; Rosenthal, P. J.; Barteau, S.; Streckfuss, J.;
28 Kameni-Tcheudji, J.; Chibale, K. Synthesis and in vitro and in vivo pharmacological
29 evaluation of new 4-aminoquinoline-based compounds. *ACS Med. Chem. Lett.* **2013**, *4*, 1198-
30 1202.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4 (33) Singh, K.; Kaur, H.; Smith, P.; de Kock, C.; Chibale, K.; Balzarini, J. Quinoline–
5
6 pyrimidine hybrids: synthesis, antiplasmodial activity, SAR, and mode of action studies. *J.*
7
8 *Med. Chem.* **2014**, *57*, 435-448.
- 9
10 (34) a) Madrid, P. B.; Liou, A. P.; DeRisi, J. L.; Guy, R. K. Incorporation of an
11
12 Intramolecular Hydrogen-Bonding Motif in the Side Chain of 4-Aminoquinolines Enhances
13
14 Activity against Drug-Resistant *P. falciparum*. *J. Med. Chem.* **2006**, *49*, 4535-4543. b) Ray,
15
16 S.; Madrid, P. B.; Catz, P.; LeValley, S. E.; Furniss, M. J.; Rausch, L. L.; Guy, R. K.; DeRisi,
17
18 J. L.; Iyer, L. V.; Green, C. E.; Mirsalis, J. C. Development of a New Generation of 4-
19
20 Aminoquinoline Antimalarial Compounds Using Predictive Pharmacokinetic and Toxicology
21
22 Models. *J. Med. Chem.* **2010**, *53*, 3685-3695.
- 23
24 (35) Kumar, D.; Khan, S.I.; Tekwani, B.L.; Ponnann, P.; Rawat, D.S. 4-Aminoquinoline-
25
26 Pyrimidine hybrids: Synthesis, antimalarial activity, heme binding and docking studies. *Eur.*
27
28 *J. Med. Chem.* **2015**, *89*, 490-502.
- 29
30 (36) Figueiras, M.; Coelho, L.; Wicht, K.J.; Santos, S.A.; Lavrado, J.; Gut, J.; Rosenthal, P.J.;
31
32 Nogueira, F.; Egan, T.J.; Moreira, R.; Paulo, A. N10,N11-di-alkylamine indolo[3,2-
33
34 b]quinolines as hemozoin inhibitors: Design, synthesis and antiplasmodial activity. *Bioorg.*
35
36 *Med. Chem.* **2015**, *23*, 1530–1539.
- 37
38 (37) a) Šolaja, B. A.; Opsenica, D.; Smith, K. S.; Milhous, W. K.; Terzić, N.; Opsenica, I.;
39
40 Burnett, J. C.; Nuss, J.; Gussio, R.; Bavari, S. Novel 4-Aminoquinolines Active against
41
42 Chloroquine-Resistant and Sensitive *P. falciparum* Strains that also Inhibit Botulinum
43
44 Serotype A. *J. Med. Chem.* **2008**, *51*, 4388-4391. b) For short-chain in vitro 4-aminoquinoline
45
46 data see also: Stocks, P.A.; Raynes, K.J.; Bray, P.G.; B. Park, K.; O'Neill, P.M.; Ward, S.A.
47
48 Novel Short Chain Chloroquine Analogues Retain Activity Against Chloroquine Resistant K1
49
50 *Plasmodium falciparum*. *J. Med. Chem.* **2002**, *45*, 4975-4983.
- 51
52
53
54
55
56
57
58
59
60

- 1
2
3
4 (38) Opsenica, I.M.; Verbić, T.Ž.; Tot, M.; Sciotti, R.J.; Pybus, B.S.; Djurković-Djaković,
5
6 O.; Ksenija Slavić, K.; Šolaja, B.A. Investigation into novel thiophene- and furan-based 4-
7
8 amino-7-chloroquinolines afforded antimalarials that cure mice. *Bioorg. Med. Chem.* **2015**,
9
10 23, 2176–2186.
- 11
12 (39) Roe, A.; Hawkins, G.F. The preparation of Heterocyclic Fluorine Compounds by the
13
14 Schiemann Reaction. II. The Monofluoroquinolines. *J. Am. Chem. Soc.* **1948**, 71, 1785-1786.
- 15
16 (40) Arzel, E.; Rocca, P.; Marsais, F.; Godard, A.; Quéguiner. First Total Synthesis of
17
18 Cryptomisine. *Tetrahedron*, **1999**, 55, 12149-12156.
- 19
20 (41) Milhous, W. K.; Weatherly, N. F.; Bowdre, J. H.; Desjardins, R. E. In Vitro Activities of
21
22 and Mechanisms of Resistance to Antifol Antimalarial Drugs. *Antimicrob. Agents Chemother.*
23
24 **1985**, 27, 525–530.
- 25
26 (42) Opsenica, I.; Burnett, J. C.; Gussio, R.; Opsenica, D.; Todorović, N.; Lanteri, C. A.;
27
28 Sciotti, R. J.; Gettayacamin, M.; Basilico, N.; Taramelli, D.; Nuss, J. E.; Wannner, L.; Panchal,
29
30 R. G.; Šolaja, B. A.; Bavari, S. A Chemotype That Inhibits Three Unrelated Pathogenic
31
32 Targets: The Botulinum Neurotoxin Serotype A Light Chain, *P. falciparum* Malaria, and the
33
34 Ebola Filovirus. *J. Med. Chem.* **2011**, 54, 1157–1169 and references cited therein.
- 35
36 (43) a) Ploemen, I.H.; Prudêncio, M.; Douradinha, B.G.; Ramesar, J.; Fonager, J.; van
37
38 Gemert, G.J.; Luty, A.J.; Hermsen, C.C.; Sauerwein, R.W.; Baptista, F.G.; Mota, M.M.;
39
40 Waters, A.P.; Que, I.; Lowik, C.W.; Khan, S.M.; Janse, C.J.; X Franke-Fayard, C.J.
41
42 Visualisation and quantitative analysis of the rodent malaria liver stage by real time imaging.
43
44 *PLoS ONE*, **2009**, 4, e7881. b) Prudêncio, M.; Mota, M. M.; Mendes, A. M. A toolbox to
45
46 study liver stage malaria. *Trends Parasitol.* **2011**, 27, 565–574.
- 47
48 (44) Opsenica, I.; Terzić, N.; Opsenica, D.; Angelovski, D.; Lehnig, M.; Eilbracht, P.; Tinant,
49
50 B.; Juranić, Z.; Smith, K. S.; Yang, Y. S.; Diaz, D. S.; Smith, P. L.; Milhous, W. K.; Doković,
51
52
53
54
55
56
57
58
59
60

D.; Šolaja, B. A. Tetraoxane Antimalarials and Their Reaction with Fe(II). *J. Med. Chem.*

2006, *49*, 3790–3799.

(45) Opsenica, I.; Opsenica, D.; Smith, K. S.; Milhous, W. K.; Šolaja, B. A. Chemical

Stability of the Peroxide Bond Enables Diversified Synthesis of Potent Tetraoxane

Antimalarials. *J. Med. Chem.* **2008**, *51*, 2261–2266.

(46) Opsenica, I.; Opsenica, D.; Lanteri, C.A.; Anova, L.; Milhous, W.K.; Smith, K.S.;

Šolaja, B.A. New Chimeric Antimalarials with 4-Aminoquinoline Moiety Linked to a

Tetraoxane Skeleton. *J. Med. Chem.* **2008**, *51*, 6216–6219.

(47) Šolaja, B. A.; Terzić, N.; Pocsfalvi, G.; Gerena, L.; Tinant, B.; Opsenica, D.; Milhous,

W. K. Mixed Steroidal 1,2,4,5-Tetraoxanes: Antimalarial and Antimycobacterial Activity. *J.*

Med. Chem. **2002**, *45*, 3331–3336.

(48) Opsenica, I.; Terzić, N.; Opsenica, D.; Milhous, W. K.; Šolaja, B. A. 7,8,15,16-

Tetraoxa-dispiro[5.2.5.2]hexadecane-3-carboxylic acid derivatives and their antimalarial

activity. *J. Serb. Chem. Soc.* **2004**, *69*, 919–922.

(49) Terzić, N.; Opsenica, D.; Milić, D.; Tinant, B.; Smith, K. S.; Milhous, W. K.; Šolaja, A.

B. Deoxycholic Acid-Derived Tetraoxane Antimalarials and Antiproliferatives. *J. Med.*

Chem. **2007**, *50*, 5118–5127.

(50) D'Alessandro, S.; Silvestrini, F.; Dechering, K.; Corbett, Y.; Parapini, S.; Timmerman,

M.; Galastri, L.; Basilico, N.; Sauerwein, R.; Alano, P.; Taramelli, D. A *Plasmodium*

falciparum screening assay for anti-gametocyte drugs based on parasite lactate dehydrogenase

detection. *J. Antimicrob. Chemother.* **2013**, *68*, 2048–2058.

(51) a) Madrid, P.B.; Sherrill, J.; Liou, A.P.; Weisman, J.L.; DeRisi, J.L.; Guy, R.K.

Synthesis of ring-substituted 4-aminoquinolines and evaluation of their antimalarial activities.

Bioorg. Med. Chem. Lett. **2005**, *15*, 1015–1018. b) Kaschula, C.H.; Egan, T.J.; Hunter, R.;

Basilico, N.; Parapini, S.; Taramelli, D.; Pasini, E.; Monti, D. Structure-Activity

1
2
3
4 Relationships in 4-Aminoquinoline Antiplasmodials. The Role of the Group at the 7-Position.

5
6 *J. Med. Chem.* **2002**, *45*, 3531-3539. c) De, D.; Krogstad, F.M.; Byers, L.D.; Krogstad, D.J.

7
8 Structure-Activity Relationships for Antiplasmodial Activity among 7-Substituted 4-

9
10 Aminoquinolines. *J. Med. Chem.* **1998**, *41*, 4918-4926.

11
12 (52) a) Surrey, A.R.; Cutler, R.A. The Synthesis of Some 3-Nitro- and 3-Amino-4-

13
14 dialkylaminoquinoline Derivatives. *J. Am. Chem. Soc.* **1951**, *73*, 2413-2416. b) Surrey, A.R.;

15
16 Cutler, R.A. The Preparation of 3-Halo-4-dialkylaminoquinoline derivatives. *J. Am. Chem.*

17
18 *Soc.* **1946**, *68*, 2570-2571.

19
20 (53) Geary, T. G.; Divo, A. A.; Jensen, J. B. Activity of quinoline-containing antimalarials

21
22 against chloroquine-sensitive and -resistant strains of *Plasmodium falciparum* in vitro. *Trans.*

23
24 *R. Soc. Trop. Med. Hyg.* **1987**, *81*, 499-503.

25
26 (54) Kumar, Y. K.; Kumar, G. R.; Reddy, T. J.; Sridhar, B.; Reddy, M. S. Synthesis of 3-

27
28 Sulfonylamino Quinolines from 1-(2-Aminophenyl) Propargyl Alcohols through a Ag(I)-

29
30 Catalyzed Hydroamination, (2 + 3) Cycloaddition, and an Unusual Strain-Driven Ring

31
32 Expansion. *Org. Lett.* **2015**, *17* (9), 2226-2229.

33
34 (55) Pou, S.; Winter, R.W.; Nilsen, A.; Kelly, J.X.; Li, Y.; J. Doggett, S.; Riscoe, E.W.;

35
36 Wegmann, K.W.; Hinrichs, D.J.; Riscoe, M.K. Sontochin as a Guide to the Development of

37
38 Drugs against Chloroquine-Resistant Malaria. *Antimicrob. Agents Chemother.* **2012**, *56*,

39
40 3475-3480, and references cited therein.

41
42 (56) QikProp, version 4.4, Schrödinger, Inc., New York, NY, 2015. All compound structures

43
44 were previously prepared by LigPrep (LigPrep version 3.4, Schrödinger, LLC, New York,

45
46 NY, 2015) and were used for the calculation of pharmacokinetic properties. The

47
48 pharmacokinetic profile of the test compounds showing good antimalarial activity were

49
50 predicted by using program QikProp v.4.4 (Schrödinger Suite 2015-2).

- 1
2
3
4 (57) Parapini, S.; Basilico, N.; Pasini, E.; Egan, T. J.; Olliaro, P.; Taramelli, D.; Monti, D.
5
6 Standardization of the physicochemical parameters to assess in vitro the beta-hematin
7
8 inhibitory activity of antimalarial drugs. *Exp. Parasitol.* **2000**, *96*, 249-256.
9
10 (58) Traebert, M.; Dumotier, B.; Meister, L.; Hoffmann, P.; Dominguez-Estevez, M.; Suter,
11
12 W. Inhibition of hERG K⁺ currents by antimalarial drugs in stably transfected HEK293 cells.
13
14 *Eur. J. Pharmacol.* **2004**, *484* 41–48.
15
16 (59) Sanders, N. G.; Meyers, D.J.; Sullivan, D.J. Antimalarial Efficacy of
17
18 hydroxyethylapoquinine (SN-119) and Its Derivatives. *Antimicrob. Agents Chemother.* **2014**,
19
20 *58*, 820–827.
21
22 (60) Gans, P.; O'Sullivan B. GLEE, a new computer program for glass electrode calibration,
23
24 *Talanta* **2000**, *51*, 33-37.
25
26 (61) a) Musonda, C. C.; Gut, J.; Rosenthal, P. J.; Yardley, V.; Carvalho de Souza, R. C.;
27
28 Chibale, K. Application of Multicomponent Reactions to Antimalarial Drug Discovery. Part
29
30 2: New Antiplasmodial and Antitrypanosomal 4 Aminoquinoline Gamma- and Delta-Lactams
31
32 via a “Catch and Release” Protocol. *Bioorg. Med. Chem.* **2006**, *14*, 5605–5615. b) Peck, R.
33
34 M.; Preston, R. K.; Creech, H. J. Nitrogen Mustard Analogs of Antimalarial Drugs. *J. Am.*
35
36 *Chem. Soc.* **1958**, *81*, 3984–3989. c) Price, C. C.; Leonard, N. J.; Peel, E. W.; Reitsema, R. H.
37
38 Some 4-Amino-7-chloroquinoline Derivatives. *J. Am. Chem. Soc.* **1946**, *68*, 1807–1808. d)
39
40 Singh, C.; Malik, H.; Puri, S. K. Synthesis and Antimalarial Activity of a New Series of
41
42 Trioxaquinines. *Bioorg. Med. Chem.* **2004**, *12*, 1177–1182.
43
44 (62) Margolis, B. J.; Long, K. A.; Laird, D. L. T.; Ruble, J. C.; Pulley, S. R. Assembly of 4-
45
46 Aminoquinolines via Palladium Catalysis: A Mild and Convenient Alternative to S_NAr
47
48 Methodology. *J. Org. Chem.*, **2007**, *72*, 2232–2235.
49
50 (63) Kim, S.; Oh, C. H.; Oh, Ko, J. S.; Ahn, K. H.; Kim, Y. J. Zinc-modified
51
52 cyanoborohydride as a selective reducing agent. *J. Org. Chem.*, 1985, *50* (11), 1927–1932.
53
54
55
56
57
58
59
60

1
2
3
4 (64) Arzel, E.; Rocca, P.; Marsais, F.; Godard, A.; Queguiner, G. First Total Synthesis of
5
6 Cryptomisine. *Tetrahedron*, **1999**, *55*, 12149-12156.

7
8 (65) Desjardins, R.E.; Canfield, C.J.; Haynes, D.E.; Chulay, J.D. Quantitative assessment of
9
10 antimalarial activity in vitro by a semiautomated microdilution technique. *Antimicrob. Agents*
11
12 *Chemother.* **1979**, *16* (6), 710-718.


13 (66) Opsenica, I. M.; Tot, M.; Gomba, L.; Nuss, J. E.; Sciotti, R. J.; Bavari, S.; Burnett, J. C.;
14
15 Šolaja, B. A. 4-Amino-7-chloroquinolines: Probing Ligand Efficiency Provides Botulinum
16
17 Neurotoxin Serotype A Light Chain Inhibitors with Significant Antiprotozoal Activity. *J.*
18
19 *Med. Chem.* **2013**, *56*, 5860-5871.

20
21 (67) Cevenini, L.; Camarda, G.; Michelini, E.; Siciliano, G.; Calabretta, M. M.; Bona, R.;
22
23 Kumar, T. R. S.; Cara, A.; Branchini, B. R.; Fidock, D. A.; Roda, A.; Alano, P. Multicolor
24
25 Bioluminescence Boosts Malaria Research: Quantitative Dual-Color Assay and Single-Cell
26
27 Imaging in Plasmodium falciparum Parasites. *Anal Chem.* **2014**, *86*, 8814-8821.

28
29 (68) Prudêncio, M.; Rodrigues, C.D.; Hannus, M.; Martin, C.; Real, E.; Gonçalves, L.A.;
30
31 Carret, C.; Dorkin, R.; Röhl, I.; Jahn-Hoffmann, K.; Luty, A.J.; Sauerwein, R.; Echeverri,
32
33 C.J.; Mota, M.M.; Kinome-wide RNAi screen implicates at least 5 host hepatocyte kinases in
34
35 Plasmodium sporozoite infection. *PLoS Pathog.* **2008**, *4*:e1000201.

36
37 (69) Gans, P.; Sabatini, A.; Vacca, A. Investigation of equilibria in solution. Determination of
38
39 equilibrium constants with the HYPERQUAD suite of programs, *Talanta* **1996**; *43*, 1739-
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
1753.

TABLE OF CONTENTS GRAPHIC




P. berghei

RBC (Thompson) 6/6 cure @ 80 mg/kg/day

Liver stage IC_{50} = 310 nM

Lliver load reduction 92%@80 mg/kg



Liver stage IC_{50} = 330 nM

P.b. liver load reduction 91%@100 mg/kg