


Reaction of $[\text{Os}_4(\mu\text{-Cl})(\text{CO})_{13}]^-$ with the $[\text{Au}(\text{PPh}_3)]^+$ Cation involving Insertion of an $\text{Os}(\text{CO})_4$ Fragment into the Gold–Phosphorus Bond

Christopher Cathey, Jack Lewis,* Paul R. Raithby and M. Carmen Ramírez de Arellano
University Chemical Laboratory, Lensfield Road, Cambridge CB2 1EW, UK

The reaction of the cluster anion $[\text{Os}_4(\mu\text{-Cl})(\text{CO})_{13}]^-$ **1** with $[\text{Au}(\text{PPh}_3)]^+$ affords the cluster $[\text{Os}_3\{\mu\text{-AuOs}(\text{CO})_4(\text{PPh}_3)\}(\mu\text{-Cl})(\text{CO})_{10}]^-$ **2** which represents the first example in which an $\text{Os}(\text{CO})_4$ fragment has been inserted into a gold–phosphine bond; the structure of **2** has been established by a single-crystal X-ray analysis.

It has been found that the reaction between $[\text{Ru}_3(\text{CO})_{12}]$ and $[\text{N}(\text{PPh}_3)_2]\text{Cl}$ is the best preparative route to the salt $[\text{N}(\text{PPh}_3)_2][\text{Ru}_4(\mu\text{-Cl})(\text{CO})_{13}]^-$.¹ We now report an extension of this mode of synthesis to give the corresponding osmium anion $[\text{Os}_4(\mu\text{-Cl})(\text{CO})_{13}]^-$ **1** as its $[\text{N}(\text{PPh}_3)_2]^+$ salt in a yield of 50%. The reaction of the cluster **1** (50 mg, 0.029 mol) with $[\text{Au}(\text{PPh}_3)][\text{NO}_3]$ (15.3 mg, 0.029 mmol) yields the new cluster $[\text{Os}_3\{\mu\text{-AuOs}(\text{CO})_4(\text{PPh}_3)\}(\mu\text{-Cl})(\text{CO})_{10}]^-$ **2** (20%) together with the known complexes $[\text{Os}_3\{\mu\text{-Au}(\text{PPh}_3)\}(\mu\text{-Cl})(\text{CO})_{10}]^-$ **3** (40%) and $[\text{Os}_4\text{H}_3\{\mu\text{-Au}(\text{PPh}_3)\}(\text{CO})_{12}]^3$ **4** (30%) (Scheme 1). The formation of **2** is thought to involve the cleavage of a gold–phosphorus bond of the $\text{Au}(\text{PPh}_3)$ unit and the novel insertion of an $\text{Os}(\text{CO})_4$ fragment. Alternatively **2** may be viewed as arising from the oxidative-addition of the $[\text{Au}(\text{PPh}_3)]^+$ group to the $\text{Os}(\text{CO})_4$ fragment.

The compounds **1–4** have been isolated as air stable solids. The mass and IR spectra are consistent with the proposed molecular formulae,[†] and in the cases of clusters **3** and **4** with the previously reported^{2,3} spectroscopic values. The ³¹P NMR spectra exhibit resonances in the range δ –63.6 to –62.0 which may be assigned to the co-ordinated phosphines. In order to establish the geometry of **2** its crystal structure has been determined.[‡]



Scheme 1 (i) $[\text{N}(\text{PPh}_3)_2]\text{Cl}$, tetrahydrofuran (thf), heat; (ii) $[\text{Au}(\text{PPh}_3)][\text{NO}_3]$, thf, heat

[†] Spectroscopic data. The $[\text{N}(\text{PPh}_3)_2]^+$ salt of **1**: IR $\nu(\text{CO})$ (*n*-hexane) 2069w, 2034s, 2012s, 1971m, 1805w cm^{-1} ; FAB mass spectrum M^+ (obs.) m/z 1160. (calc.) 1160; IR $\nu(\text{CO})$ (*n*-hexane) 2088w, 2042vs, 2019(sh), 2006s, 1976w cm^{-1} ; FAB mass spectrum M^+ (obs.), m/z 1647 (calc.) 1647; ¹H NMR (CDCl_3) δ 6.7 (m, 18 H, PPh₃). **4**: IR $\nu(\text{CO})$ (*n*-hexane) 2095m, 2071s, 2034vs, 2008s, 1956m cm^{-1} ; FAB mass spectrum M^+ (obs.) m/z 1558, (calc.) 1559; ¹H NMR (CDCl_3) δ 6.4 (m, 18 H, PPh₃), –21.9 (d, 0.8 Hz, 3 H, MH).

[‡] Crystal data for **2**. $\text{C}_{32}\text{H}_{15}\text{AuClO}_{14}\text{Os}_4\text{P}$, $M = 1647.6$, triclinic, space group $P\bar{1}$ (no. 2), $a = 9.553(5)$, $b = 13.499(6)$, $c = 15.693(5)$ Å, $\alpha = 108.29(3)$, $\beta = 93.18(4)$, $\gamma = 97.86(4)^\circ$, $U = 1893.0(14)$ Å³, $Z = 2$, $D_c = 2.89$ gm^{-3} , $F(000) = 1468$, $\mu(\text{Mo-K}\alpha) = 174.1$ cm^{-1} , crystal dimensions $0.10 \times 0.18 \times 0.25$ mm, 4987 unique diffractometer data. Structure solved by direct methods (SHELXTL PLUS⁴) and Fourier difference techniques, refined by full-matrix least-squares analysis on F^2 (all non-hydrogen atoms anisotropic, phenyl-H atoms placed in idealised positions and allowed to ride on the relevant C atoms, $U_H = 0.08$ Å²) (SHELXL 93⁵) to $R_1 = 0.0495$ ($wR_2 = 0.092$) for 4987 reflections [$I > 2\sigma(I)$], $w^{-1} = (\sigma F^2 + 0.035P)$ where $P = [(0 \text{ or } F_o^2) + 2F_c^2]/3$. Atomic coordinates, thermal parameters and bond lengths and angles have been deposited at the Cambridge Crystallographic Data Centre. See Instructions for Authors, *J. Chem. Soc., Dalton Trans.*, 1994, Issue 1, pp. xxiii–xxviii.

The molecular structure of **2** is shown in Fig. 1 together with some important bond parameters. The structure may be described as an osmium triangle with one of the osmium–osmium edges $[\text{Os}(1)\text{–Os}(3)]$ bridged by both a chlorine ligand and a gold ligand. The fourth osmium atom is incorporated into the gold ligand forming an almost linear $\text{Au}\text{–Os}(4)\text{–P}$ [$177.6(2)^\circ$] unit. Within the osmium triangle the length of the Os–Os edge that is bridged by the chlorine and the gold ligand is slightly longer $[\text{Os}(1)\text{–Os}(3)$ 2.908(2) Å] than is observed in the cluster $[\text{Os}_3(\text{CO})_{10}\{\text{Au}(\text{PPh}_3)\}(\mu\text{-Cl})]$ (Os–Os 2.86 Å).⁶ Within the osmium triangle the two osmium atoms involved in the bridged edge are both co-ordinated to three terminal carbonyl ligands while the third osmium atom is co-ordinated to four terminal carbonyl ligands two axial and two equatorial. A similar arrangement of the terminal carbonyl ligands has been found in $[\text{Os}_3(\text{CO})_{10}\{\text{Au}(\text{PPh}_3)\}(\mu\text{-Cl})]$.⁶

The chlorine ligand is symmetrically bridging the $\text{Os}(1)\text{–Os}(3)$ edge $[\text{Os}\text{–Cl}$ 2.457(6) and 2.445(6) Å] and the same is observed for the gold ligand with similar gold–osmium distances $[2.746(2)$ and $2.744(2)$ Å]. In the cluster $[\text{Os}_3(\text{CO})_{10}\{\text{Au}(\text{PPh}_3)\}(\mu\text{-Cl})]$ both the chlorine and the gold




Fig. 1 The molecular structure of $[\text{Os}_3\{\mu\text{-AuOs}(\text{CO})_4(\text{PPh}_3)\}(\mu\text{-Cl})(\text{CO})_{10}] \mathbf{2}$ showing the atomic numbering scheme. Selected bond lengths (Å) and angles ($^\circ$): Os(1)–Os(2) 2.842(2), Os(1)–Os(3) 2.908(2), Os(2)–Os(3) 2.852(2), Os(1)–Cl 2.445(6), Os(3)–Cl 2.457(6), Os(1)–Au 2.746(2), Os(3)–Au 2.744(2), Os(4)–Au 2.706(2), Os(4)–P 2.385(6); Os(1)–Cl–Os(3) 72.8(2), Os(4)–Au–Os(1) 145.25(5), Os(4)–Au–Os(3) 150.75(5), Os(3)–Au–Os(1) 63.97(5), P–Os(4)–Au 177.6(2)

ligand bridges could also be considered symmetric (Os–Au 2.77 and 2.75 Å, Os–Cl 2.41 and 2.43 Å).⁶ The Os–Au distances in **2** are slightly shorter than the Os–Au distances [2.802(2)–2.814(2) Å] in the anion $[\{\text{Os}_3\text{H}(\text{CO})_{10}\}_2\text{Au}]^-$ where the central Au atom links two Os₃ triangles by bridging Os–Os edges.⁷ The angle subtended at the gold atom by the two bridgehead osmium atoms is 63.97(5) $^\circ$. The gold atom makes an angle of 54 $^\circ$ to the plane of the osmium triangle which is smaller than the equivalent angle of 59 $^\circ$ for the $[\text{Os}_3(\text{CO})_{10}\{\text{Au}(\text{PPh}_3)\}(\mu\text{-Cl})]$ cluster.⁶

Within the gold ligand, Os(4) adopts an approximately octahedral geometry with four terminal carbonyl ligands in the same plane and the gold atom and the triphenylphosphine ligand *trans* to each other. To the best of our knowledge this is the first example of an Os(CO)₄ unit co-ordinated in this manner although the osmium triangle is virtually identical to the known compound $[\text{Os}_3(\text{CO})_{10}\{\text{Au}(\text{PPh}_3)\}(\mu\text{-Cl})]$.⁶ Consideration of the 18 electron rule for the Os(4) atom within the gold ligand suggests that it forms a dative bond to the gold atom. The Au–Os(4) bond length [2.706(2) Å] is shorter than

the observed osmium–gold distances in the osmium triangle.

The formation of the Au–Os(CO)₄–P linkage appears to be unprecedented. There are previous examples of an Au atom linking two metal clusters such as in the anion $[\{\text{Os}_3\text{H}(\text{CO})_{10}\}_2\text{Au}]^-$.⁷ Examples are also known where Au(PR₃) groups form terminal gold linkages to a variety of transition metals,⁸ such as in the dinuclear complexes $[\text{Mn}(\text{CO})_5\{\text{Au}(\text{PPh}_3)\}]$ and $[\text{V}(\text{CO})_6\{\text{Au}(\text{PPh}_3)\}]$, and in the Au-linked trimetal anion $[\text{Au}\{\text{Mn}(\text{CO})_5\}_2]^-$.⁹ In these systems the Au atom may be viewed as participating in a conventional two-centre two-electron covalent bond, in contrast to **2** where the 18 electron Os(4) centre may donate two electrons to the gold atom forming a dative covalent bond. Work is continuing to understand better the formation of **2** and to explore the chemistry of the new linkage.

Acknowledgements

We thank the SERC for financial support to (C. C.) and the European Economic Community for a Human Capital and Mobility Grant (to M.C.R.A.).

References

- G. R. Steinmetz, A. D. Harley and G. L. Geoffroy, *Inorg. Chem.*, **19**, 2985.
- C. W. Bradford, W. van Bronswijk, R. J. Clark and R. S. Nyholm, *J. Chem. Soc. A*, 1970, 2889.
- B. F. G. Johnson, D. A. Kaner, J. Lewis, P. R. Raithby and M. J. Taylor, *J. Chem. Soc., Chem. Commun.*, 1982, 314.
- SHELXTL PLUS, Program Version 4.0, Siemens Analytical X-Ray Instruments, Madison, WI, 1990.
- SHELXL 93, G. M. Sheldrick, Program for crystal structure refinement, University of Göttingen, 1993.
- C. M. Hay and P. R. Raithby, unpublished work.
- B. F. G. Johnson, D. A. Kaner, J. Lewis and P. R. Raithby, *J. Chem. Soc., Chem. Commun.*, 1981, 753.
- D. M. P. Mingos and M. J. Watson, *Adv. Inorg. Chem.*, 1993, **39**, 327.
- P. Braunstein and J. Dehand, *J. Organomet. Chem.*, 1975, **88**, C24.

Received 30th August 1994; Communication 4/05255B