


Synthesis and Structure-Activity Relationships of Potent New Angiotensin Converting Enzyme Inhibitors Containing Saturated Bicyclic Amino Acids¹

C. J. Blankley,* J. S. Kaltenbronn,* D. E. DeJohn, A. Werner, L. R. Bennett, G. Bobowski, U. Krolls, D. R. Johnson, W. M. Pearlman, M. L. Hoefle, A. D. Essenburg, D. M. Cohen, and H. R. Kaplan

Departments of Chemistry and Pharmacology, Warner-Lambert/Parke-Davis Pharmaceutical Research, Ann Arbor, Michigan 48105. Received November 14, 1986

The synthesis of a series of novel, potent angiotensin converting enzyme (ACE) inhibitors containing saturated bicyclic amino acids in place of proline is described. Octahydroindole-2-carboxylic acid, octahydroisindole-1-carboxylic acid, and octahydro-3-oxoisindole-1-carboxylic acid can replace proline in both sulfhydryl and non-sulfhydryl ACE inhibitors to give compounds equipotent to captopril and enalapril both in vitro and in vivo. Structure-activity relationships are discussed. Compound 11a (CI-907, indolapril) has advanced to clinical evaluation.


The development of inhibitors of angiotensin converting enzyme [EC 3.4.15.1] (ACE) has resulted in major new agents for the treatment of hypertension. The story of captopril (1a) has already become a classic in the field of rational drug design,^{2a,b} and this has inspired others to seek additional agents with the same mechanism of action. Several other structurally novel, potent ACE inhibitors have been reported, among them "keto-ACE"³ (1e) and enalapril⁴ (1c).



- 1a. R = HSCH₂CH(CH₃)CO
 b. R = HSCH₂CH₂CO
 c. R = PhCH₂CH₂CH(CO₂Et)NHCH(CH₃)CO
 d. R = PhCH₂CH₂CH(CO₂H)NHCH(CH₃)CO
 e. R = PhCH₂CH(NHCOPh)COCH₂CH₂CO

Compounds 1a-e contain L-proline as the C-terminal portion. Several structure-activity relationship (SAR) studies^{2a,b} have indicated that, although this amino acid may not be optimal for in vitro ACE inhibition, it, or other cyclic secondary amino acids, such as thiazolidine-2-carboxylic acid,⁵ is most effective for providing useful levels of in vivo activity. Studies on both captopril analogues² and simple dipeptide ACE inhibitors⁶ showed the enhanced in vitro activity of compounds containing certain lipophilic C-terminal amino acids such as phenylalanine and tryptophan, but these analogues proved to be inferior to the proline derivative in in vivo studies.² The suscep-


Scheme I^a



^a (a) H₂, Rh/C, HOAc. (b) Aqueous HCl. (c) *t*-BuOH, H₂SO₄, dioxane.

tibility of the former analogues to nonspecific amidases, in contrast to the proline derivatives, has been cited as the likely reason for diminished oral effectiveness.²

We earlier reported⁷ the result of a strategy of "cyclization" of the captopril molecule to provide a hydrolytically stable analogue with the essential functional group requirements intact. This resulted in 2, a compound in which both in vitro and in vivo ACE inhibition was retained, albeit weakly. Encouraged by this result, we



- (1) This paper has been presented in part; see *Abstracts of Papers*, 185th National Meeting of the American Chemical Society, Seattle, WA; American Chemical Society: Washington, DC, 1983; MEDI 66.
 (2) (a) Cushman, D. W.; Ondetti, M. A. *Prog. Med. Chem.* **1980**, *17*, 42-104. (b) Petrillo, E. W., Jr.; Ondetti, M. A. *Med. Res. Rev.* **1982**, *2*, 1-41.
 (3) (a) Almquist, R. G.; Chao, W. R.; Willis, M. E.; Johnson, H. R. *J. Med. Chem.* **1980**, *23*, 1392. (b) Meyer, R. F.; Nicolaides, E. D.; Tinney, F. J.; Lunney, E. A.; Holmes, A.; Hoefle, M. L.; Smith, R. D.; Essenburg, A. D.; Kaplan, H. R.; Almquist, R. G. *J. Med. Chem.* **1981**, *24*, 964. (c) Meyer, R. F.; Essenburg, A. D.; Smith, R. D.; Kaplan, H. R. *J. Med. Chem.* **1982**, *25*, 996.
 (4) Patchett, A. A.; Harris, E.; Tristram, E. W.; Wyvratt, M. J.; Wu, M. T.; Taub, R.; Peterson, E. R.; Ikeler, T. J.; Broeke, J. T.; Payne, L. G.; Ondeyka, D. L.; Thursett, E. D.; Greenlee, W. J.; Lohr, N. S.; Hoffsommer, R. D.; Joshua, H.; Ruyle, W. V.; Rothrock, W.; Aster, S. D.; Maycock, A. L.; Robinson, F. M.; Hirschman, R.; Sweet, C. S.; Ulm, E. H.; Gross, D. M.; Vassil, T. C.; Stone, C. A. *Nature (London)* **1980**, *288*, 280.
 (5) (a) Funae, Y.; Komori, T.; Sasaki, D.; Yomamoto, K. *Jpn. J. Pharmacol.* **1978**, *28*, 925. (b) Mita, L.; Iwao, J.; Masayuka, O.; Takehisa, C.; Tadashi, I. *Chem. Pharm. Bull.* **1978**, *26*, 1333.
 (6) Cheung, H. S.; Wang, F. L.; Ondetti, M. A.; Sabo, E. F.; Cushman, D. W. *J. Biol. Chem.* **1980**, *255*, 401.

sought to apply a similar reasoning to the proline terminus itself with the idea of converting potent in vitro inhibitors into compounds with equivalent in vivo potency. The success of one such attempt, the transformation of phenylalanine to 1,2,3,4-tetrahydroisoquinoline-3-carboxylic

- (7) Klutchko, S. L.; Hoefle, M. L.; Smith, R. D.; Essenburg, A. D.; Parker, R. B.; Nemeth, V. L.; Ryan, M. J.; Dugan, D. H.; Kaplan, H. R. *J. Med. Chem.* **1981**, *24*, 104.

acid and conversion of the latter to highly active in vitro and in vivo ACE inhibitors such as quinapril (3) is being described separately.⁸ In this paper we report further results along these lines where we sought to retain the favorable five-membered ring moiety of proline in an enhanced lipophilic environment. To this end, we examined the suitability of octahydroindole-2-carboxylic acid (4a) and octahydroisindole-1-carboxylic acids 5a and 6a as alternatives to proline in active ACE inhibitors.

Chemistry

At the time this work was initiated, neither 4a, 5a, or 6a had been described in the literature.⁹ We prepared racemic 4a by the method of Scheme I. Octahydro ethyl ester 4d showed no chromatographic evidence for the presence of more than one diastereoisomer, and in subsequent reactions, other derivatives of 4a behaved as single compounds. The identity of what is at least the highly predominant isomer was ascertained by conversion of 4a to its 3-bromobenzoyl derivative 7a, under Schotten-Baumann conditions. X-ray analysis of 7a established the all-cis disposition of hydrogen atoms at stereocenters around the five-membered ring.¹⁰

Racemic 4a was converted to sulfhydryl ACE inhibitor analogues 8a,b and 9a-d (Table I) by methods described in the literature^{11a} and outlined in Scheme IV. The two diastereoisomers of the α -methyl compounds 9a,b could be separated by fractional crystallization.


Resolution of 4a was accomplished initially via separation of α -methylbenzylamine salts of benzoyl derivative 7b, but subsequently the tartrate salts of the *tert*-butyl ester 4e also proved effective for this purpose. Since high activity was found to reside in the levorotatory isomer, resolved 4b was used to prepare 8c,d and 9e,f.

In most experiments, racemic 3-(acetylthio)-2-methylpropanoyl chloride was used to install the required side chain. Resolution of 3-(benzoylthio)-2-methylpropanoic acid by a literature method¹² provided the preferred *S* enantiomer of the side chain for the preparation of 9e and 9f.

The various stereoisomers of non-sulfhydryl compounds 11a-h (Table II) were prepared by coupling resolved *tert*-butyl esters 4f and 4g with the *S,S* and *R,S* diastereoisomers of 14 (Scheme V). The preparation of the optically pure isomers of 14 has been described.¹³ Noteworthy was the fact that steric hindrance of the amine group of 14 obviated the need to protect it during activation and coupling.


Compound 13 was obtained as a mixture of isomers by methods previously described.^{3b} Cyclization of 11a gave

Scheme II^a




^a (a) SOCl₂. (b) Br₂, Δ . (c) *t*-BuOH, Py. (d) PhCH₂NH₂. (e) H₂, Pd/C. (f) H₂, Rh/C, HOAc.

Scheme III^a



^a (a) Zn/HCl. (b) EtOH/HCl. (c) H₂, Rh/C. (d) NaOH.

Scheme IV




12, a crystalline material suitable for X-ray analysis.¹⁰ This established the absolute configuration of all chiral centers of 12, and hence of active isomer 11a as *S*, in analogy with enalapril.⁴ Also confirmed was the relative stereochemistry at the ring junction and carboxyl centers of 11a.

The octahydroisindole-1-carboxylic acid derivatives 5a-c served as precursors for ACE inhibitors 16 and 17, which were prepared as outlined in Scheme II. Compounds 5a-c again behaved as single diastereoisomers, and it was tentatively presumed that the hydrogens present at the chiral centers have a *cis* relationship in analogy with 4a,d,e. These compounds were employed only in racemic form. Non-sulfhydryl analogues 17a,b were obtained as predominantly single optical isomers by coupling (*S,S*)-14 with 5c and separating the resulting diastereoisomers. Assignments of configuration were made from the activity results and established SAR in analogous systems.⁴


Finally, it was of interest to prepare octahydro-3-oxo-isindole-1-carboxylic acid derivatives 18a-d and 19 in view of the high activity reported for analogues where pyroglutamic acid replaces proline.^{11a,b} The requisite precursor 6 was prepared as shown in Scheme III and again behaved as a single isomer. It was used in racemic form. The pyroglutamic acid analogues 20a,b^{11a} and 21 were prepared for comparison.¹⁴

- (8) Klutchko, S.; Blankley, C. J.; Fleming, R. W.; Hinkley, J. M.; Werner, A. E.; Nordin, I.; Holmes, A.; Hoefle, M. L.; Cohen, D. M.; Essenburg, A. D.; Kaplan, H. D. *J. Med. Chem.* 1986, 29, 1953.
- (9) A recent paper describes the synthesis of (-)-4b and (-)-4c by a different procedure. Vincent, M.; Remond, G.; Portevin, B.; Serkiz, B.; Lambie, M. *Tetrahedron Lett.* 1982, 23, 1677.
- (10) Determined by Dr. A. MacPhail, Duke University. X-ray crystal results on 7a and 12 are included in the supplementary material for this paper.
- (11) (a) Imaki, K.; Sakuyama, S.; Okada, T.; Toda, M.; Hayashi, M.; Miyamoto, T.; Kawasaki, A.; Okegawa, T. *Chem. Pharm. Bull.* 1981, 29, 2210. (b) Condon, M. E.; Petrillo, E. W., Jr.; Ryono, D. E.; Reid, J. A.; Neubeck, R.; Puar, M.; Heikes, J. E.; Sabo, E. F.; Losee, K. A.; Cushman, D. W.; Ondetti, M. A. *J. Med. Chem.* 1982, 25, 250.
- (12) Iwao, J.; Ooys, M.; Kato, E.; Watanabe, T. *Jpn. Kokai Tokkyo Koho* 79, 151, 912; *Chem. Abstr.* 1980, 92, 215076q.
- (13) Kaltenbronn, J. S.; DeJohn, D.; Krolls, U. *Org. Prep. Proc. Int.* 1983, 15, 35.

Table I. Physical Properties and in Vitro ACE Inhibitory Activity of Sulfhydryl Analogues


compd	R ₁	R ₂	X	synth ^a method	recryst solvent	mp, °C	config ^b	[α] ²³ _D , ^c deg	formula ^d	IC ₅₀ ^j , μM
8a	CH ₃ CO	H	A	A	EtOAc	131–133	racemic		C ₁₄ H ₂₁ NO ₄ S	0.56
8b	H	H	A	C	MeCN	145–146	racemic		C ₁₂ H ₁₉ NO ₃ S	0.013
8c	CH ₃ CO	H	A	A	EtOAc	110–112	S	-51.0	C ₁₄ H ₂₁ NO ₄ S	0.48
8d	H	H	A	C	EtOAc	168.5–170	S	-68.5	C ₁₂ H ₁₉ NO ₃ S	0.0064
9a	CH ₃ CO	CH ₃	A	A	EtOAc	172–173	(R,R + S,S)		C ₁₅ H ₂₃ NO ₄ S	0.16
9b	CH ₃ CO	CH ₃	A	A	EtOAc	151.5–153.5	(R,S + S,R)		C ₁₅ H ₂₃ NO ₄ S	35.0
9c	H	CH ₃	A	C	EtOAc	155–156	(R,R + S,S)		C ₁₃ H ₂₁ NO ₃ S	0.0077
9d	H	CH ₃	A	C	EtOAc	141–142	(R,S + S,R)		C ₁₃ H ₂₁ NO ₃ S	1.10
9e	PhCO	CH ₃	A	A	EtOAc	184.5–185.5	(S,S)	-135.6	C ₂₀ H ₂₅ NO ₄ S	0.54
9f	H	CH ₃	A	C	MeCN	145–148	(S,S)	-53.5	C ₂₅ H ₄₄ N ₂ O ₃ S	0.0052
10	m				EtOAc	148–152	racemic		C ₁₂ H ₁₇ NO ₂ S	0.007
16a	CH ₃ CO	H	B	B	EtOAc	140–145	racemic		C ₁₄ H ₂₁ NO ₄ S	1.2
16b	H	H	B	C		170–175 dec	racemic		C ₁₂ H ₁₆ N ₂ O ₃ S·C ₁₂ H ₂₃ N ^{e,i}	0.024
16c	CH ₃ CO	CH ₃	B	B		oil	mixture		C ₁₅ H ₂₃ NO ₄ S·0.3CHCl ₃ ^f	0.42
16d	H	CH ₃	B	C		119–129	mixture		C ₁₃ H ₂₁ NO ₃ S ^g	0.028
18a	CH ₃ CO	H	C	B	MeCN	198–203	racemic		C ₁₄ H ₁₉ NO ₅ S·C ₁₂ H ₂₃ N ⁱ	0.66
18b	H	H	C	C	MeCN	188–208	racemic		C ₁₂ H ₁₇ NO ₄ S·C ₁₂ H ₂₃ N ⁱ	0.013
18c	CH ₃ CO	CH ₃	C	B	EtOAc	185–188	mixture		C ₁₅ H ₂₁ NO ₅ S·C ₁₂ H ₂₃ N ⁱ	0.3
18d	H	CH ₃	C	C	MeCN	191–200 dec	mixture		C ₁₃ H ₁₉ NO ₄ S·C ₁₂ H ₂₃ N ^{h,i}	0.014
20a	H	H	n	B	MeCN	185–191 then 225–227	S	-31.7	C ₂₀ H ₃₄ N ₂ O ₄ S	0.96 ^j
20b	H	CH ₃	n	B	MeCN	188–191.5	(S,S)	-58.6	C ₂₁ H ₃₆ N ₂ O ₄ S	0.0093 ^j
23a	H	H	D				racemic			0.14 ^k
23b	H	CH ₃	D				(S,S)			0.0037 ^k
captopril (1a)	H	CH ₃	proline				(S,S)		C ₁₂ H ₁₇ NO ₂ S	0.012
1b	H	H	proline				S			0.82

^a See Experimental Section. ^b "Mixture" refers to an unseparated mixture of diastereomers. ^c c 1.0, 1:1 MeOH/0.1 N HCl. ^d Analyses for C, H, N were within ±0.4% except as noted. ^e C: calcd, 65.71; found, 65.24. ^f C: calcd, 52.62; found, 51.95. ^g C: calcd, 57.53; found, 56.17. ^h N: calcd, 6.50; found, 5.89. ⁱ Isolated as the dicyclohexylamine salt. ^j Molar concentration required for 50% inhibition. ^k Reference 15. ^l See reference 11a. The authors found IC₅₀ values for 20a of 0.06 μM and for (S,S)-20b 0.0036 μM. ^m See the structure for 10. ⁿ Pyroglutamic acid.

Table II. Physical Properties and in Vitro ACE Inhibitory Activity of Non-Sulfhydryl Analogues


compd	R ₁	X	synth ^a method	mp, °C	config ^b	[α] ²³ _D , ^c deg	formula ^d	IC ₅₀ ^h , μM
11a	C ₂ H ₅	A	D	amorph	SSS	-28.8	C ₂₄ H ₃₄ N ₂ O ₅ ·HCl·0.5H ₂ O	0.080
11b	H	A	E	165–166 dec	SSS	-34.0	C ₂₂ H ₃₀ N ₂ O ₅	0.0024
11c	C ₂ H ₅	A	D	amorph	RSS	-66.8	C ₂₄ H ₃₄ N ₂ O ₅ ·0.5H ₂ O ^e	1.30
11d	H	A	E	amorph	RSS		C ₂₂ H ₃₀ N ₂ O ₅	0.011
11e	C ₂ H ₅	A	D	amorph	SSR	+28.8	C ₂₄ H ₃₄ N ₂ O ₅ ·0.25H ₂ O ^f	57.0
11f	H	A	E	147–149 dec	SSR	+41.9	C ₂₂ H ₃₀ N ₂ O ₅ ·H ₂ O ^g	2.4
11g	C ₂ H ₅	A	D	amorph	RSR	-1.9	C ₂₄ H ₃₄ N ₂ O ₅ ·HCl·H ₂ O ^h	>100
11h	H	A	E	124–126 dec	RSR	+8.9	C ₂₂ H ₃₀ N ₂ O ₅ ·0.5H ₂ O	6.4
12	m			138–141	SSS	-53.5	C ₂₄ H ₃₂ N ₂ O ₄	>100
13	m			amorph	mixture		C ₂₈ H ₃₂ N ₂ O ₅ ·0.5H ₂ O ⁱ	0.011
17a	C ₂ H ₅	B	D	198–216 dec	SSS	-33.6	C ₂₄ H ₃₄ N ₂ O ₅ ·HCl	0.17
17b	C ₂ H ₅	B	D	204–214 dec	SSR	+24.8	C ₂₄ H ₃₄ N ₂ O ₅ ·HCl·0.8H ₂ O	7.50
17c	H	B	E	186–190	SSS	-36.5	C ₂₂ H ₃₀ N ₂ O ₅ ·1.2H ₂ O	0.0026
17d	H	B	E	145–163	SSR	+33.2	C ₂₂ H ₃₀ N ₂ O ₅ ·1.5H ₂ O	0.018
19	C ₂ H ₅	C	D	amorph	(SSS + SSR)	+9.5	C ₂₄ H ₃₂ N ₂ O ₅ ·HCl·1.2H ₂ O ^j	0.34
21	C ₂ H ₅	n	D	amorph	SSS	-17.1 ^l	C ₂₀ H ₂₆ N ₂ O ₆ ·HCl·1.1H ₂ O	0.063
enalapril (1c)	C ₂ H ₅	o			SSS			0.14
enalaprilat (1d)	H	o			SSS			0.0023
1e	m							0.0032

^a See Experimental Section. ^b "Mixture" refers to an unseparated mixture of diastereomers. The S and R configurations refer to the optical centers present as one proceeds from left to right in the structure as indicated. The bicyclic amino acid portion is considered as a unit and is designated as either S or R. ^c c 1.0, 1:1 MeOH/0.1 N HCl. ^d Analyses for C, H, N were within ±0.4% except as noted. ^e H: calcd, 7.62; found, 7.13. ^f H: calcd, 7.59; found, 6.80. ^g H: calcd, 7.67; found, 7.15. ^h H: calcd, 7.69; found, 6.97. ⁱ N: calcd, 5.77; found, 5.28. ^j C: calcd, 57.35; found, 56.85. ^k Molar concentration required for 50% inhibition. ^l Reference 14 reports [α]²³_D = -31.8° (c 0.96, EtOH). ^m See text. ⁿ Pyroglutamic acid. ^o Proline.

Table III. Summary of Effects of Certain ACE Inhibitors on Blood Pressure in the Conscious Renal (2 Kidney/1 Clip) Hypertensive Rat

compd	dose, ^a mg/kg po	no. tested	mean aortic blood pressure	
			base line, ^b mmHg	max change, ^c mmHg
8b	3	5	198 ± 5	-73 (at 8 h)
	10	5	198 ± 5	-89 (at 3 h)
	30	5	198 ± 5	-91 (at 5 h)
9c	0.1	4	184 ± 7	-16 (at 6 h)
	0.3	4	185 ± 14	-19 (at 8 h)
	1.0	4	178 ± 9	-72 (at 5 h)
	3.0	4	190 ± 5	-81 (at 5 h)
	30	4	197 ± 6	-115 (at 10 h)
9f	30	3	193 ± 6	-72 (at 6 h)
16b	30	3	184 ± 3	-76 (at 6 h)
16d	30	4	152 ± 3	-50 (at 6 h)
18b	30	5	205 ± 6	-57 (at 8 h)
18d	30	3	192 ± 19	-100 (at 8 h)
20b	30	3	185 ± 13	-48 (at 8 h)
11a	0.1	4	185 ± 5	-52 (at 10 h)
	0.3	4	191 ± 10	-73 (at 5 h)
	1.0	4	190 ± 5	-72 (at 3 h)
	10.0	4	197 ± 7	-86 (at 6 h)
	30.0	4	185 ± 5	-102 (at 6 h)
11b	30	4	198 ± 5	-104 (at 5 h)
13	30	4	169 ± 2	-27 (at 8 h)
17a	3	4	199 ± 12	-87 (at 6 h)
1a (captopril)	0.3	4	189 ± 6	-25 (at 2 h)
	3.0	4	192 ± 7	-93 (at 6 h)
	30.0	4	186 ± 4	-101 (at 6 h)
1c (enalapril)	0.3	4	201 ± 9	-39 (at 5 h)
	1.0	4	175 ± 2	-24 (at 8 h)
	3.0	4	200 ± 5	-56 (at 5 h)
	10.0	4	199 ± 5	-56 (at 4 h)


^a All compounds were dissolved/suspended in 4% gum acacia with the exception of 11b, which was dissolved in a 0.1% aqueous solution of methocel, and 13, which was dissolved in 5% EtOH. ^b All values are the mean ± 1 SEM. ^c Average result for three to five animals. Vehicle controls average 183 ± 5 mmHg with maximum decreases about 16–20 mmHg, or 10%. Hence, blood pressure decreases in excess of 20 mmHg are taken as significant for the purposes of this comparison.

Table IV lists the physical constants of the new amino acids and derivatives used in this study.

Biological Results and Discussion

The assay for in vitro ACE inhibition was performed as previously described⁷ with Hip-Gly-Gly as the synthetic substrate. In vitro results are reported in Tables I and II. It is clear that, allowing for racemic mixtures in some cases, all of the bicyclic analogues are at least as potent in vitro as their monocyclic counterparts. In the sulfhydryl series, the derivative with the octahydroindole-2-carboxylic acid in place of proline (compound 9f) actually improves the IC₅₀ by 2.3-fold. A recent study has reported a similar activity enhancement of a related proline replacement, indoline-2-carboxylic acid¹⁵ (IC₅₀ = 0.0037 μM for 23b). It is quite clear though that the additional geometrical features present in 4b can impart enhanced in vitro activity to analogues with less than optimal side chains, at least when compared with the proline series. For example, compare 8d (IC₅₀ = 0.006 μM), 1b (IC₅₀ = 0.82 μM),¹⁶ and 23a (IC₅₀ = 0.14 μM). This represents an activity en-

Scheme V



hancement of 136-fold for 8d relative to 1b and 23-fold for 8d over 23a. Data given for the isomeric octahydroindole analogues suggest that similar activity enhancements are obtained in these series as well. Quantitation is not possible, however, due to the unavailability of pure chiral diastereoisomers. Cyclic analogue 10 was prepared as a possible prodrug form of 8b. Surprisingly this compound showed in vitro activity comparable to that of 8b. It was not determined, however, whether 8b was actually formed under the assay conditions.

With regard to the non-sulfhydryl analogues, a similar pattern is observed. Optimal isomers 11b and 1d have nearly identical in vitro activities, while suboptimal 11d shows a 75-fold activity enhancement over the corresponding diastereoisomer of 1d (IC₅₀ = 0.82 μM).⁴ Again, the octahydroindole derivatives give comparable activity to their monocyclic analogues, reflecting further a parallelism in the SAR for the captopril and enalapril types. Although the data of Table II does not allow a definitive judgment on the effect of octahydroindole vs. proline in the "keto-ACE" type of inhibitor, we have observed the substantial activity enhancing effect of the octahydroindole-2-carboxylic acid in other series of non-sulfhydryl ACE inhibitors.¹⁷

Oral activity of selected analogues in 2-kidney-1-clip Goldblatt hypertensive rats is reported in Table III. All of the optimum diastereoisomers show potent activity in this model. Additional pharmacology of 11a (CI-907, SCH 31846, indolapril) has been reported elsewhere,^{18a-c,19} as well

- (14) Compound 21 was recently reported. See: Johnson, A. L.; Price, W. A.; Wong, P. C.; Vavala, R. F.; Stump, J. M. *J. Med. Chem.* 1985, 28, 1596.
 (15) Kim, D. H.; Guinosso, C. J.; Buzby, G. C., Jr.; Herbst, D. R.; McCaully, R. J.; Wicks, T. C.; Wendt, R. L. *J. Med. Chem.* 1983, 26, 394.
 (16) Reference 2 gives IC₅₀ values of 1d as 0.20 μM and 1a as 0.023 μM; Hip-His-Leu as substrate.

- (17) Roark, W. H.; Tinney, F. J.; Cohen, D.; Essenburg, A. D.; Kaplan, H. R. *J. Med. Chem.* 1985, 28, 1291.
 (18) (a) Kaplan, H. R.; Burmeister, W.; Essenburg, A.; Major, T. C.; Mertz, T.; Randolph, A. *Pharmacologist* 1982, 24, 176 (Abstr 446). (b) Ryan, M. J.; Boucher, D. M.; Cohen, D. M.; Olszewski, B. J.; Singer, R. M.; Smith, R. D.; Kaplan, H. R. *Pharmacologist* 1982, 24, 176 (Abstr 447). (c) Ryan, M. J.; Boucher, D. M.; Cohen, D. M.; Essenburg, A. D.; Major, T. C.; Mertz, T. E.; Olszewski, B. J.; Randolph, A. E.; Singer, R. M.; Kaplan, H. R. *J. Pharmacol. Exp. Ther.* 1984, 228, 312. (d) Borondy, P. E.; Michniewicz, B. M.; Yakatan, G. J. *Pharmacologist* 1982, 24, 95 (Abstr 14).

Table IV. New Amino Acids and Derivatives

compd	R ₁	R ₂	salt	mp, °C	[α] ²³ _D , ^a deg	formula ^b
4a	H	H		238–240	racemic	C ₉ H ₁₅ NO ₂
4b	H	H		268–270 dec	–47.9	C ₉ H ₁₅ NO ₂
4c	H	H		267–268 dec	+48.4	C ₉ H ₁₅ NO ₂ ^c
4d	C ₂ H ₅	H		oil	racemic	C ₁₁ H ₁₉ NO ₂ ^d
4e	<i>t</i> -C ₄ H ₉	H		oil	racemic	C ₁₃ H ₂₃ NO ₂ ^c
4f	<i>t</i> -C ₄ H ₉	H		oil	–29.4	C ₁₃ H ₂₃ NO ₂ ^c
4g	<i>t</i> -C ₄ H ₉	H		oil	+29.9	C ₁₃ H ₂₃ NO ₂ ^c
5a	H	H	HCl	257–262 dec	racemic	C ₉ H ₁₅ NO ₂ ·HCl·0.2H ₂ O
5b	C ₂ H ₅	H	HCl	174–176	racemic	C ₁₁ H ₁₉ NO ₂ ·HCl
5c	<i>t</i> -C ₄ H ₉	H	HCl	170–176 dec	racemic	C ₁₃ H ₂₃ NO ₂ ·HCl·0.15H ₂ O
6	H	H		205–213 dec	racemic	C ₈ H ₁₃ NO ₃
7a	H	3-BrPhCO		171–174	racemic	C ₁₆ H ₁₈ BrNO ₃
7b	H	PhCO		191–193	racemic	C ₁₆ H ₁₉ NO ₃
7c	H	PhCO		169–171	–48.9	C ₁₆ H ₁₉ NO ₃
7d	H	PhCO		169–171	+50.5	C ₁₆ H ₁₉ NO ₃ ^c
15	<i>t</i> -C ₄ H ₉	H	HCl	148–154 dec	racemic	C ₁₃ H ₁₇ NO ₂ ·HCl·0.2H ₂ O

^a c 1, MeOH. ^b Analyses for C, H, N were within ±0.4% except as noted. ^c Not analyzed. ^d C: calcd, 66.97; found, 66.54.

as pharmacokinetic studies preparatory to clinical development.^{18d}

In summary, we have described the potent ACE inhibitory and blood pressure lowering effects of a number of saturated bicyclic amino acid derivatives. These not only confirm the salutary effect on ACE activity of additional lipophilic character at the proline site in the model proposed by Ondetti et al.^{2a,b} but also add information about bulk and shape tolerance in this region. The question of real interest now is to what extent these altered geometrical and physicochemical properties will manifest themselves in a physiological setting.

Experimental Section

Melting points were determined in a Thomas-Hoover capillary melting point apparatus or a Mel-Temp apparatus. Infrared (IR) data were recorded on a Beckman IR-9 or IR-7 prism grating instrument on a Digilab FTS-14 interferometer. Nuclear magnetic resonance measurements (NMR) were made on a Bruker WH-90 pulsed Fourier transform instrument. IR and NMR were compatible with the assigned structures. Homogeneity of the products was determined by ascending thin-layer chromatography (TLC) on precoated TLC sheets (silica gel 60 F 254, Merck), using principally the solvent system HOAc–MeCN–toluene (1:9:10). HPLC analyses were carried out on a C18 reverse phase Alltech column using 40% acetonitrile/60% 0.005 M Pic A as the solvent. Peaks were detected by UV at 210 nm.

Coupling of Bicyclic Amino Acids with Sulfur-Containing Side Chains Leading to Analogues of the Sulfhydryl Type. A General Procedure Illustrated with the Preparation of (2 α ,3 $\alpha\beta$,7 $\alpha\beta$)-1-[3-(Acetylthio)-2-methyl-1-oxopropyl]octahydro-1H-indole-2-carboxylic Acid (9a and 9b). Method A. A mixture of 3.0 g (0.02 mol) of the amino acid 4a and 3.5 g (0.02 mol) of hexamethyldisilazane in 15 mL of CH₃CN was treated with a few drops of chlorotrimethylsilane and heated at reflux for 3 h. The resulting solution was then cooled in an ice bath and treated dropwise with a solution of 3.2 g (0.02 mol) of 3-(acetylthio)-2-methylpropionyl chloride in 5 mL of CH₃CN. After 1 h at 0 °C, the solution was allowed to warm to room temperature and then warmed to reflux, allowing the major portion of the

chlorotrimethylsilane and CH₃CN to escape. The solution was then recooled to 0 °C and 0.5 mL of H₂O added, and the solution was warmed briefly to reflux and filtered through Celite. Removal of the solvent under reduced pressure left an oil as a mixture of diastereomers.

Trituration of the oil with Et₂O caused separation of the less soluble diastereomer 9a. Several recrystallizations from EtOAc gave pure 9a. HPLC analysis showed only one diastereomer.

Extensive fractional recrystallization of the material obtained from the mother liquors and soluble in cyclohexane gave a sample of the more soluble isomer 9b. HPLC analysis showed less than 8% of 9a present.

Alternate Coupling Leading to Analogues of the Sulfhydryl Type. A General Procedure Illustrated with the Preparation of 2-[3-(Acetylthio)-1-oxopropyl]octahydro-1H-indole-1-carboxylic Acid (16a). Method B. A mixture of 2.0 g (9.7 mmol) of 5a and 2.3 g (29 mmol) of pyridine in 25 mL of THF was cooled in ice and treated dropwise with 1.8 g (10.8 mmol) of 3-(acetylthio)propionyl chloride. After 2 h in the cold, the mixture was allowed to warm to room temperature over 1 h and then concentrated under reduced pressure. The residue was taken up in EtOAc and washed twice with dilute HCl. Drying of the EtOAc over MgSO₄ and removal of the solvent under reduced pressure left an oil which solidified. Two recrystallizations from EtOAc gave 1.3 g (45%) of 16a.

Removal of the Acetyl Group, a General Procedure Illustrated with the Preparation of (2 α ,3 $\alpha\beta$,7 $\alpha\beta$)-Octahydro-1-(3-mercapto-2-methyl-1-oxopropyl)-1H-indole-2-carboxylic Acid (9c). Method C. Under N₂, a solution of 1.0 g (3 mmol) of 9a in 10 mL of a 5 N solution of NH₃/MeOH was stirred at room temperature for 2.5 h. The mixture was concentrated under reduced pressure and taken up in H₂O. The pH was brought to 2.0 with 10% NaHSO₄ and extracted twice with EtOAc. The EtOAc was washed with saturated NaCl and dried over MgSO₄. Removal of the solvent under reduced pressure left 0.8 g of a white solid. Recrystallization from EtOAc gave pure 9c.

Coupling of Bicyclic Amino Acids Leading to Analogues of the Non-Sulfhydryl Type (Table II). A General Procedure Illustrated with the Preparation of [2S-[1-[R*(R*)],2 α ,3 $\alpha\beta$,7 $\alpha\beta$]]-1-[2-[[1-(Ethoxycarbonyl)-3-phenylpropyl]amino]-1-oxopropyl]octahydro-1H-indole-2-carboxylic Acid Hydrochloride (11a). Method D. A solution of 8.0 g (28.6 mmol) of (S,S)-14,¹³ 6.62 g (28.6 mmol) of 4f, and 3.87 g (28.6 mmol) of 1-hydroxybenzotriazole in 120 mL of THF was cooled in ice and treated dropwise with a solution of 5.91 g (28.6 mmol) of N,N'-dicyclohexylcarbodiimide in 15 mL of THF. The solution was stirred at 0 °C for 1 h and then at room tem-

(19) Baum, T.; Sybertz, E. J.; Ahn, H. S.; Watkins, R. W.; Powell, M. L.; LaRocca, P. T. In *New Drugs Annual: Cardiovascular Drugs*; Scriabine, A., Ed.; Raven: New York, 1985; Vol. 3, p 43.

perature overnight. The dicyclohexylurea was filtered off and the solvent removed under reduced pressure. The residue was taken up in EtOAc and washed with saturated NaHCO₃ and then saturated NaCl. After drying over MgSO₄, the solvent was removed under reduced pressure. The residue was taken up in hexane, charcoaled, and filtered through Celite, and the solvent was removed under reduced pressure to give 13.9 g of the crude diester as an oil; $[\alpha]^{23}_D$ -69.3° (c, 1, MeOH). HPLC analysis showed the diester to be 97% pure.

A solution of 13.9 g (28.6 mmol) of the diester in 135 mL of CH₂Cl₂ was cooled to 0 °C and saturated with HCl gas. After the solution was allowed to stand at 0 °C overnight, the solvent was removed under reduced pressure. Additional CH₂Cl₂ was added and the solvent removed again. The residue was then taken up in 100 mL of CH₂Cl₂, charcoaled, filtered through Celite, and concentrated under reduced pressure to a foam. After trituration with Et₂O, the solid was dissolved in H₂O, filtered, and lyophilized to give 12.1 g (93% yield) of 11a as an amorphous solid. HPLC analysis showed this to be 99% pure.

Hydrolysis to the Diacid. A General Procedure Illustrated with the Preparation of [2*S*-[1-[*R-(*R**)],2*α*,3*αβ*,7*αβ*]]-1-[2-[(1-Carboxy-3-phenylpropyl)-amino]-1-oxopropyl]octahydro-1*H*-indole-2-carboxylic Acid (11*b*). Method E.** A solution of 10.0 g (21.4 mmol) of 11a in 50 mL of H₂O was treated with 50 mL (70.7 mmol) of 1.4 N NaOH and stirred at room temperature overnight. The pH was brought to 4.5 with dilute HCl, a small amount of insoluble material filtered off, and the filtrate lyophilized. The residue was treated with warm EtOH to remove NaCl and the EtOH then removed under reduced pressure. The residue was purified by ion-exchange chromatography on Amberlite IR-120, eluting with 2 N NH₄OH. Removal of the solvent under reduced pressure and trituration of the residue with H₂O gave 6.61 g (76.7%) of 11b as a white solid.

(6*αα*,10*αα*,11*αα*)-Decahydro-1*H*,5*H*-[1,4]thiazepino[4,3-*a*]indole-1,5-dione (10). A solution of 0.4 g (1.9 mmol) of *N,N*-dicyclohexylcarbodiimide and 0.2 mL of pyridine in 35 mL of CH₂Cl₂ was cooled in ice and treated dropwise with a solution of 0.5 g (1.9 mmol) of 8b in 5 mL of CH₂Cl₂ over 20 min. The mixture was then stirred for 2 days at room temperature, filtered, and concentrated under reduced pressure. The product was isolated by chromatography on silica gel, eluting with CHCl₃/MeOH/HOAc (196/3/1). Combination of the appropriate fractions and recrystallization from EtOAc gave pure 10.

[3*S*-[2(*R),3*α*,5*αβ*,9*αβ*,10*αβ*]]-Decahydro-3-methyl-1,4-dioxo-*α*-(2-phenylethyl)pyrazino[1,2-*a*]indole-2(1*H*)-acetic Acid Ethyl Ester (12).** A solution of 9.4 g (20 mmol) of 11a and 3.0 g (20 mmol) of 1-hydroxybenzotriazole in 200 mL of THF was cooled to 0 °C and treated with 21 mL (21 mmol) of a 1 M solution of *N,N*-dicyclohexylcarbodiimide in THF. After the mixture was stirred at room temperature for 16 h, the dicyclohexylurea was filtered off and the filtrate concentrated under reduced pressure. The residue was taken up in EtOAc and washed with saturated NaHCO₃ and then saturated NaCl. After drying over MgSO₄ and removal of the solvent under reduced pressure, the crude product was obtained. Recrystallization from MeOH gave 4.2 g of 12, mp 143–146 °C.

1-[5-(Benzoylamino)-1,4-dioxo-6-phenylhexyl]octahydro-1*H*-indole-2-carboxylic Acid (13). To a slurry of 16.2 g (0.05 mol) of 6-phenyl-5-(benzoylamino)-4-oxohexanoic acid in 100 mL of CH₃CN was added in portions 12.0 g (0.074 mol) of carbonyldiimidazole. On warming to 30 °C, solution soon occurred and the solution was kept at this temperature for 0.5 h. The solution was cooled in ice and treated with a solution of 8.46 g (0.05 mol) of 4a in 100 mL of CH₃CN. The solution was then allowed to stir at room temperature overnight. The solvent was removed under reduced pressure, and the residue taken up in CH₂Cl₂ and washed with dilute HCl and then with water. After drying over MgSO₄, the solvent was removed under reduced pressure, leaving 20.8 g of the crude product as a foam. A portion of this was chromatographed on silica gel, eluting with EtOAc. Combining the appropriate fractions gave 13 as a mixture of diastereomers.

(2*α*,3*αβ*,7*αβ*)-Octahydro-1*H*-indole-2-carboxylic Acid Ethyl Ester (4*d*). A solution of 100 g (0.53 mol) of indole-2-carboxylic acid ethyl ester²⁰ in 1 L of absolute EtOH and 32 mL of con-

centrated H₂SO₄ was hydrogenated in a Parr apparatus with 4.0 g of a 10% Rh/C catalyst. After the required amount of H₂ had been taken up, the catalyst was removed by filtration, and the filtrate was evaporated in vacuo. The syrupy residue was dissolved in ice water and first neutralized with K₂CO₃ and then made basic with KHCO₃. The mixture was extracted twice with Et₂O and the Et₂O washed with saturated NaCl. Drying over Na₂SO₄ and removal of the solvent under reduced pressure gave 78.5 g of the ester as a colorless oil of high purity.

(2*α*,3*αβ*,7*αβ*)-Octahydro-1*H*-indole-2-carboxylic Acid (4*a*). A solution of 2.0 g (0.01 mol) of 4*d* in 25 mL of 15% HCl was heated at reflux for 4 h and then evaporated to dryness in vacuo. Recrystallization from CH₃CN/EtOAc gave 1.7 g of hydrochloride salt, mp 186–187 °C dec.

The free acid was obtained by dissolving 1.2 g of the hydrochloride in 10 mL of H₂O and adding 2 N NaOH to pH 6.5. The resulting solution was evaporated to dryness under reduced pressure and the residue then refluxed with 50 mL of CH₃CN and filtered hot. Concentration of the filtrate to 10 mL and cooling gave 0.5 g of pure 4a.

(2*α*,3*αβ*,7*αβ*)-1-Benzoyloctahydro-1*H*-indole-2-carboxylic Acid (7*b*). A solution of 58.32 g (0.344 mol) of 4a in 290 mL of H₂O was cooled in ice and treated dropwise simultaneously over a 1-h period with 2 N NaOH and 42.0 mL (0.36 mol, a 5% XS) of benzoyl chloride, while the pH was maintained between 10 and 11 and the temperature between 5 and 10 °C. When the addition was complete, the mixture was allowed to stir at 5 °C for 3 h while base was added to maintain the pH at 10–11. Approximately 350 mL of 2 N NaOH was used. The pH was brought to 1.8 with dilute HCl and the solid collected and washed with water. Recrystallization from EtOH/H₂O gave 91.0 g (96.6%) of pure acid, mp 191–193 °C.

(2*α*,3*αβ*,7*αβ*)-1-(3-Bromobenzoyl)octahydro-1*H*-indole-2-carboxylic Acid (7*a*). By a procedure similar to the above, but using 3-bromobenzoyl chloride, there was obtained 7a as a crystalline solid.

Resolution of 7*b* To Give 7*c*. A solution of 92.93 g (0.7668 mol) of (*S*)-(-)-*α*-methylbenzylamine in 1670 mL of MeOH was treated with 209.6 g (0.7668 mol) of 7b and the mixture swirled. Almost all dissolved. The solution was warmed slightly on the steam bath and filtered. The filtrate was diluted with 3.0 L of EtOAc and seeded. Crystallization started at once. After swirling occasionally at room temperature for 1 h, the mixture was placed in the cold room overnight. The solid was collected and washed with cold EtOAc. There was obtained 114.4 g (75.6%) of the salt; mp 210–213 °C; $[\alpha]^{23}_D$ -48.3° (c 1.04, MeOH).

Recrystallization of 109.4 g of this salt from 1.1 L of MeOH and 2.2 L of EtOAc gave 87.87 g of pure salt; mp 210–214 °C; $[\alpha]^{23}_D$ -50.0° (c 1.02, MeOH).

A suspension of 87.87 g of the salt in 1617 mL of H₂O and 650 mL of MeOH was brought to pH 1.5 with dilute HCl. Almost all dissolved and another solid formed. After being stirred at room temperature for 1 h, the mixture was diluted with 878 mL of H₂O and cooled. The solid was collected and washed with H₂O. There was obtained 52.63 g of pure 7c.

Resolution of 7*b* To Give 7*d*. By a procedure similar to the above but using (*R*)-(+)-*α*-methylbenzylamine as the resolving agent there was obtained 7d as a crystalline solid.

Hydrolysis of 7*c* To Give 4*b*. A suspension of 66.28 g (0.242 mol) of 7c in 660 mL of concentrated HCl and 660 mL of H₂O was heated at reflux for 4 h. All material was in solution after approximately 15 min. At the end of the heating period the solution was diluted with 660 mL of H₂O and cooled. The precipitated benzoic acid was filtered off and the filtrate extracted twice with CHCl₃. The pH was brought to 6.4 with 50% NaOH and the material concentrated to a slush on a rotary evaporator. This material was transferred to a tray and dried in a vacuum oven, leaving 516 g of a white solid. This material was digested with two 3-L portions of boiling anhydrous EtOH. Removal of the EtOH under reduced pressure left 52.5 g of the crude product. This material was desalted by passing over an Amberlite IR-120 column, eluting with 2 N NH₄OH. Removal of the solvent under

(20) Noland, W. E.; Baude, F. J. *Organic Syntheses*; Wiley: New York, 1973; Collect. Vol. V, p 567.

reduced pressure left 41 g of a white solid. Two recrystallizations from MeOH/anhidrous EtOH gave 36.01 g (87.8%) of resolved product, **4b**.

Hydrolysis of 7d To Give 4c. Hydrolysis of **7d** followed by a workup similar to the above gave **4c**.

(2 α ,3 $\alpha\beta$,7 $\alpha\beta$)-Octahydro-1*H*-indole-2-carboxylic Acid 1,1-Dimethylethyl Ester (4e). A solution of 14.23 g (0.084 mol) of **4a** in 150 mL of dioxane contained in a pressure vessel was treated with 15 mL of concentrated H₂SO₄ and 84 g of isobutylene and kept at 20 °C for 20 h with stirring. The mixture was then poured into ice water containing 45 mL (0.86 mol) of 50% NaOH solution and the mixture extracted three times with ether. The ether was washed with H₂O and then saturated NaCl solution. Drying over MgSO₄ and removal of the ether under reduced pressure gave 14.4 g (76% yield) of the ester as an oil. The ester is sufficiently pure for use in subsequent reactions.

Resolution of 4e To Give 4f. A solution of 3.3 g (0.022 mol) of *l*-tartaric acid (unnatural tartaric acid) in 20 mL of hot THF was treated with 5.0 g (0.022 mol) of **4e** in 10 mL of THF and the resulting solution was allowed to stand at room temperature overnight. The precipitated solid was collected and dried, giving 3.9 g of salt; mp 147–148 °C; [α]_D²⁵ -26.9° (c 1.09, MeOH). Recrystallization from 80 mL of EtOAc/MeOH (1:1) while the temperature was maintained at 0 °C overnight gave 3.11 g of pure salt; mp 152–153 °C; [α]_D²⁵ -31.7° (c 1.1, MeOH).

A suspension of 3.06 g of this salt in 30 mL of Et₂O was shaken with 30 mL of 5% Na₂CO₃ solution. The layers were separated, and the aqueous phase was extracted two times with Et₂O. The combined Et₂O layers were washed with H₂O and dried over MgSO₄. Removal of the Et₂O under reduced pressure gave 1.82 g of **4f** as an oil. The product was sufficiently pure for use in subsequent reactions.

Resolution of 4e To Give 4g. A solution of 17.17 g (0.114 mol) of *d*-tartaric acid (natural tartaric acid) in 130 mL of hot THF was treated with a solution of 25.78 g (0.114 mol) of **4e** (recovered from previous resolutions and enriched in the (+)-rotating isomer) in 85 mL of THF and allowed to stand at room temperature overnight. The precipitated solid was collected and dried, giving 36.6 g of salt. Recrystallization from 750 mL of EtOAc/MeOH (1:1) gave 31.55 g of pure salt; mp 149–150 °C dec; [α]_D²⁵ +31.5° (c 1.02, MeOH).

A suspension of this salt in 320 mL of Et₂O was shaken with 320 mL of 5% Na₂CO₃ solution. The layers were separated, and the aqueous phase was extracted two times with Et₂O. The combined Et₂O layers was washed with H₂O and dried over MgSO₄. Removal of the solvent under reduced pressure gave 18.57 g of **4g** as an oil. The product was sufficiently pure for use in subsequent reactions.

Octahydro-1*H*-isoindole-1-carboxylic Acid Hydrochloride (5a) and Esters 5b and 5c. A solution of 2.14 g of methyl 2,3-dihydro-1*H*-isoindole-1-carboxylate hydrochloride²¹ in 100 mL of MeOH and 5 mL of HOAc was hydrogenated in a Parr apparatus using 0.5 g of a 10% Rh/C catalyst. When the required amount of hydrogen had been taken up, the catalyst was filtered off and the filtrate concentrated under reduced pressure. Trituration of the residue with Et₂O gave a solid. Recrystallization from MeOH/Et₂O gave pure methyl octahydro-1*H*-isoindole-1-carboxylate hydrochloride, mp 181–183 °C.

A solution of 6.25 g of this compound was dissolved in 55 mL of 10% HCl and heated at reflux for 4 h. The solvent was removed under reduced pressure and the residue recrystallized from H₂O to give **5a**.

The esters **5b** and **5c** were prepared by hydrogenating the corresponding esters of 2,3-dihydro-1*H*-isoindole-1-carboxylic acid hydrochloride. These in turn were prepared according to ref 21 (see Scheme II) by substituting EtOH or *t*-BuOH for the MeOH used in the esterification of α,α' -dibromo-2-methylbenzeneacetyl chloride. Cyclization with benzylamine gave the 2,3-dihydro-2-(phenylmethyl)-1*H*-isoindole-1-carboxylic acid esters in 35–40% yield. Debonylation was accomplished by hydrogenation with 20% Pd/C in MeOH.

Octahydro-3-oxo-1*H*-isoindole-1-carboxylic Acid (6). A solution of 15 g of ethyl 2,3-dihydro-3-oxo-1*H*-isoindole-1-carboxylic acid ethyl ester,²² **22b** in 110 mL of THF, and 110 mL of absolute EtOH was hydrogenated in a Parr apparatus using 1.0 g of 10% Rh/C as a catalyst. After the required amount of hydrogen had been absorbed, the catalyst was filtered off and the filtrate concentrated under reduced pressure. Recrystallization of the residue from EtOH gave octahydro-3-oxo-1*H*-isoindole-1-carboxylic acid ethyl ester, mp 149–157 °C.

A solution of 5 g of this ester in 10 mL of EtOH and 48 mL of 1 N NaOH was stirred at room temperature overnight. After removal of the EtOH under reduced pressure and adjustment of the pH to 2.0, the crude acid precipitated. Recrystallization from H₂O gave **6**.

Biological Methods. A. In Vitro ACE Inhibition. The in vitro ACE inhibitory activity was determined by a radioassay procedure reported previously.⁷ Activity is reported as the IC₅₀, which is the approximate molar concentration of test compound causing a 50% inhibition of the control converting-enzyme activity. The test solutions were prepared by dissolving 2–5 mg of test compound in 1 mL of Me₂SO and diluting to the desired concentration with a pH 8 buffer of 0.05 M Hepes (Calbiochem), 0.1 M NaCl, and 0.6 M Na₂SO₄ in H₂O.

B. Blood Pressure and Heart Rate Test in the Conscious Rat. Hypertension of renal origin was produced in 4-week-old Sprague–Dawley male albino rats (Charles River; Wilmington, MA) by placing a silver clip (0.2-mm gap) around the left renal artery near the aorta and leaving the contralateral kidney intact. Hypertension was allowed to develop for 4–8 weeks. The rats were then cannulated for blood pressure monitoring as described previously.²³ Only rats with mean aortic blood pressure of > 160 mmHg were used. At the time of cannulation the rats weighed 280–320 g. The rats were given free access to a standard lab chow (5012 Purina; Richmond, IN) and tap water and were maintained on a 12 h dark/12 h light cycle. One minute running average values of heart rate and aortic blood pressure (mean, systolic, and diastolic) for each rat were recorded every 30th min by means of a computer-assisted data capture system.

Acknowledgment. We thank Dr. F. A. MacKellar and associates for the analytical and spectral data, D. Schweiss for the preparation of compound **12**, and Dr. R. Meyer for the preparation of compound **13**. We also thank Prof. A. MacPhail of Duke University for the X-ray crystallographic determinations.

Supplementary Material Available: X-ray data, fractional coordinates and thermal parameters, and interatomic distances and angles for compounds **12** and **10** (19 pages). Ordering information is given on any current masthead page.

(21) Cignarella, G.; Cerri, R.; Grella, G.; Sanna, P. *Gazz. Chim. Ital.* 1976, 106, 65.

(22) Darapsky, A.; Heinrichs, P. *J. Prakt. Chem.* 1936, 146, 307.

(23) Smith, R. D.; Wood, T. J.; Tessman, D. K.; Olszewski, B. J.; Currier, G.; Kaplan, H. R. *DHEW Publ. (NIH) (U.S.)* 1980, 41, 1473.