

This article was downloaded by: [Dalhousie University]

On: 27 November 2012, At: 04:48

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uopp20>

AN EFFICIENT SYNTHESIS OF 3, 5-bis(2-CYANOISOPROPYL)TOLUENE

Yu-Hua Mei ^a, Yu Luo ^a & Wei Lu ^a

^a Institute of Medicinal Chemistry, Department of Chemistry, East China Normal University, Shanghai, 200062, PR CHINA E-mail:

Version of record first published: 18 Feb 2009.

To cite this article: Yu-Hua Mei, Yu Luo & Wei Lu (2008): AN EFFICIENT SYNTHESIS OF 3, 5-bis(2-CYANOISOPROPYL)TOLUENE, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 40:5, 487-489

To link to this article: <http://dx.doi.org/10.1080/00304940809458112>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.


10. (a) S. Kenso, W. Masami and K. Masashi, *J. Chem. Soc., Chem. Commun.*, **9**, 534 (1989).
 (b) S. Kenso, W. Masami and K. Masashi, *Bull. Chem. Soc. Jpn*, **62**, 2124 (1989).
11. W. J. Houlihan, US Patent 3898275 (1975); *Chem. Abstr.*, **83**,147305a (1975).
12. D. L. Lee (Martinez, CA), US Patent 4741769 (1988); *Chem. Abstr.*, **109**,144620u (1988).

AN EFFICIENT SYNTHESIS OF 3,5-bis(2-CYANOISOPROPYL)TOLUENE


Submitted by Yu-hua Mei, Yu Luo and Wei Lu*
(01/17/08)

*Institute of Medicinal Chemistry, Department of Chemistry,
East China Normal University, Shanghai 200062, P. R. CHINA
e-mail: wlu@chem.ecnu.edu.cn*

3,5-bis(2-Cyanoisopropyl)toluene (**4**) is a key intermediate for the preparation of *anastrozole*, an anti-tumor drug for the treatment of breast cancer. So far, the only synthetic route to **4** involves three steps from mesitylene (*Scheme 1*).^{1,2} However, this patented approach suffers from low yield, environmentally unfriendly chemicals (CCl₄ and benzoyl peroxide) and high cost (expensive iodomethane).



As a part of our exploration of a facile synthesis of *anastrozole*, we have developed an efficient preparation of 3,5-bis(2-cyanoisopropyl)toluene (**4**) from commercially available 5-methylisophthalic acid (**5**) (*Scheme 2*).



a) H_2SO_4 , CH_3OH , reflux, 90%; b) KBH_4 , LiCl , THF , 75%; c) PBr_3 , CH_2Cl_2 , 90%;
 d) KCN , TBAB , CH_2Cl_2 , H_2O 78%; e) NaH , DMF , methyl *p*-toluenesulfonate, 86%

Scheme 2

Reduction of **6**, obtained from compound **5** via esterification in methanol, gave diol **7** in 75% yield.³ Treatment of **7** with PBr_3 afforded **2** in 90% yield.^{4,5} Reaction of **2** with KCN proceeded smoothly to give compound **3** in 78% yield. The final step was modified slightly, using methyl *p*-toluenesulfonate instead of expensive iodomethane as the methylating reagent. After crystallization from ethanol, 3,5-bis(2-cyanoisopropyl)toluene (**4**) was obtained in 86% yield, in better than 99% purity.

In conclusion, we have developed an efficient synthesis of 3,5-bis(2-cyanoisopropyl)toluene (**4**) in 41% overall yield. The advantages of this procedure include mild conditions, simple work-up procedures and avoidance of toxic solvent and expensive reagent.

EXPERIMENTAL SECTION

Melting points were determined on a Büchi 510 melting point apparatus. ^1H NMR spectra were recorded on a Bruker DRX-500 (500 MHz). ^{13}C NMR spectra were obtained on a JNM-EX100 (400 MHz). Mass spectra (MS) were determined on a Finnigan MAT-95 mass spectrometer.

Methyl 5-Methyl-1,3-benzenedicarboxylate (6).- A solution of 5-methylisophthalic acid (**5**) (180 g, 1.0 mol), sulfuric acid (8 mL) in methanol (400 mL) was refluxed for 16 h. The mixture was then evaporated to remove methanol. The residue was poured into 5% aqueous Na_2CO_3 (300 mL) to precipitate a white solid, that was collected, washed with water, dried with Na_2SO_4 , and evaporated to give compound **6** (186 g, 90% yield), mp. 96–98°C. ^1H NMR (500 MHz, CDCl_3): δ 2.45 (s, 3H), 3.99 (s, 6H), 8.04 (s, 2H), 8.48 (s, 1H). ^{13}C NMR (100 MHz, CDCl_3): δ 21.1, 52.3, 127.9, 130.5, 134.4, 138.6, 166.4. MS (EI): m/z = 208.

Anal. Calcd. for $\text{C}_{11}\text{H}_{12}\text{O}_4$: C, 63.45; H, 5.81. Found: C, 63.36; H, 5.73

3,5-bis(Hydroxymethyl)toluene (7).- A mixture of **6** (20.8 g, 0.10 mol), KBH_4 (37.1 g, 0.70 mol), LiCl (29.7 g, 0.70 mol) in THF (100 mL) was heated to reflux for 6 h, then cooled to room temperature and quenched with saturated NH_4Cl solution (40 mL). After stirring for 1 h, the mixture was filtered to remove insoluble solids. Then, the filtrate was extracted with ethyl

acetate, washed with brine, dried with MgSO_4 , and concentrated to give **7** as a white solid (11.4 g, 75% yield), mp. 41-43°C. ^1H NMR (500 MHz, CDCl_3): δ 2.27 (s, 3H), 4.35 (s, 2H), 4.41 (s, 4H), 6.93 (s, 2H), 7.00 (s, 1H). ^{13}C NMR (100 MHz, CDCl_3): δ 21.1, 64.4, 122.5, 126.7, 137.9, 140.9. MS (EI): m/z = 152.

Anal. Calcd. for $\text{C}_9\text{H}_{12}\text{O}_2$: C, 71.03; H, 7.95. Found: C, 70.76; H, 7.94.

3,5-bis(Bromomethyl)toluene (2).- To a solution of compound **7** (11.46 g, 75 mmol) in CH_2Cl_2 (65 mL) at 0°C, was added PBr_3 dropwise over a period of 30 min. Then the mixture was stirred for 4 h at room temperature and poured into ice water (100 mL). The solution was extracted with CH_2Cl_2 , washed with brine, dried with Na_2SO_4 , and evaporated to dryness to give solid **3** (18.6 g, 89% yield), mp. 64-66°C, (*lit.*² mp. 66°C). ^1H NMR (500 MHz, CDCl_3): δ 2.34 (s, 3H), 4.44 (s, 4H), 7.14 (s, 2H), 7.26 (s, 1H). ^{13}C NMR (100 MHz, CDCl_3): δ 21.1, 32.9, 126.7, 129.8, 138.2, 139.2. MS (EI): m/z = 278. HPLC: 99.7%.

3,5-bis(2-Cyanoisopropyl)toluene (4).- To a mixture of NaH (6.32 g, 160 mmol) in DMF (30 mL) under a nitrogen atmosphere at 0°C was added a solution of compound **4** (6 g, 35 mmol) and methyl *p*-toluenesulfonate (26.4 g, 140 mmol) in DMF (30 mL). Then, the mixture was stirred for 30 min, poured into ice water (100 mL) to precipitate a white solid, which was collected, washed with water and dried to give crude **4**. Crystallization from ethanol gave pure solid **4** (6.8 g, 86% yield), mp. 126-127°C, (*lit.*¹ mp. 126-127°C), purity > 99% (HPLC analysis). ^1H NMR (500 MHz, CDCl_3): δ 1.73 (s, 12H), 2.40 (s, 3H), 7.26 (s, 2H), 7.32 (s, 1H). ^{13}C NMR (100 MHz, CDCl_3): δ 21.5, 28.9, 29.0, 37.13, 118.6, 124.2, 125.4, 139.5, 142.1. MS (EI): m/z = 226.

Acknowledgement.- We gratefully acknowledge the financial support for New Century Excellent Talents in university (NCET).

REFERENCES

1. P. N. Edwards and M. S. Large, EP0296749, 28 Dec 1988; *Chem. Abstr.*, **111**, 7410 (1989).
2. Z. M. Ge, J. L. Cui, T. M. Cheng and R. T. Li, *Chin. J. Med. Chem.*, **13**, 146 (2003); *Chem. Abstr.*, **142**, 397462 (2005).
3. B. Núbia and C. S. Jorge, *Tetrahedron Lett.*, **45**, 6021 (2004).
4. C. R. Noller, R. Dinsmore, D. E. Badertscher and A. R. Lux, *Org. Synth., Coll. Vol.* **2**, 358 (1943).
5. L. H. Smith, *Org. Synth., Coll. Vol.* **3**, 793 (1955).