


Reduction of  $\beta$ -Hydroxyketones by  
 $\text{SmI}_2/\text{H}_2\text{O}/\text{Et}_3\text{N}$ Todd A. Davis,<sup>†</sup> Pramod R. Chopade,<sup>†</sup> Göran Hilmersson,<sup>‡</sup> and  
Robert A. Flowers, II<sup>\*,§</sup>Department of Chemistry & Biochemistry, Texas Tech University,  
Lubbock, Texas 79409, Department of Chemistry, Göteborg University,  
SE-412 96 Göteborg, Sweden, and Department of Chemistry, Lehigh University,  
Bethlehem, Pennsylvania 18015

rof2@lehigh.edu

Received October 20, 2004

## ABSTRACT



Reduction of a series of  $\beta$ -hydroxyketones by  $\text{SmI}_2/\text{H}_2\text{O}/\text{Et}_3\text{N}$  provided 1,3-diols in quantitative yields. The reactions were exceedingly clean with no byproduct formation, negating the need for further purification. Most reactions provided moderate to excellent diastereoselectivity with *syn*-diols as the major isomer in most instances.


The pioneering discovery by Inanaga<sup>1</sup> describing the use of HMPA in accelerating  $\text{SmI}_2$ -mediated reactions has resulted in widespread applications of the  $\text{SmI}_2$ –HMPA mixture. This combination is utilized in several reduction and reductive coupling reactions. Due to its ability to enhance reaction outcomes, HMPA is the preferred cosolvent in  $\text{SmI}_2$ -mediated reactions. There is, however, one serious drawback to the use of HMPA. It is a suspected human carcinogen and, therefore, extreme caution needs to be exercised during its use. Several nontoxic alternatives to HMPA have been reported recently. Cabri<sup>2</sup> and co-workers have explored the use of 1,1,3,3-tetramethylguanidine (TMG), 1,1,3,3-tetramethylurea (TMU) and  $\text{Et}_3\text{N}$  as cosolvents in halide-olefin coupling reactions. Curran<sup>3</sup> has reported the effects of water and DMPU in the reduction of ketones. These additives have proven to be useful in several reactions but unfortunately do not have the broad applicability of HMPA, and as a consequence, the search for an alternative is ongoing.

In this regard, the work reported by Hilmersson<sup>4</sup> and co-workers is of considerable significance. They have reported

the use of an  $\text{SmI}_2/\text{H}_2\text{O}/\text{Et}_3\text{N}$  mixture in the reduction of ketones. These reactions are instantaneous and provide excellent yields of reduced products. A comparison of the  $\text{H}_2\text{O}/\text{Et}_3\text{N}$  method and the HMPA/alcohol method in reduction of ketones indicates that  $\text{H}_2\text{O}/\text{Et}_3\text{N}$  is approximately 100 times faster. This method has also been applied in the reduction of  $\alpha,\beta$ -unsaturated esters, imines, and conjugated olefins with excellent results.<sup>5,6</sup> These examples clearly show the utility of  $\text{SmI}_2/\text{H}_2\text{O}/\text{amine}$  mixtures in reduction of several functional groups. Moreover, they provide better yields and require much less time than the HMPA/alcohol systems. The workup and subsequent purification of products are straightforward since the byproducts precipitate during the course of the reaction. Therefore, the combination of amine and  $\text{H}_2\text{O}$  provides an excellent alternative to HMPA in  $\text{SmI}_2$ -based reactions.

Initial mechanistic studies show that water and  $\text{Et}_3\text{N}$  do not accelerate the reactions separately. The acceleration is a result of the  $\text{Et}_3\text{N}$ – $\text{H}_2\text{O}$  mixture. Other amines such as  $N,N,N',N'$ -tetramethylethylenediamine (TMEDA) and  $N,N,N',N'',N''$ -pentamethyldiethylenetriamine (PMDTA) have the same effect as  $\text{Et}_3\text{N}$ , while replacement of water by alcohols has a deleterious impact on the rates of reduction. It has been

<sup>†</sup> Texas Tech University.<sup>‡</sup> Göteborg University.<sup>§</sup> Lehigh University.(1) Inanaga, J.; Ishikawa, M.; Yamaguchi, M. *Chem. Lett.* **1987**, 1485.(2) Cabri, W.; Candiani, I.; Colombo, M.; Franzoi, L.; Bedeschi, A. *Tetrahedron Lett.* **1995**, 36, 949.(3) Hasegawa, E.; Curran, D. P. *J. Org. Chem.* **1993**, 58, 5008.(4) Dahlen, A.; Hilmersson, G. *Tetrahedron Lett.* **2002**, 43, 7197.(5) Dahlen, A.; Hilmersson, G. *Tetrahedron Lett.* **2003**, 44, 2661.(6) Dahlen, A.; Hilmersson, G. *Chem. Eur. J.* **2003**, 9, 1123.

**Table 1.** Reduction of  $\beta$ -Hydroxyketones by  $\text{SmI}_2/\text{H}_2\text{O}/\text{Et}_3\text{N}$ 

entry	R <sub>1</sub>	R <sub>2</sub>	ds (syn:anti) <sup>a</sup>
1	<i>t</i> -Bu	Me	12:1
2	<i>t</i> -Bu	Et	29:1
3	<i>t</i> -Bu	<i>i</i> -Pr	99:1
4	<i>t</i> -Bu	<i>t</i> -Bu	4:1
5	<i>t</i> -Bu	Ph	1.5:1
6	Ph	Me	99:1
7	Ph	Et	>99:1
8	Ph	<i>i</i> -Pr	>99:1
9	Ph	<i>t</i> -Bu	1:41
10	Ph	Ph	2.6:1
11	Me	<i>t</i> -Bu	99:1
12	Me	Ph	1:2

<sup>a</sup> Substrate ( $\beta$ -hydroxyketone, 1 mmol) was placed in a flame-dried round-bottom flask and dissolved in 10 mL of anhydrous THF, and the solution was cooled to 0 °C. A mixture of  $\text{SmI}_2$  (2.5 equiv) in THF, triethylamine (5 equiv), and deaerated water (6.25 equiv) was added dropwise to the reaction mixture. The reaction mixture was allowed to stir at 0 °C until the color of the reaction changed from blue to yellow.

proposed that rapid precipitation of  $\text{Sm}(\text{OH})_3$  and a tertiary ammonium salt,  $\text{Et}_3\text{NH}^+ \text{I}^-$ , provides the driving force for the reduction.<sup>6</sup>

To expand the applicability of the  $\text{SmI}_2/\text{H}_2\text{O}/\text{Et}_3\text{N}$  reagent and to determine its general utility in important single-electron-transfer-promoted reactions, the reduction of  $\beta$ -hydroxyketones to 1,3-diols was studied. The seminal work of Keck has shown that the reduction of  $\beta$ -hydroxyketones is sensitive to proton donor source and temperature.<sup>7</sup> Recent work in our laboratory has shown that solvation also plays an important role in determining the stereochemical outcome of these reductions.<sup>8</sup> A limited survey of the effects of substitution and solvent on reduction of  $\beta$ -hydroxyketones in three solvents showed that, in most cases, DME provided superior diastereoselectivities over THF (both solvents leading to the syn product), whereas reductions in  $\text{CH}_3\text{CN}$  led to *anti*-1,3-diols. Since the nature of substitution in substrates is known to influence outcomes in  $\text{SmI}_2/\text{H}_2\text{O}/\text{Et}_3\text{N}$ -mediated reactions,<sup>6</sup> a series of  $\beta$ -hydroxyketones were synthesized to examine the role of substituents in detail.

Table 1 contains the series of  $\beta$ -hydroxyketones examined in this study. The  $\beta$ -hydroxyketones were synthesized by the aldol reaction of ketones and aldehydes or the L-proline-catalyzed asymmetric aldol reaction.<sup>9</sup> Substrates were subjected to reduction using  $\text{SmI}_2/\text{H}_2\text{O}/\text{Et}_3\text{N}$ . These reactions were performed at 0 °C in an inert atmosphere of  $\text{N}_2$ , and all reactions were completed within 5 min of addition of the

$\text{Sm}(\text{II})$  reductant. The diastereoselectivity was determined by gas chromatography, and the stereochemistry of the 1,3-diols was determined utilizing the protocol described by Rychnovsky.<sup>10</sup> The results are shown in Table 1.

All reactions were quantitative, and the precipitation of byproducts  $\text{Sm}(\text{OH})_3$  and  $\text{HNEt}_3^+ \text{I}^-$  made purification quite simple. Filtration of the precipitate and rotary evaporation of solvent provided clean product, and no further purification was necessary. Inspection of the results in Table 1 shows a number of interesting trends. Most reactions provided the syn diastereomer predominantly, although two cases (entries 9 and 12) provided the anti diastereomer as the major product. When  $\text{R}_2$  was Ph, the diastereoselectivity decreased dramatically, whereas Me, Et, and *i*-Pr substituents resulted in good to excellent diastereoselectivity.

Comparison with other reported protocols using  $\text{SmI}_2$  reveals some interesting trends and differences among  $\text{SmI}_2$ -based approaches to the reduction of  $\beta$ -hydroxyketones. When  $\text{R}_1 = t\text{-Bu}$  and  $\text{R}_2 = \text{Ph}$  (entry 5, Table 1), poor diastereoselectivities were obtained using  $\text{SmI}_2/\text{H}_2\text{O}/\text{Et}_3\text{N}$  and  $\text{SmI}_2/\text{MeOH}$  in THF<sup>7</sup> or DME<sup>8</sup> suggesting that this substitution pattern may be inherently difficult to reduce diastereoselectively. When  $\text{R}_1 = \text{Me}$  and  $\text{R}_2 = \text{Ph}$  (entry 12, Table 1), excellent diastereoselectivities providing the anti diastereomer were obtained with  $\text{SmI}_2/\text{MeOH}$  in THF and DME, whereas reduction by  $\text{SmI}_2/\text{H}_2\text{O}/\text{Et}_3\text{N}$  provided only modest selectivity for the anti diastereomer. Substrates in entries 1, 6, and 11 of Table 1 are reduced with greater diastereoselectivity by  $\text{SmI}_2/\text{H}_2\text{O}/\text{Et}_3\text{N}$  than  $\text{SmI}_2$  in THF/MeOH.

While ease of reduction, workup, and in some cases higher diastereoselectivities are obtained in reductions of  $\beta$ -hydroxyketones with  $\text{SmI}_2/\text{H}_2\text{O}/\text{Et}_3\text{N}$  in THF compared to  $\text{SmI}_2/\text{MeOH}$ , it is important to assess various mechanistic scenarios responsible for reaction outcomes so that practitioners can make judicious choices best suited to their system of interest. The reduction of ketones by  $\text{SmI}_2$  in the presence of proton sources likely proceeds through a House-type mechanism,<sup>11</sup> and recent mechanistic work has shown that the rate-limiting step is the first proton transfer to the initially formed ketyl radical anion.<sup>12,13</sup> While reduction of  $\beta$ -hydroxyketones may be somewhat more complex since the presence of an internal proton source ( $\beta$ -OH) may also act as a donor, initial rate studies are also consistent with the initial proton transfer to the ketyl radical anion as the rate-limiting step.<sup>14</sup> The radical produced after protonation of the ketyl is reduced to a carbanion by a second equivalent of  $\text{Sm}(\text{II})$ . The stereo-defining step in the present reductions is likely to be the final proton transfer to the carbanion.

There is strong synthetic and mechanistic evidence to support the presence of chelation along the reaction coordinate of  $\text{Sm}(\text{II})$ -mediated reduction and reductive coupling

(10) Rychnovsky, S. D.; Yang, G.; Powers, J. P. *J. Org. Chem.* **1993**, *58*, 5251.

(11) House, H. O. In *Modern Synthetic Reactions*, 2nd ed.; W. A. Benjamin: Menlo Park, CA, 1972.

(12) Dahlén, A.; Hilmersson, G. *Tetrahedron Lett.* **2001**, *42*, 5565.


(13) Chopade, P. R.; Prasad, E.; Flowers, R. A., II. *J. Am. Chem. Soc.* **2004**, *126*, 44.

(14) Chopade, P. R. Ph.D. Dissertation, Texas Tech University, Lubbock, TX, 2004.

(7) Keck, G. E.; Wager, C. A.; Sell, T.; Wager, T. T. *J. Org. Chem.* **1999**, *64*, 2172.

(8) Chopade, P. R.; Davis, T. A.; Prasad, E.; Flowers, R. A., II. *Org. Lett.* **2004**, *6*, 2685.

(9) List, B.; Lerner, R. A.; Barbas, C. F., III. *J. Am. Chem. Soc.* **2000**, *122*, 239.




**Figure 1.** Electrostatic interaction between lone pairs on carbanion intermediate and neighboring oxygen.

reactions of  $\beta$ -substituted ketones, and this is likely to be important in the reductions described above.<sup>7,15–17</sup> The majority of the reactions examined in the present case provide the *syn* diastereomer as the major product. After analysis of numerous chelation models, the results of the reductions described herein are consistent with the importance of electrostatic interactions in defining the final stereochemistry of the 1,3-diol product (Figure 1). Reaction of a  $\beta$ -hydroxyketone with  $\text{SmI}_2/\text{H}_2\text{O}/\text{Et}_3\text{N}$  through two electron transfers (from 2 equiv of  $\text{SmI}_2$ ) and one proton transfer (from  $\text{H}_2\text{O}$ ) leads to two potential intermediates shown as **A** and **B** in Figure 1. Protonation of **A** leads to the formation of the *syn*-1,3-diol, whereas protonation of **B** leads to the formation of the *anti*-1,3-diol. In **A**, the electron lone pair of the oxygen and the lone pair of the adjacent carbanion are nearly antiperiplanar and as a result the electrostatic repulsion between them is not significant. Conversely, in **B**, the lone pairs are only separated by approximately  $60^\circ$ , leading to increased electrostatic repulsion. On the basis of this analysis, the transition state (TS) from **A** is of lower energy than the TS from **B**.

A different explanation for the *syn* stereoselectivity is possible as well. The axially oriented radical precursor of **A** is expected to be stabilized due to favorable overlap of the singly occupied orbital and the nonbonding orbital on the neighboring oxygen. Upon reduction of the radical, coplanarity of the nonbonding orbitals on the carbon and neighboring oxygen, while Coulombically favorable, leads to an unfavorable orbital interaction.<sup>18</sup> This intermediate (resembling **A**) could be protonated quickly by water in the

system before configurational change takes place at the carbon center. Alternatively, distortion of the intermediate due to the interaction between the lone pair of the carbanion and the samarium (as shown for **A** in Figure 1) could move the nonbonding orbitals on carbon and oxygen out of plane, providing stabilization as well.

The mechanistic scenarios described above do not explain the change in diastereoselectivity when  $\text{R}_2 = \text{Ph}$ . After careful analysis, it is our hypothesis that the electron-deficient  $\text{Sm}^{3+}$  cation could possibly interact with the  $\pi$ -system of the benzene ring (Figure 2). Since this interaction is only



**Figure 2.** Depiction of possible intermediate leading to poor stereoselectivity or reversal of stereoselectivity when  $\text{R}_2 = \text{Ph}$ .

possible in **B**, it could act to stabilize **B** depending on the substituent at  $\text{R}_1$  and as a result provide an alternative

(15) Molander, G. A.; Harris, C. R. *J. Org. Chem.* **1998**, *63*, 812.

(16) Kawatsura, M.; Hosaka, K.; Matsuda, F.; Shirahama, H. *Synlett* **1995**, 729.

(17) Prasad, E.; Flowers, R. A., II. *J. Am. Chem. Soc.* **2002**, *124*, 6357.

(18) Skrydstrup, T.; Jarretton, O.; Mazéas, D.; Urban, D.; Beau, J.-M. *Chem. Eur. J.* **1998**, *4*, 655.

mechanistic (stereodifferentiating) pathway for the reduction. Additional detailed mechanistic studies on a variety of substrates will be necessary to further test this hypothesis.

In conclusion, reduction of a series of  $\beta$ -hydroxyketones by  $\text{SmI}_2/\text{H}_2\text{O}/\text{Et}_3\text{N}$  in THF provided good to excellent selectivity for the *syn*-diol with a wide range of substrates. All reductions provided quantitative yields and were completed within a few minutes. Furthermore, precipitation of byproducts eliminates the need for further purification, thus simplifying the workup compared to standard  $\text{Sm}(\text{II})$ -based reductions. Analysis of various transition state models proceeding through chelation control are consistent with the importance of electrostatic interactions controlling the diastereoselectivity of the 1,3-diol in most instances. Regardless of the exact mechanistic details of the present reductions, the data presented herein show the utility and ease of the  $\text{SmI}_2/\text{H}_2\text{O}/\text{Et}_3\text{N}$  reducing system in the reduction of  $\beta$ -hy-

droxyketones. Examination of the use of this novel reducing system in reductive coupling reactions is currently underway, and the results of these studies will be reported in due course.

**Acknowledgment.** R.A.F. is grateful to the National Science Foundation (CHE-0413845 and CHE-0423972) for support of this work. G.H. is grateful to the Swedish Research Council (2001-1713 and 2003-2442) for financial support. We thank Professor David Birney for many fruitful discussions related to this work. We also thank reviewers A and B for their insightful comments.

**Supporting Information Available:** General methods, experimental protocols, and spectroscopic data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL047835P