

Design and Synthesis of New Naphthalenic Derivatives as Ligands for 2-[¹²⁵I]Iodomelatonin Binding Sites

Michel Langlois,*[†] Béatrice Brémont,[†] Shuren Shen,[†] Annie Poncet,[†] Jean Andrieux,[†] Sames Sicsic,[†] Isabelle Serraz,[†] Monique Mathé-Allainmat,[†] Pierre Renard,[‡] and Philippe Delagrangé[§]


CNRS-BIOCIS, Faculté de Pharmacie, 5 rue J.B. Clément, 92296 Châtenay-Malabry, France, ADIR, 1 rue C. Hebert, 92415 Courbevoie, France, and Institut de Recherche Internationale Servier, Place des Pleiades, 92415 Courbevoie, France

Received September 14, 1994[®]

New melatonin-like agents were designed from the frameworks of 2,5-dimethoxyphenethylamine, an important structural moiety for the 5-HT receptor, and (2-methoxynaphthyl)ethylamine. The compounds were synthesized by classical methods and evaluated in binding assays with chicken brain membranes using 2-[¹²⁵I]iodomelatonin as the radioligand. Preliminary studies on the series of *N*-acyl-disubstituted phenethylamines showed the favorable role of the methoxy group in the ortho position of the side chain on the affinity for the receptor ($K_i = 8 \pm 0.2$ nM) for *N*-[2-(2-methoxy-5-bromophenyl)ethyl]propionamide (**3o**). This effect was confirmed in a series of the naphthalene derivatives, a bioisosteric moiety of the indole ring, and several potent ligands for melatonin binding sites were prepared such as *N*-[2-(2-methoxynaphthyl)ethyl]propionamide (**4b**) ($K_i = 0.67 \pm 0.05$ nM) and *N*-[2-(2,7-dimethoxynaphthyl)ethyl]cyclopropylformamide ($K_i = 0.05 \pm 0.004$ nM) (**4k**). Structure–activity relationships are discussed with regard to melatonin and bioisosteric naphthalenic compound **2**. The K_i value for **4b** was affected to a similar extent to that of melatonin by GTP- γ -S or Mn²⁺ in competition experiments, suggesting an agonist profile for this compound.

Introduction

There is no longer any doubt that melatonin (**1**) is a pineal hormone which modulates a variety of endocrinological, neurophysiological, and behavioral functions in vertebrates.¹ It regulates reproduction in photoperiodic species² but is also implicated in circadian rhythms and plays a key role in a number of disorders such as anxiety, seasonal depression, and delayed sleep phase syndrome.^{3,4} More recently, the antitumoral properties of melatonin and its implication in immune system responsiveness have been described.⁵ Furthermore, the hydroxyl radical scavenger properties of melatonin have been demonstrated, and a role for this hormone in the aging process has been suggested.⁶ The synthesis of melatonin, which follows a circadian rhythm, occurs in the pinealocytes of the pineal gland and in the retina, and the maximum level is reached during the night.⁷ However, the pattern of the nocturnal rise depends upon the animal species. Therefore, melatonin has been described as “the chemical expression of darkness”.⁸ It



is synthesized from serotonin (5-HT) in two enzymatic steps: 5-HT is *N*-acetylated by *N*-acetyltransferase (NAT) to produce *N*-acetylserotonin and subsequently *O*-methylated to give melatonin.⁹ Melatonin is highly

lipophilic, can diffuse out of the pineal gland, and is then released into the blood. Its effects seem to be mediated through membrane receptors¹⁰ located in different regions of the brain, particularly in the suprachiasmatic nucleus (SCN) and the pars tuberalis where it regulates reproductive function¹¹ in photoperiodic species. However, the distribution of melatonin receptors varies greatly among mammals, and contradictory data have been reported.¹² The receptor is doubtless G-protein-coupled since several experiments have demonstrated inhibition of the high-affinity specific binding of 2-[¹²⁵I]iodomelatonin by guanine nucleotides in cerebral tissues of several species and a dose-dependent inhibition of forskolin-stimulated cAMP formation by melatonin.¹³

The design and availability of potent, specific agonists or antagonists for melatonin receptors are important to elucidate totally its role in the physiological events in which it is implicated. These compounds could find therapeutic applications in a number of disorders such as jet lag,¹⁴ anxiety,¹⁵ depression, or disturbances of the circadian rhythms.¹⁶

Several indolic analogues of melatonin have been synthesized as agonists,¹⁷ and structure–activity studies have shown the favorable role of 2-halogeno substitution and enabled the development of 2-[¹²⁵I]iodomelatonin as a radioligand for binding studies and autoradiography.¹⁸ Recently, several papers reported the synthesis and agonist properties of several nonindolic compounds. One group¹⁹ designed compounds such as **2** using the bioisosteric properties of naphthalene with regard to the indole ring, while another group²⁰ used the analogy between 2-amido-8-methoxytetralin and the pharmacophore groups of the melatonin moiety.


We present here our initial results on the design of new nonindolic melatonin-like compounds, emphasizing particularly the importance of the presence of the methoxy group in the ortho position of the ethylaceta-

[†] CNRS-BIOCIS.

[‡] ADIR.


[§] Institut de Recherche Internationale Servier.

[®] Abstract published in *Advance ACS Abstracts*, May 1, 1995.



mido chain to obtain a compound with marked affinity for melatonin receptors. They were designed from the framework of serotonergic agonists because of the structural analogy between melatonin and serotonin. The recent demonstration of 2-amido-8-methoxytetralin²⁰ as a melatonin agonist derived from 8-OH-DPAT, a potent 5-HT_{1A} agonist, agrees with this hypothesis.


As a preliminary approach, we used 2,5-dimethoxyphenethylamine, an important structural base for 5-HT₂ receptor agonists,²¹ to prepare amido derivatives **3** (R₁ = 5-MeO, R₂ = MeO). They were evaluated in binding assays (Table 1) with 2-[¹²⁵I]iodomelatonin radioligand using chicken brain membranes according to the method already reported for pigeon brain,²² and their affinities were compared to those of several *N*-acetyl-monosubstituted phenethylamines **3** (R₂ = H). The favorable role of the (2-methoxyphenethyl)amido moiety having been demonstrated by these preliminary structure-activity studies, we then investigated the preparation of the corresponding naphthalene derivatives **4**, a more suitable aromatic system for the design of indolic bioisosteres.



Chemistry

Compounds **3** and **4** (Tables 1 and 2) were obtained by acylation reactions of the corresponding amines which were commercially available or synthesized by classical methods. The amines were prepared according to the routes shown in Scheme 1. **14** and **16–18** were prepared by LiAlH₄ reduction of the nitroethylene derivatives **9–12** obtained through the Knoevenagel condensation of the corresponding formyl derivatives with nitromethane or nitroethane (method A).^{17b,23} Amines **13** and **15** were synthesized through the cyano derivatives **7** and **8** which were hydrogenated in the presence of Raney Ni (method B). 5-Ethylaldehyde (**5**) and 2,7-dimethoxy-1-naphthaldehyde (**6**), the intermediates for the synthesis of **14** and **18**, respectively, were prepared by a formylation reaction with dichloromethyl methyl ether²⁴ in the presence of TiCl₄, and the cyano derivatives **7** and **8** were synthesized routinely from 2-methoxy-5-methylbenzyl alcohol and 2-methoxy-5-bromobenzyl alcohol which were transformed into the corresponding chlorides to react with KCN in DMSO.

The amides **3a–o** and **4a–e,g,i–k** (Tables 1 and 2) were prepared by reaction of the amines with the acyl chlorides in CH₂Cl₂ in the presence of Et₃N (method C) or reaction of the acetic anhydride under Schotten–



13 R = Me

14 R = Et

15 R = Br

16 R₁ = H R₂ = H

17 R₁ = Me R₂ = H

18 R₁ = H R₂ = OMe


Baumann conditions (method D), while the trifluoroacetyl derivatives **4f,h,l** were synthesized directly with trifluoroacetic anhydride in CH₂Cl₂ in the presence of pyridine (method E).

Biochemistry

The affinities values (*K_i*) of the amido derivatives **3** and **4** for melatonin binding sites are shown in Tables 1 and 2, respectively. They were evaluated *in vitro* by binding assays²² using 2-[¹²⁵I]iodomelatonin and chicken brain membranes. Specific binding of the ligand was stable, saturable, and of high affinity (*K_d* = 0.05 nM), and nonspecific binding was less than 10% of the total binding. Scatchard analyses gave linear plots, and the values of the Hill coefficients were close to 1, suggesting a single class of binding sites for the ligand. *B_{max}* values for specific 2-[¹²⁵I]iodomelatonin binding sites in preparations from chickens killed at 12 a.m. and 6 p.m. were about 20% higher than the number of binding sites at 12 p.m. (Figure 1). However, no significant variations in the binding affinities were observed at the different time points. These data suggest that the *in vivo* rhythm of the melatonergic system is regulated by variations in the number of binding sites available rather than by an increase in their affinity.²⁵ In the inhibition studies with the new compounds described above, brain homogenates from the chicken killed at 12 a.m. were used and incubated at 25 °C for 60 min with the compounds under investigation. The affinity of melatonin was assessed in this preparation by competition experiments and agreed (*K_i* = 0.64 nM) with the value already reported.²⁰


Discussion

The preliminary approach, which consisted of using the hypothetical analogy between the structural elements of the receptor binding sites of serotonin and melatonin, prompted us to prepare amido derivatives of 2,5-dimethoxyphenethylamine, an important framework in the serotonergic field. The relatively good affinity of compound **3h** (*K_i* = 69.9 nM), obtained in competition studies with 2-[¹²⁵I]iodomelatonin, confirmed the soundness of this hypothesis. In our hands, this derivative was more potent than 2-acetamido-8-methoxytetralin (*K_i* = 206 nM), another template used recently by Copinga²⁰ to design new melatonergic ligands. Structural variations of the amido group of compound **3h** brought about a clear increase in the affinity of the propionyl derivative **3i**, but a marked drop was observed with cyclopropyl and butyl substitutions (compounds **3j,k**). These results differed from those obtained by Guardiola¹⁹ in a series of naphthalenic derivatives where the optimum binding affinity was attained with propyl, butyl, or cyclopropyl chains in

Scheme 1^a

^a (a) MeNO₂ or EtNO₂, AcONH₄; (b) SOCl₂, pyridine, toluene, KCN, DMSO, rt; (c) LiAlH₄, THF, 40 °C, 24 h; (d) NH₄OH, H₂, Ra Ni, EtOH; (e) RCOCl, CH₂Cl₂, Et₃N; (f) (Ac)₂O, AcONa, H₂O; (g) (CF₃CO)₂O, CH₂Cl₂, pyridine.

Table 1. Displacement of 2-[¹²⁵I]Iodomelatonin Binding from Melatonin Receptors in Chicken Brain Membranes by the Compounds **3**




compd	R ₁	R ₂	R	K _i ± SEM, nM ^a
3a	4-Cl	H	CH ₃	>1000
3b	3-Cl	H	CH ₃	>1000
3c	2-Cl	H	CH ₃	>1000
3d	3-F	H	CH ₃	>1000
3e	4-MeO	H	CH ₃	>1000
3f	3-MeO	H	CH ₃	253 ± 57
3g	H	MeO	CH ₃	213 ± 23
3h	5-MeO	MeO	CH ₃	69.9 ± 5.1
3i	5-MeO	MeO	C ₂ H ₅	17.5 ± 3.2
3j	5-MeO	MeO	c-C ₃ H ₅	386 ± 28
3k	5-MeO	MeO	n-C ₄ H ₉	>1000
3l	5-CH ₃	MeO	CH ₃	192 ± 23
3m	5-C ₂ H ₅	MeO	CH ₃	217 ± 23
3n	5-Br	MeO	CH ₃	42 ± 2.8
3o	5-Br	MeO	C ₂ H ₅	8 ± 0.2
melatonin				0.67 ± 0.04
2-acetamido-8-methoxytetralin				206 ± 20

^a K_i values are expressed in nM ± the standard error of the mean (SEM) and were calculated using the Cheng-Prusoff equation from IC₅₀ values obtained from competition curves; the data are the results of one or two separate determinations.

binding assays using the ovine pars tuberalis. In particular, the affinity of the latter substituent was increased by 2 orders of magnitude with regard to the parent compound. The influence of the position of the methoxy group on the phenyl ring was evaluated with the simple derivatives **3e-g** which emphasized the prevalence of the meta or ortho positions over the para position, and these results agree with a recent report.²⁰ Although these data concern compounds with a moderate level of activity, they illustrate, compared to the results obtained for the other monosubstituted compounds (**3a-d**), the superiority of the methoxy substitution for recognition by the melatonin receptor site. At this stage of the study, it was postulated that the relatively good affinity of compound **3i** for melatonin receptors was due to the melatonin-like position of the 5-methoxy group of this compound, while the 2-methoxy group could mimic the pyrrole ring of the indole. The contribution of the 5-methoxy group to the improved affinity of compound **3i** for melatonin receptors, and in particular that of the oxygen atom, was clearly demonstrated by the weak activity seen with the deoxy compounds **3l,m**. These data could suggest that, as with serotonergic receptors, a hydrogen bond²⁶ may be implicated in the binding of the ligand to the receptor.

Table 2. Displacement of 2-[¹²⁵I]Iodomelatonin Binding from Melatonin Receptors in Chicken Brain Membranes by the Compounds **4**



compd	R ₁	R ₂	R	K _i ± SEM, nM ^a
4a	H	H	CH ₃	2.68 ± 0.3
4b	H	H	C ₂ H ₅	0.67 ± 0.05
4c	H	H	n-C ₃ H ₇	1.37 ± 0.22
4d	H	H	c-C ₃ H ₅	19.4 ± 2.7
4e	H	H	n-C ₄ H ₉	237 ± 50
4f	H	H	CF ₃	0.54 ± 0.05
4g	CH ₃	H	CH ₃	421 ± 48
4h	CH ₃	H	CF ₃	>1000
4i	H	OMe	CH ₃	0.1 ± 0.04
4j	H	OMe	C ₂ H ₅	0.07 ± 0.004
4k	H	OMe	c-C ₃ H ₇	0.05 ± 0.004
4l	H	OMe	CF ₃	0.4 ± 0.04
19				321 ± 33

2b

0.54 ± 0.03

^a K_i values are expressed and were calculated as described in Table 1; the data are the results of one or two separate determinations. ^b K_i = 0.1 ± 0.03 nM determined in a receptor binding assay using ovine pars tuberalis membranes; see ref 19.

However, a significant increase in the affinity was obtained with the substitution of a bromine atom for the 5-methoxy group, and the *N*-propionyl derivative **3o** showed a marked affinity (K_i = 8 nM) for 2-[¹²⁵I]-iodomelatonin binding sites, being only 12-fold less active than melatonin itself (K_i = 0.67 nM). These data suggested that the presence of both methoxy groups is not necessary to obtain compounds with potent affinity, but moreover, they emphasized the essential role of the methoxy group in the ortho position which could give rise to productive interactions with the melatonin receptor site. Therefore, we prepared and evaluated in binding assays compounds such as **4a** (Table 2) where the phenyl ring was substituted by a naphthalene moiety, a well-known bioisosteric group of the indole.²⁷ Compound **4a** had nanomolar affinity for 2-[¹²⁵I]-iodomelatonin binding sites, and comparison of its affinity to that of the unsubstituted derivative **19** demonstrated the efficient role of substitution in the ortho position of the side chain. These results are different from those reported previously in the naphthalene series¹⁹ with




Figure 1. Scatchard plots of the saturation curves of the specific binding of 2-[¹²⁵I]iodomelatonin (0.02–0.8 nM) in brain membranes from chickens killed by decapitation at 12 a.m., 6 p.m., and 12 p.m., respectively. The data were analyzed by linear regression and are the results of one or two separate determinations.

compounds derived from **2** and indole derivatives where the methoxy group needed to be located in the same position as in melatonin to provide potent compounds.

We investigated the influence of the length of the acyl chain with compounds **4b–e**, and as described above for compound **3h**, only the propionyl group increased the affinity (compound **4b**, $K_i = 0.67$ nM) to be equipotent to melatonin. We demonstrated again the failure of the cyclopropyl substituent (compound **4d**, $K_i = 19.4$ nM) to enhance the activity, and the favorable effect of the trifluoroacetyl group (**4f**) to maintain the affinity was also noted. On the other hand, introduction of a methyl group in the α position of the acetamido function resulted in a dramatic drop in the activity of compounds **4g,h**. The different results obtained with the variations of the acyl chain on the affinity of compounds derived from **4a** and **2** for melatonin receptors could be explained by a different positioning of the molecules in the receptor site. This could be due to an interaction with one or other methoxy groups, involving a better or worse fit with the putative lipophilic pocket located near the amido function. However, it is also possible that the steric hindrance of the ortho methoxy group in **4a** brings about, with regard to **2**, some important modifications in the conformation of the flexible side chain and consequently alters the parameters of the pharmacophore implicated in interactions with the receptor site. Nevertheless, the role of the methoxy groups in the 2 or 7 position of the naphthalene ring as anchorage points on the receptor site was demonstrated by the important drop in the affinity observed with the unsubstituted naphthalene acetamido derivative **19** ($K_i = 321 \pm 33$ nM). However, it is possible that compound **4a** could be superimposed with melatonin in two ways corresponding to the different positions of the methoxy group in the receptor site (Figure 2). In position A, the orientation of the side chain of **4a** would be the mirror image of that of melatonin or the bioisosteric derivative **2**, and the different methoxy groups would be superimposed. In this overlay, although the acetamido groups would be located in the same spatial positions in the receptor, the alkyl chain of the acyl group would interact with a different part of the receptor. Consequently, this interaction could explain the difference in affinity

observed with the propyl and cyclopropyl substituents of compounds **4c,d**, respectively, with regard to those of the corresponding derivatives of **2**. On the other hand, superimposition with melatonin or compound **2** according to position B implicates a similar orientation of the side chains for the different compounds and suggests the existence of productive interactions between the methoxy group of **4a** and a region of the receptor located in the neighborhood of the side chain capable of interacting with an ortho substituent. The existence of this additional binding site agrees with earlier structure–activity relationship (SAR) studies²⁸ which demonstrated the favorable role on the affinity values of 2-halogen substitution in melatonin and, particularly, the very high affinity of the 2-iodo derivative. However, the present data reported for **4a** do not indicate which of the two superimpositions with melatonin is correct; further structural variations on the 2 and 7 positions of the naphthalene ring are required to draw definite conclusions. Nevertheless, the existence of the accessory binding site was confirmed in the present study by the enhancement by 1 order of magnitude of the affinity of the 2,7-dimethoxy derivatives (compounds **4i–l**) with regard to the monomethoxy compounds **4a** and **2**. Thus, compounds **4j,k** ($K_i = 0.07$ and 0.05 nM, respectively) were equipotent to 2-iodomelatonin. Moreover, in contrast to the results with the 2-methoxy derivatives, we did not observe a drop in the affinity with the cyclopropyl substitution which provided compound **4k** with high affinity for melatonin binding sites. Consequently, it seems clear that, for these compounds, the 7-methoxy group mimics the methoxy group of melatonin and the 2-methoxy group complements the site occupied by the iodine atom in 2-iodomelatonin.

This study has confirmed the suitable bioisosteric properties of the naphthalene ring for recognition by the melatonin receptor and the existence of an additional binding site capable of giving productive interactions. However, the structural parameters of the melatonin-ergic pharmacophore and, particularly, the relative spatial position of the aromatic system and the amido function in the active conformer are still unclear. The previously reported²⁹ conformational analysis of melatonin demonstrated, as expected for such molecules, a large number of conformers in a relatively small energy range. It is essential to design a potent, conformationally restricted ligand in order to obtain a good insight into the structural parameters of the melatonin-ergic pharmacophore. The structure of the rigid 2-amido-8-methoxytetralin reported recently²⁰ does not seem to be a useful starting framework because of the weak activity, and this point was confirmed recently by the data reported by Garrat³⁰ on a series of the tricyclic indole derivatives related to melatonin, demonstrating the superiority of the compounds with the cis configuration of the ethyl chain equipotent to melatonin while the structurally, closely related tetralin derivatives were markedly less potent.

In the course of this work, it seemed to us worthwhile to evaluate the potential agonist or antagonist properties of the synthesized compounds. Therefore, as a preliminary approach, the influence of guanine nucleotides and cations on the affinity values of melatonin and compound **4b** for the melatonin receptor site was




Figure 2. Putative positions of the compound **4a** in the melatonin receptor. In position A, the 2-methoxy group mimics the melatonergic methoxy; in position B, it interacts with an accessory binding site.

Table 3. Effects of GTP- γ -S and Mn²⁺ on the Affinities of Melatonin and Compound **4b** for Chicken Brain Receptors

compd	K_i^a	K_i	
		+ 3 $\times 10^{-5}$ /M GTP- γ -S ^b	+ 1 mM MnCl ₂ ^b
melatonin	0.31 \pm 0.04	1.08 \pm 0.5	0.176 \pm 0.048
4b	0.68 \pm 0.04	2.86 \pm 0.48	0.297 \pm 0.04

^a K_i values are expressed and were calculated as described in Table 1. ^b The experimental conditions were similar to those reported in the Experimental Section, but the assays were carried out in the presence of GTP- γ -S or MnCl₂.

studied. There is evidence that this receptor is linked to a G-protein³¹ and the addition of GTP or a stable analogue, such as GTP- γ -S, to a membrane preparation decreases the number of high-affinity binding sites available to agonists, giving a shift to the right of the competition curve. In contrast, antagonists bind equally well to the G-protein-coupled or uncoupled receptors, and no shift in the competition curve is seen. Alternatively, Hamon³² reported that divalent cations such as Mn²⁺ increase the apparent affinity of agonists for their binding sites, whereas the affinity of antagonists is unaffected. A simple binding assay for identifying new agonists and antagonists was carried out by evaluating the ratio of the K_i values in the presence of GTP or Mn²⁺. An agonist should have a ratio close to 10, whereas it would not differ from unity for an antagonist.³³ In the present study, the K_i values for melatonin and compound **4b** were measured in the presence of GTP- γ -S (30 μ M) and compared to the values in the presence of Mn²⁺ (1 mM), and the results are shown in Table 3. The affinity of **4b** was affected by both GTP- γ -S and Mn²⁺ to a similar extent to that of melatonin with a ratio close to 10, suggesting an agonist profile for this compound. The marked affinity of compound **4b** for melatonin receptors was confirmed in another binding assay using ovine pars tuberalis membranes^{34,35} ($K_i = 0.61$ nM ($n = 3$)). Furthermore, the agonist profile of compound **4b** was evaluated by measuring its influence on the cAMP synthesis. It is well-known that melatonin inhibits forskolin-stimulated cAMP synthesis¹³ and the agonist or antagonist activity of new compounds can be evaluated by this method. An assay has been developed using cultured ovine pars tuberalis

cells, and the cAMP index was calculated according to the method recently reported by Depreux et al.^{19b} Compound **4b** displayed an activity index of 1.02 and fully mimicked the activity of melatonin, indicating that it can be considered as a full agonist for melatonin receptors.

Conclusion

These data confirm the interest of the new naphthalene derivatives **4** as nonindolic, melatonergic agents, the agonist profile of which should be demonstrated in further pharmacological studies. They also emphasize the important role played by the substituent located on the ortho position of the amido chain on the value of the affinity of the compound for melatonin receptors. On the other hand, these derivatives did not provide complete structural information on the melatonergic pharmacophore, and for this purpose, the development of high-affinity, conformationally locked compounds is essential. Further studies are in progress to evaluate the pharmacological potential of compound **4b** as a nonindolic, melatonergic agent.

Experimental Section

Chemistry. Melting points were determined on a Mettler FP 61 melting point apparatus and are uncorrected. ¹H and ¹³C NMR spectra were recorded on a Bruker AC 200 spectrometer at 200 and 50 MHz, respectively. Chemical shifts are reported in parts per million (δ) relative to tetramethylsilane as an internal standard, and signals are quoted as s (singlet), ds (dedoubled singlet), d (doublet), dd (dedoubled doublet), t (triplet), dt (dedoubled triplet), q (quartet), br s (broad singlet), or m (multiplet). Elemental analyses were performed at the CNRS's analysis services in Vernaison or Gif sur Yvette (France) and are within 0.4% of the theoretical values.

Materials. Tetrahydrofuran (THF) was distilled from sodium/benzophenone. The column chromatography was performed on Merck silica gel 60 (70/230 mesh). 2-(4-Chlorophenyl)ethylamine, 2-(3-chlorophenyl)ethylamine, 2-(2-chlorophenyl)ethylamine, 2-(3-fluorophenyl)ethylamine, 2-(4-methoxyphenyl)ethylamine, 2-(3-methoxyphenyl)ethylamine, 2-(2-methoxyphenyl)ethylamine, 2-(2,5-dimethoxyphenyl)ethylamine, 4-ethylphenol, 5-bromoanisaldehyde, 5-methylsalicylic acid, 2-methoxy-1-naphthaldehyde, and 2,7-dimethoxynaphthalene were purchased from Aldrich (France).

Synthesis of Formyl Compounds 5 and 6. The method is described for the synthesis of 5-ethylanisaldehyde (5). To

a stirred solution of 4-ethylphenol (3.06 g, 25.05 mmol) and K₂CO₃ (7.78 g, 56.29 mmol) in anhydrous acetone (100 mL) under an Ar atmosphere was added dropwise dimethyl sulfate (2.32 mL, 24.56 mmol). After the addition, the reaction mixture was refluxed for 24 h. Excess of (CH₃O)₂SO₂ was decomposed by dropwise addition of water (1.4 mL). The solids were removed by filtration and washed with CH₂Cl₂. The organic layers were dried (MgSO₄), and the solvent was evaporated under reduced pressure. The crude product was purified by chromatography (silica gel, CH₂Cl₂) to yield 2.64 g (19.36 mmol, 77%) of 4-ethylanisole as a colorless liquid: bp 90 °C (13 mmHg); ¹H NMR (CDCl₃) δ 1.28 (t, 3H, CH₃, *J* = 7.6 Hz), 2.66 (q, 2H, CH₂, *J* = 7.6 Hz), 3.84 (s, 3H, MeO), 6.89 (d, 2H, H₂ and H₆, *J* = 8.6 Hz), 7.17 (d, 2H, H₃ and H₅, *J* = 8.6 Hz). To 4-ethylanisole (2.35 g, 17.26 mmol) in anhydrous CH₂Cl₂ (50 mL) under an inert atmosphere and at 0 °C were added dropwise TiCl₄ (99.9%, 2.65 mL, 24.17 mmol) and dichloromethyl methyl ether (2.35 mL, 25.98 mmol). The reaction mixture was stirred at 0 °C for 1 h and at room temperature for 4 h. The mixture was poured into 1 N HCl (50 mL) while stirring, and the two phases were separated. The H₂O layer was washed with CH₂Cl₂. The CH₂Cl₂ layers were washed with H₂O (20 mL) and a saturated NaHCO₃ solution (2 × 20 mL) and then dried over MgSO₄. After evaporation of the solvent under reduced pressure, the crude liquid was purified by column chromatography (silica gel, CH₂Cl₂) to yield 1.76 g (10.72 mmol, 62%) of **5** as a yellow liquid: ¹H NMR (CDCl₃) δ 1.21 (t, 3H, CH₃, *J* = 7.6 Hz), 2.61 (q, 2H, CH₂, *J* = 7.6 Hz), 3.90 (s, 3H, MeO), 6.90 (d, 1H, H₃, *J* = 8.6 Hz), 7.38 (dd, 1H, H₄, *J* = 2.3, 8.6 Hz), 7.65 (ds, 1H, H₆, *J* = 2.3 Hz), 10.44 (s, 1H, CHO); ¹³C NMR (CDCl₃) δ 15.47 (CH₃), 27.66 (CH₂), 55.62 (MeO), 110.60 (C₃), 124.47 (C₁), 127.25 (C₆), 135.46 (C₄), 136.38 (C₅), 160.08 (C₂), 189.94 (CHO).

2,7-Dimethoxy-1-naphthaldehyde (6). This compound was obtained from 2,7-dimethoxynaphthalene. The product was purified by recrystallization from EtOH to yield pure **6** as a yellow crystalline solid: yield 72%; mp 98 °C; ¹H NMR (CDCl₃) δ 3.96 (s, 3H, MeO), 4.00 (s, 3H, MeO), 7.01–7.08 (m, 2H, H₃ and H₆), 7.62 (d, 1H, H₄ or H₅, *J* = 9 Hz), 7.92 (d, 1H, H₄ or H₅, *J* = 9 Hz), 8.82 (ds, 1H, H₈, *J* = 2.4 Hz), 10.86 (s, 1H, CHO); ¹³C NMR (CDCl₃) δ 55.10 (MeO), 56.05 (MeO), 103.13 (C₈), 109.10 (C₃), 115.31 (C₁), 117.05 (C₆), 123.67 (C₁₀), 129.42 (C₅), 133.16 (C₉), 137.04 (C₄), 161.17 (C₇), 164.51 (C₂), 191.73 (CHO).

Synthesis of Benzyl Cyanides 7 and 8. The method is described for the synthesis of 1-(cyanomethyl)-2-methoxy-5-methylbenzene (**7**). It was prepared from 5-methylsalicylic acid according to the following process: to a stirred solution of 5-methylsalicylic acid (5.07 g, 33.32 mmol) and K₂CO₃ (anhydrous, 22.33 g, 161.6 mmol) in anhydrous acetone (175 mL) under an Ar atmosphere was added dropwise dimethyl sulfate (7.25 mL, 76.62 mmol). After the addition, the reaction mixture was refluxed for 6 h. Water (5 mL) was added dropwise to destroy the excess (CH₃O)₂SO₂. After the usual workup, the crude product was purified by chromatography (silica gel, CH₂Cl₂) to yield 5.47 g (30.36 mmol, 91%) of methyl 2-methoxy-5-methylbenzoate: bp 130 °C (12 mmHg); ¹H NMR (CDCl₃) δ 2.27 (s, 3H, CH₃), 3.84 (s, 3H, MeO), 3.85 (s, 3H, MeO), 6.86 (d, 1H, H₃, *J* = 8.4 Hz), 7.25 (dd, 1H, H₄, *J* = 2.4, 8.4 Hz), 7.59 (sd, 1H, H₆, *J* = 2.4 Hz).

The previous compound (4.99 g, 27.69 mmol) in THF (150 mL) was added to a stirred suspension of LiAlH₄ (1.56 g, 41.08 mmol) in THF (100 mL) under an Ar atmosphere. After the addition, the reaction mixture was refluxed for 20 h. After cooling at 0 °C, water (1.3 mL), NaOH (15%, 1.3 mL), ether (40 mL), and water (5 mL) were added dropwise successively to destroy the excess hydride. The mixture was filtered, and the filtrate was dried over MgSO₄. After evaporation of the solvents under reduced pressure, the crude alcohol was purified by distillation to yield 3.68 g (24.18 mmol, 87%) of 2-methoxy-5-methylbenzyl alcohol: bp 131 °C (12 mmHg); ¹H NMR (CDCl₃) δ 2.3 (s, 3H, CH₃), 2.40 (br s, 1H, OH), 3.84 (s, 3H, MeO), 4.65 (s, 2H, CH₂), 6.78 (d, 1H, H₃, *J* = 8.6 Hz), 7.07 (d, 1H, H₄, *J* = 8.6 Hz), 7.09 (s, 1H, H₆).

To a stirred solution of 2-methoxy-5-methylbenzyl alcohol (3.23 g, 21.22 mmol) in anhydrous toluene (50 mL) under an

Ar atmosphere and at 0 °C were added dropwise pyridine (1.7 mL, 21.10 mmol) and then thionyl chloride (3.5 mL, 47.98 mmol). The reaction mixture was allowed to stand at room temperature for 24 h. The mixture was poured into an ice bath and stirred for 1 h. After the usual workup, the crude product was purified by distillation to yield 2.5 g (14.65 mmol, 69%) of 2-methoxy-5-methylbenzyl chloride: bp 118 °C (13 mmHg); ¹H NMR (CDCl₃) δ 2.30 (s, 3H, CH₃), 3.86 (s, 3H, MeO), 4.64 (s, 2H, CH₂), 6.80 (d, 1H, H₃, *J* = 8.7 Hz), 7.11 (dd, 1H, H₄, *J* = 1.9, 8.7 Hz), 7.17 (sd, 1H, H₆).

To a stirred mixture of 2-methoxy-5-methylbenzyl chloride (2.44 g, 14.30 mmol) in anhydrous DMSO (50 mL) were added KCN (1.85 g, 28.41 mmol) and a catalytic amount of KI, and the reaction mixture was stirred at room temperature for 20 h. After evaporation of the DMSO under reduced pressure, the residue was dissolved in CH₂Cl₂ (50 mL) and water (20 mL). The two phases were separated and the aqueous portion was washed with CH₂Cl₂ (20 mL). The combined organic extracts were washed with water (20 mL) and dried over MgSO₄, and the solvent was evaporated under reduced pressure. The crude product was purified by distillation to yield 2.11 g (13.09 mmol, 91%) of **7**: bp 138 °C (16 mmHg); ¹H NMR (CDCl₃) δ 2.30 (s, 3H, CH₃), 3.65 (s, 2H, CH₂), 3.83 (s, 3H, MeO), 6.77 (d, 1H, H₃, *J* = 8.3 Hz), 7.10 (d, 1H, H₄, *J* = 8.3 Hz), 7.17 (s, 1H, H₆); ¹³C NMR (CDCl₃) δ 18.33 (CH₂), 20.21 (CH₃), 55.37 (OCH₃), 110.23 (C₃), 118.04, 118.14 (C₁ and C_N), 129.54, 129.64, 129.96 (C₄, C₅, and C₆), 154.5 (C₂).

1-(Cyanomethyl)-2-methoxy-5-bromobenzene (8). It was prepared from 5-bromo-*o*-anisaldehyde which was reduced by LiAlH₄ in 5-bromo-2-methoxybenzyl alcohol which gave 5-bromo-2-methoxybenzyl chloride according to the process used previously for 5-methyl-2-methoxybenzyl alcohol. It was obtained after purification by chromatography (silica gel, ether–pentane, 3:7); ¹H NMR (CDCl₃) δ 3.78 (s, 3H, MeO), 4.51 (s, 2H, CH₂), 6.69 (d, 1H, H₃, *J* = 8.6 Hz), 7.24–7.41 (m, 2H, H₄ and H₆).

The cyano derivative **8** was obtained according to the above procedure described for **7**. The product was purified by chromatography (silica gel, ether–pentane, 8:2) and then by recrystallization from hexane/AcOEt to yield the pure compound as a yellow crystalline solid: yield 35%; mp 61 °C; ¹H NMR (CDCl₃) δ 3.58 (s, 2H, CH₂), 3.78 (s, 3H, MeO), 6.70 (d, 1H, H₃, *J* = 8.7 Hz), 7.34 (dd, 1H, H₄, *J* = 2.3, 8.7 Hz), 7.41 (ds, 1H, H₆, *J* = 2.3 Hz). Anal. (C₉H₈NOBr).

Synthesis of the Nitro Compounds 9–12. The method is described for the synthesis of 1-nitro-2-(2-methoxynaphthyl)ethylene (**9**). A solution of 2-methoxy-1-naphthaldehyde (1.52 g, 8.16 mmol) and ammonium acetate (0.4 g, 5.19 mmol) in nitromethane (20 mL) was refluxed for 2 h. After evaporation of the solvent under reduced pressure, the residue was dissolved in CH₂Cl₂ (30 mL) and H₂O (30 mL) was added. The aqueous portion was washed with CH₂Cl₂ (2 × 20 mL), and the combined organic layers were washed with H₂O (20 mL). The CH₂Cl₂ solution was dried (MgSO₄), and the solvent was evaporated under reduced pressure to afford 1.77 g (7.73 mmol, 95%) of **9** as a yellow crystalline solid: mp 138 °C; ¹H NMR (CDCl₃) δ 4.04 (s, 3H, MeO), 7.25 (d, 1H, ArH, *J* = 9.7 Hz), 7.37 (dt, 1H, ArH, *J* = 7.2 Hz), 7.56 (dt, 1H, ArH, *J* = 7.2 Hz), 7.76 (d, 1H, ArH, *J* = 8 Hz), 7.92 (d, 1H, ArH, *J* = 9.2 Hz), 8.12 (d, 1H, ArH, *J* = 8.6 Hz), 8.09 (d, 1H, CHNO₂, *J* = 13.4 Hz), 8.79 (d, 1H, ArCH, *J* = 13.4 Hz).

2-Nitro-1-(2-methoxynaphthyl)propene (10). The reaction was carried out with 2-methoxynaphthaldehyde in nitroethane as described in the above procedure. Two isomers, *Z* and *E*, were obtained (2/1): yield 95%; mp 72 °C; ¹H NMR *Z* component (CDCl₃) δ 2.1 (s, 3H, CH₃), 3.97 (s, 3H, MeO), 7.23–7.96 (m, 6H, ArH), 8.33 (s, 1H, ArCH); ¹H NMR *E* component (CDCl₃) δ 2.52 (s, 3H, CH₃), 3.89 (s, 3H, MeO), 6.92 (s, 1H, CH=C–NO₂), 7.23–7.96 (m, 6H, ArH).

1-Nitro-2-(2-methoxy-5-ethylphenyl)ethylene (11). The reaction was carried out with 5-ethylanisaldehyde (**5**) in nitromethane as described in the above procedure. The crude product was purified by column chromatography (silica gel, CH₂Cl₂) to afford **11** as a yellow liquid: yield 82%; bp 155 °C (0.9 mmHg); ¹H NMR (CDCl₃) δ 1.23 (t, 3H, CH₃, *J* = 7.6 Hz), 2.61 (q, 2H, CH₂, *J* = 7.6 Hz), 3.92 (s, 3H, MeO), 6.89 (d, 1H,

H₃, $J = 8.6$ Hz), 7.26 (s, 1H, H₆), 7.28 (d, 1H, H₄, $J = 8.6$ Hz), 7.88 (d, 1H, CHNO₂, $J = 13.6$ Hz), 8.13 (d, 1H, ArCH, $J = 13.6$ Hz); ¹³C NMR (CDCl₃) δ 15.50 (CH₃), 27.61 (CH₂), 55.64 (OMe), 111.32 (C₃), 118.85 (C₁), 131.51 (C₆), 132.83 (C₄), 136.80 (C₅), 135.57, 138.06 (ArCH and CHNO₂), 157.67 (C₂).

1-Nitro-2-(2,7-dimethoxynaphthyl)ethylene (12). The reaction was carried out with 2,7-dimethoxy-1-naphthaldehyde (6) in nitromethane as described in the above procedure. The crude product was purified by column chromatography (silica gel, CH₂Cl₂) to afford **12** as a yellow crystalline solid: yield 95%; mp 140 °C; ¹H NMR (CDCl₃) δ 3.93 (s, 3H, MeO), 4.02 (s, 3H, MeO), 7.04 (dd, 1H, H₆, $J = 2.4, 8.9$ Hz), 7.06 (d, 1H, H₃, $J = 9.1$ Hz), 7.24 (ds, 1H, H₅, $J = 2.4$ Hz), 7.66 (d, 1H, H₄ or H₅, $J = 8.9$ Hz), 7.82 (d, 1H, H₄ or H₅, $J = 9.1$ Hz), 8.07 (d, 1H, CHNO₂, $J = 13.2$ Hz), 8.62 (d, 1H, ArCH, $J = 13.2$ Hz); ¹³C NMR (CDCl₃) δ 55.51 (OMe), 56.21 (OMe), 101.52 (C₃), 109.53 (C₃ or C₆), 110.68 (C₁), 116.72 (C₃ or C₆), 124.41 (C₉ or C₁₀), 130.76, 131.03, 134.24 (ArCH, C₄, and C₅), 135.28 (C₉ or C₁₀), 139.69 (CHNO₂), 159.9 (C₂ and C₇).

Synthesis of Amines 14 and 16–18. Method A. The method is described for the synthesis of 2-(2-methoxynaphthyl)ethylamine (16). To a stirred suspension of LiAlH₄ (1.74 g, 45.82 mmol) in anhydrous THF (50 mL) under an Ar atmosphere and at 0 °C was added dropwise a solution of 1-nitro-2-(2-methoxynaphthyl)ethylene (9) (2.05 g, 8.94 mmol) in THF (75 mL). After the addition, the reaction mixture was heated at 40 °C for 24 h. After cooling at 0 °C, water (1.6 mL), 15% NaOH solution (1.6 mL), ether (37 mL), and water (4.7 mL) were added. The mixture was filtered, and the filtrate was dried over MgSO₄. After evaporation of the solvents under reduced pressure, the crude oil was purified by column chromatography (silica gel, AcOEt–MeOH–Et₃N, 15:3:2) to yield 1.42 g (7.06 mmol, 79%) of **16** as an orange oil: ¹H NMR (CDCl₃) δ 1.71 (br s, 2H, NH₂), 2.98 (t, 2H, ArCH₂, $J = 7.1$ Hz), 3.26 (t, 2H, CH₂NH₂, $J = 7.1$ Hz), 3.94 (s, 3H, MeO), 7.27 (d, 1H, H₁, $J = 9.1$ Hz), 7.34 (dt, 1H, ArH, $J = 7.8$ Hz), 7.48 (dt, 1H, ArH, $J = 1.2, J = 7.8$ Hz), 7.75 (d, 1H, ArH, $J = 9.1$ Hz), 7.79 (d, 1H, ArH, $J = 8.5$ Hz), 7.98 (d, 1H, ArH, $J = 8.5$ Hz).

The oil was dissolved in ether (10 mL), and 2 equiv of a solution of 4 N chlorhydric ether was added. After evaporation of the solvent under reduced pressure, the solid was recrystallized from ethanol/ether to afford the pure HCl salt of the amine **16**: yield 100%; mp >220 °C. Anal. (C₁₃H₁₆NOCl).

1-(2-Methoxynaphthyl)-2-aminopropane (17). The reaction was carried out with 2-nitro-1-(2-methoxynaphthyl)propane (10) as described in the above procedure. The crude oil was purified by chromatography (silica gel, AcOEt and then AcOEt–MeOH–Et₃N, 15:3:2) to afford **17** as an orange oil: yield 59%; ¹H NMR (CDCl₃) δ 1.19 (d, 3H, CH₃, $J = 6.7$ Hz), 1.44 (br s, 2H, NH₂), 3.14 (d, 2H, CH₂, $J = 6.7$ Hz), 3.28 (m, 1H, CH, $J = 6.7$ Hz), 3.94 (s, 3H, MeO), 7.28 (d, 1H, ArH, $J = 8.8$ Hz), 7.34 (dt, 1H, ArH, $J = 1.2, 7.8$ Hz), 7.47 (dt, 1H, ArH, $J = 1.2, 7.8$ Hz), 7.76 (d, 1H, ArH, $J = 8.8$ Hz), 7.79 (d, 1H, ArH, $J = 7$ Hz), 7.99 (d, 1H, ArH, $J = 8.5$ Hz); ¹³C NMR (CDCl₃) δ 23.83 (CH₃), 35.23 (ArCH₂), 47.85 (CHN), 56.25 (OMe), 113.08 (C₃), 121.04 (C₁), 123.11, 123.45 (C₈ and C₆), 126.15, 127.91, 128.44 (C₅, C₇, and C₄), 129.13 (C₁₀), 133.37 (C₉), 154.92 (C₂).

2-(2-Methoxy-5-ethylphenyl)ethylamine (14). The reaction was carried out with 1-nitro-2-(2-methoxy-5-ethylphenyl)ethylene (11) as described in the above procedure. The crude compound was dissolved in ether (10 mL), and 2 equiv of a solution of 4 N chlorhydric ether was added. After evaporation of the solvent under reduced pressure, the solid was recrystallized from ethanol/ether to afford the pure HCl salt of the amine **14**: yield 28%; mp 173 °C; ¹H NMR (CD₃OD) δ 1.19 (t, 3H, CH₃, $J = 7.6$ Hz), 2.56 (q, 2H, ArCH₂, $J = 7.6$ Hz), 2.92 (t, 2H, ArCH₂, $J = 7.6$ Hz), 3.11 (t, 2H, CH₂N, $J = 7.6$ Hz), 3.82 (s, 3H, MeO), 6.89 (d, 1H, H₃, $J = 8.3$ Hz), 7.02 (ds, 1H, H₆, $J = 2.1$ Hz), 7.10 (dd, 1H, H₄, $J = 2.1, 8.3$ Hz); ¹³C NMR (CD₃OD) δ 16.40 (CH₃), 28.86 (ArCH₂), 29.97 (ArCH₂), 40.87 (CH₂N), 59.81 (MeO), 111.68 (C₃), 125.61 (C₁), 128.80 (C₆), 131.14 (C₄), 137.40 (C₅), 157.01 (C₂). Anal. (C₁₁H₁₆NOCl).

2-(2,7-Dimethoxynaphthyl)ethylamine (18). The reaction was carried out with 1-nitro-2-(2,7-dimethoxynaphthyl)-

ethylene (**12**) as described in the above procedure. The crude compound was dissolved in ether (10 mL), and 2 equiv of a solution of 4 N chlorhydric ether was added. After evaporation of the solvent under reduced pressure, the solid was recrystallized from ethanol/ether to afford the pure HCl salt of the amine **18**: yield 63%; mp >222 °C dec; ¹H NMR (CD₃OD) δ 3.14 (t, 2H, ArCH₂, $J = 7.8$ Hz), 3.41 (t, 2H, CH₂N, $J = 7.8$ Hz), 3.94 (s, 3H, MeO), 3.97 (s, 3H, MeO), 7.02 (dd, 1H, H₆, $J = 2.4, 8.9$ Hz), 7.23 (ds, 1H, H₅, $J = 2.4$ Hz), 7.24 (d, 1H, H₃, $J = 9$ Hz), 7.72 (d, 1H, H₄ or H₅, $J = 8.8$ Hz), 7.76 (d, 1H, H₄ or H₅, $J = 8.6$ Hz); ¹³C NMR (CD₃OD) δ 24.16 (ArCH₂), 40.27 (CH₂N), 55.88 (MeO), 56.58 (MeO), 101.84 (C₃), 103.62, 111.22 (C₃ and C₆), 121.45 (C₁), 126.21 (C₁₀), 130.17, 131.50 (C₄ and C₅), 135.39 (C₉), 157.04 (C₇), 160.39 (C₂). Anal. (C₁₄H₁₈NO₂Cl).

Synthesis of Amines 13 and 15. Method B. The method is described for the synthesis of 2-(2-methoxy-5-methylphenyl)ethylamine (13). Ammoniaque (3.8 mL) was added to a vigorously stirred solution of **7** (1.7 g, 10.54 mmol) in ethanol (38 mL). A catalytic amount of Raney Ni was added to the mixture, and the reaction mixture was heated at 40–50 °C for 48 h under a H₂ atmosphere. After cooling, the mixture was filtered on Celite, and the solid was washed with ethanol. After evaporation of the solvent under reduced pressure, the residue was dissolved in ether (30 mL) and dried over MgSO₄, and the solvent was evaporated. To a solution of the crude amine **13** in ether (10 mL) was added a solution of 4 N chlorhydric ether. After evaporation of the solvent under reduced pressure, the solid was recrystallized from a mixture of ethanol/ether to afford 1.07 g (5.31 mmol, yield 50%) of the pure HCl salt of the amine **13**: mp 170 °C; ¹H NMR (CD₃OD) δ 2.25 (s, 3H, CH₃), 2.91 (t, 2H, ArCH₂, $J = 7.4$ Hz), 3.11 (t, 2H, CH₂N, $J = 7.4$ Hz), 3.81 (s, 3H, MeO), 6.85 (d, 1H, H₃, $J = 8.3$ Hz), 7.01 (ds, 1H, H₆), 7.06 (dd, 1H, H₄, $J = 1.7, 8.3$ Hz); ¹³C NMR (CD₃OD) δ 20.49 (CH₃), 29.83 (ArCH₂), 40.91 (CH₂N), 55.89 (MeO), 111.69 (C₃), 125.56 (C₁), 130.06 (C₆), 131.13 (C₄), 132.29 (C₅), 156.90 (C₂). Anal. (C₁₀H₁₆NOCl).

2-(2-Methoxy-5-bromophenyl)ethylamine (15). The reaction was carried out with **8** as described in the above procedure. The crude amine was used directly in the next reaction without further purification: ¹H NMR (CDCl₃) δ 1.39 (br s, 2H, NH₂), 2.66 (m, 2H, ArCH₂), 2.83 (m, 2H, CH₂N), 3.71 (s, 3H, MeO), 6.64 (d, 1H, ArH, $J = 8.5$ Hz), 7.16–7.23 (m, 2H, ArH).

Synthesis of Amides 3a–k,n,o, 4a–e,g,j,k. Method C. The method is described for the synthesis of *N*-[2-(2-methoxynaphthyl)ethyl]propionamide (**4b**). To a stirred solution of **16** (1.40 g, 6.96 mmol) in anhydrous CH₂Cl₂ (20 mL) under an Ar atmosphere and at 0 °C were added dropwise Et₃N (1.5 mL, 10.4 mmol) and then propionyl chloride (0.7 mL, 8.06 mmol). After these additions, the reaction mixture was stirred at room temperature for 30 min. The mixture was poured into water (20 mL) and extracted with CH₂Cl₂ (3 × 20 mL). The combined organic extracts were dried over MgSO₄, and the solvent was evaporated under reduced pressure. The crude solid was recrystallized from pentane/AcOEt to yield 0.70 g (2.72 mmol, 39%) of **4b** as a white crystalline solid: mp 103 °C; ¹H NMR (CDCl₃) δ 1.08 (t, 3H, CH₃, $J = 7.2$ Hz), 2.11 (q, 2H, COCH₂, $J = 7.2$ Hz), 3.33 (t, 2H, ArCH₂, $J = 6.7$ Hz), 3.53 (t, 2H, CH₂N, $J = 6.7$ Hz), 3.96 (s, 3H, CH₃O), 5.82 (br s, 1H, NH), 7.28 (d, 1H, ArH, $J = 8.7$ Hz), 7.34 (dt, 1H, ArH, $J = 1.3, 7.8$ Hz), 7.49 (dt, 1H, ArH, $J = 1.3, 7.8$ Hz), 7.76 (d, 1H, ArH, $J = 9.0$ Hz), 7.78 (d, 1H, ArH, $J = 7.5$ Hz), 8.02 (d, 1H, ArH, $J = 8.6$ Hz); ¹³C NMR (CDCl₃) δ 9.63 (CH₃), 24.46 (ArCH₂), 29.67 (COCH₂), 39.85 (CH₂N), 56.41 (MeO), 112.89 (C₃), 120.26 (C₁), 123.10, 123.47 (C₆ and C₈), 126.69, 128.35, 128.47 (C₄, C₅, and C₇), 129.22 (C₁₀), 133.24 (C₉), 154.58 (C₂), 173.82 (CO). Anal. (C₁₆H₁₉NO₂).

***N*-[2-(4-Chlorophenyl)ethyl]acetamide (3a).** Acylation of 2-(4-chlorophenyl)ethylamine with acetyl chloride in the above procedure produced **3a** as a white crystalline solid purified by recrystallization from hexane/AcOEt: yield 84%; mp 96 °C; ¹H NMR (CDCl₃) δ 1.77 (s, 3H, CH₃), 2.61 (t, 2H, ArCH₂, $J = 7.0$ Hz), 3.29 (q, 2H, CH₂N, $J = 6.6$ Hz), 5.66 (br s, 1H, NH), 6.93 (d, 2H, H₂ and H₆, $J = 8.4$ Hz), 7.09 (d, 2H, H₃ and H₅, $J = 8.4$ Hz); ¹³C NMR (CDCl₃) δ 23.17 (CH₃), 35.02 (ArCH₂),

40.68 (CH₂NH), 128.78 (C₃ and C₅), 130.11 (C₂ and C₆), 132.41 (C₄), 137.34 (C₁), 170.32 (CO). Anal. (C₁₀H₁₂NOCl).

N-[2-(3-Chlorophenyl)ethyl]acetamide (3b). Acylation of 2-(3-chlorophenyl)ethylamine with acetyl chloride in the above procedure produced **3b** as a yellow oil purified by column chromatography (silica gel, CHCl₃-MeOH, 95:5): yield 89%; ¹H NMR (CDCl₃) δ 1.87 (s, 3H, CH₃), 2.72 (t, 2H, ArCH₂, *J* = 7.0 Hz), 3.41 (q, 2H, CH₂N, *J* = 6.7 Hz), 5.66 (br s, 1H, NH), 6.99–7.03 (dd, 1H, H₆, *J* = 8.1 Hz), 7.11–7.20 (m, 3H, H₂, H₄, and H₅). Anal. (C₁₀H₁₂NOCl).

N-[2-(2-Chlorophenyl)ethyl]acetamide (3c). Acylation of 2-(2-chlorophenyl)ethylamine with acetyl chloride in the above procedure produced **3c** as an oil purified by column chromatography (silica gel, CHCl₃-MeOH, 95:5): yield 61%; ¹H NMR (CDCl₃) δ 1.87 (s, 3H, CH₃), 2.88 (t, 2H, ArCH₂, *J* = 7.0 Hz), 3.44 (q, 2H, CH₂N, *J* = 6.6 Hz), 5.77 (br s, 1H, NH), 7.06–7.16 (m, 3H, H₄, H₅, and H₆), 7.27 (d, 1H, H₃, *J* = 8.2 Hz). Anal. (C₁₀H₁₂NOCl).

N-[2-(3-Fluorophenyl)ethyl]acetamide (3d). Acylation of 2-(3-fluorophenyl)ethylamine with acetyl chloride in the above procedure produced **3d** as an oil purified by column chromatography (silica gel, CH₂Cl₂-MeOH, 95:5): yield 58%; ¹H NMR (CDCl₃) δ 1.86 (s, 3H, CH₃), 2.73 (t, 2H, ArCH₂, *J* = 7.1 Hz), 3.40 (q, 2H, CH₂N, *J* = 6.6 Hz), 5.86 (br s, 1H, NH), 6.78–6.90 (m, 3H, H₂, H₄, and H₆), 7.16 (t, 1H, H₅, *J* = 8.0 Hz); ¹³C NMR (CDCl₃) δ 23.23 (CH₃), 35.49 (ArCH₂), 40.62 (CH₂NH), 113.48 (d, 1C, C₄, *J* = 21 Hz), 115.66 (d, 1C, C₂, *J* = 21 Hz), 124.52 (C₆), 130.16 (d, 1C, C₅, *J* = 8 Hz), 141.66 (d, 1C, C₁, *J* = 7 Hz), 160.62, 165.51 (d, 1C, C₃, *J* = 250 Hz), 170.47 (CO). Anal. (C₁₀H₁₂NOF).

N-[2-(4-Methoxyphenyl)ethyl]acetamide (3e). Acylation of 2-(4-methoxyphenyl)ethylamine with acetyl chloride in the above procedure produced **3e** as a white crystalline solid purified by recrystallization from hexane/AcOEt: yield 41%; mp 86 °C; ¹H NMR (CDCl₃) δ 1.84 (s, 3H, CH₃), 2.67 (t, 2H, ArCH₂, *J* = 6.9 Hz), 3.39 (q, 2H, CH₂N, *J* = 6.8 Hz), 3.71 (s, 3H, MeO), 5.58 (br s, 1H, NH), 6.76 (d, 2H, H₃ and H₅, *J* = 8.6 Hz), 7.02 (d, 2H, H₂ and H₆, *J* = 8.5 Hz). Anal. (C₁₁H₁₅NO₂).

N-[2-(3-Methoxyphenyl)ethyl]acetamide (3f). Acylation of 2-(3-methoxyphenyl)ethylamine with acetyl chloride in the above procedure produced **3f** as an oil purified by column chromatography (silica gel; CHCl₃-MeOH, 9:1): yield 60%; ¹H NMR (CDCl₃) δ 1.84 (s, 3H, CH₃), 2.69 (t, 2H, ArCH₂, *J* = 7.0 Hz), 3.40 (q, 2H, CH₂N, *J* = 6.6 Hz), 3.69 (s, 3H, OCH₃), 5.74 (br s, 1H, NH), 6.64–6.70 (m, 2H, H₄ and H₆), 7.12 (t, 1H, H₅, *J* = 7.7 Hz); ¹³C NMR (CDCl₃) δ 23.26 (CH₃), 35.68 (ArCH₂), 40.68 (CH₂NH), 55.25 (OMe), 111.91 (C₄), 114.56 (C₂), 121.12 (C₆), 129.70 (C₅), 140.54 (C₁), 159.92 (C₃), 170.35 (CO). Anal. (C₁₁H₁₅NO₂).

N-[2-(2-Methoxyphenyl)ethyl]acetamide (3g). Acylation of 2-(2-methoxyphenyl)ethylamine with acetyl chloride in the above procedure produced **3g** as a white crystalline solid purified by recrystallization from hexane/AcOEt: yield 53%; mp 76 °C; ¹H NMR (CDCl₃) δ 1.87 (s, 3H, COCH₃), 2.78 (t, 2H, ArCH₂, *J* = 6.9 Hz), 3.53 (t, 2H, CH₂N, *J* = 6.9 Hz), 3.72 (s, 3H, MeO), 6.36 (br s, 1H, NH), 6.65–6.76 (m, 3H, ArH), 7.18 (t, 1H, ArH, *J* = 7.9 Hz). Anal. (C₁₁H₁₅NO₂).

N-[2-(2,5-Dimethoxyphenyl)ethyl]acetamide (3h). Acylation of 2-(2,5-dimethoxyphenyl)ethylamine with acetyl chloride in the above procedure produced **3h** as a white crystalline solid purified by recrystallization from hexane/AcOEt: yield 69%; mp 96 °C; ¹H NMR (CDCl₃) δ 1.86 (s, 3H, COCH₃), 2.73 (t, 2H, ArCH₂, *J* = 6.6 Hz), 3.40 (t, 2H, CH₂N, *J* = 6.6 Hz), 3.69 (s, 3H, MeO), 3.73 (s, 3H, MeO), 5.64 (br s, 1H, NH), 6.64–6.76 (m, 3H, ArH). Anal. (C₁₂H₁₇NO₂).

N-[2-(2,5-Dimethoxyphenyl)ethyl]propionamide (3i). Acylation of 2-(2,5-dimethoxyphenyl)ethylamine with propionyl chloride in the above procedure produced **3i** as a white crystalline solid purified by recrystallization from hexane/AcOEt: yield 81%; mp 115 °C; ¹H NMR (CDCl₃) δ 1.05 (t, 3H, CH₃, *J* = 7.6 Hz), 2.08 (q, 2H, COCH₂), 2.73 (t, 2H, ArCH₂, *J* = 6.6 Hz), 3.40 (m, 2H, CH₂N), 3.69 (s, 3H, MeO), 3.73 (s, 3H, MeO), 5.69 (br s, 1H, NH), 6.64–6.76 (m, 3H, ArH). Anal. (C₁₃H₁₉NO₂).

N-[2-(2,5-Dimethoxyphenyl)ethyl]cyclopropylformamide (3j). Acylation of 2-(2,5-dimethoxyphenyl)ethyl-

amine with cyclopropanecarbonyl chloride in the above procedure produced **3j** as a white crystalline solid purified by recrystallization from hexane/AcOEt: yield 84%; mp 118 °C; ¹H NMR (CDCl₃) δ 0.56–0.65 (m, 2H, cyclopropyl), 0.87–0.94 (m, 2H, cyclopropyl), 1.13–1.23 (m, 1H, cyclopropyl), 2.75 (t, 2H, ArCH₂, *J* = 6.7 Hz), 3.42 (m, 2H, CH₂N), 3.69 (s, 3H, MeO), 3.74 (s, 3H, MeO), 5.79 (br s, 1H, NH), 6.66–6.76 (m, 3H, ArH). Anal. (C₁₄H₁₉NO₂).

N-[2-(2,5-Dimethoxyphenyl)ethyl]-*n*-valeramide (3k). Acylation of 2-(2,5-dimethoxyphenyl)ethylamine with *n*-valeryl chloride in the above procedure produced **3k** as a white crystalline solid purified by recrystallization from hexane/AcOEt: yield 68%; mp 80 °C; ¹H NMR (CDCl₃) δ 0.82 (t, 3H, CH₃, *J* = 7.2 Hz), 1.10–1.32 (m, 2H, CH₂), 1.42–1.47 (m, 2H, CH₂), 2.05 (t, 2H, COCH₂, *J* = 7.2 Hz), 2.73 (t, 2H, ArCH₂, *J* = 6.6 Hz), 3.40 (m, 2H, CH₂N), 3.69 (s, 3H, MeO), 3.73 (s, 3H, MeO), 5.66 (br s, 1H, NH), 6.64–6.75 (m, 3H, ArH). Anal. (C₁₅H₂₃NO₂).

N-[2-(2-Methoxy-5-bromophenyl)ethyl]acetamide (3n). Acylation of the crude amine **15** with acetyl chloride in the above procedure produced **3n** as a white crystalline solid purified by recrystallization from hexane/AcOEt: yield 39%; mp 103 °C; ¹H NMR (CDCl₃) δ 1.87 (s, 3H, COCH₃), 2.71 (t, 2H, ArCH₂, *J* = 6.8 Hz), 3.34–3.44 (m, 2H, CH₂N), 3.75 (s, 3H, MeO), 5.53 (br s, 1H, NH), 6.67 (d, 1H, ArH, *J* = 8.6 Hz), 7.17–7.27 (m, 2H, ArH). Anal. (C₁₁H₁₄NO₂Br).

N-[2-(2-Methoxy-5-bromophenyl)ethyl]propionamide (3o). Acylation of the crude amine **15** with propionyl chloride in the above procedure produced **3o** as a white crystalline solid purified by recrystallization from hexane/AcOEt: yield 30%; mp 106 °C; ¹H NMR (CDCl₃) δ 1.06 (t, 3H, CH₃, *J* = 7.6 Hz), 2.09 (q, 2H, CH₂, *J* = 7.6 Hz), 2.72 (t, 2H, ArCH₂, *J* = 7.6 Hz), 3.34–3.44 (m, 2H, CH₂N), 3.75 (s, 3H, MeO), 5.55 (br s, 1H, NH), 6.66 (d, 1H, ArH, *J* = 8.7 Hz), 7.15–7.27 (m, 2H, ArH). Anal. (C₁₂H₁₆NO₂Br).

N-[2-(2-Methoxynaphthyl)ethyl]acetamide (4a). Acylation of **16** with acetyl chloride in the above procedure produced **4a** as a white crystalline solid purified by recrystallization from hexane/ethyl acetate: yield 77%; mp 139 °C; ¹H NMR (CDCl₃) δ 1.81 (s, 3H, COCH₃), 3.25 (t, 2H, ArCH₂, *J* = 6.8 Hz), 3.48 (t, 2H, CH₂N, *J* = 6.8 Hz), 3.90 (s, 3H, CH₃O), 5.69 (br s, 1H, NH), 7.19–7.31 (m, 2H, ArH), 7.39–7.47 (m, 1H, ArH), 7.68–7.74 (m, 2H, ArH), 7.94 (d, 1H, ArH, *J* = 8.6 Hz). Anal. (C₁₅H₁₇NO₂).

N-[2-(2-Methoxynaphthyl)ethyl]-*n*-butyramide (4c). Acylation of **16** with butyryl chloride in the above procedure produced **4c** as a white crystalline solid purified by recrystallization from pentane/AcOEt: yield 48%; mp 76 °C; ¹H NMR (CDCl₃) δ 0.86 (t, 3H, CH₃, *J* = 7.5 Hz), 1.57 (m, 2H, CH₂, *J* = 7.5 Hz), 2.03 (t, 2H, COCH₂, *J* = 7.5 Hz), 3.30 (t, 2H, ArCH₂, *J* = 6.5 Hz), 3.52 (t, 2H, CH₂N, *J* = 6.5 Hz), 3.95 (s, 3H, CH₃O), 5.74 (br s, 1H, NH), 7.26 (d, 1H, ArH, *J* = 9.4 Hz), 7.33 (dt, 1H, ArH, *J* = 7.4 Hz), 7.48 (dt, 1H, ArH, *J* = 1.0, 7.4 Hz), 7.76 (d, 1H, ArH, *J* = 9.0 Hz), 7.77 (d, 1H, ArH, *J* = 7.5 Hz), 8.03 (d, 1H, ArH, *J* = 8.5 Hz). Anal. (C₁₇H₂₁NO₂).

N-[2-(2-Methoxynaphthyl)ethyl]cyclopropylformamide (4d). Acylation of **16** with cyclopropanecarbonyl chloride in the above procedure produced **4d** as a beige crystalline solid purified by column chromatography (silica gel, CH₂Cl₂-MeOH, 99.8:0.2) followed by recrystallization from hexane/AcOEt: yield 39%; mp 127.5 °C; ¹H NMR (CDCl₃) δ 0.56–0.65 (m, 2H, cyclopropyl), 0.85–0.92 (m, 2H, cyclopropyl), 1.08–1.18 (m, 2H, cyclopropyl), 3.24 (t, 2H, ArCH₂, *J* = 6.3 Hz), 3.49 (t, 2H, CH₂N, *J* = 6.3 Hz), 3.90 (s, 3H, CH₃O), 5.86 (br s, 1H, NH), 7.21 (d, 1H, ArH, *J* = 8.8 Hz), 7.30 (dt, 1H, ArH, *J* = 1.3, 7.8 Hz), 7.44 (dt, 1H, ArH, *J* = 1.3, 7.8 Hz), 7.71 (d, 1H, ArH, *J* = 9.0 Hz), 7.73 (d, 1H, ArH, *J* = 7.5 Hz), 7.96 (d, 1H, ArH, *J* = 8.6 Hz). Anal. (C₁₇H₁₉NO₂).

N-[2-(2-Methoxynaphthyl)ethyl]-*n*-valeramide (4e). Acylation of **16** with *n*-valeryl chloride in the above procedure produced **4e** as a white crystalline solid purified by recrystallization from pentane/AcOEt: yield 47%; mp 72 °C; ¹H NMR (CDCl₃) δ 0.88 (t, 3H, CH₃, *J* = 7.3 Hz), 1.27 (m, 2H, CH₂, *J* = 7.3 Hz), 1.54 (m, 2H, CH₂, *J* = 7.3 Hz), 2.08 (t, 2H, COCH₂, *J* = 7.6 Hz), 3.33 (t, 2H, ArCH₂, *J* = 6.4 Hz), 3.54 (t, 2H, CH₂N, *J* = 6.4 Hz), 3.97 (s, 3H, CH₃O), 5.77 (br s, 1H, NH), 7.28 (d, 1H,

ArH, $J = 9.3$ Hz), 7.33 (t, 1H, ArH, $J = 7.7$ Hz), 7.50 (dt, 1H, ArH, $J = 1.2, 7.7$ Hz), 7.77 (d, 1H, ArH, $J = 9.1$ Hz), 7.79 (d, 1H, ArH, $J = 7.8$ Hz), 8.03 (d, 1H, ArH, $J = 8.6$ Hz). Anal. ($C_{18}H_{23}NO_2$).

N-[1-Methyl-2-(2-methoxynaphthyl)ethyl]acetamide (4g). Acylation of **17** with acetyl chloride in the above procedure produced **4g** as a white crystalline solid purified by column chromatography (silica gel, CH_2Cl_2 -MeOH, 98:2): yield 90%; mp 151 °C; 1H NMR ($CDCl_3$) δ 1.24 (d, 3H, CH_3 -CH, $J = 6.5$ Hz), 1.84 (s, 3H, COCH₃), 3.13–3.38 (m, 2H, ArCH₂, $J_{A-B} = 13.7$ Hz, $J_{A-X} = 6.3$ Hz, $J_{B-X} = 8$ Hz), 3.97 (s, 3H, CH₃O), 4.14–4.34 (m, 1H, CH, $J_{A-X} = 6.3$ Hz, $J_{B-X} = 8$ Hz), 6.02 (d, 1H, NH, $J = 5.7$ Hz), 7.28 (d, 1H, ArH, $J = 7.8$ Hz), 7.34 (dt, 1H, ArH, $J = 7.8$ Hz), 7.50 (dt, 1H, ArH, $J = 1.2, 7.3$ Hz), 7.76 (d, 1H, ArH, $J = 9.1$ Hz), 7.78 (d, 1H, ArH, $J = 8.1$ Hz), 8.07 (d, 1H, ArH, $J = 8.6$ Hz); ^{13}C NMR ($CDCl_3$) δ 20.71 (CH₃CO), 23.45 (CH₃CH), 31.22 (ArCH₂), 47.0 (CHN), 56.45 (MeO), 112.92 (C₃), 119.93 (C₁), 123.40, 123.51 (C₆ and C₈), 126.66, 128.43, 128.43 (C₄, C₅, and C₇), 129.32 (C₁₀), 133.3 (C₉), 154.62 (C₂), 169.48 (CO). Anal. ($C_{16}H_{19}NO_2$).

N-[2-(2,7-Dimethoxynaphthyl)ethyl]propionamide (4j). Acylation of the free amine **18** with propionyl chloride in the above procedure produced **4j** as a white crystalline solid purified by recrystallization from CH_2Cl_2 /petroleum ether: yield 74%; mp 126 °C; 1H NMR ($CDCl_3$) δ 1.00 (t, 3H, CH_3 , $J = 7.6$ Hz), 2.04 (q, 2H, COCH₂, $J = 7.6$ Hz), 3.20 (t, 2H, ArCH₂, $J = 6.8$ Hz), 3.46 (q, 2H, CH₂N, $J = 6.4$ Hz), 3.87 (s, 3H, MeO), 3.88 (s, 3H, MeO), 5.78 (br s, 1H, NH), 6.98 (dd, 1H, H₆, $J = 2.5, 8.9$ Hz), 7.03 (d, 1H, H₃, $J = 8.9$ Hz), 7.26 (ds, 1H, H₈, $J = 2.5$ Hz), 7.59 (d, 2H, H₄ and H₅, $J = 8.9$ Hz); ^{13}C NMR ($CDCl_3$) δ 9.64 (CH₃), 24.73 (ArCH₂), 29.74 (CH₂), 39.52 (CH₂N), 55.38 (MeO), 56.27 (MeO), 101.65 (C₈), 110.22, 116.31 (C₃ and C₆), 119.04 (C₁), 124.71 (C₁₀), 128.02, 129.97 (C₄ and C₅), 134.65 (C₉), 155.18 (C₇), 158.53 (C₂), 173.84 (CO). Anal. ($C_{17}H_{21}NO_3$).

N-[2-(2,7-Dimethoxynaphthyl)ethyl]cyclopropylformamide (4k). Acylation of the free amine **18** with cyclopropanecarbonyl chloride in the above procedure produced **4k** as a beige crystalline solid purified by column chromatography (silica gel, CH_2Cl_2 -MeOH, 98:2): yield 61%; mp 140 °C; 1H NMR ($CDCl_3$) δ 0.62–0.72 (m, 2H, cyclopropyl), 0.89–0.97 (m, 2H, cyclopropyl), 1.15–1.25 (m, 2H, cyclopropyl), 3.28 (t, 2H, ArCH₂, $J = 6.8$ Hz), 3.56 (q, 2H, CH₂N, $J = 6.3$ Hz), 3.94 (s, 3H, MeO), 3.96 (s, 3H, MeO), 5.94 (br s, 1H, NH), 6.91 (dd, 1H, H₆, $J = 2.4, 8.9$ Hz), 7.12 (d, 1H, H₃, $J = 8.9$ Hz), 7.30 (ds, 1H, H₈, $J = 2.4$ Hz), 7.68 (d, 1H, H₄ or H₅, $J = 8.9$ Hz), 7.69 (d, 1H, H₄ or H₅, $J = 8.9$ Hz); ^{13}C NMR ($CDCl_3$) δ 6.79 (2C, CH₂ cyclopropyl), 14.77 (CH cyclopropyl), 24.92 (ArCH₂), 39.74 (CH₂N), 55.34 (MeO), 56.28 (MeO), 101.70 (C₈), 110.28, 116.25 (C₃ and C₆), 119.14 (C₁), 124.70 (C₁₀), 127.99, 129.95 (C₄ and C₅), 134.63 (C₉), 155.24 (C₇), 158.50 (C₂), 173 (CO). Anal. ($C_{18}H_{21}NO_3$).

Synthesis of Acetamides 3l,m and 4i. Method D. The method is described for the synthesis of *N*-[2-(2-methoxy-5-methylphenyl)ethyl]acetamide (**3l**). To a stirred solution of the hydrochloride salt **13** (0.31 g, 1.54 mmol) in water (15 mL) were added successively AcONa (1.6 g, 19.51 mmol) and acetic anhydride (3.6 mL, 38.15 mmol). After these additions, the reaction mixture was stirred at room temperature for 30 min. The amide was extracted from water with CH_2Cl_2 (3 × 20 mL). This organic extract was washed with a saturated NaHCO₃ solution (30 mL) and water (30 mL) and then dried (MgSO₄). After evaporation of the solvent under reduced pressure, the crude product was purified by column chromatography (silica gel, CH_2Cl_2 -MeOH, 95:5) to yield 0.25 g (1.21 mmol, 79%) of **3l**: mp 76 °C; 1H NMR ($CDCl_3$) δ 1.91 (s, 3H, COCH₃), 2.25 (s, 3H, CH₃), 2.77 (t, 2H, ArCH₂, $J = 6.8$ Hz), 3.44 (q, 2H, CH₂N, $J = 6.4$ Hz), 3.79 (s, 3H, MeO), 5.95 (br s, 1H, NH), 6.74 (d, 1H, H₃, $J = 8.1$ Hz), 6.94 (dd, 1H, H₄, $J = 1.8, 8.1$ Hz), 7.01 (ds, 1H, H₆, $J = 1.8$ Hz); ^{13}C NMR ($CDCl_3$) δ 20.41 (ArCH₃), 23.27 (COCH₃), 30.07 (ArCH₂), 40.13 (CH₂NH), 55.47 (OCH₃), 110.47 (C₃), 127.25 (C₁), 128.07 (C₆), 129.96 (C₄), 131.37 (C₅), 155.46 (C₂), 170.07 (CO). Anal. ($C_{12}H_{17}NO_2$).

N-[2-(2-Methoxy-5-ethylphenyl)ethyl]acetamide (3m). Acetylation of the chlorhydrate of the amine **14** in the above procedure produced **3m**. The crude amide was purified by

column chromatography (silica gel, CH_2Cl_2 -MeOH, 98:2): yield 87%; mp 82 °C; 1H NMR ($CDCl_3$) δ 1.13 (t, 3H, CH₃, $J = 7.6$ Hz), 1.85 (s, 3H, COCH₃), 2.50 (q, 2H, C₂CH₂, $J = 7.6$ Hz), 2.73 (t, 2H, C₁CH₂, $J = 6.7$ Hz), 3.39 (q, 2H, CH₂N, $J = 6.3$ Hz), 3.74 (s, 3H, MeO), 5.69 (br s, 1H, NH), 6.72 (d, 1H, H₃, $J = 8.3$ Hz), 6.90 (ds, 1H, H₆), 6.96 (dd, 1H, H₄, $J = 8.3$ Hz); ^{13}C NMR ($CDCl_3$) δ 15.99 (CH₃), 23.43 (COCH₃), 28.07 (CH₂), 30.26 (ArCH₂), 40.25 (CH₂NH), 55.59 (OCH₃), 110.60 (C₃), 126.98 (C₆), 127.39 (C₁), 130.37 (C₄), 136.71 (C₅), 155.71 (C₂), 170.10 (CO). Anal. ($C_{13}H_{19}NO_2$).

N-[2-(2,7-Dimethoxynaphthyl)ethyl]acetamide (4i). Acylation of the chlorhydrate of the amine **18** in the above procedure produced **4i**. The crude amide was purified by column chromatography (silica gel, CH_2Cl_2 -MeOH, 98:2): yield 64%; mp 110 °C; 1H NMR ($CDCl_3$) δ 1.91 (s, 3H, COCH₃), 3.29 (t, 2H, ArCH₂, $J = 7$ Hz), 3.52 (q, 2H, CH₂N, $J = 6.5$ Hz), 3.95 (s, 3H, MeO), 3.97 (s, 3H, MeO), 5.79 (br s, 1H, NH), 7.02 (dd, 1H, H₆, $J = 2.4, 8.8$ Hz), 7.12 (d, 1H, H₃, $J = 8.9$ Hz), 7.35 (ds, 1H, H₈, $J = 2.4$ Hz), 7.68 (d, 1H, H₄ or H₅, $J = 9$ Hz), 7.69 (d, 1H, H₄ or H₅, $J = 8.8$ Hz); ^{13}C NMR ($CDCl_3$) δ 23.33 (CH₃), 24.79 (ArCH₂), 39.68 (CH₂N), 55.39 (MeO), 56.31 (MeO), 101.56 (C₈), 110.25, 116.38 (C₃ and C₆), 118.98 (C₁), 124.69 (C₁₀), 128.02, 129.95 (C₄ and C₅), 134.67 (C₉), 155.19 (C₇), 158.53 (C₂), 170.25 (CO). Anal. ($C_{16}H_{19}NO_3$).

Synthesis of Trifluoroacetamides 4f,h,l. Method E. The method is described for the synthesis of *N*-[1-methyl-2-(2-methoxynaphthyl)ethyl]trifluoroacetamide (**4h**). To a stirred solution of **17** (0.31 g, 1.44 mmol) in anhydrous CH_2Cl_2 (20 mL) under an Ar atmosphere and at 0 °C were added dropwise pyridine (0.12 mL, 1.44 mmol) and then trifluoroacetic anhydride (0.245 mL, 1.73 mmol). After these additions, the reaction mixture was allowed to stir at room temperature for 30 min. The mixture was poured into water (20 mL), the two phases were separated, and the aqueous portion was washed with CH_2Cl_2 (2 × 20 mL). The combined organic extracts were washed with a saturated NaHCO₃ solution (40 mL) and water (40 mL) and dried over MgSO₄. After evaporation of the solvent under reduced pressure, the crude solid was purified by column chromatography (silica gel, CH_2Cl_2 -MeOH, 98:2) and recrystallization from pentane/AcOEt to yield 0.15 g (0.482 mmol, 34%) of **4h** as a white crystalline solid: mp 123 °C; 1H NMR ($CDCl_3$) δ 1.41 (d, 3H, CH₃, $J = 6.4$ Hz), 3.22–3.46 (m, 2H, ArCH₂, $J_{A-B} = 14.3$ Hz, $J_{A-X} = 4.7$ Hz, $J_{B-X} = 9.5$ Hz), 4.00 (s, 3H, CH₃O), 4.21–4.26 (m, 1H, CH, $J_{A-X} = 4.7$ Hz, $J_{B-X} = 9.5$ Hz), 7.30 (d, 1H, ArH, $J = 9$ Hz), 7.38 (dt, 1H, ArH, $J = 7.5$ Hz), 7.53 (dt, 1H, ArH, $J = 7.5$ Hz), 7.82 (d, 2H, ArH, $J = 8.8$ Hz), 7.92 (d, 1H, ArH, $J = 8.5$ Hz); ^{13}C NMR ($CDCl_3$) δ 20.40 (CH₃), 30.69 (ArCH₂), 48.46 (CHN), 56.10 (MeO), 112.34 (C₃), 118.49 (C₁), 122.64, 123.73 (C₆ and C₈), 127.0, 128.75, 129.09 (C₄, C₅, and C₇), 129.45 (C₁₀), 133.2 (C₉), 154.27 (C₂). Anal. ($C_{16}H_{19}NO_2F_3$).

N-[2-(2-Methoxynaphthyl)ethyl]trifluoroacetamide (4f). Acylation of **16** with trifluoroacetic anhydride in the above procedure produced **4f** as a white crystalline solid purified by recrystallization from hexane/AcOEt: yield 29%; mp 87 °C; 1H NMR ($CDCl_3$) δ 3.34 (t, 2H, ArCH₂, $J = 6.8$ Hz), 3.55–3.64 (m, 2H, CH₂N), 3.91 (s, 3H, CH₃O), 6.97 (br s, 1H, NH), 7.18–7.49 (m, 3H, ArH), 7.72–7.87 (m, 3H, ArH). Anal. ($C_{15}H_{14}NO_2F_3$).

N-[2-(2,7-Dimethoxynaphthyl)ethyl]trifluoroacetamide (4l). Acylation of the free amine **18** with trifluoroacetic anhydride in the above procedure produced **4l** as a white crystalline solid purified by column chromatography (silica gel, CH_2Cl_2): yield 48%; mp 132 °C; 1H NMR ($CDCl_3$) δ 3.29 (t, 2H, ArCH₂, $J = 6.6$ Hz), 3.58 (q, 2H, CH₂N, $J = 6.1$ Hz), 3.88 (s, 3H, MeO), 3.89 (s, 3H, MeO), 6.96 (br s, 1H, NH), 6.96 (dd, 1H, H₆, $J = 2.5, 8.9$ Hz), 7.06 (d, 1H, H₃, $J = 8.9$ Hz), 7.12 (ds, 1H, H₈, $J = 2.5$ Hz), 7.63 (d, 1H, H₄ or H₅, $J = 8.9$ Hz), 7.65 (d, 1H, H₄ or H₅, $J = 8.9$ Hz); ^{13}C NMR ($CDCl_3$) δ 23.76 (ArCH₂), 40.32 (CH₂N), 55.36 (MeO), 56.09 (MeO), 101.10 (C₈), 109.88, 116.45 (C₃ and C₆), 115.91 (q, 1C, CF₃, $J = 288$ Hz), 118.77 (C₁), 124.87 (C₁₀), 128.73, 130.34 (C₄ and C₅), 134.28 (C₉), 155.14 (C₇), 157.37 (q, 1C, CO, $J = 37$ Hz), 158.88 (C₂). Anal. ($C_{16}H_{19}NO_3$).

Melatonin Binding Assays. Chickens (Redbro, 4 months, 3–4 kg; Cellu-bio, France) were decapitated at 12 a.m., 6 p.m.,

and 12 p.m., respectively. The whole brains were stored at -80°C . They were homogenized with a Polytron homogenizer in 10 volumes of ice-cold Tris-HCl buffer (50 mM, pH 7.4) and washed twice by centrifugation (4400g, 25 min, 4°C). The resulting pellet was resuspended in 10 volumes of the same buffer to a final concentration of 5 mg of protein/mL determined by the method of Lowry.³⁶ The membrane aliquots were stored at -80°C until subsequent use. The binding assays were performed according to the method of Dubocovich with modifications.^{37,22,18} Membrane aliquots (20 μL) were incubated in a total volume of 0.25 mL of Tris-HCl buffer (50 mM, pH 7.4) with eight concentrations (0.02–0.8 nM) of the labeled ligand 2-[¹²⁵I]iodomelatonin (2000 Ci/mmol; purchased from NEN or amersham) for the saturation experiments; 0.05 nM 2-[¹²⁵I]iodomelatonin and seven concentrations of the compound under test were used for the competition studies. Each binding assay was performed in triplicate. The incubation (25 $^{\circ}\text{C}$, 60 min) was stopped by the addition of 3 mL of ice-cold buffer and immediate vacuum filtration through glass fiber filters (GF/B Whatman strips) presoaked in 0.1% poly(ethylenimine), using a Brandel cell harvester. The filters were washed (3×4 mL) with buffer, dried, and counted on a γ -counter (Crystal-Packard).

Nonspecific binding was defined with 10 μM of 2-iodomelatonin and represented 10% of the total binding. Competition studies with brain homogenates from the chicken killed at 12 a.m., were also performed in the presence of 30 μM GTP- γ -S or 1 mM MnCl_2 ³⁸ to evaluate the agonist or antagonist profile of the tested compounds.

Data Analysis. Both saturation and competition experiments were analyzed using nonlinear least squares curve fitting or linear regression programs in LOTUS 1.2.3 and GraphPad. K_i values were determined using the Cheng-Prussoff equation.³⁹

References

- Reiter, R. J. Pineal melatonin: Cell biology of its synthesis and of its physiological interactions. *Endocrinol. Rev.* **1991**, *12*, 151–180.
- Reiter, R. J. The pineal and its hormone in the control of the reproduction in the mammals. *Endocrinol. Rev.* **1980**, *1*, 109–131.
- (a) Reiter, R. J. The melatonin rhythm: both a clock and a calendar. *Experientia* **1993**, *49*, 654–664. (b) Pierpaoli, W. The pineal gland: a circadian or seasonal aging clock? *Aging* **1991**, *3*, 99–101. (c) Blehar, M. C.; Lewy, A. J. Seasonal mood disorders: Consensus and controversy. *Psychopharmacol. Bull.* **1990**, *4*, 465–482.
- Waldhauser, F.; Ehrhart, B.; Förster, E. Clinical aspects of the melatonin action: impact of development, aging, and puberty, involvement of melatonin in psychiatric disease and importance of neuroimmunoenocrine interactions. *Experientia* **1993**, *49*, 671–681.
- (a) Maestroni, G. J. M. The immunoneuroendocrine role of melatonin. *J. Pineal Res.* **1993**, *14*, 1–10. (b) Ying, S.-W.; Niles, L. P.; Crocker, C. Human malignant melanoma cells express high-affinity receptors for melatonin: antiproliferative effects of melatonin and 6-chloromelatonin. *Eur. J. Pharmacol. Mol. Pharmacol. Sect.* **1993**, *246*, 89–96.
- (a) Pöggeler, B.; Reiter, R. J.; Tan, D.-X.; Chen, L.-D.; Manchester, L. C. Melatonin, hydroxyl radical-mediated oxidative damage, and aging: A hypothesis. *J. Pineal Res.* **1993**, *14*, 151–158. (b) Grad, B. R.; Rozenzweig, R. The role of melatonin and serotonin in aging: Update. *Psychoneuroendocrinology* **1993**, *18*, 283–295. (c) Schmid, H. A. Decreased melatonin biosynthesis, calcium flux, pineal gland calcification and aging: A hypothetical framework. *Gerontology* **1993**, *39*, 189–199.
- Zawilska, J. B.; Nowak, J. Z. Regulatory mechanisms in melatonin biosynthesis in retina. *Neurochem. Int.* **1992**, *20*, 23–36.
- (a) Utiger, R. D. Melatonin – the hormone of darkness. *N. Engl. J. Med.* **1992**, *327*, 1377–1379. (b) Reiter, R. J. Melatonin: the chemical expression of darkness. *Mol. Cell. Endocrinol.* **1991**, *79*, C153–C158.
- (a) Deguchi, T. Physiology and molecular biology of arylamine N-acetyltransferases. *Biomed. Res.* **1992**, *13*, 231–242. (b) Zatz, M. Does the circadian pacemaker act through cyclic AMP to drive the melatonin rhythm in chick pineal cells. *J. Biol. Rhythms* **1992**, *7*, 301–311.
- (a) Stankov, B.; Fracchini, F.; Reiter, R. J. Melatonin binding sites in the central nervous system. *Brain Res.* **1991**, *16*, 245–256. (b) Krause, D. N.; Dubocovich, M. L. Melatonin receptors. *Annu. Rev. Pharmacol. Toxicol.* **1991**, *31*, 549–568.
- (a) Skene, D. J.; Masson-Pévet, M.; Pévet, P. Characterisation of melatonin binding sites in the pars tuberalis of the European hamster. *J. Neuroendocrinol.* **1992**, *4*, 189–192. (b) Helliwell, R. J. A.; Williams, L. M. Melatonin binding sites in the ovine brain and pituitary: Characterisation during the oestrous cycle. *J. Neuroendocrinol.* **1992**, *4*, 287–294.
- Kennaway, D. J.; Hugel, H. M. Melatonin binding sites: are they receptors? *Mol. Cell. Endocrinol.* **1992**, *88*, C1–C9.
- (a) Morgan, P. J.; Lawson, W.; Davidson, G.; Howell, H. E. Guanine nucleotides regulate the affinity of melatonin receptors on the ovine pars tuberalis. *Neuroendocrinology* **1989**, *50*, 359–362. (b) Chong, N. W. S.; Sugden, D. Guanine nucleotides regulate 2-[¹²⁵I]iodomelatonin binding sites in chick retinal pigment epithelium but not in neuronal retina. *J. Neurochem.* **1991**, *57*, 685–689. (c) Pang, C. S.; Brown, G. M.; Tang, P. L.; Pang, S. F. G-protein linked binding sites in the chicken lung. *Neurosci. Lett.* **1993**, *162*, 17–20.
- Redfern, P. H. Can pharmacological agents be used effectively in the alleviation of Jet-lag? *Drugs* **1992**, *43*, 146–153.
- Guardiola-Lemaître, B.; Lenègre, A.; Porsolt, R. D. Combined effects of diazepam and melatonin in two tests for anxiolytic activity in the mouse. *Pharmacol. Biochem. Behav.* **1992**, *41*, 405–408.
- Durlach-Misteli, C.; Van Ree, J. M. Dopamine and melatonin in the nucleus accumbens may be implicated in the mode of action of antidepressant drugs. *Eur. J. Pharmacol.* **1992**, *217*, 15–21.
- (a) Heward, C. B.; Hadley, M. E. Structure activity relationships of melatonin and related indoleamines. *Life Sci.* **1975**, 1167–1178. (b) Spadoni, G.; Stankov, B.; Duranti, A.; Biella, G.; Lucini, V.; Salvatori, A.; Fracchini, F. 2-Substituted 5-methoxy-N-acetyltryptamines: synthesis, binding affinity for the melatonin receptor, and evaluation of the biological activity. *J. Med. Chem.* **1993**, *36*, 4069–4074. (c) Garrat, P. J.; Vonhoff, S. Mapping the melatonin receptor. 1. The 5-methoxyl group of melatonin is not an essential requirement for biological activity. *BioMed. Chem. Lett.* **1994**, *4*, 1555–1558.
- (a) Dubocovich, M. L.; Shankar, G.; Miockel, M. 2-[¹²⁵I]iodomelatonin labels sites with identical pharmacological characteristics in chicken brain and chicken retina. *Eur. J. Pharmacol.* **1989**, *162*, 289–299. (b) Siuciak, J. A.; Krause, D. N.; Dubocovich, M. L. Quantitative pharmacological analysis of 2-[¹²⁵I]iodomelatonin binding sites in discrete areas of the chicken brain. *J. Neurosci.* **1991**, *11*, 2855–2864.
- (a) Yous, S.; Andrieux, J.; Howell, H. E.; Morgan, P. J.; Renard, P.; Pfeiffer, B.; Lesieur, D.; Guardiola-Lemaître, B. Novel naphthalenic ligands with high affinity for the melatonin receptor. *J. Med. Chem.* **1992**, *35*, 1484–1486. (b) Depreux, P.; Lesieur, D.; Mansour, H. A.; Morgan, P.; Howell, F.; Renard, P.; Caignard, D.-H.; Pfeiffer, B.; Delagrè, P.; Guardiola, B.; Yous, S.; Demarque, A.; Adam, G.; Andrieux, J. Synthesis and structure-activity relationships of novel naphthalenic and bioisosteric related amidic derivatives as melatonin receptor ligands. *J. Med. Chem.* **1994**, *37*, 3231–3239.
- Copinga, S.; Tepper, P. G.; Grol, C. J.; Horn, A. S.; Dubocovich, M. L. 2-Amido-8-methoxytetralins: A series of nonindolic melatonin-like agents. *J. Med. Chem.* **1993**, *36*, 2891–2898.
- Glennon, R. A.; McKenney, J. D.; Lyon, R. A.; Titeler, M. 5-HT₁ and 5-HT₂ binding characteristics of 1-(2,5-dimethoxyphenyl)-2-aminopropane analogues. *J. Med. Chem.* **1986**, *29*, 194–199.
- Yuan, H.; Pang, S. F. 2-[¹²⁵I]iodomelatonin-binding sites in the pigeon brain: binding characteristics, regional distribution and diurnal variation. *J. Endocrinol.* **1991**, *128*, 475–482.
- (a) Glennon, R. A.; Liebowitz, S. M.; Anderson, G. M. Serotonin receptor affinities of psychoactive phenylalkylamine analogues. *J. Med. Chem.* **1980**, *23*, 294–299. (b) Zweig, J. S.; Castagnoli, N., Jr. In vitro O-Demethylation of the psychomimetic amine 1-(2,5-dimethoxy-4-methylphenyl)-2-aminopropane. *J. Med. Chem.* **1977**, *20*, 414–420. (c) Monte, A. P.; Marona-Lewicka, D.; Cozzi, N. V.; Nichols, D. E. Synthesis and pharmacological examination of benzofuran, indan and tetralin analogues of 3,4-(methylenedioxy)amphetamine. *J. Med. Chem.* **1993**, *36*, 3700–3706.
- Einhorn, J.; Demerseman, P.; Rene, L.; Royer, R.; Cavier, R. Recherches sur les dérivés nitrés d'intérêt biologiques- XXVIII-Synthèse et activités contre les microorganismes d'analogues hydrogénés des nitro-2-naphthofuranes. (Study on nitro-derivatives of biological interest. XXVIII. Synthesis of hydrogenated analogs of 2-nitronaphthofurans and activities against microorganisms.) *Eur. J. Med. Chem.* **1983**, *18*, 79–84.
- Brooks, D. S.; Cassone, V. M. Daily and circadian regulation of 2-[¹²⁵I]iodomelatonin in the chick brain. *Endocrinology* **1992**, *131*, 1297–1304.
- Findlay, J.; Eliopoulos, E. Three-dimensional modelling of G protein-linked receptors. *TIPS* **1990**, *11*, 492–499.

- (27) Langlois, M.; Soulier, J. L.; Yang, D.; Bremont, B.; Florac, C.; Rampillon, V.; Giudice, A. Structural analysis of 5-HT₃ receptor antagonists: synthesis and pharmacological activity of various aromatic esters or amides derived from tropane and 1,2,6-trisubstituted piperidine. *Eur. J. Med. Chem.* **1993**, *28*, 869–880.
- (28) Duranti, E.; Stankov, B.; Spadoni, G.; Duranti, A.; Lucini, V.; Capsoni, S.; Biella, G.; Frascini, F. 2-bromomelatonin: synthesis and characterization of a potent melatonin agonist. *Life Sci.* **1992**, *51*, 479–485.
- (29) Jansen, J. M.; Karlén, A.; Grol, C. J.; Hacksell, U. Conformational properties of melatonin and two conformationally restricted agonists: a molecular mechanics and NMR spectroscopy study. *Drug design and discovery*; Harwood Academic Publishers GmbH: New York, 1993; Vol. 10, pp 115–133.
- (30) Garrat, P. J.; Vonhoff, S. Mapping the melatonin receptor. 2. Synthesis and biological activity of indole derived melatonin analogues with restricted conformations of the C-3 amidoethane side-chain. *BioMed. Chem. Lett.* **1994**, *4*, 1559–1564.
- (31) (a) Laitinen, J. T.; Flugge, G.; Saavedra, J. M. Characterization of melatonin receptors in the rat area postrema: modulation of affinity with cations and guanine nucleotides. *Neuroendocrinology* **1990**, *51*, 619–624. (b) Rivkees, S. A.; Carlson, L. L.; Reppert, S. M. G-protein regulation of melatonin receptors in lizard brain. *Proc. Natl. Acad. Sci. U.S.A.* **1989**, *86*, 3882–3886. (c) Chong, N. W. S.; Sugden, D. Thermodynamic analysis of agonist and antagonist binding to the chicken brain melatonin receptor. *Br. J. Pharmacol.* **1994**, *111*, 295–301.
- (32) Hamon, M.; Goetz, C.; Gozlan, H. *CNS Receptor Subtypes: Pharmacological Significance and Clinical Implications*; Mandel, P., DeFeudis, F. V., Eds.; Raven Press: New York, 1983; pp 349–359.
- (33) Langlois, M.; Gaudy, F.; Shen, S.; Brémont, B. Synthesis of new derivatives of 8-OH-DPAT: influence of substitution on the aromatic ring on the pharmacological profile. *BioMed. Chem. Lett.* **1993**, *3*, 2035–2038.
- (34) Morgan, P.; Williams, L.; Davidson, G.; Laxson, W.; Howell, E. Melatonin receptors of the ovine pars tuberalis: characterization and autoradiographical localization. *J. Neuroendocrinol.* **1989**, *1*, 1–4.
- (35) Morgan, P. J.; Lawson, W.; Davidson, G.; Howell, H. E. Melatonin inhibits cyclic AMP production in cultured ovine pars tuberalis cells. *J. Mol. Endocrinol.* **1989**, *3*, R5–R8.
- (36) Lowry, O. H.; Rosebrough, N. J.; Farr, A. L.; Randall, R. J. Protein measurement with the Folin phenel reagent. *J. Biol. Chem.* **1951**, *193*, 265–275.
- (37) Duncan, M. J.; Takahashi, J. B.; 2-[¹²⁵I]iodomelatonin binding sites in hamster brain membranes: Pharmacological characteristics and regional distribution. *Endocrinology* **1988**, *122*, 1825–1833.
- (38) Hall, M. D.; Gozlan, H.; Emerit, M. B.; El Mestikawy, S.; Pichat, L.; Hamon, M. Differentiation of pre- and post-synaptic high affinity serotonin receptor binding sites using physico-chemical parameters and modifying agents. *Neurochem. Res.* **1986**, *11*, 891–912.
- (39) Cheng, Y. C.; Prusoff, W. H. Relation between the inhibition constant K_i and the concentration of inhibitor which causes fifty per cent inhibition (IC₅₀) of an enzymatic reaction. *Biochem. Pharmacol.* **1973**, *22*, 3099–3108.

JM940616X