

RSC Advances



This article can be cited before page numbers have been issued, to do this please use: Z. Li, X. xu, X. Jiang, Y. Li, Z. Yu and X. Zhang, *RSC Adv.*, 2015, DOI: 10.1039/C5RA01649E.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

COMMUNICATION

Facile reduction of aromatic nitro compounds to aromatic amines catalysed by support-free nanoporous silver

Cite this: DOI: 10.1039/x0xx00000x

Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

Zhiwen Li,^a Xiaohong Xu,^a Xiaojian Jiang,^a Yingchun Li,^b Zhixin Yu,^c and Xiaomei Zhang^{a*}

Nanoporous silver was used as the catalyst for the reduction of aromatic nitro compounds even in the presence of some sensitive functional groups under mild conditions with excellent yields. Lesser amount of NaBH₄ was used. The reaction kinetics was studied with the help of UV-visible spectrophotometry.

Nanoporous metals (np-M) made by dealloying have attracted increasing interest due to their broad application in fuel cell technologies, biomolecular sensing, surface-enhanced Raman scattering (SERS), electrocatalysis, and heterogeneous catalysis. Their unique structural properties of mechanical rigidity, nanoscale three-dimensional bicontinuous porous structure, easy functionalization and high corrosion resistance make these unsupported materials to be a good choice for green, sustainable catalysts.¹⁻³ To date, some of np-M have been used in gas phase,^{4,5} liquid phase⁶⁻⁸ and electrochemical^{9,10} catalytic reactions and exhibited effective selectivity and catalytic activity. Compared with gas phase and electrochemical catalytic reactions, investigation about np-M-catalyzed liquid phase catalytic reactions was developed lately (in 2010). Therefore relative works were rare and most of them limited in nanoporous gold (np-Au). Although other np-M was used in a small number of liquid phase catalytic reactions, the results were exciting.¹¹⁻¹⁴ Searching efficient, high-selective np-M-catalyzed reaction systems are necessary due to their unique structural properties mentioned above.

Aromatic amines are important substances in synthetic chemistry and chemical engineering.^{15,16} Reduction of aromatic nitro compounds is one of the main paths to get aromatic amines. Different reductants have been used for this reaction with transition metals as the catalysts. But high temperature and pressure were always required.¹⁷⁻¹⁹ Sodium borohydride (NaBH₄) was a common reductant for hydrogenation. When it was used for reduction reactions with the help of transition metal catalysts, although normal pressure and room temperature was available, substantial molar

excess of NaBH₄ relative to the substrates, usually exceeding 100-fold excess, was needed.²⁰⁻²³ It is important to search suitable catalyst which could increase the catalytic efficiency and decrease the amount of NaBH₄.²⁴

In the present work, we comparatively investigated the catalytic activity of different nanoporous metals for the reduction of aromatic nitro compounds and found that nanoporous silver (np-Ag) exhibiting the highest activity. The aromatic nitro compounds with not only electron-withdrawing groups but also electron-donating groups could be hydrogenated to the corresponding aromatic amines with good yield and high selectivity at room temperature. Compared with other transition metal catalysts, when np-Ag was used as catalyst, only 25-fold excess of NaBH₄ was needed. In addition, np-Ag could be reused at list seven times without the loss of catalytic activity.




Figure 1. SEM image of dealloyed np-Ag a) before catalytic reaction and b) after being used for seven times.

The previously reported fabrication method for np-Ag was by etching magnesium from Ag-Mg alloy.²⁵ In our present case, as

COMMUNICATION

aluminium can melt with Au, Ag, Cu, Pd, therefore different aluminum-base ($\text{Al}_{75}\text{M}_{25}$) alloys were prepared firstly by melting aluminum with other metals (Au, Ag, Cu, Pd). $\text{Al}_{75}\text{Ag}_{25}$ was chose as a representative example to be introduced. Sodium hydroxide solution (NaOH, 10 wt. %) was used to etch the aluminum from the alloy. After the dealloying process the nanoporous silver was got, as showed in Fig. 1a, nanoscale three-dimensional bicontinuous porous structure was obtained and the average ligament size was around 30 nm. In order to check the chemical state of Ag, XPS of nanoporous Ag before the reaction was detected. As can be seen from the XPS spectrum of Ag 3d region, Fig. 2a, two peaks occurred at 368.55 and 374.55 eV were well corresponded with Ag $3d_{5/2}$ and Ag $3d_{3/2}$ binding energies, indicating the metallic nature of np-Ag.²⁶




Figure 2. XPS spectra of nanoporous Ag before (a) and after (b) the reaction.

Nanoporous silver was first used as the catalysts for the reduction of 4-nitrophenol (4-NP) by using NaBH_4 as reductant. Ultraviolet and visible (UV/Vis) spectrophotometer was used to analyze the reaction process. As shown in Fig. 3a, the characteristic absorption peak of 4-NP and 4-aminophenol (4-AP) in the presence of NaBH_4 was appeared at 400 and 300 nm, respectively. During the reaction, the intensity of the characteristic peak of 4-NP decreased, while the absorption peak at 300 nm attributing to 4-AP increased gradually, indicating the successful reduction of 4-NP to 4-AP. In this reaction, NaBH_4 was used in 25-fold excess, so the reaction rate would follow pseudo-first-order kinetics. A linear correlation between $\ln(A_t/A_0)$ and reaction time, which A_t and A_0 represented the absorbance at the fixed intervals and the absorbance at the initial stage, respectively, were got, Fig. 3b. This result further confirmed the reaction is a pseudo-first-order and its rate constant (k) was calculated from the slope to be 0.036 min^{-1} .




Figure 3. Time-dependent UV-visible absorption spectra for the reduction of 4-nitrophenol over nanoporous silver in aqueous media at room temperature, Fig. 2a and plot of $\ln(A_t/A_0)$ versus time for the reduction of 4-nitrophenol over nanoporous silver, Fig. 2b. Reactions were performed using np-Ag 0.05 mmol, 4-NP 20 mM and NaBH_4 500 mM in 10 mL aqueous solution at room temperature.

Table 1. The expressions of different nanoporous metal catalysts in the reaction.

Entry	Catalyst	k / min^{-1}	BET / m^2/g	TOF / min^{-1}
1	np-Ag-Al	0.036	10	1.91
2	np-Ag-Mg	0.019	14	1.02
3	np-Pd-Al	0.033	40	0.48
4	np-Au-Al	0.028	15	1.03
5	np-Cu-Al	0.017	8	1.19

^{a)} Reactions were performed using catalyst 0.05 mmol, 4-NP 20 mM and NaBH_4 500 mM in 10 mL aqueous solution at room temperature.

Then other nanoporous metal catalysts were used to reduce the 4-NP under the same conditions (Table 1, Entry 2-5). Corresponding rate constant (k) of the catalysts was also obtained, and a plot of $\ln(A_t/A_0)$ versus time were shown, (Figs. S1-S4, ESI). Meanwhile, the turnover frequency (TOF) of each catalyst was calculated, (the calculation was showed in ESI). Compared with np-Ag-Al, the nanoporous silver fabricated from the magnesium-silver alloy (np-Ag-Mg) show lower activity. This result would most originate from the difference in their sacrificial metal (Al vs. Mg), which effort the structure and the surface state of the catalysts. At the same mol percent, the nanoporous palladium (np-Pd) exhibited the similar activity to np-Ag-Al, but the TOF was lower as the specific surface area was large. Due to the rapid decomposition of NaBH_4 on the surface of nanoporous gold (np-Au), the activity of np-Au ($k=0.028 \text{ min}^{-1}$) was lower than np-Ag. The nanoporous copper (np-Cu) showed limited activity ($k=0.017 \text{ min}^{-1}$). np-Ag-Al exhibited highest activity among the nanoporous materials we used.


Table 2. The result of 4-nitrophenol reduction catalyzed by nanoporous silver under different reaction conditions

Entry	$C_{4\text{-NP}} / \text{mM}$	$C_{\text{NaBH}_4} / \text{mM}$	k / min^{-1}
1	20	200	0.035
2	20	500	0.036
3	20	1000	0.034
4	20	2000	0.037
5	10	500	0.083
6	5	500	0.150

^{a)} Reactions were performed using catalyst 0.05 mmol, 4-NP and NaBH_4 in 10 mL aqueous solution at room temperature.

To better understand the behaviour of np-Ag-Al and the mechanism of the reaction, different concentration of 4-NP and NaBH_4 were used in the reaction. At the same concentration of 4-NP (20 mM), changing the dosage of NaBH_4 didn't significantly impact the activity. Compared with other Ag-based catalysts,²⁷⁻³⁰ even when NaBH_4 was 10-fold excess, the reaction could also occur smoothly. This result confirmed the higher catalytic activity of np-Ag towards the reduction of 4-nitrophenol. As in the previous research, 100-fold excess of NaBH_4 was usually used. On the other hand, when the concentration of NaBH_4 (500 mM) remained unchanged, the

reaction rate constant decreased with the increase of 4-NP concentration, implying that the reaction rate was in direct proportion to the concentration of 4-NP. In common, the reduction of 4-nitrophenol contains two processes, namely the decomposition of NaBH_4 to produce hydrogen and the hydrogenation of aromatic nitro compounds. According to the above experimental results, it was clearly that the hydrogenation of aromatic nitro compounds was the rate limiting step.



Scheme 1. Proposed reaction pathway for the reduction of nitroarenes.

The way of the hydrogenation was further investigated, to understand the reduction mechanism. Usually, there are two accepted routes for the reduction of aromatic nitro compounds based on the electrochemical model as presented by Haber, namely direct route and condensation route.³¹ As shown in Scheme 1, the direct route was a stepwise hydrogenation process. During this process, the intermediate products, nitrosobenzene (**1aa**) and phenylhydroxylamine (**1aaa**), were occurred. However, in the condensation route, intermediate product was azobenzene. As the two accepted routes had different intermediate products, detecting the intermediate products would be an effective method to explore the possible mechanism. Then, the following experiment was done. When the reduction of nitrobenzene run 0.5 h, the catalyst was separated from the solution and the solution was analyzed by gas chromatography-mass spectrometer (GC-MS). Nitrobenzene (**1a**), nitrosobenzene (**1aa**) and aniline (**1b**) were observed, indicating np-Ag catalyzed nitroarenes reduction following the direct route. It is noteworthy, as showed in scheme 1, the intermediate products in direct route contained nitrosobenzene (**1aa**) and phenylhydroxylamine (**1aaa**). However, in our present case, no phenylhydroxylamine (**1aaa**) was detected. It therefore deduced that the hydrogenation of nitrobenzene (**1a**) to nitrosobenzene (**1aa**) would be the rate limiting step.

Except the hydrogenation, electron transfer was also speculated. According to the work before, the reaction of 4-NP to 4-AP was a thermodynamically feasible process considering the BH_4^- (-1.33 V vs. NHE) and 4-NP (-0.76 V vs. NHE). However, it is a kinetically restricted process, (the reaction could not happen even in two days' time) in the absence of a catalyst.^{27, 28} Usually it is believed that the electron transfer from BH_4^- to 4-NP was relayed by the metal surface.^{29, 30} As the NaBH_4 dissociate on the surface of np-Ag, electron would transfer on the surface of np-Ag. When the aromatic nitro compounds was adsorbed on the surface of np-Ag, the electron would further transferred from the silver surface to the nitro compounds.

The substrate generality of the catalyst was test by using a series of aromatic nitro compounds (Table 3). Nitrobenzene without substituent group could be reduced into aniline easily (Entry 1). Aromatic nitro compounds with electron-withdrawing group and

electron-donating group could also be reduced with high selectivity. For nitrophenol with the same substituent at different position, the 3-nitrophenol showed higher activity than 2-nitrophenol and 4-nitrophenol. The reason may be from the stability of the intermediate product nitrophenolate ion. As the resonance effect of 3-nitrophenol is lowest among the three nitrophenols, so the 3-NP was easier reduced. These results were in accordance with the previous literatures.^{32, 33} Similar results were also found for the reduction of 2, 3, and 4-toluidine (Entry 5-7). The amino on the benzene ring did not affect the reduction of nitro group and the 4-phenylenediamine was achieved in 96% yield (Entry 8). np-Ag can also be applied to reduction of nitrobenzene containing halogen or unsaturated substitutes. For example, 1-iodo-4-nitrobenzene was chemoselectively reduced to 4-iodoaniline with high selectivity (Entry 9). No dehalogenation side-product was detected. The nitroarenes with methoxycarbonyl and nitrile substitutes were reduced to methyl-4-aminobenzoate and 4-aminobenzonitrile, respectively, in high yield (Entry 10, 11).

Table 3. Reduction of different aromatic nitro compounds catalyzed by nanoporous silver


Entry	Substrate	Product	Time / min	Yield / %
1			90	99
2			105	99
3			60	98
4			60	99
5			420	91
6			180	95
7			240	95
8			360	96
9			720	90
10			480	95
11			360	95

^{a)} Reactions were performed using catalyst 0.05 mmol, aromatic nitro compounds 20 mM and NaBH_4 500 mM in 10 mL aqueous solution at room temperature.

As the high stability was one of the most important properties of heterogeneous catalysts, the recycling capacity and reusability of np-Ag were also tested. After the reaction, np-Ag was separated from the reaction mixture by filtrating and the catalyst was washed by

ethanol and water. Then np-Ag was reused without further purification. From scheme 2, it can be seen that the catalyst could be reused even seven times without reducing the catalytic activity. Fig. 1b is the SEM image of the recovered catalyst after seven uses. Compared to the fresh one (Fig. 1a), little change in porous morphology and feature dimension was observed, but the ligament size coarsened from 30 nm before the reaction to around 40 nm. After the reaction the XPS spectrum of np-Ag was also detected, Fig. 2b. The binding energies of Ag 3d_{5/2} and Ag 3d_{3/2} were appeared at 368.5 and 374.5 eV, which implied that the surface status did not change after the reaction. To clarify whether the np-Ag catalyst had been leached into the reaction mixture or not, the reaction solution was also tested by inductivity coupled plasma (ICP-AES) analysis when the conversion of 4-NP was about 50%. The rate was lower than the detection limits (<0.02 ppm). No leaching of silver was detected. These evidences proved that np-Ag was a green and sustainable catalyst for the reduction of aromatic nitro compounds.

Table 4. Reusability studies of np-Ag catalyst.



Reaction Cycle ^{a)}	1	2	3	4	5	6	7
Yield/%	99	99	98	99	97	98	97
Time/min	120	120	120	120	120	135	135

^{a)} Reactions were performed using catalyst 0.1 mmol, 4-NP (1 mmol) and NaBH₄ (25 mmol) in 10 mL aqueous solution at room temperature.

Conclusions

Briefly conclusion, we realized the reduction of nitroarenes with lesser amount of NaBH₄ reductant under the mild reaction condition by using nanoporous silver as a high active and selective catalyst. The aromatic nitro compounds containing both electro-withdrawing and electro-donating substituents could be reduced with high yields. The catalyst can be reused several times without evident loss of its catalytic activity. As a cost-effective nanoporous material, the well reduction property exhibited by nanoporous silver here will further arouse the investigated interest of nanoporous silver in wider green organic synthesis field.

Acknowledgements

Financial support from the National Science Foundation of China (21472117 and 21176144) was acknowledged.

Notes and references

^{a)} School of Chemistry and Chemical Engineering, Shandong University, Jinan 250100, China.

^{b)} School of Pharmacy, Shihezi University, Shihezi 832000, China.

^{c)} Department of Petroleum Engineering, University of Stavanger, Stavanger 4036, Norway.

† Footnotes should appear here. These might include comments relevant to but not central to the matter under discussion, limited experimental and spectral data, and crystallographic data.


Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/c3000000x/

- 1 Y. Ding, M. Chen, *MRS Bull.* 2009, **34**, 569-576.
- 2 X. Zhang, Y. Ding, *Catal. Sci. Technol.* 2013, **3**, 2862-2868.
- 3 Y. Yamamoto, *Tetrahedron* 2014, **70**, 2305-2317.
- 4 C. Xu, J. Su, X. Xu, P. Liu, H. Zhao, F. Tian, Y. Ding, *J. Am. Chem. Soc.* 2007, **129**, 42-43.
- 5 A. Wittstock, V. Zielasek, J. Biener, C. M. Friend, M. Bäumer, *Science* 2010, **327**, 319-322.
- 6 H. Yin, C. Zhou, C. Xu, P. Liu, X. Xu, Y. Ding, *J. Phys. Chem. C* 2008, **112**, 9673-9678.
- 7 N. Asao, Y. Ishikawa, N. Hatakeyama, Y. Yamamoto Menggenbateer, M. Chen, W. Zhang, A. Inoue, *Angew. Chem. Int. Ed.* 2010, **49**, 10093-10095.
- 8 M. Yan, T. Jin, Y. Ishikawa, T. Minato, T. Fujita, L. Chen, M. Bao, N. Asao, M. Chen, Y. Yamamoto, *J. Am. Chem. Soc.* 2012, **134**, 17536-17542.
- 9 R. Wang, C. Wang, W. Cai, Y. Ding, *Adv. Mater.* 2010, **22**, 1845-1848.
- 10 R. Wang, C. Xu, X. Bi, Y. Ding, *Energ. Environ. Sci.* 2012, **5**, 5281-5286.
- 11 S. Tanaka, T. Kaneko, N. Asao, Y. Yamamoto, M. Chen, W. Zhang, A. Inoue, *Chem. Commun.* 2011, **47**, 5985-5987.
- 12 T. Kaneko, S. Tanaka, N. Asao, Y. Yamamoto, M. Chen, W. Zhang, A. Inoue, *Adv. Synth. Catal.* 2011, **353**, 2927-2932.
- 13 Z. Li, S. Lin, L. Ji, Z. Zhang, X. Zhang, Y. Ding, *Catal. Sci. Technol.* 2014, **4**, 1734-1737.
- 14 Z. Li, C. Zhang, J. Tian, Z. Zhang, X. Zhang, Y. Ding, *Catal. Commun.* 2014, **53**, 53-56.
- 15 S. Lawrence, *Amines: Synthesis, properties and applications*, Cambridge University Press, Cambridge, 2004.
- 16 E. Burkhardt, K. Matos, *Chem. Rev.* 2006, **106**, 2617-2650.
- 17 I. Nakamura, Y. Yamanoi, T. Imaoka, K. Yamamoto, H. Nishihara, *Angew. Chem. Int. Ed.* 2011, **50**, 5830-5833.
- 18 N. Sakai, K. Fujii, S. Nabeshima, R. Ikeda, T. Konakahara, *Chem. Commun.* 2010, **46**, 3173-3175.
- 19 R. V. Jagadeesh, G. Wienhöfer, F. A. Westerhaus, A.-E. Surkus, H. Junge, K. Junge, M. Beller, *Chem. Eur. J.* 2011, **17**, 14375-14379.
- 20 H. Wei and Y. Lu, *Chem. Asian J.* 2012, **7**, 680-683.
- 21 K. Kuroda, T. Ishida, M. Haruta, *J. Mol. Catal. A-Chem.* 2009, **298**, 7-11.
- 22 K. Layek, M. Lakshmi-Kantam, M. Shirai, D. Nishio-Hamane, T. Sasaki, M. Maheswaran, *Green Chem.* 2012, **14**, 3164-3174.
- 23 S. Saha, A. Pal, S. Kundu, S. Basu, T. Pal, *Langmuir* 2010, **26**, 2885-2893.
- 24 Z. Zhang, C. Shao, Y. Sun, J. Mu, M. Zhang, P. Zhang, Z. Guo, P. Liang, C. Wang, Y. Liu, *J. Mater. Chem.* 2012, **22**, 1387-1395.
- 25 T. Kou, D. Li, C. Zhang, Z. Zhang, H. Yang, *J. Mol. Catal. A-Chem.* 2014, **382**, 55-63.
- 26 F. Kelly, J. Johnston, *ACS Appl. Mater. Interfaces*, 2011, **3**, 1083-1092.
- 27 S. Saha, A. Pal, S. Kundu, S. Basu, T. Pal, *Langmuir* 2010, **26**, 2885-2893.

Journal Name

- 28 J. Huang, S. Vongehr, S. Tang, H. Lu, X. Meng, *J. Phys. Chem. C* 2010, **114**, 15005-15010.
- 29 P. Zhang, C. Shao, Z. Zhang, M. Zhang, J. Mu, Z. Guo, Y. Liu, *Nanoscale* 2011, **3**, 3357-3363.
- 30 Z. Zhang, C. Shao, Y. Sun, J. Mu, M. Zhang, P. Zhang, Z. Guo, P. Liang, C. Wang, Y. Liu, *J. Mater. Chem.* 2012, **22**, 1387-1395.
- 31 F. Haber, *Z. Elektrochem.* 1898, **22**, 506-514.
- 32 Y. Choi, H. Bae, E. Seo, S. Jang, K. H. Park, B. Kim, *J. Mater. Chem.* 2011, **21**, 15431-15436.
- 33 S. Jana, S. Ghosh, S. Nath, S. Pande, S. Praharaj, S. Panigrahi, S. Basu, T. Endo, T. Pal, *Appl. Catal. A-gen.* 2006, **313**, 41-48.

Facile reduction of aromatic
nitro compounds to aromatic
amines catalysed by
support-free nanoporous silver



RSC Advances

Zhiwen Li, Xiaohong Xu,
Xiaojian Jiang, Yingchun Li,
Zhixin Yu, and Xiaomei Zhang*

Reduction of aromatic nitro compounds to aromatic amines was realized with excellent yield by using nanoporous silver as sustainable, heterogeneous catalyst.