

of picric acid. The solid was isolated as described below (yields).

*Anal.* Calcd. for  $C_{13}H_{12}O_7N_4$ : N, 16.7. Found: N, 16.7.

**Wolff-Kishner.**—A yield of 7.8 g. (83.5%) of the semicarbazide hydrochloride of 2-acetylpyridine was obtained by the method of Woodward, *et al.*,<sup>6</sup> using ten equivalents of hydrochloric acid, m. p. 202° (uncor.).

*Anal.* Calcd. for  $C_8H_{10}ON_4 \cdot HCl$ : N, 26.2; Cl, 16.6. Found: N, 26.6; Cl, 17.0, 16.9.

The reduction was carried out by heating the semicarbazide hydrochloride with eight equivalents of sodium ethylate in an oil-bath at 180°. The distillate was caught in a 50% alcoholic solution of picric acid.

**Huang-Minlon.**<sup>7</sup>—A mixture of 2-acetylpyridine (0.06 mole), 5 ml. of 85% hydrazine hydrate, 5 g. of sodium hydroxide, and 80 ml. of diethylene glycol was heated for six hours. No attempt was made to remove the water. The solution was cooled, extracted with benzene, and converted into the picrate.

**Yields.**—No attempt was made to obtain maximum yields. The solid picrates were isolated from the alcoholic solution, dissolved in hot acetone, treated with decolorizing carbon, crystallized by cooling and finally recrystallized from alcohol. From the weights of the crude products, and the purified picrates the following estimates of yields were made: Clemmensen 80%, Wolff-Kishner 50% and Huang-Minlon 65%.

**Melting Points and Fusion Analysis.**—Using a copper block<sup>8</sup> and raising the temperature at a rate no faster than 1–2° per minute each picrate, and all possible combinations of picrate mixtures melted at 107–107.5° (uncor.). When the picrate of the Clemmensen reduction product was mixed with (a) the picrate of 2-acetylpyridine a depression of 20° in the melting point was noted; (b) the picrate of the hydrogenation product of Gregg and Craig the melting point was 107.5°. A fusion analysis showed these last pair to be identical also, for no eutectic melt was noted at the boundary. This in contrast with the eutectic melt shown in the fusion analysis of the first pair.

**Acknowledgments.**—I should like to acknowledge thanks to Robert Seiwald for the analytical data, and Dr. Luthy for a sample of the picrate of the hydrogenation product of 2-vinylpyridine and for the fusion analysis.

(6) C. F. Woodward, A. Eisner and P. G. Hains, *THIS JOURNAL*, **66**, 911 (1944).

(7) Huang-Minlon, *ibid.*, **68**, 2487 (1946).

(8) F. W. Bergstrom, *Ind. Eng. Chem., Anal. Ed.*, **9**, 340 (1937).

UNIVERSITY OF SAN FRANCISCO  
SAN FRANCISCO 17, CALIFORNIA RECEIVED MAY 14, 1949

## The Reaction of Propyl Disulfide with Decyl Mercaptan

By GEORGE GORIN,<sup>1</sup> GREGG DOUGHERTY AND ARTHUR V. TOBOLSKY

It is known that thioglycolic acid reacts with cystine in solution to give cysteine<sup>2</sup>; indeed, a quantitative study of the reaction has been made.<sup>3</sup> We wish to report some results which indicate that a similar reaction occurs between simple alkyl disulfides and mercaptans.

Mixtures of propyl disulfide and decyl mercaptan were heated in sealed Pyrex glass tubes for varying lengths of time. The samples were then

(1) Thiokol Corporation Fellow 1946–1948.

(2) Goddard and Michaelis, *J. Biol. Chem.*, **106**, 605 (1934).

(3) Bersin and Steudel, *Ber.*, **71B**, 1015 (1938).

cooled, the mercaptans were titrated with standard silver nitrate solution to an amperometric end-point, and the precipitated silver mercaptides weighed.<sup>4</sup> It was found that the number of moles of total mercaptan did not change during the reaction, but that the weight of the precipitate decreased continuously to an equilibrium value. It was thus indicated that there had taken place a mole per mole exchange of propyl for decyl mercaptan. The data obtained in a run at 138–139° are given in Table I.

TABLE I  
MIXTURES OF PROPYL DISULFIDE AND DECYL MERCAPTAN  
HEATED AT 138–139°

Time, hr.	Moles SH found	Av. mol. wt. of AgSR	% PrSH
0		280	1
3	275	268	13
6	282	260	21
9	276	253	29
12	278	248	34
36	277	243	39
62	277	226	56

At 180° the reaction proceeded more quickly, and after a few hours heating considerable amounts of propyl mercaptan could be isolated directly by distillation of the cooled reaction mixture.

We believe the reaction proceeds by a stepwise exchange which at first gives rise to a mixed disulfide. We made no attempt to isolate this compound but hope to obtain evidence on this question in future work.

### Experimental

**Material.**—The propyl disulfide was an Eastman Kodak Co. "white label" product which had been redistilled at least once, b. p. 93.3–94.6° at 30 mm. The decyl mercaptan was a Connecticut Hard Rubber Co. product, which had been redistilled at least once; b. p. 102–106° at 15 mm.,  $n_D^{20}$  1.4534, molecular weight<sup>4</sup> 171 (calculated 174).

**Approximate Rate Measurements.**—Equal volumes (1.00 ml.) of a mixture of one mole of propyl disulfide and one mole decyl mercaptan were sealed in several tubes of nearly equal volumes (1.5 ml.) and heated in a vapor bath for some time (*p*-xylene used at 138–139°; *p*-cymene at 176°). At the end of the appropriate period each tube was withdrawn and cooled, and the contents washed into 375 ml. of 95% alcohol containing 25 ml. of 0.2 *M* alcoholic ammonium acetate as supporting electrolyte. The solution was then titrated with standard 0.1 *N* alcoholic silver nitrate to an amperometric end-point. The precipitate was transferred to Gooch-type filters consisting of a layer of asbestos placed over a medium Pyrex glass fritted filter disk, and dried *in vacuo* at 60–70° to constant weight.

**Acknowledgments.**—The authors are indebted to Dr. F. O. Davis and Mr. E. M. Fettes of the Thiokol Corporation for their active interest and valuable advice.

FRICK CHEMICAL LABORATORY  
PRINCETON UNIVERSITY  
PRINCETON, NEW JERSEY

RECEIVED MAY 7, 1949

(4) Laitinen, O'Brien and Nelson, *Ind. Eng. Chem., Anal. Ed.*, **18**, 471 (1946).