

### Preliminary communication

---

## AN ORGANOMETALLIC ROUTE TO ( $\pm$ )-3-*p*-TOLUENESULFINYL-2-PYRONE

GARY H. POSNER\* and WAYNE HARRISON

*Department of Chemistry, The Johns Hopkins University, Baltimore, Maryland 21218 (U.S.A.)*

(Received July 17th, 1984)

### Summary


3-Bromo-2-pyrone reacts with dimethylcopperlithium and then with *p*-tolyl *p*-toluenethiosulfonate to form 3-*p*-toluenethio-2-pyrone. Organocopper species are likely intermediates in this reaction. Oxidation of the sulfide produces ( $\pm$ )-3-*p*-toluenesulfinyl-2-pyrone. The overall process involves an important and useful umpolung, a reversal of reactivity at C(3) of the pyrone ring from an electrophilic center to a nucleophilic center.

---

As part of our research program on the use of enantiomerically pure unsaturated ketone and lactone sulfoxides for asymmetric carbon—carbon bond formation with various organometallic reagents [1], we required (+)- or (–)-3-*p*-toluenesulfinyl-2-pyrone (3). Because even racemic pyrone sulfoxide 3 had not been made before, we chose to prepare the racemate and to investigate its chemistry before attempting the more difficult task of preparing enantiomerically pure (+)- or (–)-3. We report here a reliable and very short (3-step, 2-pot) synthesis of ( $\pm$ )-3 involving organocopper intermediates.

3-Bromo-2-pyrone (1a) was prepared via bromination—dehydrobromination of 2-pyrone [2] using a slightly modified (40 h instead of 24 h reflux, 1.4/1 molar ratio of pyrone/carbon tetrachloride; 59% overall yield from 2-pyrone) literature procedure [3]. Attempts at direct bromine  $\rightarrow$  lithium exchange [4], (i.e. at formation of 3-lithiopyrone 1b) using *n*-butyllithium at  $-78^\circ\text{C}$  failed, giving only what appeared to be *n*-butyl addition products [5]. In contrast, bromine  $\rightarrow$  copper exchange was highly successful.

Treating 3-bromo-2-pyrone (1a) with 2 equivalents of dimethylcopperlithium in diethyl ether at  $-78^\circ\text{C}$  led, after 2 h, to complete bromine  $\rightarrow$  copper exchange; hydrolytic work-up gave 2-pyrone in good yield, with less than 5% of the coupling product [6] 3-methyl-2-pyrone being formed. A mechanistic




(1a, X = Br; 1b, X = Li;

2, X = *p*-TolS;

3, X = *p*-TolS )


rationalization of this result is shown in Scheme 1 involving transient copper(III) intermediate 4 which apparently undergoes selective reductive elimination of ethane (rather than of 3-methyl-2-pyrone) to form a 3-cuprio-2-pyrone (e.g. 5); the excess dimethylcopperlithium might react with cuprio-pyrone 5 to form a more reactive cuprate species (e.g. 6) and possibly methylcopper (a yellow-green precipitate is initially observed) [7]. Similar bromine  $\rightarrow$  copper exchanges instead of carbon-carbon couplings are known during reaction of organocopper reagents with some  $\alpha$ -bromoketones [8],  $\alpha$ -bromoacrylate esters [9], and alkenyl bromides [10]; it has been possible in these cases to intercept the newly-formed organocopper species with a limited number of electrophiles either intermolecularly [8,9] or intramolecularly [10].



SCHEME 1

When 3-bromo-2-pyrone (1a) reacts with dimethylcopperlithium at  $-78^{\circ}\text{C}$  for 3 h and then with *p*-tolyl *p*-toluenethiosulfonate [11] at  $-30^{\circ}\text{C}$  for 2 h, a coupling reaction occurs to form 3-*p*-tolylthio-2-pyrone (2,  $^1\text{H}$ , NMR ( $\text{CDCl}_3$ ):  $\delta$  6.54 (dd, 1H,  $H_4$ )) isolated after preparative TLC in 49% yield (based on bromopyrone 1a, eq. 1). Standard oxidation of sulfide 2 produces the desired, racemic, crystalline, lactone sulfoxide 3 (m.p.  $112\text{--}113^{\circ}\text{C}$ ;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  8.08 (dd, 1H,  $J_{4,5}$  6.7 Hz,  $J_{4,6}$  2.1 Hz,  $H_4$ ). Analysis: found: C, 61.52; H, 4.30; S, 13.69;  $\text{C}_{12}\text{H}_{10}\text{O}_3\text{S}$  calcd.: C, 61.60; H, 4.31; S, 13.61%).

It is noteworthy that a C(3)-tolylthio group causes an upfield chemical shift of H(4) comparable to that caused by a C(3)-methoxy group, whereas a C(3)-tolylsulfinyl group causes a downfield chemical shift of H(4) comparable to that caused by a C(3)-ethoxycarbonyl group [12].



These significant results represent an important and useful umpolung [13] or reversal of reactivity at C(3) of bromopyrone 1a from an electrophilic center to a nucleophilic center, thereby allowing regiospecific attachment of a sulfur substituent as an electrophile. Indeed C(3) attachment of other electrophiles should be possible as well; we have had some initial success with introducing an allyl group.

We are actively studying the chemistry of 3-sulfinyl-2-pyrone (3) as well as the reactivity of the intermediate organocopper species (e.g. 5, 6) toward various electrophiles, and a full report will appear in due course.

**Acknowledgement.** We thank the National Science Foundation for financial assistance in the form of a research grant (CHE-83-12161) and a grant (PCM83-03176) toward purchase of a departmental 400 MHz NMR spectrometer. We also thank the NIH for contributing toward purchase of the NMR spectrometer (grant 1 S10 RR01934).

## References

- (a) G.H. Posner, T.P. Kogan, S.R. Haines, and L. Frye, *Tetrahedron Lett.*, 25 (1984) 2627; (b) G.H. Posner, T.P. Kogan, and M. Hulce, *ibid.*, 25 (1984) 383; (c) G.H. Posner and M. Hulce, *ibid.*, 25 (1984) 379; (d) G.H. Posner, L.L. Frye, and M. Hulce, *Tetrahedron*, 40 (1984) 401.
- Prepared from commercially available 5,6-dihydro-2-pyrone according to M. Nakagawa, M. Tonozuka, M. Obi, M. Kinchi, T. Hino, and Y. Ban, *Synthesis*, (1974) 510.
- W.H. Pirkle and M. Dines, *J. Org. Chem.*, 34 (1969) 2239.
- R.G. Jones and H. Gilman, *Org. Reactions*, 6 (1951) 339.

- 5 For a study of addition of Grignard reagents to 2-pyrones see P. Lhoste, M. Moreau and J. Dreux, *Tetrahedron*, 40 (1984) 1551, and ref. therein.
- 6 G.H. Posner, *Org. Reactions*, 22 (1975) 253.
- 7 G.H. Posner, *An Introduction to Synthesis Using Organocopper Reagents*, John Wiley, New York, 1980.
- 8 (a) G.H. Posner and J.J. Sterling, *J. Am. Chem. Soc.*, 95 (1973) 3076; see also (b) A.E. Greene, J.-P. Lansard, J.-L. Luche, and C. Petrier, *J. Org. Chem.*, 48 (1983) 4763; (c) G.R. Clark, J. Lin and M. Nikaido, *Tetrahedron Lett.*, 25 (1984) 2645.
- 9 (a) J.P. Marino and D.M. Floyd, *J. Am. Chem. Soc.*, 96 (1974) 7138; (b) J.P. Marino and D.M. Floyd, *Tetrahedron Lett.*, (1975) 3897, 3901; (c) See also J. Klein and R. Levene, *J. Am. Chem. Soc.*, 94 (1972) 2520.
- 10 (a) E.J. Corey and I. Kuwajima, *J. Am. Chem. Soc.*, 92 (1970) 395; (b) E.J. Corey, M. Narisada, T. Hiroaka, and R.A. Ellison, *ibid.*, 92 (1970) 396.
- 11 Prepared by *m*-chloroperbenzoic acid oxidation of di-*p*-tolyl disulfide.
- 12 W.H. Pirkle and W.V. Turner, *J. Org. Chem.*, 40 (1975) 1616.
- 13 D. Seebach, *Angew. Chem. Int. Ed. Engl.*, 18 (1979) 239.