

Note

Reductive elimination of C–CN compounds from nickel complexes as a reverse reaction for oxidative addition of benzonitriles to nickel

Takakazu Yamamoto*, Isao Yamaguchi, Mahmut Ablá

Chemical Resources Laboratory, Tokyo Institute of Technology, 4259 Nagatsuta, Midori-ku, Yokohama 226-8503, Japan

Received 18 November 2002; received in revised form 5 February 2003; accepted 5 February 2003

Abstract

Treatment of (bpy)Ni(CN)(C₆H₄CN-*o* or -*m*) with acrylonitrile led to reductive elimination of CN–C₆H₄CN from the complex. The reductive elimination proceeded much faster for the C₆H₄CN-*m* complex.

© 2003 Elsevier Science B.V. All rights reserved.


Keywords: Nickel; Reductive elimination; Oxidative addition; C–CN compound

1. Introduction

Oxidative addition of nitriles to transition metal complexes is recently calling attention [1–3]; we have also been concerned with the oxidative addition [2].

On the other hand, reductive elimination is one of most fundamental organometallic processes. In contrast to many reports on reductive elimination of R–R [4,5], Ar–Ar [6], and C–X (X = hetero-atom [7]), report on reductive elimination of C–CN compounds, the reverse reaction of the oxidative addition of nitrile, is limited, although it is considered a key step in organic synthetic reactions concerned with nitriles [3,8]. Recently, Lewis-acid-accelerated reductive elimination of R–CN from (diphosphine)Pd(R)(CN) has been reported [9]. However, reductive elimination of Ar–CN has not been reported due to less examples of the cyano(aryl)metal complex, LnM(CN)(Ar). In addition, the M–Ar bond has usually higher stability than the M–R bond, and reversibility of the oxidative addition and reductive elimination of ArCN to and from metal complexes has not been clarified. We have previously reported preparation of cyanoaryl nickel complexes **1** and **2** according to oxidative addition of the corresponding benzonitriles to a zerovalent nickel complex, Ni(bpy)(cod)

(bpy = 2,2'-bipyridyl; cod = 1,5-cyclooctadiene) [2].



It has been reported that coordination of electron-withdrawing molecules such as acrylonitrile [5] and protic acids [6] to Ni promote reductive elimination of R–R from (bpy)NiR₂ [5] and Ar–Ar from (bpy)NiAr₂ [6], respectively. We now report similar effect of acrylonitrile and hydrogen chloride on the reductive elimination reaction of the (bpy)Ni(CN)(aryl) type complexes.



2. Results and discussion

We first checked the generality of the oxidative addition of benzonitriles to the Ni(0)-bpy complex by examining reactions of benzonitrile and *m*-dicyanobenzene.

Reaction of Ni(bpy)(cod) with benzonitrile and *m*-dicyanobenzene also caused oxidative addition of CN–C₆H₄R (R = H, *m*-CN) to give (bpy)Ni(CN)(C₆H₄R) (**3** and **4**) as orange solids in 84% yields, respectively Eq. (1).

* Corresponding author. Tel.: +81-459245220; fax: +81-459245276.

E-mail address: tyamamot@res.titech.ac.jp (T. Yamamoto).



IR spectrum of **3** showed a peak due to $\nu(\text{C}\equiv\text{N})$ at a lower frequency (2110 cm^{-1}) than that of benzonitrile (2228 cm^{-1}). $\nu(\text{C}\equiv\text{N})$ peaks of **4** appeared at 2219 and 2116 cm^{-1} , which were shifted to a lower frequency from that of *m*-dicyanobenzene (2235 cm^{-1}), similar to the cases of **1** and **2**.

Fig. 1 shows X-ray crystal structure of **3**. The bond distances between Ni- and bpy-nitrogen ($1.944(4)$ – $1.959(4)$) are comparable with those found in the other square-planar nickel complexes [10]. The N–Ni–N bond angle of **3** (82.5°) is similar to those of **1** (83.0°) and **2** (82.5°). Complexes **1**, **2**, and **4** were stable under air, however, **3** was somewhat unstable under air and was stored under N_2 .

Reaction of **1** with acrylonitrile (excess) caused reductive elimination of *o*-dicyanobenzene (isolated yield = 0.3 mol mol^{-1} **1**) that was isolated by using HPLC (Eq. (2)). Determination of the yield by gas chromatography was not possible due to its high boiling point of the product.

Treatment of **4** with acrylonitrile also leads to reductive elimination to provide *m*-dicyanobenzene (isolated yield = 0.3 mol mol^{-1} **4**). The nickel complex is considered to be converted into Ni(acrylonitrile)(bpy) after the reductive elimination, similar to the case of reductive elimination reaction of $\text{NiR}_2(\text{bpy})$ caused by acrylonitrile [5a].

According to the reductive elimination, the UV–vis spectrum of an acetone solution containing **1** and acrylonitrile changes with time as depicted in Fig. 2. The change of the absorbance at 407 nm with time t obeys the first-order rate law with a rate constant of $29 \times 10^{-6}\text{ s}^{-1}$ at 35°C . The pseudo-first-order rate constant k_{obs} increases proportionally with the concentration of acrylonitrile, indicating that the rate of the reductive elimination can be expressed by the second-order rate law.

$$\text{Rate} = k[\mathbf{1}][\text{CH}_2=\text{CHCN}] \quad (3)$$

The k value of $29 \times 10^{-6}\text{ s}^{-1}\text{M}^{-1}$ at 35°C is by




Fig. 1. Molecular structure of $(\text{bpy})\text{Ni}(\text{CN})(\text{Ph})$ (**3**). Selected bond lengths (\AA) and angles ($^\circ$): Ni(1)–N(2) $1.944(4)$, Ni(1)–N(3) $1.959(4)$, Ni(1)–C(1) $1.845(6)$, Ni(1)–C(2) $1.890(5)$, N(1)–C(1) $1.139(6)$, N(2)–Ni(1)–N(3) $82.5(2)$, N(2)–Ni(1)–C(1) $175.0(2)$, N(3)–Ni(1)–C(1) $97.3(3)$, N(3)–Ni(1)–C(1) $172.2(2)$, Ni(3)–C(2)–C(3) $121.7(5)$.




Fig. 2. Change in UV-vis spectrum during the reaction of **1** with acrylonitrile in acetone at 35 °C. $[\text{CH}_2=\text{CHCN}] = 1.0 \text{ M}$. Under N_2 .

about 10^3 times smaller than that obtained for the reductive elimination reaction of $\text{NiEt}_2(\text{bpy})$ enhanced by electron-withdrawing olefins [5,6], due to higher stability of **1** than $\text{NiEt}_2(\text{bpy})$.

The temperature dependence of the k value for the reaction with acrylonitrile gives an Arrhenius activation energy of 107 kJ mol^{-1} .

Similar reaction of **4** with acrylonitrile caused a shift of the UV-vis peak to a longer wavelength when the reductive elimination of $m\text{-CN-C}_6\text{H}_4\text{CN}$ proceeded, as shown in Fig. 3. In this case, the UV-vis spectrum changes did not obey first-order rate law.

Scheme 1 shows a plausible reaction pathway for the acrylonitrile promoted reductive elimination of $\text{CN-C}_6\text{H}_4\text{CN}$ from $(\text{bpy})\text{Ni}(\text{CN})(\text{C}_6\text{H}_4\text{CN})$.

The rate of the reductive elimination of $\text{CN-C}_6\text{H}_4\text{CN}$ from **4** was by about two orders of magnitude faster than that from **1** as judged from the UV-vis change.




Fig. 3. Changes in UV-vis spectrum during the reaction of **4** with acrylonitrile in acetone at 35 °C. $[\text{CH}_2=\text{CHCN}] = 1.0 \text{ M}$. Under N_2 .

Slower reductive elimination in **1** may be related to known *o*-substitution effect for stabilizing Ni-aryl compounds [11] and/or blocking of the coordination site for acrylonitrile by the *o*-CN group. Preliminary experiments indicated that hydrochloric acid also led to the reductive elimination of $o\text{-CN-C}_6\text{H}_4\text{CN}$ from **1** (yield = 70%), similar to the case of $\text{Ni}(\text{aryl})_2(\text{bpy})$ [6].


3. Experimental

All reactions and manipulations were carried out under inert gas using standard Schlenk techniques. All solvents were dried and distilled prior to use. IR and NMR spectra were recorded on a JASCO-IR 810 spectrophotometer and JEOL EX-400 spectrometer, respectively. Elemental analyses were carried out with a Yanagimoto Type MT-2 CHN autocorder.

3.1. Synthesis of complex 3

A yellow THF solution (15 ml) containing $\text{Ni}(\text{cod})_2$ (0.23 g, 0.82 mmol) and 0.1 ml of cod turned to violet when bpy (0.13 g, 0.82 mmol) was added. After stirring for 2 h at room temperature (r.t.), benzonitrile (0.40 cm^3 , 3.9 mmol) was added, and the reaction mixture was stirred for 4 days. The color of the reaction mixture turned to pale orange with formation of an orange precipitate. After addition of hexane (20 cm^3), the solid was separated by filtration, washed with diethyl ether, and dried under vacuum to give **3** as an orange powder (0.22 g, 84%). Crystals for microanalysis and X-ray crystallographic analysis were obtained by recrystallization from acetone at $-20 \text{ }^\circ\text{C}$. Element anal. Found: C, 64.00; H, 3.82; N, 13.51. Calc. for $(\text{bpy})\text{Ni}(\text{CN})(\text{C}_6\text{H}_5)$: C, 64.21; H, 4.12; N, 13.21.

Complex **4** was synthesized in a similar manner by using 0.57 g (2.1 mmol) of $\text{Ni}(\text{cod})_2$, 0.51 ml of cod, 0.33 g (2.1 mmol) of bpy, and 0.27 g (2.1 mmol) of *m*-dicyanobenzene in 5 ml of DMSO at r.t. An orange powder of **4** was obtained in 84% yield. $^1\text{H-NMR}$ ($\text{DMSO-}d_6$): $\delta = 8.93$ (1H, 6-H of bpy); 8.48 (2H, 3- and 3'-H of bpy); 8.26 and 8.20 (2H, 4- and 4'-H of bpy); 7.84 (m, 2H, *m*- $\text{C}_6\text{H}_4\text{CN}$); 7.78 (1H, 5-H of bpy); 7.50 (1H, 5'-H of bpy); 7.34 (d, 1H, 6-H of *m*- $\text{C}_6\text{H}_4\text{CN}$); 7.18 (t, 1H, 5-H of *m*- $\text{C}_6\text{H}_4\text{CN}$); 7.13 (sh, 1H, 6'-H of bpy). The $^1\text{H-NMR}$ pattern of bpy was similar to that of reported **1**, including splitting of 6- and 6'-H [2]. The assignment of the peaks of *m*- $\text{C}_6\text{H}_4\text{CN}$ is tentative, however, the $^1\text{H-NMR}$ pattern of *m*- $\text{C}_6\text{H}_4\text{CN}$ is much changed from that of *m*-dicyanobenzene: $\delta = 8.47$ (s, 1H, 2-H); 8.19 (m, 2H, 4- and 6-H); 7.79 (t, 1H, 5-H).



Scheme 1. A plausible reaction pathway for the reductive elimination.

3.2. Reaction of **1** with acrylonitrile

To an acetone (20 ml) solution of **1** (34 mg, 0.10 mmol) was added acrylonitrile (0.66 cm³, 10 mmol). After the reaction mixture was stirred for 10 h, the solvent was removed by evaporation to give a light brown solid. The solid was extracted with chloroform, and removal of solvent by evaporation provided a light brown solid which was purified by using HPLC to give *o*-dicyanobenzene (4 mg).

Similar reaction of **4** with acrylonitrile gave *m*-dicyanobenzene.

3.3. Reaction of **1** with HCl

To an acetone (40 ml) solution of **1** (68 mg, 0.20 mmol) was added conc. HCl (3.3 cm³). After the reaction mixture was stirred for 15 h, the solvent was removed by evaporation to give a light yellow solid. The solid was extracted with chloroform, and removal of solvent by evaporation provided a light brown solid which was purified by using HPLC to give *o*-dicyanobenzene (18 mg).

3.4. Crystal data for **3**

C₁₇H₁₃N₃Ni, *M* = 318.01, monoclinic, space group *P*2₁/*a* (No. 14), *a* = 8.948(6) Å, *b* = 18.441(4) Å, *c* = 9.266(6) Å, β = 113.62(5)°, *V* = 1400(1) Å³, *Z* = 4, *D*_{calc} = 1.508 Mg m⁻³, *F*(0 0 0) = 656.00. The diffraction data were collected with a Rigaku AFC5R diffractometer at ambient temperature (23 °C) using the ω scan mode (2θ ≤ 55°). Correction for Lorentz and polarization effects and an empirical absorption correction (ψ scan) were applied. The structure was solved by a common combination of direct method (SAPI91) and subsequent Fourier techniques. The positional and thermal parameters of non-hydrogen atoms were refined anisotropically, while hydrogen atoms were located by assuming the ideal geometry.

Acknowledgements

This work was partly supported by a Grant-in-Aid for Scientific Research and the 21st Century COE program from the Ministry of Education, Science, Culture and Sports, Japan.

References

- [1] (a) F.L. Taw, P.S. White, R.G. Bergman, M. Brookhart, *J. Am. Chem. Soc.* 124 (2002) 4192; (b) J.J. Garcia, W.D. Jones, *Organometallics* 19 (2000) 5544; (c) D. Churchill, J.H. Shin, T. Hascall, J.M. Hahn, B.M. Bridgewater, B.M. Parkin, *Organometallics* 18 (1999) 2403.
- [2] M. Abila, T. Yamamoto, *J. Organomet. Chem.* 532 (1997) 267.
- [3] T. Morimoto, *Organomet. News* 105 (2002) 105.
- [4] J.P. Collman, L.S. Hegeudus, J.R. Norton, R.G. Finke, *Principles and Applications of Organotransition Metal Chemistry*, University Science Books, Mill Valley, CA, 1987.
- [5] (a) T. Yamamoto, A. Yamamoto, S. Ikeda, *J. Am. Chem. Soc.* 93 (1971) 3350; (b) B. Åkermark, H. Johnsen, B. Ross, U. Wahlgren, *J. Am. Chem. Soc.* 100 (1979) 5876; (c) K. Tatsumi, A. Nakamura, S. Komiya, T. Yamamoto, A. Yamamoto, *J. Am. Chem. Soc.* 106 (1984) 8181.
- [6] (a) Y. Murakami, T. Yamamoto, *Inorg. Chem.* 36 (1997) 5682; (b) T. Yamamoto, Y. Murakami, M. Abila, *Chem. Lett.* (1999) 419; (c) T. Yamamoto, M. Abila, Y. Murakami, *Bull. Chem. Soc. Jpn* 75 (2002) 1997 (The bar A in Figure 3 in this reference should be shortened).
- [7] (a) A.H. Roy, J.F. Hartwig, *J. Am. Chem. Soc.* 123 (2001) 1232; (b) K. Koo, G.L. Hillhouse, *Organometallics* 14 (1995) 4421; (c) M.S. Driver, J.F. Hartwig, *J. Am. Chem. Soc.* 118 (1996) 7217.
- [8] (a) R.J. McKinney, C. Roe, *J. Am. Chem. Soc.* 108 (1986) 5167; (b) J.A. Miller, *Tetrahedron Lett.* 42 (2001) 6991.
- [9] (a) J.E. Marccone, K.G. Moloy, *J. Am. Chem. Soc.* 120 (1998) 8527; (b) J. Huang, C.M. Haar, S.P. Nolan, *Organometallics* 18 (1999) 297.
- [10] (a) J. Kaiser, J. Sieler, U. Braum, L. Golic, E. Dinjus, D. Wather, *J. Organomet. Chem.* 224 (1982) 81; (b) D. Walther, E. Dinjus, L. Sieler, L. Anderson, O. Lindqvist, *J. Organomet. Chem.* 276 (1984) 99.
- [11] N. Seno, S. Tsuchiya, M. Hidai, Y. Uchida, *Bull. Chem. Soc. Jpn* 49 (1976) 1184.