


Table I. Nmr Data of the Methyl and Nickel Complexes^a

Complex	Solvent	Temp, °C	δ , chemical shifts in ppm ^b		
			CH ₃ -Ni	CH ₃ (acac)	CH(acac)
CH ₃ Ni(acac)(PPh ₃) ₂	Benzene	25	0.09 (3, s)	1.39 (3, s) 1.89 (3, s)	5.30 (1, s)
	Pyridine	25	0.04 (3, s)	1.78 (6, s)	5.40 (1, s)
CH ₃ CH ₂ Ni(acac)(PPh ₃)	Acetone- <i>d</i> ₆	25	CH ₃ CH ₂ Ni	CH ₃ (acac)	CH(acac)
			0.18 (5, s)	1.42 (3, s) 1.89 (3, s)	5.37 (1, s)
	Benzene	25	0.77 (5, s)	1.43 (3, s)	5.30 (1, s)
				1.93 (3, s)	5.40 (1, s)
	Pyridine ^c (220 MHz)	64	-36	0.60 (3, t) ^d	1.76 (6, s)
0.89 (2, q) ^d				1.62 (3, br, s)	
			0.6 (3, br, s)	1.82 (3, br, s)	
			0.9 (2, br, s)		

^a Chemical shifts are referenced to internal TMS. When TMS interferes with the sample peaks, an appropriate peak of the known chemical shift relative to TMS was used as the standard. The chemical shift values with an external TMS varied depending on the concentration of the sample. ^b Figures in parentheses mean peak intensity and the multiplicity: s, singlet; t, triplet; q, quartet; br, broad. ^c Peaks due to ethane and the coordinated ethylene which were formed by decomposition of **1** are observed at δ 0.72 and 2.5–2.8, respectively. ^d $J = 7.4$ Hz.

at -36° . The peaks due to the ethyl protons in **1** show striking variation depending on solvent. The ethyl peaks are observed as a singlet in benzene, acetone, toluene, and tetrahydrofuran whereas in more basic solvents such as pyridine and triethylamine they are observed as a multiplet at 100 MHz and as a pair of a triplet and a quartet at 220 MHz. The singlet ethyl peak of **1** in toluene is broadened by lowering the temperature, but neither the splitting of the peak nor the appearance of a hydride peak was observed at -100° . Examination of the ³¹P nmr spectrum of **1** revealed that the coordinated triphenylphosphine ligand exchanges quite rapidly with the added triphenylphosphine in pyridine even at -40° where the methyl protons in the acetylacetonato ligand are observed as nonequivalent. On the other hand the exchange of the triphenylphosphine ligand in benzene or toluene is much slower. These results suggest that a partial dissociation of the acetylacetonato ligand (bidentate \rightarrow monodentate) and recoordination may be operative as a mechanism to make the methyl protons of the acetylacetonato ligand equivalent. Lewis bases enhance the ligands dissociation whereas they suppress the proton interchange of the ethyl group bonded to nickel. Although the β elimination is considered as a possible mechanism to explain the rapid interconversion in less basic solvents, the ethylene-coordinated nickel hydride which is assumed as the intermediate may not be present as a separate entity of a measurable lifetime, since no hydride peak was observed and attempts to cause the isomerization of butene-1 with **1** failed.

By enhancing the ligand exchange rate, Lewis bases cause another complication; pyridine (py) causes the disproportionation of **1** as follows



Ni(acac)₂·2py was isolated and identified after leaving the pyridine solution of **1** at room temperature for 2 weeks or after heating the solution at 70° for 2 hr. The presence of ethane formed by the disproportionation reaction is observed as a singlet peak at δ 0.72 and the peak of the coordinated ethylene, which is present in the equilibrium $(\text{PPh}_3)_2\text{NiC}_2\text{H}_4 \rightleftharpoons (\text{PPh}_3)_2\text{Ni}$

+ C₂H₄, is observed in the range δ 2.5–2.8.^{7,9} Methane is formed by treating **2** in pyridine, and a similar disproportionation of **2** to eq 1 appears to be taking place.

The thermal stability of the present ethyl complex **1** which has an ethyl group having β hydrogens readily abstractable by transition metal is noteworthy in discussing the theory to explain the stability of the transition metal alkyls.¹⁰

Acknowledgment. The authors are grateful to Professor I. Morishima of Kyoto University for the measurement of the 220-MHz nmr spectrum.

(9) P.-T. Cheng, C. D. Cook, S. C. Nyburg, and K. Y. Wan, *Inorg. Chem.*, **10**, 2210 (1971).

(10) W. Mowat, A. Shortland, G. Yagupsky, N. J. Hill, M. Yagupsky, and G. Wilkinson, *J. Chem. Soc., Dalton Trans.*, 533 (1972); G. M. Whitesides, J. E. Gaasch, and E. R. Stedronsky, *J. Amer. Chem. Soc.*, **94**, 5258 (1972); P. S. Braterman and R. J. Cross, *J. Chem. Soc., Dalton Trans.*, 657 (1972).

Akio Yamamoto,* Takakazu Yamamoto
Toshio Saruyama, Yoshiyuki Nakamura

Tokyo Institute of Technology
Research Laboratory of Resources Utilization
O-okayama, Meguro, Tokyo 152, Japan
Received February 27, 1973

Nucleic Acid Related Compounds. VII. Conversion of Ribonucleoside 2',3'-Ortho Esters into Deoxy, Epoxy, and Unsaturated Nucleosides^{1,2}

Sir:


We wish to report the conversion of nucleoside 2',3'-ortho esters into the corresponding 3'-halo-3'-deoxy-xylo ester intermediates which are versatile precursors of epoxy, deoxy, and unsaturated nucleosides.³

Usual means of access to these modified nucleosides involve prior construction of a suitable carbohydrate derivative followed by coupling with a blocked and/or

(1) This work was generously supported by Grant No. A5890 from the National Research Council of Canada, Grant No. 9530-108 from the Defense Research Board, and The University of Alberta.

(2) For the previous paper in this series see: M. J. Robins and E. M. Trip, *Biochemistry*, **12**, 2179 (1973).

(3) R. Mengel and M. J. Robins, 161st National Meeting of the American Chemical Society, Los Angeles, Calif., March-April, 1971, Abstract CARB-002.



activated base.^{4a-d} Alternatively, intramolecular cyclonucleoside participation which requires specific base structure and functionality has been used.^{4a,d,e} We are interested in developing general transformations of intact nucleosides which are applicable to nucleoside antibiotics readily available by fermentation but practically inaccessible by coupling reactions.⁵

The elegant pioneering work of Winstein and Meerwein on acyloxonium ions has been reviewed recently⁶ along with numerous subsequent applications. Our observation of the facile conversion of 3',4'-*O*-ethoxymethylidene-*psicofuranine* to the 1',3',4'-*O*-orthoformate with boron trifluoride etherate⁷ led us to explore entry into acyloxonium ion mediated reactions of nucleosides^{6,8,9} via 2',3'-ortho esters.

Treatment of 2',3'-*O*-methoxyethylideneadenosine¹⁰ (**1**, Z = N) with either boron trifluoride etherate or antimony pentachloride in the presence of iodide gave mixtures of 2'-*O*- and 3'-*O*-acetyladenosine even under "anhydrous" conditions. Pyridine hydrohalide was weakly effective in promoting the desired reaction. Heating **1** (Z = N) at reflux with pivalic acid chloride in pyridine for 2 hr gave a mixture from which could be isolated 6-*N*-pivalamido-9-(3-chloro-3-deoxy-2-*O*-acetyl-5-*O*-pivalyl- β -D-xylofuranosyl)purine¹¹ [**3**, Z = N; X = Cl, R = COC[CH₃]₃; R' = CH₃ (70% yield); uv max (MeOH) 272 nm (ϵ 17,000); nmr (CDCl₃, TMS internal) δ 1.23 (s, 9, 5'-OCOC[CH₃]₃), 1.40 (s, 9, N⁶-

COC[CH₃]₃), 2.18 (s, 3, 2'-OCOC[CH₃]₃), 4.48 (m, 3, H_{3'}, H_{5',5''}), 4.65 (m, 1, H_{4'}), 5.63 (d of d, $J_{2'-1'} = 2$ Hz, $J_{2'-3'} = 1.5$ Hz, 1, H_{2'}), 6.31 (d, $J_{1'-2'} = 2$ Hz, 1, H_{1'}), 8.41 (s, 1, H₈), 8.56 (br s, 1, N⁶-*H*-pivalyl), 8.76 (s, 1, H₂); mass spectrum calcd for C₂₂H₃₀ClN₅O₆, 495.1885; found, 495.1905] and 6-*N*-pivalamido-9-(3-chloro-3-deoxy-5-*O*-pivalyl-2-*O*-[4,4-dimethyl-3-pivalyloxypent-2-enyl]- β -D-xylofuranosyl)purine¹¹ [**3**, Z = N; X = Cl; R = COC[CH₃]₃; R' = CH=C(OCOC[CH₃]₃)C[CH₃]₃ (11% yield); uv max (MeOH) 272 nm (ϵ 17,800); nmr (CDCl₃, TMS internal) δ 1.16 (s, 9, CH=C(OPv)C[CH₃]₃), 1.23 and 1.25 (s and s, 9 and 9, 5'-OCOC[CH₃]₃ and CH=C(OCOC[CH₃]₃)C[CH₃]₃), 1.41 (s, 9, N⁶-COC[CH₃]₃), 4.47 (m, 3, H_{3'}; H_{5',5''}), 4.65 (m, 1, H_{4'}), 5.57 (d of d, $J_{2'-1'} = 2$ Hz, $J_{2'-3'} = 1.5$ Hz, 1, H_{2'}), 5.76 (s, 1, CH=C(OPv)(*t*-Bu)), 6.33 (d, $J_{1'-2'} = 2$ Hz, H_{1'}), 8.41 (s, 1, H₈), 8.53 (br s, 1, N⁶-*H*-pivalyl), 8.76 (s, 1, H₂); mass spectrum calcd for C₃₂H₄₆ClN₅O₈, 663.3035; found, 663.3017]. Treatment of **1** (Z = N) with pivalyl chloride and excess sodium iodide for 10 min in refluxing pyridine gave **3**¹¹ (Z = N; X = I, R = COC[CH₃]₃; R' = CH=C(OCOC[CH₃]₃)C[CH₃]₃; mass spectrum calcd for C₃₂H₄₆IN₅O₈, 755.2391; found, 755.2424) as the major product in 60% yield with none of the 2'-*O*-acetyl derivative observed.

Methanolic sodium methoxide converted the various products **3** (Z = N) into the known 2',3'-anhydroadenosine^{11,12} (**4**, B = adenine). This confirms the trans 2',3' relationship and the "down" configuration of the oxygen function in **3** as well as providing convenient access to the useful synthetic intermediate **4**.

Hydrogenolysis of iodo enol ester **3** gave the corresponding 3'-deoxyadenosine derivative¹¹ (**3**, Z = N; X = H; R = COC[CH₃]₃; R' = CH=C(OCOC[CH₃]₃)C[CH₃]₃; mp 92.5–93.5°; mass spectrum calcd for C₃₂H₄₇N₅O₈, 629.3425; found, 629.3409) with the hindered double bond intact. A minor quantity of the 2'-deoxyadenosine derivative,¹¹ mp 127–129°, was isolated by fractional crystallization indicating some attack of iodide at C₂; of the "symmetrical" 1,3-dioxolan-2-ylum intermediate (**2**, R = COC[CH₃]₃; R' = CH=C(OCOC[CH₃]₃)C[CH₃]₃). Chromatography¹³ of the aqueous phase of the deblocked orig-

(12) (a) C. D. Anderson, L. Goodman, and B. R. Baker, *J. Amer. Chem. Soc.*, **81**, 3967 (1959); (b) A. Benitez, O. P. Crews, Jr., L. Goodman, and B. R. Baker, *J. Org. Chem.*, **25**, 1946 (1960).

(13) C. A. Dekker, *J. Amer. Chem. Soc.*, **87**, 4027 (1965).

(4) For comprehensive reviews, see: (a) A. M. Michelson, "The Chemistry of Nucleosides and Nucleotides," Academic Press, New York, N. Y., 1963; (b) C. A. Dekker and L. Goodman in "The Carbohydrates," 2nd ed, Vol. IIA, W. Pigman and D. Horton, Ed., Academic Press, New York, N. Y., 1970, Chapter 29; (c) J. A. Montgomery and H. J. Thomas, *Advan. Carbohyd. Chem.*, **17**, 301 (1962); (d) J. J. Fox and I. Wempen, *ibid.*, **14**, 283 (1959); (e) M. Ikehara, *Accounts Chem. Res.*, **2**, 47 (1969).

(5) R. J. Suhadolnik, "Nucleoside Antibiotics," Wiley-Interscience, New York, N. Y., 1970.

(6) C. U. Pittman, Jr., S. P. McManus, and J. W. Larsen, *Chem. Rev.*, **72**, 357 (1972).

(7) J. R. McCarthy, Jr., R. K. Robins, and M. J. Robins, *J. Amer. Chem. Soc.*, **90**, 4993 (1968).

(8) L. Goodman, *Advan. Carbohyd. Chem.*, **22**, 109 (1967).

(9) (a) S. Greenberg and J. G. Moffatt, *J. Amer. Chem. Soc.*, **95**, 4016 (1973); A. F. Russell, S. Greenberg, and J. G. Moffatt, *ibid.*, **95**, 4025 (1973); (b) M. J. Robins, J. R. McCarthy, Jr., R. A. Jones, and R. Mengel, *Can. J. Chem.*, **51**, 1313 (1973).

(10) H. P. M. Fromageot, B. E. Griffin, C. B. Reese, and J. E. Sulston, *Tetrahedron*, **23**, 2315 (1967).

(11) All compounds had satisfactory elemental analyses for C, H, N, and where applicable Cl or I. Spectral data were comparable with the examples given or with literature and/or values determined on known samples.

inal hydrogenolysis mixture gave 3'-deoxyadenosine (cordycepin)^{11,14} (5, B = adenine) and 2'-deoxyadenosine^{11,12a} in a ratio of 9:1.

Treatment of the above iodo enol ester 3 with 1,5-diazabicyclo[4.3.0]nonene-5 (DBN) and other nonsaponifying bases gave the blocked (3-deoxy-β-D-glycero-pent-3-enofuranosyl) heterocycle plus the corresponding heterocycle-substituted furan derivative. Deblocking gave 6,¹¹ mp 228–230°, which was hydrogenated to 5^{11,14} plus its 4' epimer.^{11,15}

Analogous reaction of 2',3'-O-methoxyethylidene-tubercidin¹¹ (1, Z = CH) gave 3¹¹ (Z = CH; X = I; R = COC[CH₃]₃; R' = CH=C(OCOC[CH₃]₃)C-[CH₃]₃; mass spectrum calcd for C₃₃H₄₇N₄O₈, 754.2339; found, 754.2376). Transformations of this material to give 4,¹¹ mp 167° dec, 5,^{9b,11} and 6,¹¹ mp 190–192° (B = 4-aminopyrrolo[2,3-d]pyrimidine) proceeded similarly with the exception that no 2'-deoxytubercidin was detected in the hydrogenolysis.

Isolation of intermediates involved in characterizing the interesting acyloxonium ion diacylation mechanism of enol ester formation, details of various other products formed, and applications of these useful intermediates in nucleoside chemistry will be reported in detail.

(14) W. W. Lee, A. Benitez, C. D. Anderson, L. Goodman, and B. R. Baker, *J. Amer. Chem. Soc.*, **83**, 1906 (1961); E. A. Kaczka, E. L. Dulaney, C. O. Gitterman, H. B. Woodruff, and K. Folkers, *Biochem. Biophys. Res. Commun.*, **14**, 452 (1964).

(15) K. L. Nagpal and J. P. Horwitz, *J. Org. Chem.*, **36**, 3743 (1971).

(16) University of Alberta Postdoctoral Fellow, 1969–1971.

Morris J. Robins,* Rudolf Mengel,¹⁶ Roger A. Jones

Department of Chemistry, The University of Alberta
Edmonton, Alberta, Canada T6G 2G2

Received March 13, 1973

Nitrosyl Transfer Reactions

Sir:

The recent literature contains several examples of reactions involving transfer of carbon monoxide from one metal atom to another.¹ We wish to report the first observations relating to nitrosyl transfer reactions.

Methanolic solutions of CoD₂PPh₃² rapidly absorb NO to yield a mononitrosyl adduct with ν_{NO} = 1710 cm⁻¹ (CHCl₃ solution). The solution precipitates a solid of composition CoNOD₂(MeOH),³ with ν_{NO} = 1639 cm⁻¹ (KBr); coordinated phosphine is not present. The similarity of the equatorial ligands and the NO stretching frequency of CoNOD₂(MeOH) to those of Co(en)₂NOCl⁺ClO₄⁻ 4 and CoNO(tet)⁵ suggests the Co–N–O moiety is bent in the dimethylglyoximate complex. Proton nmr of CoNOD₂(MeOH) in CDCl₃ exhibits a methoxy resonance at a chemical shift identical with that of uncoordinated methanol; azeotropic distillation of methanol from a benzene solution of

(1) J. Alexander and A. Wojcicki, *Inorg. Chem.*, **12**, 74 (1973); B. Booth, M. Else, R. Fields, H. Goldwhite, and R. Hazeldine, *J. Organometal. Chem.*, **14**, 417 (1968).

(2) G. Schrauzer and R. Windgassen, *Chem. Ber.*, **99**, 602 (1966); D = monoanion of dimethylglyoxime.


(3) Compounds were characterized by elemental analyses and all applicable spectroscopic methods. Compare M. Tamaki, I. Masuda, and K. Shinra, *Bull. Chem. Soc. Jap.*, **45**, 171 (1972).

(4) D. Snyder and D. Weaver, *Inorg. Chem.*, **9**, 2760 (1970).



(5) R. Wiest and R. Weiss, *Rev. Chim. Miner.*, **9**, 655 (1972); tet = tetradentate ligands, e.g., N,N'-ethylenebis(acetylacetoniminate) or N,N'-ethylenebis(benzoylacetoniminate).

CoNOD₂(MeOH) yields unsolvated CoNOD₂. All of the observations imply a large trans effect for NO, consistent with previous observations on bent nitrosyls.⁶


CoNOD₂(MeOH) reacts with CoCl₂L₂ and L (L = PPh₃) (2:1:2 molar ratio) in ethanol to yield CoClD₂L (1 mol), CoD₂L (1 mol), and an equilibrium mixture⁷



NaBPh₄ displaces this equilibrium to the left by quantitatively precipitating Co(NO)₂L₂⁺BPh₄⁻. The overall reaction (1) involves the transfer of two nitrosyl




groups and a chlorine atom. Since the NO donor reagent is a mononitrosyl, it is natural to consider a stepwise process. The intermediacy of a mononitrosyl in reaction 1 is suggested by the observation that Co(NO)Cl₂L₂⁸ reacts with CoNOD₂(MeOH) and L (1:1:1 mol ratio) to form the dinitrosyl (2). No Ph₃PO is de-



tected after these reactions, indicating the absence of free NO. An alternative mechanism involving initiation of the reaction by catalytic amounts of the halogen acceptor⁹ CoD₂ is ruled out by the observation that neither CoD₂ nor CoD₂L will reduce CoCl₂L₂.

Square-pyramidal cobalt complexes with CH₃ or bent NO in the apical position exhibit many similarities. Foremost is the common ambiguity in assignment of oxidation states: CH₃(+1), CH₃, or CH₃(-1) vs. NO(+1), NO, or NO(-1). Both groups have very high trans effects, sometimes allowing isolation of the complex with the trans position unoccupied.^{5,10} Co-CH₃D₂ is dimeric,¹¹ resonances of nonequivalent dimethylglyoximate methyl groups being apparent below -12°. The proton nmr of CoNOD₂(MeOH) shows only one resonance for dimethylglyoximate methyl groups even at -90°, implying an even stronger trans effect for bent NO than for CH₃. Finally the nitrosyl transfer reaction observed here mimics the known alkyl transfer reactions of alkyl cobalt Schiff base complexes.¹²

Although simple nitrosyl transfer must occur at some stage in reaction 1, it seems likely that the efficacy of CoNOD₂(MeOH) as a nitrosyl source is related to the fact that the CoD₂ produced can also function as a halogen acceptor. Consistent with this idea, we find that nitrosyl-halogen interchange appears to be a rather general reaction. For example



(6) C. Brock, B. Coyle, and J. Ibers, *J. Chem. Soc. A*, 2146 (1971).

(7) T. Bianco, M. Rossi, and L. Uva, *Inorg. Chim. Acta*, **3**, 443 (1969). We write here Co(NO)₂(PPh₃)₂Cl to indicate this equilibrium mixture.

(8) J. Collman, P. Farnham, and G. Dolcetti, *J. Amer. Chem. Soc.*, **93**, 1788 (1971).

(9) A. Adin and J. Espenson, *Inorg. Chem.*, **11**, 686 (1972); L. Marzilli, J. Salerno, and L. Epps, *ibid.*, **11**, 2050 (1972).

(10) G. Costa, G. Mestroni, G. Tauzher, and L. Stafani, *J. Organometal. Chem.*, **6**, 181 (1966); C. Floriani, M. Poppis, and F. Calderazzo, *ibid.*, **12**, 209 (1968).

(11) A. Herlinger and T. Brown, *J. Amer. Chem. Soc.*, **94**, 388 (1972).

(12) A. v. Bergen and B. West, *Chem. Commun.*, 52 (1971).