

Synthesis, anti-GABA activity and preferred conformation of bicuculline and norbicuculline enantiomers

J Kardos¹, T Blandl¹, ND Luyen¹, G Dörnyei¹, E Gács-Baitz¹, M Simonyi¹,
DJ Cash², G Blaskó³, Cs Szántay¹

¹Central Research Institute for Chemistry, The Hungarian Academy of Sciences, 1525 Budapest, PO Box 17, Hungary;

²Biochemistry Department, School of Medicine, University of Missouri, Columbia, MO, 65201 USA;

³EGIS Pharmaceuticals, 1475 Budapest, PO Box 100, Hungary

(Received 19 October 1995; accepted 18 March 1996)

Summary — Synthesis of *erythro*-(±)-[1*SR*,9*RS*]-norbicuculline and *threo*-(±)-[1*SR*,9*SR*]-noradlumidine from piperonal was performed using Bischler–Napieralski cyclization as a key step. Resolution gave rise to (+)-[1*S*,9*R*]-norbicuculline ([1*S*,9*R*] norBIC) and (–)-[1*R*,9*S*]-norbicuculline ([1*R*,9*S*] norBIC) in >99.5% enantiomeric purity. Bicuculline enantiomers were readily obtained by methylation of the latter products. [1*S*,9*R*]BIC was about 70 times more potent than [1*R*,9*S*] BIC as an inhibitor of GABA_A receptor binding and was about 100 and 900 times more potent than [1*S*,9*R*] norBIC at pH 7.1 and 5.0 respectively. Similarly, [1*S*,9*R*] norBIC was much less potent than [1*S*,9*R*] BIC as an inhibitor of GABA-specific ³⁶Cl[–] ion flux. The observed increase of about two orders of magnitude in the in vitro biological activity caused by *N*-2-CH₃ substitution in [1*S*,9*R*] norBIC was attributed to different conformations for *erythro*- and *nor-erythro*-bicucullines indicated by ¹H nuclear Overhauser enhancements of [1*S*,9*R*] BIC and [1*S*,9*R*] norBIC.

bicuculline enantiomer / norbicuculline enantiomer / Bischler–Napieralski cyclization / ¹H nuclear Overhauser enhancement / [³H]GABA binding / ³⁶Cl[–] ion flux


Since the announcement by Curtis et al [1] that the phthalideisoquinoline (PIQ) alkaloid, [1*S*,9*R*] BIC possessed potent inhibitory activity against the depressant action of GABA in the central nervous system, it has been of interest to study the effect of its antipode, [1*R*,9*S*] BIC. The effect of [1*R*,9*S*] BIC, however, has not been unequivocally reported because of the scarcity of this natural product and the confusion caused by the reversal of the sign of optical rotation for quaternary bicuculline derivatives [2]. Based on the available structural data and in vitro biological activity measurements of 39 PIQ derivatives, a good correlation was found [3] between activities and preferred conformations of *erythro* and *threo* PIQs and analogs.

We report herein a synthetic route for bicuculline and norbicuculline enantiomers. The effect of *N*-methyl substitution on GABA_A receptor activity and preferred conformations were evaluated by [³H]GABA binding, GABA specific ³⁶Cl[–] ion flux and ¹H nuclear Overhauser enhancement (nOe) measurements respectively.

Chemistry

[1*S*,9*R*] BIC **11a**, first encountered as a constituent of *Dicentra cucullaria* [4] and subsequently found in other species of genera *Rhoeadales* [5, 6], was first totally synthesized by Groenewoud and Robinson [7] as early as 1936. Reissert synthesis of [1*SR*,9*RS*] norBIC has been cited [8] as an unpublished result of Kerekes [8], however, total synthesis of [1*SR*,9*RS*] norBIC **9** has not been described up to now.

As demethylation of PIQ alkaloids results in decomposition of the molecule, a linear approach to [1*SR*,9*RS*] BIC via [1*SR*,9*RS*] norBIC had to be elaborated to meet our demand to produce [1*S*,9*R*] BIC and [1*S*,9*R*] norBIC as well as [1*R*,9*S*]BIC and [1*R*,9*S*] norBIC. For this purpose the Bischler–Napieralski route, applied to the synthesis of (±)-[1*SR*,9*RS*]- α -narcotine [9–11], seemed to be appropriate. The two building blocks needed for Bischler–Napieralski cyclization, ie, carboxylic acid **4** (scheme 1; stereostructures are represented according to [12]) and homopiperonylamine **6** [13], were both prepared from pipe-



Scheme 1.

ronal **1**. According to Ziegler and Fowler [14], **1** gave a Schiff base **2** with cyclohexylamine. This was treated with butyllithium at $-78\text{ }^{\circ}\text{C}$, then with CO_2 in THF/hexane solution to produce carboxylic acid **3** after hydrolysis. Phthalide-3-carboxylic acid **4** was obtained from **3** in two steps; addition of HCN was followed by aqueous sulfuric acid hydrolysis [9]. Compound **4** was then converted to acyl chloride **5** with thionyl chloride, which, without isolation, was reacted with homopiperonylamine **6** obtained from piperonal **1** in two steps via nitrostyrol [13], to supply amide **7**. POCl_3 cyclization of **7** led to $\Delta^{1,9}$ -norBIC **8** containing a double bond between the two rings ('enamine form') even in the protonated salt form (no C9-H

signal could be observed in the $^1\text{H-NMR}$ spectra of **8** in D_2O solution). Sodium borohydride reduction of **8** resulted in a diastereomeric mixture of *erythro*-[1*SR*, 9*RS*] norBIC **9** and its *threo* epimer (\pm)-[1*SR*, 9*SR*]-noradlumidine ([1*SR*, 9*SR*] norADLD, **10**) in a ratio of about 5:1. The relative stereochemistry of the chiral atoms C1 and C9 in the two compounds was deduced from the characteristic chemical shifts of the C2'-H protons (5.90 and 6.93 ppm for the major and minor epimers respectively) and by comparing the different $n\text{Oe}$ values of protons C1-H, C8-H, C9-H and C2'-H (table I) with other *erythro* and *threo* PIQ alkaloids [15].

After separating the two diastereomeric racemates on the column, the major *erythro* base **9** was resolved in acetone with (-)- and (+)-*O,O*-dibenzoyltartaric acid and the enantiomeric purity was checked on a chiral HPLC column (Chiralcel OD) showing a baseline separation of **9a** and **9b**. After repeated recrystallization of the two diastereomeric salt-pairs, [1*S*, 9*R*] norBIC **9a** and [1*R*, 9*S*] norBIC **9b** were obtained in more than 99.5% optical purity. Eschweiler-Clark or methyl iodide methylation of the optically pure [1*S*, 9*R*] norBIC and [1*R*, 9*S*] norBIC gave [1*S*, 9*R*] BIC and [1*R*, 9*S*] BIC **11a** and **11b** respectively, the

Table I. Nuclear Overhauser enhancement data (%) in $^1\text{H-NMR}$.

Proton irradiated	Proton observed				
	C2'-H	C1-H	C8-H	C9-H	N2-CH ₃
10^a					
H8	—	4.4	—	15.4	—
H9	2.4	6.3	15.3	—	—
H1	5.6	—	5.5	7.4	—
12^b					
H8	—	6.8	—	3.5	—
H9	2.2	7.3	4.4	—	4.4
H1	2.0	—	9.8	8.2	6.4 ^c
9a^d					
H8	1.9	3.4	—	15.0	—
H9	2.3	6.4	17.8	—	—
H1	1.0	—	5.4	6.8	—
11a^e					
H8	—	7.6	—	10.7	—
H9	3.0	8.3	12.0	—	0.8
H1	3.6	—	10.3	8.9	6.4 ^c

^a $J_{\text{C1-H,C9-H}} = 3.2\text{ Hz}$; $\delta_{\text{C4-H}_{\text{ax}}} = 2.74\text{ ppm}$. ^b $J_{\text{C1-H,C9-H}} = 3.3\text{ Hz}$; $\delta_{\text{C4-H}_{\text{ax}}} = 2.75\text{ ppm}$. ^cShare of C3-H is higher in [1*S*, 9*SR*] ADLD than in [1*S*, 9*R*] BIC. ^d $J_{\text{C1-H,C9-H}} = 4.0\text{ Hz}$; $\delta_{\text{C4-H}_{\text{ax}}} = 2.43\text{ ppm}$. ^e $J_{\text{C1-H,C9-H}} = 4.1\text{ Hz}$; $\delta_{\text{C4-H}_{\text{ax}}} = 2.24\text{ ppm}$; no $n\text{Oe}$ was observed between N2-CH₃ and C4-H_{ax} protons suggesting the following configuration: N2-CH₃ is equatorial (α); lone pair electrons at N2 (N2-LPE) are axial (β).

former being identical in every respect (mp, TLC, IR, ^1H - and ^{13}C -NMR, MS) with the natural sample, thus providing a final proof of the *erythro* assignment for [1*S*,9*RS*] norBIC (**9**).

The *threo* racemate ([1*S*,9*SR*] norADLD, **10**) and its methylated product ([1*S*,9*SR*] ADLD, **12**) were also obtained but not biologically evaluated.

Biological data

Enantiomers of **9** and **11** were initially evaluated for inhibition of [^3H]GABA binding in membranes from the rat cerebral cortex. As shown in table II, GABA_A receptors expressed greater enantioselectivity towards **11a** and **11b** than **9a** and **9b**, a result that is consistent [3] with the higher potency of **11a**. As racemic bicuculline and norbicuculline appeared to have identical activities and so were thought to possess similar conformations [3], we were quite surprised to find that $\text{N}2\text{-H} \rightarrow \text{N}2\text{-CH}_3$ substitution in **11a** caused an approximately 100-fold increase in inhibition of GABA_A receptor binding compared with **9a**.

The effect of $\text{N}2\text{-H} \rightarrow \text{N}2\text{-CH}_3$ substitution on [^3H]-GABA binding was further evaluated with the protonated **9a** and **11a** analogs. By protonation at pH 5.0 [16], the affinity of **11a** increased ($K_i = 0.39 \mu\text{M}$, $K_{\text{GABA}}/K_i = 0.11$), while that of **9a** decreased ($K_i = 340 \mu\text{M}$, $K_{\text{GABA}}/K_i = 0.0001$).

Substantial increase in the affinity caused by $\text{N}2\text{-CH}_3$ substitution of unprotonated ($K_{9a}/K_{11a} = 100$) and protonated ($K_{9a}/K_{11a} = 870$) analogs and the differential effect of protonation on the affinities of **11a** ($K_{11a}/K_{11a,H^+} = 2.3$) and **9a** ($K_{9a}/K_{9a,H^+} = 0.27$) suggested that the *erythro* conformations of bicuculline and norbicuculline were different. A comparison of the nOe values and the chemical shift data for **11a** and **9a** (table I) revealed that (i) nOe interactions of C1-H and C8-H in [1*S*,9*R*] BIC were twice as large as in [1*S*,9*R*] norBIC; (ii) smaller $\delta_{\text{C}4\text{-H}_{\text{ax}}}$ in [1*S*,9*R*] BIC (2.24 ppm vs 2.43 ppm) indicated that C4-H_{ax} is in the shielding

zone of the phthalide ring; (iii) the nOe effects of C8-H and both C2'-H and C9-H were larger in [1*S*,9*R*] norBIC than in [1*S*,9*R*] BIC. These nOe interactions are indicated in figure 1. The nOe data are in agreement with a conformational change of [1*S*,9*R*] norBIC which results in an increased distance between the lone-pair electrons ($\text{N}2\text{-LPE}$) and the carbonyl group (fig 1).

The inhibitory effect of these bicuculline derivatives (**9a**, **9b**, **11a** and **11b**) on bicuculline-sensitive GABA binding and GABA receptor function was compared (table II). Inhibition of the in vitro binding of [^3H] GABA is parallel to the inhibition of GABA-mediated influx of $^{36}\text{Cl}^-$ ions into the membrane vesicles (40 μM GABA, 6 s, 30 °C) by the same concentrations (46 μM) of the bicuculline derivatives. It is apparent that **9a** was a less potent antagonist of GABA_A function ($^{36}\text{Cl}^-$ flux) than **11a**.

The $\text{N}2\text{-H} \rightarrow \text{N}2\text{-CH}_3$ substitution is apparently responsible for a substantial improvement of anti-GABA activity of *erythro* PIQ derivatives. The unusually large effect was explained by the conformational change in ring B resulting in different $\text{N}2\text{-LPE}$ -carbonyl distances in nor-*erythro*- and *erythro*-bicucullines.




Fig 1. Different conformations of the bicuculline (A) and norbicuculline (B) skeleton as indicated by the nOe interactions.

Table II. [^3H] GABA binding and GABA-specific $^{36}\text{Cl}^-$ ion flux inhibition data in membranes from the rat cerebral cortex.

Compound	Binding ^a inhibition affinity			$^{36}\text{Cl}^-$ transport ^b inhibition influx	
	K_i (μM)	K_{11a}/K_i	K_{GABA}/K_i	dpm	%
GABA	0.018 ± 0.001	49.4	1.0	383 ± 48	0 ± 7
9a	91.5 ± 9.2	0.01	0.002	274 ± 59	28 ± 8
9b	365 ± 70	0.002	0.0005	360 ± 15	6 ± 4
11a	0.89 ± 0.04	1.0	0.02	5 ± 20	99 ± 5
11b	64.0 ± 6.4	0.014	0.0003	282 ± 37	26 ± 10

^aReduced χ_r^2 values were the following: 0.34, 2.81, 1.52, 7.35 and 0.94 for GABA, **9a**, **9b**, **11a** and **11b** respectively; other displacement experiments gave similar results; ^bdata \pm SEM were from six to ten determinations.

Experimental protocols

The ^1H and ^{13}C NMR spectra were taken on a UXR 400 spectrometer. Mass spectra were run on a MS 902 spectrometer. Infrared spectra were taken on a Nicolet 205 FT-I. HPLC was run on a PU 4000 with a Chiralcel OD column eluted with isopropanol. Analyses indicated by the symbols of the elements were within $\pm 0.4\%$ of theoretical values.

Synthesis

2-Formyl-5,6-methylenedioxybenzoic acid **3**

Compound **3** was prepared from piperonal **1** in three steps according to Ziegler and Fowler [14] in 63% yield, mp 164–165 °C (ref [14]; mp 165–165.5 °C).

6,7-Methylenedioxyphthalide-3-carboxylic acid **14**

To the stirred suspension of **3** (5 g, 25.7 mmol) in water (15 mL), cooled to 0–5 °C, a solution of KCN (10 g, 154 mmol) in water (25 mL) was added over 15 min and stirred for an additional 15 min. A mixture of conc HCl aq (17.5 mL) and water (17.5 mL) was added and stirred for 5 h while cooling. The precipitate was filtered off and the reaction mixture was extracted with EtOAc. The combined precipitate and oil, obtained by evaporating the dried EtOAc solution, was refluxed for 2 h in a mixture of conc H_2SO_4 (3.5 mL) and water (15 mL). After cooling, the mixture was extracted with EtOAc and the organic phase was dried and evaporated. On CH_2Cl_2 treatment the residue gave colourless crystals of **4** (4.65 g, 81%) mp 190–192 °C; anal $\text{C}_{10}\text{H}_6\text{O}_6$ (C, H, O); IR (KBr) γ -lactone, 1775 cm^{-1} , C=O, 1720 cm^{-1} ; ^1H NMR ($\text{DMSO}-d_6/\text{CDCl}_3$) δ 5.80 (s, 1H, C3-H), 6.21 (s, 2H, OCH_2O), 7.12 (m, 1H, C5-H), 7.16 (m, 1H, C4-H), 8.80 (s, 1H, COOH).

Amide **7**

To a stirred solution of carboxylic acid **4** (4.0 g, 18 mmol) in dry benzene, (40 mL) thionyl chloride (20 mL) was added dropwise at room temperature, then the mixture was refluxed for 2 h. The mixture containing acyl chloride **5** was evaporated in vacuo. The amine **6** [13] (2.9 g, 18 mmol) was dissolved in benzene (80 mL) and 2.5% aq NaOH solution (18 mL) was added with stirring. A solution of acyl chloride **5** in benzene (30 mL) was added and stirred for 1 h. The precipitate formed was filtered off. The benzene solution was dried and evaporated. The residue and the precipitate were combined and recrystallized from EtOAc to yield amide **7** (3.6 g, 54%), mp 165–166 °C; anal $\text{C}_{19}\text{H}_{15}\text{NO}_7$ (C, H, N, O); mass spectrum m/z (%) 369 (M^+ , 126), 339 (2), 177 (25), 148 (100); IR (KBr) NH, 3280 cm^{-1} , lactone C=O, 1760 cm^{-1} , amid C=O, 1660 cm^{-1} ; ^1H NMR (CDCl_3) δ 2.71 (t, 2H, benzyl- CH_2), 3.46 (m, 2H, NHCH_2), 5.68 (s, 1H, OCH), 5.90 (s, 2H, OCH_2O), 6.20 (s, 2H, OCH_2O), 6.49 (s, 1H, NH), 6.52 (s, 1H, C2-H), 6.68 (s, 1H, C5-H), 7.12 (m, 1H, C3'-H), 7.30 (m, 1H, C2'-H); ^{13}C NMR ($\text{DMSO}-d_6/\text{CDCl}_3$); 34.65 (benzyl-C), 40.32 (CH_2NH), 78.47 (OCH), 100.50 + 103.25 (OCH_2O), 106.33 (C6'), 107.85 (C5), 108.65 (C2), 114.11 (C2'), 115.89 (C3'), 121.27 (C6), 131.77 (C1), 137.36 (C1'), 144.57, 145.79, 147.27 and 149.41 (C3, C4, C4' and C-5'), 166.07 (C=O).

$\Delta^{1,9}$ -Norbicuculline **8**

Amide **7** (3.50 g, 9.5 mmol), dissolved in freshly distilled POCl_3 (35 mL), was stirred at 100 °C for 1 h under argon. The cooled mixture was poured onto ice (300 g), extracted with diethyl ether, the water phase was then neutralized with conc NH_4OH , the yellow precipitate filtered, dissolved in EtOAc and acidified with conc HCl to pH 4. The crystals were filtered to

give $\Delta^{1,9}$ -norBIC. HCl **8** (2.7 g, 76%) mp 173–174 °C (MeOH); anal $\text{C}_{19}\text{H}_{14}\text{NO}_6\text{Cl}$ (C, H, N, O, Cl); mass spectrum m/z (%) 351 ($\text{M} - \text{HCl}$, 85), 322 (100), 294 (11), 177 (91), 36 (HCl); IR (KBr) = N^+H_2 , 3290 cm^{-1} , lactone C=O, 1720 cm^{-1} ; ^1H NMR (D_2O) δ 3.15 (m, 2H, C4-H₂), 3.90 (m, 2H, C3-H₂), 6.12 + 6.22 (s-s, 2-2H, OCH_2O), 6.98 (s, 1H, C8-H), 7.02 (s, 1H, C5-H), 7.11 (d, 1H, C3'-H), 7.42 (d, 1H, C2'-H).

Norbicuculline **9** and noradlumidine **10**

Cyclized product **8** (5.4 g 13.9 mmol) was dissolved in a mixture of CH_2Cl_2 (300 mL) and AcOH (6 mL), cooled to 0–5 °C. To the stirred mixture NaBH_4 (1.62 g, 42.8 mmol) was added in small portions (~3 h) then stirred for 1 h. Excess reducing agent was then decomposed with acetone (2 mL). The mixture was washed with water, dried (MgSO_4), and evaporated in vacuo. Chromatography on 250 g 63-200 mesh Kieselgel 60 (eluent: benzene/acetone, 1:1) gave **9** (2.1 g, 43%) mp 184–185 °C and **10** (0.4 g, 8%) mp 203–204 °C. **9**: Anal $\text{C}_{19}\text{H}_{15}\text{NO}_6$ (C, H, N, O); mass spectrum m/z (%) 353 (M^+ , 1), 335 (2), 176 (100), 149 (3); IR (KBr) C=O, 1750 cm^{-1} ; ^1H NMR (CDCl_3) 1.86 (br s, 1H, NH), 2.43 (m, $J = 15.5 + 8.05 + 5.4$ Hz, 1H, C4-H_{ax}), 2.53 (m, $J = 15.5 + 5.4 + 4.5$ Hz, 1H, C4-H_{eq}), 2.73 (m, $J = 11.6 + 5.4 + 5.4$ Hz, 1H, C3-H_{eq}), 2.83 (m, $J = 11.6 + 8.0 + 4.5$ Hz, 1H, C3-H_{ax}), 4.73 (d, $J = 4$ Hz, 1H, C1-H), 5.72 (dd, $J = 4 + 1$ Hz, 1Hz, C9-H), 5.99 + 6.15 (s-s, 2-2H, OCH_2O), 5.90 (d, $J = 7 + 1$ Hz, 1H, C2'-H), 6.60 (s, 1H, C5-H), 6.74 (s, 1H, C8-H), 6.83 (d, $J = 7$ Hz, 1H, C3'-H), see table I for nOe data; ^{13}C NMR (CDCl_3) δ 29.72 (C4), 40.65 (C3), 56.76 (C1), 84.62 (C9), 100.91 + 103.15 (OCH_2O), 106.28 (C8), 109.16 (C5), 109.88 (C6'), 113.21 (C2'), 115.29 (C3'), 124.93 (C4a), 130.80 (C8a), 139.78 (C1'), 144.49 (C5'), 146.07 and 146.54 (C6 and C7), 148.95 (C4'), 167.48 (C=O). **10**: Anal $\text{C}_{19}\text{H}_{15}\text{NO}_6$ (C, H, N, O); mass spectrum m/z (%) 353 (M^+ , 1), 335 (3), 176 (100), 149 (4); IR (KBr) C=O, 1750 cm^{-1} ; ^1H NMR (CDCl_3) δ 1.80 (br s, 1H, NH), 2.55–2.80 (m, 2H, C4-H₂), 2.90–3.20 (m, 2H, C3-H₂), 4.45 (d, $J = 3.2$ Hz, 1H, C1-H), 5.77 (d, $J = 3.2$ Hz, 1H, C9-H), 5.88 + 6.16 (d, 4H, OCH_2O), 6.55 (s, 1H, C5H), 6.70 (s, 1H, C8-H), 6.93 (d, $J = 7$ Hz, 1H, C2'-H), 7.08 (d, $J = 7$ Hz, 1H, C3'-H), see table I for nOe data; ^{13}C NMR (CDCl_3) δ 29.91 (C4), 41.52 (C3), 58.04 (C1), 84.63 (C9), 100.82 + 103.35 (OCH_2O), 106.21 (C8), 109.11 (C5), 109.78 (C6'), 113.65 (C2'), 114.62 (C3'), 126.26 (C4a), 130.32 (C8a), 140.42 (C1'), 144.85 (C5'), 145.94 and 146.45 (C6 and C7), 149.23 (C4'), 167.24 (C=O).

(+)-[1S,9R]- and (-)-[1R,9S]-Norbicuculline **9a** and **9b**

To the solution of **9** (200 mg, 0.6 mmol) in acetone (12 mL), a solution of (-)-*O,O*-dibenzoyltartaric acid (DBTA, 200 mg, 0.6 mmol) in acetone (4 mL) was added. After stirring for 30 min the mixture was left to stand at room temperature for 2–3 h. The tartrate salt was filtered, recrystallized from acetone, then dissolved in water (20 mL) treated with conc NH_4OH (pH 8), and extracted with CH_2Cl_2 . Having distilled off the solvent the residue was crystallized from MeOH to produce **9a** (80 mg, 40%) mp 195–197 °C; optical rotation ($c = 1$, CHCl_3) $[\alpha]_{\text{D}}^{25} = +256^\circ$; enantiomeric purity (HPLC) >99.5%. Mother liquor treated with (+)-DBTA (100 mg, 0.3 mmol) similarly gave **9b** (90 mg, 45%); optical rotation ($c = 1$, CHCl_3) $[\alpha]_{\text{D}}^{25} = -250^\circ$; enantiomeric purity (HPLC) >99.5%.

(+)-[1S,9R]- and (-)-[1R,9S]-Bicuculline **11a** and **11b**

To **9a** or **9b** (200 mg, 0.58 mmol each) dissolved in HCOOH (2.5 mL), 37% HCHO was added (0.5 mL) and stirred at 100 °C for 15 min, then evaporated in vacuo. The residue was dissolved in 10% HCl (20 mL), neutralized with conc NH_4OH

and extracted with CH_2Cl_2 . The organic phase was dried over MgSO_4 , evaporated and purified on 20 g 63-200 mesh Kieselgel 60 column (eluent: $\text{CHCl}_3/\text{MeOH}$, 10:1) to yield **11a** identical with an authentic sample (170 mg, 81%), mp 192–193 °C: optical rotation ($c = 1$, CHCl_3) $[\alpha]_{\text{D}}^{25} = +126^\circ$ (ref [6]: $[\alpha]_{\text{D}}^{20} = +132.7^\circ$, $c = 0.49$, CHCl_3); and **11b** (175 mg, 84%): optical rotation ($c = 1$, CHCl_3) $[\alpha]_{\text{D}}^{25} = -124.8^\circ$ (ref [6]: $[\alpha]_{\text{D}}^{33} = -128^\circ$, $c = 0.27$, CHCl_3).

(+)-[1SR,9SR]-Adlumidine **12**

Compound **10** (100 mg, 0.29 mmol) was methylated as above. Purification on 10 g 63-200 mesh Kieselgel 60 column (eluent: $\text{CHCl}_3/\text{MeOH}$, 20:1) gave **12** (70 mg, 67%) mp 201–202 °C (205 °C in ref [7]); ^1H NMR nOe data are summarized in table I.

Inhibition of $[^3\text{H}]\text{GABA}$ binding and GABA-specific $^{36}\text{Cl}^-$ ion flux

Cortical membranes in 50 mM Tris-HCl pH 7.1 buffer were incubated in the dark for 40 min with 4 nM $[^3\text{H}]\text{GABA}$ at 4 °C in the presence or absence of test compounds [3]. Stock solutions (2 mM) of PIQ derivatives were freshly prepared in diluted HCl solution (pH 3) and stored in the dark on ice until use. After incubating **9a** and **11a** with membranes for 40 min at 4 °C, relative changes in UV absorbances at $\lambda_{\text{max}2} = 326$ nm (A_2) and $\lambda_{\text{max}1} = 292$ nm (A_1), A_2/A_1 , indicated less than 5–6% opening of the phthalide ring [17]. Non-specific binding was determined in the presence of 46 μM **11a**. Goodness of fit for one-site ligand analysis [18] of the displacement experiment was expressed as the reduced χ_r^2 value (table II.) Saturation data with 2–250 nM $[^3\text{H}]\text{GABA}$ indicated 0.8 pmol/mg protein density of **11a**-sensitive binding sites.

For GABA-specific $^{36}\text{Cl}^-$ transport, the fast kinetic technique [19–21] was applied. In the presence or absence of 46 μM **9a**, **9b**, **11a** or **11b**, a cortical membrane vesicle suspension [21] (0.225 mL) in 10 mM HEPES buffered physiological salt solution (HBSS, pH 7.5) was rapidly mixed with 0.225 mL of HBSS containing 15 $\mu\text{Ci/mL}$ $^{36}\text{Cl}^-$ ion, 40 μM GABA and incubated for 6 [19, 20] at 30 °C; under the conditions the A_2/A_1 ratio for **9a** and **11a** did not change.

Acknowledgment

This work was supported by EGIS Pharmaceuticals (Budapest, Hungary) and by grants OTKA 1762 (Hungary) and US-Hungarian JF 277 (JK).

References

- 1 Curtis DR, Duggan AW, Felix D, Johnston GAR (1970) *Nature (Lond)* 226, 1222–1224
- 2 Simonyi M, Blaskó G, Kardos J, Kajtar M (1989) *Chirality* 1, 178–179
- 3 Kardos J, Blaskó G, Kerekes P, Kovács I, Simonyi M (1984) *Biochem Pharmacol* 33, 3537–3545
- 4 Manske RHF (1933) *Can J Chem* 8, 142–146
- 5 Glasby JS (1975) In: *Encyclopedia of the Alkaloids Vol 1*, Plenum Press, New York
- 6 MacLean DB (1985) In: *The Alkaloids Vol 24* (Brossi A, ed) Academic Press, Orlando, pp 253–286
- 7 Groenewoud PWG, Robinson R (1936) *J Chem Soc* 199–202
- 8 Blaskó G, Kerekes P, Makleit S (1987) In: *The Alkaloids Vol 31* (Brossi A, ed) Academic Press, San Diego, pp 1–18
- 9 Perkin Jr WH, Ray JN, Robinson R (1925) *J Chem Soc* 127, 740–744
- 10 Kerekes P, Bognár R (1971) *J Prakt Chem* 313, 923–928
- 11 Varga Zs, Blaskó G, Dornyei G, Szántay Cs (1991) *Acta Chim Hung* 128, 831–837
- 12 Simonyi M, Gal J, Testa B (1989) *Trends Pharmacol Sci* 10, 349–354
- 13 Schales O (1935) *Berichte* 68, 1579–1581
- 14 Ziegler FE, Fowler KW (1976) *Org Chem* 41, 1564–1566
- 15 Kövér KE, Kerekes P (1986) *Magn Res Chem* 24, 113–122
- 16 Olsen RW (1976) In: *GABA in Nervous System Function* (Roberts E, Chase TN, Tower DB, eds) Excerpta Medica and American Elsevier Publishing Co, Amsterdam, 287–304
- 17 Olsen RW, Ban M, Miller T, Johnston GAR (1975) *Brain Res* 98, 383–387
- 18 Munson PJ, Rodbard D (1980) *Anal Biochem* 107, 220–239
- 19 Cash DJ, Subbarao K (1987) *Biochemistry* 26, 7556–7562
- 20 Cash DJ, Subbarao K (1987) *Biochemistry* 26, 7562–7570
- 21 Serfőz P, Cash DJ (1992) *FEBS Lett* 310, 55–59