

Synthesis of Fluorinated Amide Derivatives via a Radical N-Perfluoroalkylation–Defluorination Pathway

 Zhiyao Zheng,[†] Angela van der Werf,[†] Marie Deliaval, and Nicklas Selander*

 Cite This: <https://dx.doi.org/10.1021/acs.orglett.0c00768>


Read Online


ACCESS |

Metrics & More

Article Recommendations

Supporting Information

ABSTRACT: A one-pot approach to fluorinated hydroxamic acid, amide, and thioamide derivatives is reported. The reaction proceeds via an N-perfluoroalkylation of nitrosoarenes with perfluoroalkanesulfonates, resulting in labile N-perfluoroalkylated hydroxylamines. By the addition of suitable additives, a controllable oxy/thiodeflourination of the fluorinated hydroxylamine intermediates was achieved. The method highlights N-perfluoroalkylated amines as versatile intermediates for further synthesis.



N-Perfluoroalkylation - Defluorinative Functionalization:


The incorporation of fluorine into organic compounds has become increasingly important for pharmaceutical and agrochemical sciences as the presence of a fluorinated moiety can drastically change physical, chemical, and biological properties.¹ Thus, a large number of methods for the direct introduction of perfluoroalkyl groups (most commonly CF_3 groups), onto C-, O-, and S-atoms² via electrophilic, nucleophilic,⁴ and radical⁵ pathways have been reported. The corresponding N-perfluoroalkylation is, however, less investigated.⁶ One reason is the poor stability of perfluoroalkylated amines, as they are prone to fluoride elimination, assisted by the nitrogen lone pair,⁷ resulting in highly electrophilic fluoroiminium species. Although the high reactivity of these intermediates is challenging to control, the synthesis and direct utilization of perfluoroalkyl amino compounds is an attractive method for the diversification of fluorinated compounds.


Since the seminal work of Yarovenko,⁸ Ishikawa,⁹ and Petrov¹⁰ on the *in situ* preparation of fluorinated ethylamine and its use as a fluorination agent for the conversion of alcohols into alkyl fluorides, the direct functionalization of perfluoroalkyl amines has received limited attention. Recently, Leroux reported an elegant method for the activation of fluorinated ethylamines via fluoride abstraction by a Lewis acid (e.g., $\text{BF}_3 \cdot \text{OEt}_2$) to afford relatively stable iminium salts.^{7c} The nucleophilic addition of electron-rich arenes or heteroarenes enabled the synthesis of various aryl or heteroaryl fluoromethyl ketones (Scheme 1a).¹¹ Furthermore, trapping with electron-rich olefins or C–H acidic compounds, followed by cyclization with hydrazine or hydroxylamine, provided access to a wide range of fluorinated pyrazoles and isoxazoles.^{11,12}

A recent report by Baidya and co-workers disclosed an N-selective nitroso aldol reaction between *gem*-difluoroenolates and nitrosoarenes. The difluoroalkyl intermediate underwent an intramolecular nucleophilic substitution, followed by a rearrangement via C–F and N–O bond cleavage to furnish α -ketoamides (Scheme 1b).¹³

Scheme 1. Defluorination of Fluorinated Amines

 a) Leroux: Fluoroalkyl amino reagent for difluoromethylation and acylation:^{11,12}

 b) Baidya: Cascade nitroso aldol reaction and defluorination:¹³


c) This work: one-pot synthesis of fluorinated amides and thioamides:



Despite the above-mentioned progress, novel reactivity discoveries of perfluoroalkyl amines are still rare, and the direct transformation of these intermediates incorporating a


Received: February 28, 2020

controllable defluorination process is desirable. Recently, we developed a radical N-trifluoromethylation of nitrosoarenes^{6b} with sodium triflinate to obtain a range of isolable N-trifluoromethylated hydroxylamines. However, when longer-chain perfluoroalkanesulfonates^{5a,14} were used, we observed a decomposition of the perfluorinated hydroxylamine upon attempted isolation. As one of the decomposition products was identified as a hydroxamic acid derivative (formal oxydefluorination at the α -position), we set out to investigate a controllable defluorination. Notably, fluorinated amides are important motifs in biologically active molecules¹⁵ and valuable building blocks for pharmaceuticals,¹⁶ dyes,¹⁷ and functional materials.¹⁸ Only a few synthesis methods for perfluorinated amides are reported;^{15b,17,19} most of them have a limited substrate scope and rely on sensitive acid derivatives or coupling reagents.

Herein, we report a one-pot N-perfluoroalkylation reaction of nitrosoarenes for the synthesis of fluorinated hydroxamic acids, amides, and thioamides via N-perfluorinated hydroxylamine intermediate **2** (Scheme 1c).

We first investigated the scope of the perfluoroalkylation reaction with respect to various nitrosoarenes and sodium perfluoroalkanesulfonates (for details, see the SI, Table S-1). As judged by ^1H NMR, a variety of nitrosoarenes bearing either electron-withdrawing or electron-donating substituents at the *ortho*-, *meta*-, or *para*-positions were transformed into the corresponding hydroxylamines in moderate to high yields (**2a–2q**, 56–88% by NMR). In addition, the sodium sulfinate component could be varied to include both shorter (perfluoropropyl) and longer (8*H*-perfluorooctyl) perfluoroalkyl chains (**2s–2u**, 68–80%, Scheme 2).

Scheme 2. N-Perfluoroalkylation of Nitroso Compounds^a



^aGeneral Conditions: **1** (0.50 mmol), $\text{NaSO}_2\text{CF}_2\text{R}_F$ (1.5 mmol), hydroquinone (0.55 mmol), $\text{Cu}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ (1 mol %), EtOAc (4.0 mL), and $^t\text{BuOOH}$ (70% aq, 1.5 mmol). ^bAfter adding Ac_2O (1.0 mL) and NaHCO_3 (3.0 mmol), stirring for 24 h, isol. yield.

Although the hydroxylamines **2** were not isolable, electron-deficient hydroxylamine products were found to be more stable; acetylated hydroxylamine **6q** could be isolated in 61% yield after the acetylation of **2q** (Scheme 2). An aliphatic nitroso compound was a suitable substrate as well. However, the corresponding hydroxylamine was found to be the least stable; oxydefluorinated product **7r** was obtained in 54% isolated yield, instead of the expected hydroxylamine.

Intrigued by the oxydefluorination reaction observed for product **7r** and the relative stability of hydroxylamines **2** in solution, we set out to find general conditions facilitating the oxydefluorination in a one-pot process. As shown in Table 1, stirring the reaction mixture at a higher temperature for an

extended time was beneficial for the oxydefluorination, yielding hydroxamic acid **7b** in 25% yield (entry 1).

Table 1. Condition Screening for Oxydefluorination^a

1b	<i>N</i> -perfluoroalkylation	2b	Cond. for oxydefluorination	7b R = OH, 3b R = H, 4b R = OAc
			85% NMR yield (Method A) 75% NMR yield (Method B)	
1	50 °C, 24 h		25	-
2 ^c	HCl (20 equiv), rt, 6 h		36	-
3 ^{c,d}	HCl (20 equiv), HOAc, 0 °C, 6 h	41 (35) ^e	-	-
4	InCl_3 (3 equiv), 50 °C, 6 h	26/24 ^f	-	-
5	$\text{BF}_3 \cdot \text{OEt}_2$ (10 equiv), rt, 6 h	24 ^f	-	-
6 ^{c,d}	Zn/HCl (40 equiv), HOAc, 65 °C, 6 h	-	72 (70) ^e	-
7	NaOAc (6 equiv), rt, 16 h	-	-	69 (64) ^e
8 ^f	NaOAc (6 equiv), rt, 16 h	-	-	68 (64) ^e


^aMethod A: **1b** (0.10 mmol), $\text{NaSO}_2(\text{CF}_2)_3\text{CF}_3$ (0.3 mmol), hydroquinone (0.11 mmol), $\text{Cu}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ (1 mol %), EtOAc (0.8 mL), and $^t\text{BuOOH}$ (70% aq, 0.3 mmol), rt, 1 h. ^b ^{19}F NMR yields determined with HFIP as an internal standard. ^c37% HCl was used. ^d1.0 mL of HOAc was used. ^eIsolated yield (0.5 mmol). ^fMethod B: CuCl_2 (5 mol %) and $^t\text{BuOOH}$ (55% in decane, 0.3 mmol) were used instead of $\text{Cu}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ and aq. $^t\text{BuOOH}$.

We then screened a variety of Brønsted and Lewis acids for their effects on the oxydefluorination (see also the Supporting Information). Using HCl as a promoter led to an increased yield of **7b** (36%, entry 2), while AcOH as a cosolvent resulted in a slight improvement: the reaction at 0 °C afforded hydroxamic acid **7b** in 41% yield (35% isolated, entry 3). Our attempts to use InCl_3 or $\text{BF}_3 \cdot \text{OEt}_2$ as promoters were unsuccessful, both when the perfluoroalkylation was conducted under nondry conditions (Method A) and when TBHP in decane (Method B) was used (entries 4 and 5). Gratifyingly, under acidic reductive conditions (Zn/HCl in AcOH), amide **3b** was obtained in 72% yield after oxydefluorination and N–O bond reduction (entry 6).

Furthermore, we examined the effect of bases on the oxydefluorination. While K_2CO_3 and pyridine led to decomposition, intermediate **2b** was converted into the corresponding O-acetylated hydroxamic acid using NaOAc. The unexpected product **4b** was isolated in 64% yield (entries 7 and 8).

It is worth to mention that when moderate yields were obtained, no evident side products were identifiable in these one-pot reactions. If the oxydefluorination was not efficient, the hydroxylamine intermediates decomposed, and complex mixtures were obtained.

Having evaluated the oxydefluorination of hydroxylamine intermediates **2**, we examined the scope of this one-pot protocol for the synthesis of amides **3** (Scheme 3).²⁰ Nitrosobenzene was transformed into amide **3a** in 55% yield via the one-pot perfluoroalkylation/oxydefluorination under reductive conditions. High yields were generally obtained for ortho-substituted amides (**3b**, **3e**, **3g**, and **3h**, 53–70%), with the exception of **3c** (42% yield). Halogen substituents were also well tolerated (**3g–j**, 53–70%). The meta-CF₃-substituted


Scheme 3. Synthesis of Amides 3 with Zn/HCl^a

^aReaction conditions: Step (1) Method A (*cf., Table 1*), 0.50 mmol scale. Step (2) HOAc (5 mL), Zn (powder, 20 mmol), and HCl (37% aq, 20 mmol) were added to the reaction mixture, 2 h at 65 °C. Isolated yields over two steps. ^b1.0 mmol scale. ^cMethod B (*cf., Table 1*) was used for Step (1), 0.50 mmol scale. ^d0.20 mmol scale.

amide 3k was obtained in a low yield (31%, Method A), whereas dry conditions (Method B) led to a slightly higher yield (38%). A range of para-substituted amides were obtained in good yields (3l–p, 49–63%), including electron-rich (3l and 3m, 60 and 49%) and electron-poor (3n and 3o, 63 and 61%) arenes. Notably, aminoglutethimide derivative 3p was obtained in 55% yield. Amides derived from other perfluoroalkanesulfonates were isolated in similar yields (3s–u, 58–64%).

The oxydefluorination reaction of *ortho*-substituted substrates, with sodium acetate as a promoter, delivered *O*-acyl hydroxamic acids 4b–f in high yields (59–78%, Scheme 4). Electron-rich and electron-poor arenes as well as *ortho*-disubstituted substrates worked well using nondry reaction conditions (Method A). For product 4f, the yield was improved from 44 to 62% by the addition of 1,1,2,2-tetrachloroethane (TCE) as a cosolvent.

However, for substrates lacking an *ortho*-substituent (4a, 4j, 4k, and 4n), dry reaction conditions (Method B) and a cosolvent were necessary to obtain high yields. For example,

Scheme 4. Oxydefluorination with NaOAc^a

^aReaction conditions: Step (1) Method A, 0.50 mmol scale. Step (2) NaOAc (3.0 mmol) was added to the reaction mixture, 16 h at rt. Isolated yields over two steps. ^b¹⁹F NMR yields determined with HFIP as an internal standard, using Method A. ^cIsolated yields using Method B for Step (1), 0.50 mmol scale. DCM (4.0 mL) was added as a cosolvent in Step 2. ^dTCE (4.0 mL) was added as a cosolvent in Step 2.


only traces of 4k were observed using Method A, whereas 44% yield was reached using Method B (*Scheme 4*).

In analogy with NaOAc, we next explored KSAc as an additive for the defluorinating functionalization. We envisioned that thioamide derivatives could be obtained under similar conditions. Such a procedure could serve as an alternative method for the synthesis of fluorinated thioamides.²¹ Unlike the oxydefluorination with NaOAc, the thioacetate-promoted reaction led to the formation of thioamides 5, involving a concomitant cleavage of the N–O bond. Additionally, dry conditions (Method B) had to be used in order to obtain the thioamide products (*Scheme 5*).



Substrates bearing *ortho*-substituents reacted smoothly, affording thioamides 5b and 5c in 68 and 63% yield, respectively. For other substrates, a cosolvent was required (5a, 5j, 5k, and 5o). Nitrosobenzene was transformed into thioamide 5a in 57% yield, whereas the meta-substituted thioamides 5j and 5k were obtained in slightly lower yields (47–50%). Ester derivative 5o was isolated in 62% yield.

To demonstrate the applicability of our method, amide 3b was reduced by LiAlH₄ into amine 9 in 65% yield. Furthermore, thioamide 5c was efficiently transformed into benzothiazole 10 in 70% yield, via an oxidative cyclization using CAN.²² The S-phenyl thioimidate 11 could be obtained in 88% yield from 5c by reaction with a diaryliodonium salt (*Scheme 6*).²³

A proposed mechanism for the defluorinating functionalization is presented in *Scheme 7*. The defluorination of hydroxylamine intermediate 2 is likely facilitated by the nitrogen lone pair^{7,13} and hyperconjugation from the perfluoroalkyl group, yielding oxaziridine A.²⁴ Our attempts to detect intermediate A (by NMR and MS) were fruitless, as a rapid conversion into the products was observed. In the

Scheme 5. Synthesis of Thioamides 5^a

^aReaction conditions: Step (1) Method B, 0.50 mmol scale. Step (2) KSAC (3.0 mmol) was added to the reaction mixture 16 h, rt. Isolated yields over two steps. ^bDCM (4.0 mL) was added as a cosolvent in Step (2).

Scheme 6. Synthesis Applications**Scheme 7. Mechanistic Proposal**

presence of Zn/HCl, A is reduced to iminoyl fluoride B, which after hydrolysis affords amide 3 (Scheme 7).

In the presence of sodium acetate, O-acyl hydroxamic acid 4 can be obtained via a nucleophilic attack by acetate, followed by a rearrangement of intermediate C. To confirm that the acetyl group originates from sodium acetate (and not EtOAc), the reaction was performed with sodium propionate, yielding O-propionyl hydroxamic acid 8b in 44% yield (see the SI). When potassium thioacetate is used as a nucleophile, the opening of oxaziridine A by thioacetate gives intermediate D.

An analogous rearrangement to E, followed by a cleavage of the N–O and C–S bonds,²⁵ affords iminoyl fluoride B, acetate, and sulfur. The subsequent nucleophilic addition of thioacetate gives intermediate F, which then is hydrolyzed to form thioamide product 5. Upon GCMS analysis of the crude reaction mixtures for thioamides 5a and 5b, we detected the corresponding intermediates F, in support of our mechanistic hypothesis.

In summary, an efficient one-pot N-perfluoroalkylation–defluorination functionalization was developed. After perfluoroalkylation of the nitrosoarene starting materials, labile perfluoroalkyl hydroxylamines were obtained. Although not isolable, the hydroxylamine intermediates were readily converted into a variety of perfluoroalkylated amides, hydroxamic acids, and thioamides, through a controllable defluorination pathway. This work demonstrates the versatility of fluorinated hydroxylamines as intermediates for the synthesis of novel fluorine-containing targets.

■ ASSOCIATED CONTENT

SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.orglett.0c00768>.

Detailed experimental data and characterization data (PDF)

■ AUTHOR INFORMATION

Corresponding Author

Nicklas Selander – Department of Organic Chemistry, Arrhenius Laboratory, Stockholm University SE-106 91 Stockholm, Sweden; orcid.org/0000-0002-9085-0476; Email: nicklas.selander@su.se

Authors

Zhiyao Zheng – Department of Organic Chemistry, Arrhenius Laboratory, Stockholm University SE-106 91 Stockholm, Sweden; orcid.org/0000-0003-2187-675X

Angela van der Werf – Department of Organic Chemistry, Arrhenius Laboratory, Stockholm University SE-106 91 Stockholm, Sweden; orcid.org/0000-0001-6517-8879

Marie Deliaval – Department of Organic Chemistry, Arrhenius Laboratory, Stockholm University SE-106 91 Stockholm, Sweden; orcid.org/0000-0003-1396-2818

Complete contact information is available at: <https://pubs.acs.org/10.1021/acs.orglett.0c00768>

Author Contributions

[†]Z.Z. and A.v.d.W. contributed equally to this work.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank the Knut och Alice Wallenbergs Foundation (Dnr: 2018.0066) for the financial support, the Carl Trygger Foundation for a postdoc scholarship (Z.Z.), and Erika Linde for initial studies.

■ REFERENCES

- (a) Meanwell, N. A. *J. Med. Chem.* **2018**, *61*, 5822. (b) Zhou, Y.; Wang, J.; Gu, Z.; Wang, S.; Zhu, W.; Aceña, J. L.; Soloshonok, V. A.; Izawa, K.; Liu, H. *Chem. Rev.* **2016**, *116*, 422. (c) Gillis, E. P.;

- Eastman, K. J.; Hill, M. D.; Donnelly, D. J.; Meanwell, N. A. *J. Med. Chem.* **2015**, *58*, 8315. (d) Fujiwara, T.; O'Hagan, D. *J. Fluorine Chem.* **2014**, *167*, 16. (e) Jeschke, P. *Pest Manage. Sci.* **2010**, *66*, 10. (f) Smart, B. E. *J. Fluorine Chem.* **2001**, *109*, 3.
- (2) (a) Ni, C.; Hu, J. *Chem. Soc. Rev.* **2016**, *45*, 5441. (b) Yang, X.; Wu, T.; Phipps, R. J.; Toste, F. D. *Chem. Rev.* **2015**, *115*, 826. (c) Alonso, C.; Martinez de Marigorta, E.; Rubiales, G.; Palacios, F. *Chem. Rev.* **2015**, *115*, 1847. (d) Merino, E.; Nevado, C. *Chem. Soc. Rev.* **2014**, *43*, 6598. (e) Liang, T.; Neumann, C. N.; Ritter, T. *Angew. Chem., Int. Ed.* **2013**, *52*, 8214. (f) Ni, C.; Hu, M.; Hu, J. *Chem. Rev.* **2015**, *115*, 765. (g) Xu, X.-H.; Matsuzaki, K.; Shibata, N. *Chem. Rev.* **2015**, *115*, 731.
- (3) (a) Charpentier, J.; Früh, N.; Togni, A. *Chem. Rev.* **2015**, *115*, 650. (b) Barata-Vallejo, S.; Lantaño, B.; Postigo, A. *Chem. - Eur. J.* **2014**, *20*, 16806.
- (4) (a) Liu, X.; Xu, C.; Wang, M.; Liu, Q. *Chem. Rev.* **2015**, *115*, 683. (b) Dilman, A. D.; Levin, V. V. *Eur. J. Org. Chem.* **2011**, *2011*, 831.
- (5) (a) Zhao, Y.; Liu, F. *Tetrahedron Lett.* **2018**, *59*, 180. (b) Barata-Vallejo, S.; Cooke, M. V.; Postigo, A. *ACS Catal.* **2018**, *8*, 7287. (c) Song, H.-X.; Han, Q.-Y.; Zhao, C.-L.; Zhang, C.-P. *Green Chem.* **2018**, *20*, 1662. (d) Lefebvre, Q. *Synlett* **2017**, *28*, 19. (e) He, Z.; Tan, P.; Ni, C.; Hu, J. *Org. Lett.* **2015**, *17*, 1838. (f) Ji, Y.; Brueckl, T.; Baxter, R. D.; Fujiwara, Y.; Seiple, I. B.; Su, S.; Blackmond, D. G.; Baran, P. S. *Proc. Natl. Acad. Sci. U. S. A.* **2011**, *108*, 14411. (g) Fujiwara, Y.; Dixon, J. A.; O'Hara, F.; Funder, E. D.; Dixon, D. D.; Rodriguez, R. A.; Baxter, R. D.; Herlé, B.; Sach, N.; Collins, M. R.; Ishihara, Y.; Baran, P. S. *Nature* **2012**, *492*, 95. (h) Fujiwara, Y.; Dixon, J. A.; Rodriguez, R. A.; Baxter, R. D.; Dixon, D. D.; Collins, M. R.; Blackmond, D. G.; Baran, P. S. *J. Am. Chem. Soc.* **2012**, *134*, 1494.
- (6) (a) Milcent, T.; Crousse, B. C. R. *Chim.* **2018**, *21*, 771. (b) van der Werf, A.; Hriberek, M.; Selander, N. *Org. Lett.* **2017**, *19*, 2374. (c) Scattolin, T.; Deckers, K.; Schoenebeck, F. *Angew. Chem., Int. Ed.* **2017**, *56*, 221. (d) Blastik, Z. E.; Voltrová, S.; Matoušek, V.; Jurásek, B.; Manley, D. W.; Klepetářová, B.; Beier, P. *Angew. Chem., Int. Ed.* **2017**, *56*, 346. (e) Zheng, G.; Ma, X.; Li, J.; Zhu, D.; Wang, M. *J. Org. Chem.* **2015**, *80*, 8910. (f) Teng, F.; Cheng, J.; Bolm, C. *Org. Lett.* **2015**, *17*, 3166. (g) Niedermann, K.; Früh, N.; Senn, R.; Czarniecki, B.; Verel, R.; Togni, A. *Angew. Chem., Int. Ed.* **2012**, *51*, 6511. (h) Niedermann, K.; Früh, N.; Vinogradova, E.; Wiehn, M. S.; Moreno, A.; Togni, A. *Angew. Chem., Int. Ed.* **2011**, *50*, 1059. (i) Umemoto, T.; Adachi, K.; Ishihara, S. *J. Org. Chem.* **2007**, *72*, 6905. (j) Petko, K. I.; Sokolenko, T. M.; Bezduzdny, A. V.; Yagupolskii, L. M. *J. Fluorine Chem.* **2005**, *126*, 1342. (k) Jiang, H.; Chen, Y.; Chen, B.; Xu, H.; Wan, W.; Deng, H.; Ma, K.; Wu, S.; Hao, J. *Org. Lett.* **2017**, *19*, 2406. (l) Scattolin, T.; Bouayad-Gervais, S.; Schoenebeck, F. *Nature* **2019**, *573*, 102. (m) Liang, S.; Wei, J.; Jiang, L.; Liu, J.; Mumtaz, Y.; Yi, W. *Chem. Commun.* **2019**, *55*, 8536.
- (7) (a) Kloeter, G.; Seppelt, K. *J. Am. Chem. Soc.* **1979**, *101*, 347. (b) Hagooly, Y.; Gatenyo, J.; Hagooly, A.; Rozen, S. *J. Org. Chem.* **2009**, *74*, 8578. (c) Commaré, B.; Schmitt, E.; Aribi, F.; Panossian, A.; Vors, J.-P.; Pazenok, S.; Leroux, F. R. *Molecules* **2017**, *22*, 977.
- (8) Yarovenko, N.; Raksha, M. *Zh. Obshch. Khim.* **1959**, *29*, 2159.
- (9) (a) Takaoka, A.; Iwamoto, K.; Kitazume, T.; Ishikawa, N. *J. Fluorine Chem.* **1979**, *14*, 421. (b) Takaoka, A.; Iwakiri, H.; Ishikawa, N. *Bull. Chem. Soc. Jpn.* **1979**, *52*, 3377.
- (10) (a) Petrov, V. A.; Swearingen, S.; Hong, W.; Chris Petersen, W. *J. Fluorine Chem.* **2001**, *109*, 25. (b) Grieco, L. M.; Halliday, G. A.; Junk, C. P.; Lustig, S. R.; Marshall, W. J.; Petrov, V. A. *J. Fluorine Chem.* **2011**, *132*, 1198.
- (11) (a) Schmitt, E.; Rugeri, B.; Panossian, A.; Vors, J.-P.; Pazenok, S.; Leroux, F. R. *Org. Lett.* **2015**, *17*, 4510. (b) Schmitt, E.; Bouvet, S.; Pégot, B.; Panossian, A.; Vors, J.-P.; Pazenok, S.; Magnier, E.; Leroux, F. R. *Org. Lett.* **2017**, *19*, 4960.
- (12) (a) Pazenok, S.; Giornal, F.; Landelle, G.; Lui, N.; Vors, J. P.; Leroux, F. R. *Eur. J. Org. Chem.* **2013**, *2013*, 4249. (b) Giornal, F.; Landelle, G.; Lui, N.; Vors, J.-P.; Pazenok, S.; Leroux, F. R. *Org. Process Res. Dev.* **2014**, *18*, 1002. (c) Schmitt, E.; Landelle, G.; Vors, J.-P.; Lui, N.; Pazenok, S.; Leroux, F. R. *Eur. J. Org. Chem.* **2015**, *2015*, 6052.
- (13) Reddy, M. K.; Ramakrishna, I.; Baidya, M. *Org. Lett.* **2018**, *20*, 4610.
- (14) (a) Zhang, C.-P.; Chen, Q.-Y.; Guo, Y.; Xiao, J.-C.; Gu, Y.-C. *Chem. Soc. Rev.* **2012**, *41*, 4536. (b) Ma, J.; Xie, X.; Meggers, E. *Chem. - Eur. J.* **2018**, *24*, 259. (c) Wu, Z.; Wang, D.; Liu, Y.; Huan, L.; Zhu, C. *J. Am. Chem. Soc.* **2017**, *139*, 1388. (d) Wang, D.; Fang, J.; Deng, G.-J.; Gong, H. *ACS Sustainable Chem. Eng.* **2017**, *5*, 6398. (e) Torviso, M. R.; Mansilla, D.; Garcia, S.; Lantaño, B.; Barata-Vallejo, S.; Postigo, A. *J. Fluorine Chem.* **2017**, *197*, 42. (f) Ma, J.-J.; Liu, Q.-R.; Lu, G.-P.; Yi, W.-B. *J. Fluorine Chem.* **2017**, *193*, 113. (g) Cui, L.; Matusaki, Y.; Tada, N.; Miura, T.; Uno, B.; Itoh, A. *Adv. Synth. Catal.* **2013**, *355*, 2203. (h) Liu, J.-T.; Huang, W.-Y. *J. Fluorine Chem.* **1999**, *95*, 131. (i) Hu, C.-M.; Tang, X.-Q.; Qing, F.-L. *J. Fluorine Chem.* **1992**, *59*, 405. (j) Hu, C.-M.; Qing, F.-L.; Huang, W.-Y. *J. Org. Chem.* **1991**, *56*, 2801. (k) Huang, W.-Y.; Huang, B.-N.; Hu, C.-M. *J. Fluorine Chem.* **1983**, *23*, 193. (l) Huang, W.-Y.; Huang, B.-N.; Wang, W. *Acta Chim. Sinica* **1983**, *41*, 1193.
- (15) (a) Kees, K. L.; Cheeseman, R. S.; Prozialeck, D. H.; Steiner, K. E. *J. Med. Chem.* **1989**, *32*, 11. (b) Kees, K. L.; Smith, T. M.; McCaleb, M. L.; Prozialeck, D. H.; Cheeseman, R. S.; Christos, T. E.; Patt, W. C.; Steiner, K. E. *J. Med. Chem.* **1992**, *35*, 944. (c) Dubowchik, G. M.; Vrudhula, V. M.; Dasgupta, B.; Ditta, J.; Chen, T.; Sheriff, S.; Sipman, K.; Witmer, M.; Tredup, J.; Vyas, D. M.; Verdoorn, T. A.; Bollini, S.; Vinitsky, A. *Org. Lett.* **2001**, *3*, 3987. (d) Prchalová, E.; Štěpánek, O.; Smrk, S.; Kotora, M. *Future Med. Chem.* **2014**, *6*, 1201.
- (16) (a) van Oeveren, A.; Motamedi, M.; Mani, N. S.; Marschke, K. B.; López, F. J.; Schrader, W. T.; Negro-Vilar, A.; Zhi, L. *J. Med. Chem.* **2006**, *49*, 6143. (b) Kim, H. Y.; Wiles, J. A.; Wang, Q.; Pais, G. C. G.; Lucien, E.; Hashimoto, A.; Nelson, D. M.; Thanassi, J. A.; Podos, S. D.; Deshpande, M.; Pucci, M. J.; Bradbury, B. *J. J. Med. Chem.* **2011**, *54*, 3268. (c) Wong, J. C.; Tang, G.; Wu, X.; Liang, C.; Zhang, Z.; Guo, L.; Peng, Z.; Zhang, W.; Lin, X.; Wang, Z.; Mei, J.; Chen, J.; Pan, S.; Zhang, N.; Liu, Y.; Zhou, M.; Feng, L.; Zhao, W.; Li, S.; Zhang, C.; Zhang, M.; Rong, Y.; Jin, T.-G.; Zhang, X.; Ren, S.; Ji, Y.; Zhao, R.; She, J.; Ren, Y.; Xu, C.; Chen, D.; Cai, J.; Shan, S.; Pan, D.; Ning, Z.; Lu, X.; Chen, T.; He, Y.; Chen, L. *J. Med. Chem.* **2012**, *55*, 8903. (d) Fontenelle, C. Q.; Wang, Z.; Fossey, C.; Cailly, T.; Linclau, B.; Fabis, F. *Bioorg. Med. Chem.* **2013**, *21*, 7529. (e) Hirayama, S.; Iwai, T.; Higashi, E.; Nakamura, M.; Iwamatsu, C.; Itoh, K.; Nemoto, T.; Tanabe, M.; Fujii, H. *ACS Chem. Neurosci.* **2019**, *10*, 2237. (f) Koshti, N.; Reddy, G. V.; Jacobs, H.; Gopalan, A. *Synth. Commun.* **2002**, *32*, 3779. (g) Liang, F.; Cho, B. P. *J. Am. Chem. Soc.* **2007**, *129*, 12108. (h) Wang, X.; Edwards, R. L.; Ball, H.; Johnson, C.; Haymond, A.; Girma, M.; Manikkam, M.; Brothers, R. C.; McKay, K. T.; Arnett, S. D.; Osbourn, D. M.; Alvarez, S.; Boshoff, H. I.; Meyers, M. J.; Couch, R. D.; Odom John, A. R.; Dowd, C. S. *J. Med. Chem.* **2018**, *61*, 8847. (i) Cheguillaume, A.; Gillart, J.; Labar, D.; Grégoire, V.; Marchand-Brynaert, J. *Bioorg. Med. Chem.* **2005**, *13*, 1357.
- (17) Kölmel, D. K.; Rudat, B.; Braun, D. M.; Bednarek, C.; Schepers, U.; Bräse, S. *Org. Biomol. Chem.* **2013**, *11*, 3954.
- (18) (a) Mitronova, G. Y.; Belov, V. N.; Bossi, M. L.; Wurm, C. A.; Meyer, L.; Medda, R.; Moneron, G.; Bretschneider, S.; Eggeling, C.; Jakobs, S.; Hell, S. W. *Chem. - Eur. J.* **2010**, *16*, 4477. (b) Wei, Q.; Schlaich, C.; Prévost, S.; Schulz, A.; Böttcher, C.; Gradzielski, M.; Qi, Z.; Haag, R.; Schalley, C. A. *Adv. Mater.* **2014**, *26*, 7358. (c) Jbeily, M.; Schöps, R.; Kressler, J. *J. Fluorine Chem.* **2016**, *189*, 70.
- (19) (a) Ng, C.-O.; Feng, H.; Cheng, S.-C.; Xiao, Y.; Lo, L. T.-L.; Ko, C.-C. *Asian J. Org. Chem.* **2018**, *7*, 1587. (b) Xiong, W.-T.; Zhao, J.-W.; Gu, J.-W.; Zhu, S. *Tetrahedron* **2011**, *67*, 5235. (c) Rota, P.; Allevi, P.; Colombo, R.; Costa, M. L.; Anastasia, M. *Angew. Chem., Int. Ed.* **2010**, *49*, 1850. (d) Katritzky, A. R.; He, H.-Y.; Suzuki, K. *J. Org. Chem.* **2000**, *65*, 8210. (e) Kawa, H.; Ishikawa, N. *J. Fluorine Chem.* **1980**, *16*, 365.
- (20) Since hydroxamic acid **7b** was obtained in an unsatisfied yield, no further investigation of the substrate scope was conducted.

- (21) (a) Portella, C.; Muzard, M.; Bouillon, J.-P.; Brulé, C.; Grellepois, F.; Shermolovich, Y. G.; Timoshenko, V. M.; Chernega, A. N.; Sotoca-Usina, E.; Parra, M.; Gil, S. *Heteroat. Chem.* **2007**, *18*, 500.
(b) Mykhaylychenko, S. S.; Bouillon, J.-P.; Shermolovich, Y. G. *J. Fluorine Chem.* **2009**, *130*, 878.
- (22) Zhu, J.; Xie, H.; Li, S.; Chen, Z.; Wu, Y. *J. Fluorine Chem.* **2011**, *132*, 306.
- (23) Villo, P.; Kervefors, G.; Olofsson, B. *Chem. Commun.* **2018**, *54*, 8810.
- (24) (a) For the reactivity of oxaziridines, see: Arnone, A.; Foletto, S.; Metrangolo, P.; Pregnolato, M.; Resnati, G. *Org. Lett.* **1999**, *1*, 281.
(b) Petrov, V. A.; Resnati, G. *Chem. Rev.* **1996**, *96*, 1809.
- (25) (a) Huisgen, R.; Fisera, L.; Giera, H.; Sustmann, R. *J. Am. Chem. Soc.* **1995**, *117*, 9671. (b) Shangguan, N.; Katukojvala, S.; Greenberg, R.; Williams, L. *J. Am. Chem. Soc.* **2003**, *125*, 7754.
(c) Shioji, K.; Matsumoto, A.; Takao, M.; Kurauchi, Y.; Shigetomi, T.; Yokomori, Y.; Okuma, K. *Bull. Chem. Soc. Jpn.* **2007**, *80*, 743.