


RSC Advances



This article can be cited before page numbers have been issued, to do this please use: W. Xie, Z. Xia, J. Hu, Z. Shen and Q. Yao, *RSC Adv.*, 2015, DOI: 10.1039/C5RA04931H.




This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Dienone-phenol rearrangement of 4,4-disubstituted cyclohexadienones to multiple substituted phenols is achieved by employing Re₂O₇ as catalyst.

COMMUNICATION

Re₂O₇ Catalyzed Dienone-Phenol Rearrangement[†]

Cite this: DOI: 10.1039/x0xx00000x

Zilei Xia,^{a,b,||} Jiadong Hu,^{a,b,||} Zhigao Shen,^{a,b} Qizheng Yao^{*a} and Weiqing Xie^{*b,c}

Received 00th January 2012,

Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/


Dienone-phenol rearrangement of 4,4-disubstituted cyclohexadienones catalyzed by Re₂O₇ was described. Multiple substituted phenols could be efficiently obtained in good to excellent yields by employing this catalytic protocol.

Although various Rhenium complexes have been synthesized and even some of them are commercially available,¹ their catalytic abilities for organic transformations are less explored compared with other transition state metal complexes.² In recent years, increasing interests have been dedicated to Rhenium catalysts for their stability to air and moisture, unique Lewis acidity for activation unsaturated hydrocarbon bonds.^{3,4,5} Especially, the ability of activating C(Sp³)-H and C(Sp³)-H bonds by rhenium carbonyl complexes has enabled them to be amenable for C-H bond functionalization reactions.^{2b,3} Not only those low-valent Rhenium complexes, but also Re(V)⁴ and Re(VII)^{2b,5} complexes are competent catalysts in a variety of organic transformations. For example, Toste and co-workers unveiled that Re(V)-oxo complexes enable converting propargyl alcohols to functionalized intermediates via C-C,^{4b,4c} C-O,^{4e} C-N^{4g} bonds formations. MeReO₃, a Re(VII) complex, is well known for its wide usage in oxidation catalyst.^{5d} Besides its oxidation ability, Re(VII) also exhibit moderate Lewis acidity, which could be utilized for generation of carbocation intermediate under mild reaction conditions. For example, Re₂O₇ and PhSiOReO₃ are popular in stereospecific isomerization of allyl alcohols (Figure 1, eq. 1).^{5e-5h} Those Re(VII) complexes are also used for generation of oxonium ions in acetalization and Prins reaction (Figure 1, eq. 2).^{5i-5l} It's should be pointed out that the high efficiency of those reactions relies on the unique Lewis acid property of Re(VII), which could in turn stabilize the cation intermediate.


Dienone-phenol rearrangement is a rearomatization reaction of 4,4-disubstituted cyclohexadienones through bond shift, which provides a facile access to multi-substituted phenols.⁶ The mechanism

of Dienone-phenol rearrangement has been extensively studied and well employed in organic synthesis.⁷ Typically, the reaction could be promoted by acid, strong base or photo irradiation.⁶ In this context, catalytic dienone-phenol rearrangement are also well described.⁸ For example, Kim developed a domino dienone-phenol rearrangement/5-*endo-dig* cyclization of quinols catalyzed by PtCl₂ to afford benzofurans.^{8c} Fujioka also found that transformation of dienones to benzenethioethers was efficiently achieved by catalytic TfOH.^{8d} More recently, dienone-phenol rearrangement of spiro[4.5]cyclohexadienones catalyzed by Sc(OTf)₃ was described by Hamada and co-workers.^{8e} In this report, we disclose that Re₂O₇ is a robust catalyst for dienone-phenol rearrangement of 4-alkoxy-substituted cyclohexadienones, which is difficult to be furnished by previous catalysts (Figure 1, eq. 3).

isomerization of allylic alcohol: (ref. 5e-5h)



substitution of hemiacetal: (ref. 5i-5l)



dienone-phenol rearrangement: (this work)





Figure 1. Representative Re(VII) catalyzed reactions involving cationic intermediate.

COMMUNICATION

As our interest in application of dienone-phenol rearrangement in synthesis of phenol **2aa**, we found that under the effect of excessive amount of $\text{BF}_3 \cdot \text{OEt}$, cyclohexadienone **1aa** smoothly converted to phenol **2aa** in 73% yield (Table 1, entry 1). In meanwhile, only low yield was resulted when catalytic amount of $\text{BF}_3 \cdot \text{OEt}$ was employed (20%, Table 1, entry 2), which prompted us to investigate reaction conditions to find out a catalytic protocol for this reaction. As summarized in Table 1, most of the evaluated Lewis acids (e.g. $\text{Zn}(\text{OTf})_2$, $\text{Cu}(\text{TFA})_2$, AlCl_3 , entry 3-9) were ineffective for this reaction with starting material being fully recovered after 12h. Previously reported catalysts such as $\text{Sc}(\text{OTf})_3$ ^{8e} and PtCl_2 ^{8c}, which served as efficient catalysts for dienone-phenol rearrangement, were also capable of facilitating the rearrangement of **1aa** albeit in 45% and 55% yield respectively (Table 1, entry 10-11). Other type promoters (e.g. TMSOTf and TfOH^{8d}) gave unsatisfactory results (Table 1, entry 13-14). To our delight, Re_2O_7 worked very well for this rearrangement, providing phenol **2aa** in 92% yield with accelerating reaction rate and other $\text{Re}(\text{VII})$ complexes (e.g. $\text{Ph}_3\text{SiOREO}_3$, MeReO_3) alleviated this reaction (Table 1, entry 14-16). Subsequently, solvent screening showed that highest yield was observed in CH_2Cl_2 than using any other solvents (Table 1, entry 17-19). Low or higher temperature was detrimental to the reaction (Table 1, entry 20 and 21). Significantly, catalyst loading could be reduced to 5 mol% without deleterious effects on the reaction (Table 1, entry 22).


Table 1 Screening of reaction conditions for dienone-phenol rearrangement of quinol **1a**.^a




Entry	Catalyst (equiv)	T (°C)	solvent	Time (h)	Yield ^b (%)
1	$\text{BF}_3 \cdot \text{Et}_2\text{O}$ (3.0)	rt	CH_2Cl_2	4	73
2	$\text{BF}_3 \cdot \text{Et}_2\text{O}$ (0.1)	rt	CH_2Cl_2	4	20
3	$\text{Cu}(\text{TFA})_2$ (0.1)	rt	CH_2Cl_2	12	ND
4	$\text{Zn}(\text{OTf})_2$ (0.1)	rt	CH_2Cl_2	12	ND
5	Cp_2TiCl_2 (0.1)	rt	CH_2Cl_2	12	ND
6	$\text{Yb}(\text{OTf})_3$ (0.1)	rt	CH_2Cl_2	12	ND
7	InCl_3 (0.1)	rt	CH_2Cl_2	12	ND
8	AgBF_4 (0.1)	rt	CH_2Cl_2	12	ND
9	AlCl_3 (0.1)	rt	CH_2Cl_2	12	ND
10	$\text{Sc}(\text{OTf})_3$ (0.1)	rt	CH_2Cl_2	12	45
11	PtCl_2 (0.1)	rt	CH_2Cl_2	4	55
12	TMSOTf (0.1)	rt	CH_2Cl_2	4	60
13	HOTf (0.1)	rt	CH_2Cl_2	4	50
14	Re_2O_7 (0.1)	rt	CH_2Cl_2	1	92
15	$\text{Ph}_3\text{SiOREO}_3$ (0.1)	rt	CH_2Cl_2	1	83
16	MTO (0.1)	rt	CH_2Cl_2	12	ND
17	Re_2O_7 (0.1)	rt	toluene	12	22
18	Re_2O_7 (0.1)	rt	CHCl_3	12	67
19	Re_2O_7 (0.1)	rt	EtOAc	12	16
20	Re_2O_7 (0.1)	0	CH_2Cl_2	12	50
21	Re_2O_7 (0.1)	40	CH_2Cl_2	1	76
22	Re_2O_7 (0.05)	rt	CH_2Cl_2	1	93

^a Reaction conditions: quinol **1aa** (0.1 mmol) in CH_2Cl_2 (0.5 mL) was added dropwise to a solution of catalyst (0.01 mmol) in CH_2Cl_2 (0.5 mL) at rt. ^b Isolated yields.

Upon identification of reaction conditions, the substrate scope of the reaction was subsequently examined. Various cyclohexadienones were prepared by phenol oxidation and subjected to the standard reaction conditions (Scheme 1). Satisfactorily, different alkoxy groups were compatible, affording corresponding phenols in good to excellent yields (**2aa** to **2ag**). Different protecting groups were also surveyed for exploring functional groups compatibility of this reaction. Given the fact that Brønsted acids were usually generated from hydrolysis of Re_2O_7 ,⁵ acid-labile protecting groups (e.g. TBS, Boc for **2ba** and **2bb**) were not well tolerated, leading to low isolated yields due to substantial deprotection.⁹ Moderate yields could be obtained when acid-stable protecting groups were presented (e.g. Ac, Bz, Ts for **2bd-2bf**). Next, different migration groups were evaluated for this reaction. Cyclohexadienones with primary and secondary alkyl groups smoothly transferred to corresponding phenols in very high yields (**2ca** to **2ci**). However, cyclohexadienone **1d** with 4-methyl was inert under reaction conditions, owing to the low migratory aptitude of methyl and moderate Lewis acidity of Re_2O_7 . Tertiary butyl was also an unsuitable migratory group and 4-methoxyphenol was exclusively formed from **1e**, presumably via extrusion of stable tertiary carbocation (see Supporting Information). Disappointingly, only sluggish reaction mixture was obtained when cyclohexadienone **1f** was examined. Furthermore, tetra-substituted phenols could be isolated regioselectively from multiple substituted cyclohexadienone in good yields, with migratory group being attached to less hindered position (**2g** to **2i**). Additionally, the reaction could also be easily scaled up without influencing the isolated yields (**2g** and **2h**).

Scheme 1 Substrate scope of the Re₂O₇ catalyzed dienone-phenol rearrangement.

To demonstrate the efficiency of Re₂O₇ on dienone-phenol rearrangement, a direct comparison with previous protocols was made by using cyclohexadienones **1j** and **1k** as substrates (Scheme 2). In Kita's report,¹⁰ spiro-cyclohexadienone **1j** could be transferred to chroman **2j** quantitatively, promoted by excess amount of montmorillonite K10 (Scheme 2, eq. 1). Pleasingly, under the effect of catalytic Re₂O₇, chroman **2j** could be furnished in comparable yield. Furthermore, rearrangement of cyclohexadienone **1k** led to three regioisomers **2ka**, **2kb** and **2kc** in 1:2.5:1.6 ratios under the effect of Ac₂O/H₂SO₄ as a result of competitive [1,2]-shift and [3,3]-rearrangement of allyl (Scheme 2, eq. 2).¹¹ In sharp contrast, under our reaction conditions only [3,3]-shift product **2kb** and **2kc**¹¹ was obtained in 95% combined yield and 3:5 ratio, showing that reaction pathways was greatly influenced by using Re₂O₇ as catalyst.



Scheme 2 Other substrates.

Conclusions

Re₂O₇ catalyzed dienone-phenol rearrangement reaction is developed. Multiple substituted phenols could be conveniently prepared in good to excellent yields by using this catalytic procedure. In addition, [3,3]-shift product is exclusively formed under the effect of Re₂O₇ when the migratory group is allyl. The highly catalytic efficiency of Re₂O₇ on dienone-phenol rearrangement is attributed to the moderate Lewis acidity of Re₂O₇ and its ability of stabilizing putative phenyl cation intermediate of this reaction.

We are grateful for financial support from the National Natural Science Foundation of China (grand No. 21202187, 21372239).

Notes and references

^a Department of Medicinal Chemistry & State Key Laboratory of Natural Medicines, Center of Drug Discovery, China Pharmaceutical University, 24 Tongjiyaxiang, Nanjing 210009, China, E-mail: qz_yao@aliyun.com

^b Shaanxi Key Laboratory of Natural Products & Chemical Biology, College of Science, Northwest A&F University, 22 Xiong Road, Yangling 712100, Shaanxi, China, E-mail: xiewq@sic.ac.cn

^c State Key Laboratory of Bioorganic & Natural Products Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, Shanghai 200032, China

† Electronic Supplementary Information (ESI) available: General experimental procedures and spectroscopic data for the all compounds.

For ESI or other electronic format See DOI: 10.1039/c000000x/

‡ Those two authors contributed equally.

- (a) G. Wilkinson, F. G. A. Stone and E. W. Abel, *Comprehensive organometallic chemistry: the synthesis, reactions, and structures of organometallic compounds*, 1st edn., Pergamon Press, Oxford Oxfordshire; New York, 1982, Vol. 4, p. 161; (b) E. W. Abel, F. G. A. Stone and G. Wilkinson, *Comprehensive organometallic chemistry II: a review of the literature 1982-1994*, 1st edn., Pergamon, Oxford; New York, 1995 Vol. 6, p. 167; (c) D. M. P. Mingos and R. H. Crabtree, *Comprehensive organometallic chemistry III*, 1st edn., Elsevier, Amsterdam; Boston, 2007 Vol. 5, p. 855.

COMMUNICATION

- 2 For Rhenium-catalyzed reactions, see: (a) R. M. Hua and J. L. Jiang, *Curr. Org. Synth.*, 2007, **4**, 151; (b) S. Bellemin-Laponnaz, *ChemCatChem*, 2009, **1**, 357; (c) Y. Kuninobu and K. Takai, *Chem. Rev.*, 2011, **111**, 1938.
- 3 Selected examples of Re(I) catalyzed reactions: (a) H. Y. Chen and J. F. Hartwig, *Angew. Chem., Int. Edit.*, 1999, **38**, 3391; (b) Y. Kuninobu, A. Kawata and K. Takai, *J. Am. Chem. Soc.*, 2005, **127**, 13498; (c) Y. Kuninobu, T. Matsuki and K. Takai, *Org. Lett.*, 2010, **12**, 2948; (d) Y. Kuninobu, T. Nakahara, P. Yu and K. Takai, *J. Organomet. Chem.*, 2011, **696**, 348; (e) Y. Kuninobu, T. Iwanaga, M. Nishi and K. Takai, *Chem. Lett.*, 2010, **39**, 894; (f) Nishiyama, K. Kaiba and R. Umeda, *Tetrahedron Lett.*, 2010, **51**, 793-795.
- 4 (a) J. J. Kennedy-Smith, K. A. Nolin, H. P. Gunterman and F. D. Toste, *J. Am. Chem. Soc.*, 2003, **125**, 4056; (b) M. R. Luzung and F. D. Toste, *J. Am. Chem. Soc.*, 2003, **125**, 15760; (c) B. D. Sherry, A. T. Radosevich and F. D. Toste, *J. Am. Chem. Soc.*, 2003, **125**, 6076; (d) J. J. Kennedy-Smith, L. A. Young and F. D. Toste, *Org. Lett.*, 2004, **6**, 1325; (e) B. D. Sherry, R. N. Loy and F. D. Toste, *J. Am. Chem. Soc.*, 2004, **126**, 4510; (f) K. A. Nolin, R. W. Ahn and F. D. Toste, *J. Am. Chem. Soc.*, 2005, **127**, 12462; (g) R. V. Ohri, A. T. Radosevich, K. J. Hrovat, C. Musich, D. Huang, T. R. Holman and F. D. Toste, *Org. Lett.*, 2005, **7**, 2501.
- 5 For reviews see: (a) C. C. Romao, F. E. Kuhn and W. A. Herrmann, *Chem. Rev.*, 1997, **97**, 3197; (b) J. H. Espenson, *Chem. Commun.*, 1999, 479; (c) G. S. Owens, J. Aries and M. M. Abu-Omar, *Catal. Today*, 2000, **55**, 317; (d) G. Grigoropoulou, J. H. Clark and J. A. Elings, *Green Chem.*, 2003, **5**, 1; (e) S. Bellemin-Laponnaz and J. P. Le Ny, *Cr. Chim.*, 2002, **5**, 217; selected examples: (f) C. Morrill and R. H. Grubbs, *J. Am. Chem. Soc.*, 2005, **127**, 2842; (g) C. Morrill, G. L. Beutner and R. H. Grubbs, *J. Org. Chem.*, 2006, **71**, 7813-7825. (h) A. T. Herrmann, T. Saito, C. E. Stivala, J. Tom and A. Zakarian, *J. Am. Chem. Soc.*, 2010, **132**, 5962; (i) P. Ghorai and P. H. Dussault, *Org. Lett.*, 2008, **10**, 4577; (j) K. Tadpetch and S. D. Rychnovsky, *Org. Lett.*, 2008, **10**, 4839; (k) P. Ghorai and P. H. Dussault, *Org. Lett.*, 2009, **11**, 213; (l) W. Sittiwong, M. W. Richardson, C. E. Schiaffo, T. J. Fisher and P. H. Dussault, *Beilstein J. Org. Chem.*, 2013, **9**, 1526.
- 6 For reviews see: (a) B. Miller, *Acc. Chem. Res.*, 1975, **8**, 245; (b) B. M. Trost and I. Fleming, *Comprehensive organic synthesis : selectivity, strategy, and efficiency in modern organic chemistry*, 1st edn., Pergamon Press, Oxford, England ; New York, 1991, Vol. 3, p. 803; (c) L. s. Kürti and B. Czako, *Strategic applications of named reactions in organic synthesis : background and detailed mechanisms*, Elsevier Academic Press, Amsterdam ; Boston, 2005, p. 142; (d) J. J. Li, *Name reactions : a collection of detailed reaction mechanisms*, 3rd expanded edn., Springer, Berlin ; New York, 2006, p. 202.
- 7 Selected examples: (a) D. J. Hart, A. Kim, R. Krishnamurthy, G. H. Merriman and A. M. Waltos, *Tetrahedron*, 1992, **48**, 8179; (b) K. A. Parker and Y. H. Koh, *J. Am. Chem. Soc.*, 1994, **116**, 11149; (c) Z. H. Guo and A. G. Schultz, *Org. Lett.*, 2001, **3**, 1177; (d) F. Doi, T. Ogamino, T. Sugai and S. Nishiyama, *Tetrahedron Lett.*, 2003, **44**, 4877.
- 8 Catalytic hexadienone-phenol rearrangement: (a) S. Chalais, P. Laszlo and A. Mathy, *Tetrahedron Lett.*, 1986, **27**, 2627; (b) L. Moisan, M. Wagner, S. Comesse and E. Doris, *Tetrahedron Lett.*, 2006, **47**, 9093; (c) I. Kim, K. Kim and J. Choi, *J. Org. Chem.*, 2009, **74**, 8492; (d) Y. Wada, K. Otani, N. Endo, Y. Kita and H. Fujioka, *Chem. Commun.*, 2010, **46**, 797; (e) M. Yoshida, T. Nozaki, T. Nemoto and Y. Hamada, *Tetrahedron*, 2013, **69**, 9609.
- 9 No deprotection product of **2ba** was isolated.
- 10 K. Hata, H. Hamamoto, Y. Shiozaki, S. B. Cammerer and Y. Kita, *Tetrahedron*, 2007, **63**, 4052
- 11 U. Widmer, J. Zsindely, H.-J. Hansen and H. Schmid, *Helv. Chim. Acta*, 1973, **56**, 75.
- 12 The structure of **2mb** and **2mc** was determined by NMR analysis, see supporting information.