

# Selectivity in Lewis acid-mediated fragmentations of peroxides and ozonides: application to the synthesis of alkenes, homoallyl ethers, and 1,2-dioxolanes †

Patrick H. Dussault,\* Hyung-Jae Lee and Xuejun Liu

Department of Chemistry, University of Nebraska–Lincoln, Lincoln, NE 68588-0304, USA


Received (in Cambridge, UK) 21st February 2000, Accepted 1st June 2000

Published on the Web 15th August 2000

Fragmentation of dialkyl peroxides and ozonides is strongly influenced by the choice of Lewis acid.  $\text{TiCl}_4$  promotes C–O ionization ( $\text{S}_{\text{N}}1$  reaction) of tertiary peroxides while  $\text{SnCl}_4$  and  $\text{BF}_3 \cdot \text{OEt}_2$  promote O–O heterolysis (Hock reaction). The cationic intermediates are trapped with allyltrimethylsilane to afford allylated alkanes and homoallyl ethers. In the absence of a nucleophile, ozonides (1,2,4-trioxolanes) invariably undergo O–O heterolysis. However, the combination of allyltrimethylsilane and  $\text{SnCl}_4$  results in formation of 1,2-dioxolanes *via* trapping of intermediates derived from  $\text{S}_{\text{N}}1$  ionization.

## Introduction

The acid-catalyzed decomposition of peroxides is an important synthetic process for which two major modes of reaction are typically observed. Heterolysis of the O–O bond (Hock cleavage) produces oxycarbenium ion intermediates *via* migration of a neighboring substituent, while acid-catalyzed C–O ionization furnishes carbenium ions (Scheme 1). Selectivity between these



Scheme 1 Modes of Lewis acid (LA) mediated activation.


pathways is largely dependent on substrate structure,<sup>1–5</sup> with C–O ionization favored by stabilization of the resulting carbenium ion and O–O heterolysis favored by neighboring groups of high migratory aptitude. Although there have been a number of reports involving Lewis acid-mediated decomposition of peroxides, there are few data regarding reagent-directed selectivity.<sup>6,7</sup> Our interest arose from studies of Lewis acid-mediated allylation of monoperoxyacetals.  $\text{TiCl}_4$ -promoted reactions were found to proceed with selective displacement of the peroxide to form ethers, whereas  $\text{SnCl}_4$ -promoted reactions resulted in selective displacement of the alkoxide to form allylated peroxides.<sup>8,9</sup> We were interested in using similar methodology to explore the fragmentations of highly substituted peroxides and ozonides, using the relative yields of allylated products as an indicator of the ability of individual Lewis acids to dictate a particular pathway. We now report the effect of peroxide structure and Lewis acid on the fragmentation of dialkyl peroxides and ozonides.

† Experimental results for **22l**, **26** and **27** are available as supplementary data. For direct electronic access see <http://www.rsc.org/suppdata/p1/b001391i/>

## Results and discussion

### Substrate preparation

Dialkyl peroxides were prepared by alkylation of the appropriate hydroperoxide with an alkyl halide or sulfonate in the presence of  $\text{CsOH}$  (Scheme 2).<sup>10</sup> The selective saturation of




Scheme 2 Preparation of peroxide substrates. Reagents and conditions: a.  $\text{CsOH}$ , DMF, MeI or  $\text{MeO}(\text{CH}_2)_2\text{OTs}$ ; b.  $\text{PtO}_2$ ,  $\text{H}_2$ ; c. *tert*-butyl hyponitrite,  $\text{C}_6\text{H}_6$ , 60 °C.

unsaturated peroxide is based upon reported work from our group.<sup>11</sup>


Ozonides **5a–m** were prepared by co-ozonolysis of oximes and ketones,<sup>12</sup> co-ozonolysis of enol ethers and ketones,<sup>13</sup> or by ozonolysis of alkenes (Table 1).

### Fragmentation of hydroperoxides and peroxides

Reaction of 2-phenyl-2-propyl hydroperoxide with Lewis acids in the presence of allyltrimethylsilane furnished low yields of allylcumene **6** (Table 2). 2-Phenyl-2-propyl methyl peroxide (**1b**) underwent  $\text{TiCl}_4$ -mediated allylation to furnish a good yield of **6**, while the corresponding reaction with  $\text{SnCl}_4$  afforded only allylated phenyl ether **7**. Interestingly, reaction with allyltrimethylsilane and  $\text{BF}_3 \cdot \text{OEt}_2$  promoted formation of methyl ether **8b**. Methoxyethyl peroxide **1c** was anticipated to favor heterolysis based upon the ability to enter into a five-membered chelate,<sup>14,15</sup> and this outcome was observed in the presence of  $\text{SnCl}_4$ . However, both  $\text{S}_{\text{N}}1$  and Hock products were obtained in the presence of  $\text{BF}_3 \cdot \text{OEt}_2$  or  $\text{TiCl}_4$ .

**Table 1** Synthesis of ozonides


Ozonide	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	Method	Yield (%)
<b>5a</b>		-(CH <sub>2</sub> ) <sub>4</sub> -		-(CH <sub>2</sub> ) <sub>5</sub> -	A	60
<b>5b</b>		-(CH <sub>2</sub> ) <sub>5</sub> -		-(CH <sub>2</sub> ) <sub>5</sub> -	A	67
<b>5c</b>		-(CH <sub>2</sub> ) <sub>6</sub> -		-(CH <sub>2</sub> ) <sub>5</sub> -	A	63
<b>5d</b>	CH <sub>3</sub>	Ph		-(CH <sub>2</sub> ) <sub>5</sub> -	A	57
<b>5e</b>	C <sub>4</sub> H <sub>9</sub>	C <sub>4</sub> H <sub>9</sub>		-(CH <sub>2</sub> ) <sub>5</sub> -	A	47
<b>5f</b>	C <sub>8</sub> H <sub>17</sub>	H		-(CH <sub>2</sub> ) <sub>5</sub> -	B	47
<b>5g</b>	Ph	H		-(CH <sub>2</sub> ) <sub>5</sub> -	B	65
<b>5h</b>	H	H		-(CH <sub>2</sub> ) <sub>5</sub> -	C	78
<b>5i</b>	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>		-(CH <sub>2</sub> ) <sub>5</sub> -	A	53
<b>5j</b>	H	Ph	H	C <sub>3</sub> H <sub>7</sub>	C	73
<b>5k</b>	H	Ph	H	H	C	77
<b>5l</b>	CH <sub>3</sub>	Ph	H	H	C	61
<b>5m</b>	H		-(CH <sub>2</sub> ) <sub>3</sub> -	CH <sub>3</sub>	C	73

**Table 2** Acid-mediated reactions of 2-phenyl-2-propyl peroxides

Peroxide	Acid	T/°C	t/min	6 (%)	7 (%)	8 (%)
<b>1a</b>	BF <sub>3</sub> ·OEt <sub>2</sub>	RT	30	9	—	—
<b>1a</b>	TiCl <sub>4</sub>	-78	10	18	—	—
<b>1a</b>	SnCl <sub>4</sub>	-78	30	34	—	—
<b>1b</b>	BF <sub>3</sub> ·OEt <sub>2</sub>	0-RT	120	—	—	55
<b>1b</b>	TiCl <sub>4</sub>	-78	10	54	15	—
<b>1b</b>	SnCl <sub>4</sub>	-40	30	—	64	—
<b>1b</b>	SnCl <sub>4</sub>	-78	60	—	74	—
<b>1c</b>	BF <sub>3</sub> ·OEt <sub>2</sub>	RT	90	10	—	30
<b>1c</b>	TiCl <sub>4</sub>	-78	1	56	36	—
<b>1c</b>	SnCl <sub>4</sub>	-20	1	—	17	42
<b>1c</b>	SnCl <sub>4</sub>	-40	2	—	37	37
<b>1c</b>	SnCl <sub>4</sub>	-78	10	—	80	20


Octahydronaphthyl hydroperoxide **2a** underwent rapid fragmentation in the presence of SnCl<sub>4</sub> and allylsilane to afford a mixture of an elimination-prone hydroxyketone (**10a**) and enone **11** (Table 3). The corresponding reaction in the presence of EtAlCl<sub>2</sub> and allylsilane furnished **10a/11** accompanied by a small amount of the allylated S<sub>N</sub>1 adduct **9**.<sup>16</sup> The methyl peroxide, **2b**, underwent TiCl<sub>4</sub>-promoted allylation to afford only **9** whereas reaction with stoichiometric SnCl<sub>4</sub> or catalytic TMSOTf resulted in nearly exclusive (SnCl<sub>4</sub>) or exclusive (TMSOTf) formation of **10b/11**. Reaction of the methoxyethyl peroxide **2c** gave predominately (TiCl<sub>4</sub>) or exclusively (SnCl<sub>4</sub>) the rearrangement-derived ring expansion products **9c/10**. Higher yields of rearrangement product were observed in the presence of tetra-*n*-butylammonium perchlorate, with or without acetonitrile as a cosolvent. No reaction was observed in the presence of allyltributylstannane, presumably reflecting rapid metathesis with SnCl<sub>4</sub> to form a less reactive allylstannane-Lewis acid pair.<sup>17</sup>

The isomeric octahydronaphthyl peroxide (**4b**) underwent SnCl<sub>4</sub>-mediated reaction to furnish a mixture of the allylated octahydronaphthalene and the spirodecanone **12** (Scheme 3). The saturated analog **3** underwent selective consumption of the *cis* isomer to form an allylated epoxydecane (**13**) derived


**Table 3** Allylation of octahydronaphthyl peroxides

Peroxide	L. acid (equiv.)	t/min	9 (%)	[10 + 11] (%)
<b>2a</b>	SnCl <sub>4</sub> (1.0)	5	Decomp.	53
<b>2a</b>	EtAlCl <sub>2</sub> (1.0)	5	18	76
<b>2b</b>	TiCl <sub>4</sub> (1.0)	1	68	—
<b>2b</b>	SnCl <sub>4</sub> (1.0)	3	8	74
<b>2b</b>	TMSOTf (0.1)	5	—	80
<b>2c</b>	TiCl <sub>4</sub> (1.0)	1	11	45
<b>2c</b>	SnCl <sub>4</sub> (1.0)	1	—	67
<b>2c</b>	SnCl <sub>4</sub> (1.0) <sup>a</sup>	5	—	98
<b>2c</b>	SnCl <sub>4</sub> (1.0)	5	—	45
<b>2c</b>	SnCl <sub>4</sub> (1.0) <sup>ab</sup>	5	—	89

<sup>a</sup> Presence of stoichiometric *n*-Bu<sub>4</sub>NClO<sub>4</sub>. <sup>b</sup> 20% MeCN-CH<sub>2</sub>Cl<sub>2</sub> as solvent.

**Table 4** Lewis acid-mediated fragmentation of ozonides

Ozonide	Lewis acid	T/°C	Yield (%)					
			15a	15b	15c	16a	16b	16c
<b>5a</b>	TiCl <sub>4</sub>	-78	18	32	—	30	20	—
	SnCl <sub>4</sub>	0	5	39	—	45	11	—
	TMSOTf	RT	14	36	—	36	14	—
<b>5b</b>	TiCl <sub>4</sub>	-78	—	52	—	—	48	—
	SnCl <sub>4</sub>	0	—	49	—	—	51	—
	TMSOTf	RT	—	50	—	—	50	—
<b>5c</b>	TiCl <sub>4</sub>	-78	—	43	5	—	11	41
	SnCl <sub>4</sub>	0	—	44	5	—	6	45
	TMSOTf	RT	—	49	1	—	1	49

**Scheme 3** Rearrangement of octa- and decahydronaphthyl peroxides.

from migration of the central C-C bond, accompanied by smaller amounts of an allylated oxepane (**14**) derived from migration of a peripheral C-C bond.

### Fragmentation of ozonides

Table 4 summarizes Lewis acid-promoted decomposition of ozonides **5a-c**. The spiro-6,6-ozonide **5b** invariably underwent Hock-type decomposition to afford a 1:1 mixture of caprolactone (**15b**) and cyclohexanone (**16b**). The spiro-6,5- and -6,7-

**Table 5** Lewis acid-mediated allylation of ozonides

Ozonide	R <sup>1</sup>	R <sup>2</sup>	Lewis acid	T/°C	Yield (%)				
					15b	17	18	Ketones	20
5a	—	—	SnCl <sub>4</sub>	−78–0	11	50	—	16a,b (trace)	—
5b	—	—	TiCl <sub>4</sub>	−78	9	0	—	16b,c (35% combined)	—
5b	—	—	SnCl <sub>4</sub>	−78–0	17	57	—	16c (trace)	—
5b	—	—	TMSOTf	−78–0	—	NR	—	—	—
5b	—	—	SbCl <sub>5</sub>	−78–RT	Major	Decomp.	—	16b (major, as mixture)	—
5c	—	—	SnCl <sub>4</sub>	−78–0	39	24	—	16c (trace)	—
5d	CH <sub>3</sub>	Ph	SnCl <sub>4</sub>	−78–0	25	61	—	16d (93%)	—
5e	C <sub>4</sub> H <sub>9</sub>	C <sub>4</sub> H <sub>9</sub>	SnCl <sub>4</sub>	−78–0	40	14	—	16e (70%)	75
5f	H	C <sub>8</sub> H <sub>17</sub>	SnCl <sub>4</sub>	−78	—	56	—	—	50
5g	H	Ph	SnCl <sub>4</sub>	−78	—	79	—	—	13
5h	H	H	SnCl <sub>4</sub>	−78	—	10	—	—	—
5i	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>	SnCl <sub>4</sub>	−78–0	31	21	9 ( <i>cis</i> )	—	—

Ozonide	R <sup>1</sup>	R <sup>2</sup>	22 (% <i>cis</i> : <i>trans</i> )	23 (%)	Ketones (%)	20j (%)	24 (%)
5j	H	C <sub>3</sub> H <sub>7</sub>	15 (1:1)	7	39	20	—
5k	H	H	15 (1:1)	—	22	24	—
5l	CH <sub>3</sub>	H	9 (1:1)	—	43	—	2.5


analogs **5a** and **5c** underwent similar reactions, again favoring the product derived from migration of a cyclohexyl C–C bond. TiCl<sub>4</sub>-promoted reactions occurred at much lower temperatures than those promoted by SnCl<sub>4</sub> or TMSOTf.

#### Fragmentations of ozonides in the presence of allyltrimethylsilane

Reaction of ozonide **5b** with TiCl<sub>4</sub> in the presence of allyltrimethylsilane occurred at −78 °C, producing caprolactone, cyclohexanone, and cycloheptanone (Table 5). However, the corresponding reaction with SnCl<sub>4</sub> occurred only upon warming to ~0 °C and furnished spirodioxolane **17** accompanied by small amounts of cycloheptanone. No reaction was observed in the presence of TMSOTf. Application of the most favorable conditions (SnCl<sub>4</sub>, −78 to 0 °C, allyltrimethylsilane) to a series of tri- and tetrasubstituted ozonides invariably led to the predominant formation of dioxolane **17**. Decomposition of several styrene ozonides proceeded in low yield to furnish as the major product the dioxolane derived from attack on the center better able to stabilize positive charge. Reaction of **5j** also produced significant quantities of benzaldehyde and 1-phenylbut-3-enol while the reaction of **5l** furnished 43% of acetophenone and traces of 4-methyl-4-phenylhepta-1,6-diene.

#### Allylation of a bicyclic ozonide


The SnCl<sub>4</sub>-mediated allylation of a bicyclic ozonide was investigated to determine whether the S<sub>N</sub>1-type reaction proceeded *via* initial ionization of the C–O (etheral) or C–OO (peroxide) linkage. Treatment of ozonide **5m** with SnCl<sub>4</sub> and allyltrimethylsilane produced a good yield of 3,5,5-trisubstituted 1,2-dioxolane (**25**) as a single regioisomer consisting of two separate *cis* and two separate *trans* diastereomers in a 35:35:15:15 ratio (Scheme 4). The presence of a secondary

**Scheme 4** Allylation of a bicyclic ozonide.


alcohol was confirmed by the downfield shift of the H–C–O signal upon formation of acetate **26** and by the disappearance of the same signal upon oxidation to ketone **27**. The stereochemical assignment was supported by the observation of a 70:30 mixture of *cis/trans* two diastereomers upon oxidation to ketone **27**.

#### Discussion

The selective formation of allylcumene and allyloctahydro-naphthalene provides clear evidence for the ability of TiCl<sub>4</sub> to selectively promote S<sub>N</sub>1-type C–O ionization of dialkyl peroxides (Scheme 5). SnCl<sub>4</sub>-mediated allylations proceed with a strong preference for Hock rearrangement. In some cases, the intermediate oxycarbenium ion is diverted into other pathways too rapidly for trapping by allylsilane. For example, the failure to observe ring-expanded allyl ethers from **2a–c** or **4b** reflects rapid intramolecular closure of the intermediate vinyl ketals, an outcome previously observed for Cu(OTf)<sub>2</sub>-mediated rearrangement of allyl hydroperoxides.<sup>18</sup> Similarly, the conversion of 2-phenyl-2-propyl alkyl peroxides to alkyl allyl ethers results from formation and subsequent reaction of mixed acetals; the greater extent of this process in the presence of BF<sub>3</sub>·OEt<sub>2</sub> may indicate the higher nucleophilicity of alkoxyboranes relative to alkoxy-stannanes.



Scheme 5 Overview of peroxide fragmentation.



Scheme 6 Pathways for ozonide ionization.

Ozonides (1,2,4-trioxolanes) can in principle undergo acid-catalyzed decomposition through ionization of peroxide, ionization of alkoxide, or oxygen–oxygen heterolysis (Scheme 6).<sup>19</sup> Acidolysis of ozonides with  $\text{ClSO}_3\text{H}$  or  $\text{SbCl}_5$  has been reported to produce metalated or protonated carbonyl oxides through ionization of the C–OO linkage, followed by loss of carbonyl from the oxycarbenium ion intermediate.<sup>20–22</sup> Self-reaction and/or attack on another ozonide results in formation of cyclic peroxides, while trapping of the metalated carbonyl oxide with 1,1-substituted alkenes produces 1,2-dioxolanes.<sup>23</sup> Related work from our group has demonstrated  $\text{SnCl}_4$ -mediated allylation of ozonides furnishes 1,2-dioxolanes, presumably *via* similar intermediates. In the current study, both  $\text{SnCl}_4$  and  $\text{TiCl}_4$  promote O–O heterolysis. However, in the presence of allyltrimethylsilane, only  $\text{SnCl}_4$  promotes formation of 1,2-dioxolanes. This result indicates that  $\text{SnCl}_4$  is more efficient in promoting reversible ionization of the C–O linkage, that  $\text{TiCl}_4$  promotes more rapid O–O heterolysis, or some combination of these factors. The regioselectivity in allylation of unsymmetrical ozonides supports a mechanism involving ionization of both the C–O and C–OO linkages. In the case of a bicyclic ozonide, only the products corresponding to initial C–O ionization were observed.

In conclusion,  $\text{SnCl}_4$  and  $\text{TiCl}_4$  provide alternate low energy pathways for activation of dialkyl peroxides.  $\text{TiCl}_4$  tends to promote  $\text{S}_{\text{N}}1$  ionization, even when this requires activation of the more hindered oxygen. In contrast,  $\text{SnCl}_4$  generally promotes selective O–O heterolysis through activation of the less hindered peroxide oxygen. Allyltrimethylsilane is an effective trapping agent generally able to capture the carbenium or oxycarbenium intermediates. In the absence of a nucleophile, both  $\text{TiCl}_4$  and  $\text{SnCl}_4$  promote O–O heterolysis of ozonides, while the combination of  $\text{SnCl}_4$  and allyltrimethylsilane furnishes 1,2-dioxolanes through capture of an intermediate peroxy-carbenium ion. Our results demonstrate the ability to dictate the mode of peroxide and ozonide decomposition through selection of Lewis acid, as well as the ability to employ this decomposition for the controlled synthesis of alkenes and allyl ethers.

## Experimental

Standard conditions have been described.<sup>24</sup> As in any work involving peroxides, standard precautions (safety shields, avoidance of heat, light, or metal salts, use of minimal scale) should be faithfully observed.<sup>25–27</sup> NMR spectra were acquired at 300 MHz ( $^1\text{H}$ ) and 75 MHz ( $^{13}\text{C}$ ) in  $\text{CDCl}_3$  unless otherwise noted;  $J$  values are given in Hz; DEPT135 assignments are indicated with (+) or (–) where relevant. IR spectra were recorded as neat oils; C=O and O–H stretches are reported. “Standard work-up” refers to aqueous quench, two-fold ether extraction (25–100 mL, depending upon scale), drying over  $\text{Na}_2\text{SO}_4$ , and concentration *in vacuo*. Unless otherwise indicated, compounds were purified by silica flash chromatography.

### Methyl 2-phenyl-2-propyl peroxide (1b)

To a 0 °C solution of iodomethane (0.84 mL, 13.5 mmol) in DMF (30 mL) under  $\text{N}_2$  was added cumene hydroperoxide **1a** (1.2 mL, 6.8 mmol) followed by  $\text{CsOH}$  (1.7 g, 8.1 mmol). The resulting solution was stirred for 2 h at 0 °C and then submitted to a standard workup, followed by flash chromatography (2.5% ether–pentane) to afford methyl 2-phenyl-2-propyl peroxide **1b** (0.70 g, 63%) as a colorless oil:  $R_f = 0.24$  (2.5% EA–hex);  $^1\text{H}$  NMR  $\delta$  7.54–7.27 (m, 5 H), 3.82 (s, 3 H), 1.64 (s, 6 H);  $^{13}\text{C}$  NMR  $\delta$  145.3, 128.1, 127.0, 125.4, 82.9, 62.9, 26.4; Anal. Calcd for  $\text{C}_{10}\text{H}_{14}\text{O}_2$ : C, 72.26; H, 8.49. Found: C, 72.22; H, 8.41%.

### 2-Methoxyethyl 2-phenyl-2-propyl peroxide (1c)

To a 0 °C solution of 2-methoxyethanol (1.0 mL, 12.7 mmol), and pyridine (2.1 mL, 25.4 mmol) under  $\text{N}_2$  was added *p*-tolylsulfonfyl chloride (2.40 g, 12.7 mmol). After stirring for 1 h, the reaction was quenched with ice-cold 10% HCl (100 mL). The ether extracts (3 × 100 mL) were washed with aqueous saturated NaCl solution (3 × 100 mL) and dried over  $\text{Na}_2\text{SO}_4$ . After removal of solvent *in vacuo*, flash chromatography (20% EA–hex) afforded 2-methoxyethyl toluene-*p*-sulfonate ester (2.5 g, 86%) as a colorless oil:  $R_f = 0.23$  (5% EA–hex);  $^1\text{H}$  NMR  $\delta$  7.76 (d, 2 H,  $J = 8.4$ ), 7.31 (d, 2 H,  $J = 8.1$ ), 4.12 (t, 2 H,  $J = 4.5$ ), 3.54 (t, 2 H,  $J = 4.5$ ), 3.27 (s, 3 H), 2.41 (s, 3 H);  $^{13}\text{C}$  NMR  $\delta$  144.8, 132.9, 129.7, 127.8, 69.9, 69.0, 58.9, 21.5; Anal. Calcd for  $\text{C}_{10}\text{H}_{14}\text{SO}_4$ : C, 52.16; H, 6.13. Found: C, 51.99; H, 6.05%.

Alkylation of cumene hydroperoxide (1.7 mL, 9 mmol) with the methoxyethyl toluene-*p*-sulfonate (2.10 g, 9 mmol) as for **1b**, furnished, after standard work-up and flash chromatography (10% EA–hex), peroxide **1c** (0.96 g, 51%) as a colorless oil:  $R_f = 0.63$  (20% EA–hex);  $^1\text{H}$  NMR  $\delta$  7.52–7.25 (m, 5 H), 4.09 (t, 2 H,  $J = 4.8$ ), 3.59 (t, 2 H,  $J = 5.0$ ), 3.35 (s, 3 H), 1.63 (s, 6 H);  $^{13}\text{C}$  NMR  $\delta$  145.2, 128.0, 127.1, 125.5, 83.0, 74.2, 69.8, 58.9, 26.4; Anal. Calcd for  $\text{C}_{12}\text{H}_{18}\text{O}_3$ : C, 68.55; H, 8.63. Found: C, 68.36; H, 8.39%.



**4a-Hydroperoxy-1,2,3,4,4a,5,6,7-octahydronaphthalene (2a)**

Compound **2a** was prepared using a reported procedure.<sup>28</sup>

**4a-Methyldioxy-1,2,3,4,4a,5,6,7-octahydronaphthalene (2b)**

Compound **2b** was prepared in 93% yield from **2a** (142 mg, 0.85 mmol) and iodomethane (0.11 mL, 1.7 mmol) by a similar procedure as for **1b**:  $R_f = 0.50$  (0.25% EA-hex);  $^1\text{H NMR } \delta$  5.63 (app. quintet, 1 H,  $J = 2.3$ ), 3.82 (s, 3 H), 2.33–2.16 (m, 3 H), 2.04–1.85 (m, 3 H), 1.79–1.63 (m, 3 H), 1.59–1.48 (m, 2 H), 1.29–1.12 (m, 3 H);  $^{13}\text{C NMR } \delta$  136.7, 126.7, 80.7, 62.9, 35.7, 32.9, 32.3, 27.5, 25.8, 21.7, 18.7; Anal. Calcd for  $\text{C}_{11}\text{H}_{18}\text{O}_2$ : C, 72.49; H, 9.95. Found: C, 72.36; H, 9.87%.

**4a-(2-Methoxyethyl)dioxy-1,2,3,4,4a,5,6,7-octahydronaphthalene (2d)**

Compound **2d** was prepared in 61% yield from **2a** (78.8 mg, 0.47 mmol) and 2-methoxyethoxy toluene-*p*-sulfonate (108.1 mg, 0.47 mmol) as for **1d**:  $R_f = 0.37$  (5% EA-hex);  $^1\text{H NMR } \delta$  5.63 (app. quintet, 1 H,  $J = 2.2$ ), 4.11 (t, 2 H,  $J = 4.8$ ), 3.60 (t, 2 H,  $J = 4.9$ ), 3.37 (s, 3 H), 2.29–2.20 (m, 3 H), 2.03–1.84 (m, 3 H), 1.77–1.61 (m, 3 H), 1.56–1.48 (m, 2 H), 1.29–1.10 (m, 3 H);  $^{13}\text{C NMR } \delta$  136.6, 126.8, 80.7, 74.0, 70.1, 59.0, 35.7, 32.8, 32.4, 27.4, 25.9, 21.7, 18.7; Anal. Calcd for  $\text{C}_{13}\text{H}_{22}\text{O}_3$ : C, 68.99; H, 9.80. Found: C, 68.78; H, 10.00%.

**cis and trans-1-Methyldioxy-1,2,3,4,4a,5,6,7,8,8a-decahydronaphthalene (3b)**

An ethyl acetate suspension of  $\text{PtO}_2$  (81 mg, 0.36 mmol) and **2b** (443.4 mg, 2.4 mmol) was placed under an atmosphere of hydrogen for 1 h. The solution was filtered through Celite and then concentrated *in vacuo*. Filtration through silica (5% EA-hexane) and HPLC (0.1% EA-hexane) isolated **3b** (415.7 mg, 93%) as a 1:1 mixture of *cis* (5.3 min) and *trans* (7.5 min) isomers. Assignments are based on literature reports for related compounds. *trans*:  $R_f = 0.60$  (5% EA-hex);  $^1\text{H NMR}$  (500 MHz)  $\delta$  3.81 (s, 3 H), 2.04 (dq, 2 H,  $J = 13.7, 2.8$ ), 1.73–1.62 (m, 4 H), 1.50–1.44 (m, 2 H), 1.40–1.18 (m, 7 H), 1.00 (td, 2 H,  $J = 13.7, 4.0$ );  $^{13}\text{C NMR}$  (125 MHz)  $\delta$  80.8, 62.6, 45.3, 34.3, 28.4, 26.5, 21.7; HRMS Calcd for  $\text{C}_{11}\text{H}_{20}\text{O}_2$  ( $\text{M}^+$ ): 184.1463. Found: 184.1463. *cis*:  $R_f = 0.60$  (5% EA-hex);  $^1\text{H NMR}$  (500 MHz)  $\delta$  3.80 (s, 3 H), 1.76–1.70 (m, 3 H), 1.68–1.48 (m, 8 H), 1.44–1.27 (m, 6 H);  $^{13}\text{C NMR}$  (125 MHz)  $\delta$  83.6, 62.9, 37.7, 28.1, 22.7.

**1-Hydroperoxy-1,2,3,4,5,6,7,8-octahydronaphthalene (4a)<sup>29</sup>**

This was prepared from **2a** by a reported method.<sup>30</sup>

**1-Methyldioxy-1,2,3,4,5,6,7,8-octahydronaphthalene (4b)**

Alkylation of **4a** (977.4 mg, 5.8 mmol) with MeI as for **1b** furnished **4b** (819.5 mg, 77%) as a colorless oil:  $R_f = 0.50$  (2.5% EA-hex);  $^1\text{H NMR } \delta$  4.21 (br s, 1 H), 3.83 (s, 3 H), 2.27 (d, 1 H,  $J = 16.2$ ), 2.19–2.11 (m, 1 H), 1.94–1.80 (m, 5 H), 1.78–1.62 (m, 3 H), 1.59–1.41 (m, 4 H);  $^{13}\text{C NMR } \delta$  137.7, 124.8, 80.1, 62.2, 30.7, 30.6, 27.8, 27.0, 22.9, 22.7, 17.7; Anal. Calcd for  $\text{C}_{11}\text{H}_{18}\text{O}_2$ : C, 72.49; H, 9.95. Found: C, 72.69; H, 9.70%.

**Preparation of ozonides**

**Method A.** Into a 0 °C hexane solution of *O*-methyl oxime and ketone was bubbled  $\text{O}_3$ – $\text{O}_2$  until a light blue was observed. Residual  $\text{O}_3$  was sparged with  $\text{O}_2$  and solvent was removed *in vacuo*. The residue was subjected to flash chromatography.

**Method B.** Similar to method A, except a –78 °C solution of vinyl ether (1 equiv.) and cyclohexanone (2 equiv.) in 2:1 hexane– $\text{CH}_2\text{Cl}_2$  was used.

**Method C.** Similar to method A, except a –78 °C solution of alkene in hexane was used. The  $^1\text{H}$  and  $^{13}\text{C}$  NMR data of

**5a** (60%), **5b** (67%), **5c** (63%), **5d** (57%), and **5m** were identical to literature reports.<sup>12,13</sup>

**3,3-Dibutyl-1,2,4-trioxaspiro[5.4]decane (5e)**

Yield 47%;  $R_f = 0.77$  (10% EA-hex);  $^1\text{H NMR } \delta$  1.80–1.50 (m, 12 H), 1.50–1.25 (m, 10 H), 0.87 (t, 6 H,  $J = 4.5$ );  $^{13}\text{C NMR } \delta$  111.7, 108.9, 35.1, 34.7, 26.1, 25.0, 23.8, 22.9, 14.0.

**3-Octyl-1,2,4-trioxaspiro[5.4]decane (5f)**

Yield 47%;  $R_f = 0.76$  (10% EA-hex);  $^1\text{H NMR } \delta$  5.20 (t, 1 H,  $J = 4.5$ ), 1.75–1.50 (m, 8 H), 1.50–1.20 (m, 16 H), 0.87 (t, 3 H,  $J = 4.5$ );  $^{13}\text{C NMR } \delta$  109.2, 103.9, 35.5, 33.4, 32.6, 31.8, 31.2, 29.5, 29.4, 29.2, 24.9, 24.0, 23.9, 14.1, 22.7.

**3-Phenyl-1,2,4-trioxaspiro[5.4]decane (5g)**

Yield 65%;  $R_f = 0.65$  (10% EA-hex);  $^1\text{H NMR } \delta$  7.60–7.50 (m, 2 H), 7.50–7.38 (m, 3 H), 6.08 (s, 1 H), 2.00–1.80 (m, 4 H), 1.80–1.60 (m, 4 H), 1.60–1.40 (m, 2 H);  $^{13}\text{C NMR } \delta$  132.6, 130.4, 128.6, 127.9, 110.6, 103.5, 35.7, 33.6, 25.0, 24.0, 23.8.

**1,2,4-Trioxaspiro[5.4]decane (5h)**

Yield 78%;  $R_f = 0.67$  (10% EA-hex);  $^1\text{H NMR } \delta$  5.10 (s, 2 H), 1.75–1.55 (m, 8 H), 1.45–1.35 (m, 2 H);  $^{13}\text{C NMR } \delta$  108.8, 93.6, 33.8, 24.8, 23.8.

**3-tert-Butyl-3-methyl-1,2,4-trioxaspiro[5.4]decane (5i)**

Yield 53%;  $R_f = 0.75$  (10% EA-hex);  $^1\text{H NMR } \delta$  1.70–1.50 (m, 8 H), 1.50–1.30 (m, 2 H), 1.44 (s, 3 H), 1.03 (s, 9 H);  $^{13}\text{C NMR } \delta$  112.3, 110.0, 37.5, 35.4, 35.3, 26.5, 25.6, 24.5, 24.2, 20.6.

**3-Phenyl-5-propyl-1,2,4-trioxolane (5j)**

Yield 73%, 1:1 *cis*–*trans* mixture  $R_f = 0.63$  (10% EA-hex); *first eluting*:  $^1\text{H NMR } \delta$  7.62–7.40 (m, 5 H), 6.15 (s, 1 H), 5.47 (t, 1 H,  $J = 5.25$ ), 1.87 (m, 2 H), 1.60 (m, 2 H), 1.03 (t, 3 H,  $J = 7.4$ );  $^{13}\text{C NMR } \delta$  134.3, 130.1, 128.5, 127.3, 105.1, 103.30, 103.27, 34.4, 17.5, 13.9. *Second eluting*:  $^1\text{H NMR}$  same except  $\delta$  7.62–7.40, 6.09 (s, 1 H), 5.54 (t, 1 H,  $J = 4.8$ ), 1.05 (t, 3 H,  $J = 7.4$ );  $^{13}\text{C NMR } \delta$  132.6, 130.5, 128.6, 127.8, 105.6, 103.81, 103.78, 32.9, 17.2, 14.0.

**3-Phenyl-1,2,4-trioxolane (5k)**

Yield 77%;  $R_f = 0.60$  (10% EA-hex);  $^1\text{H NMR } \delta$  7.60–7.40 (m, 5 H), 6.07 (s, 1 H), 5.44 (s, 1 H), 5.36 (s, 1 H);  $^{13}\text{C NMR } \delta$  132.9, 130.4, 128.6, 127.5, 103.08, 103.05, 95.2.

**3-Methyl-3-phenyl-1,2,4-trioxolane (5l)**

Yield 61%;  $R_f = 0.62$  (10% EA-hex);  $^1\text{H NMR } \delta$  7.60–7.28 (m, 5 H), 5.28 (d, 1 H,  $J = 4.0$ ), 1.84 (s, 3 H);  $^{13}\text{C NMR } \delta$  140.7, 128.6, 128.3, 125.2, 108.3, 94.5, 24.3.

**General procedure for allylation**

To a 0 °C ( $\text{BF}_3 \cdot \text{OEt}_2$ ) or –78 °C ( $\text{SnCl}_4$  or  $\text{TiCl}_4$ ) solution of cumene hydroperoxide (0.24 mL, 1.4 mmol) and allylsilane (0.22 mL, 1.4 mmol) in  $\text{CH}_2\text{Cl}_2$  (6 mL) under  $\text{N}_2$  was added Lewis acid (1.2 mmol). After stirring for 5–30 min at the initial temperature, the reaction was brought to RT, and subjected to a standard work-up and flash chromatography. Allylations of **1b** were analyzed by  $^1\text{H NMR}$ . Allylations of **1c** were performed in the presence of 2-methylnaphthalene as an internal standard; the products were quantified by  $^1\text{H NMR}$ . Fragmentations of **2a–d** were performed as for **1b–d** except that chromatography was performed using alumina (ether–pentane) and hydroxy- or alkoxybicyclo[5.3.0]decanones **10a/b** were quantified together with bicyclodecenone **11**.

**4-Methyl-4-phenylpent-1-ene (6)**

$R_f = 0.81$  (2.5% EA–hex);  $^1\text{H NMR}$  (500 MHz)  $\delta$  7.36–7.16 (m, 5 H), 5.60–5.51 (m, 1 H), 4.98 (s, 1 H), 4.94 (d, 1 H,  $J = 8.1$ ), 2.36 (d, 2 H,  $J = 7.3$ ), 1.31 (s, 6 H);  $^{13}\text{C NMR}$  (125 MHz)  $\delta$  149.3, 135.5, 128.0, 125.8, 125.5, 116.9, 48.8, 37.6, 28.5; Anal. Calcd for  $\text{C}_{12}\text{H}_{16}$ : C, 89.94; H, 10.06. Found: C, 90.13; H, 10.04%.

**4-Methyl-4-phenoxy-pent-1-ene (7)**

$R_f = 0.48$  (2.5% EA–hex);  $^1\text{H NMR}$  (500 MHz)  $\delta$  7.27–6.98 (m, 5 H), 5.95 (m, 1 H), 5.12 (s, 1 H), 5.09 (d, 1 H,  $J = 11.3$ ), 2.41 (d, 2 H,  $J = 7.3$ ), 1.27 (s, 6 H);  $^{13}\text{C NMR}$  (125 MHz)  $\delta$  155.2, 134.5, 128.9, 124.2, 123.3, 117.7, 80.0, 46.7, 26.4; Anal. Calcd for  $\text{C}_{12}\text{H}_{16}\text{O}$ : C, 81.77; H, 9.15. Found: C, 81.75; H, 8.94%.

**4-Methoxy-4-methylpent-1-ene (8b)**

$R_f = 0.45$  (5% EA–hex);  $^1\text{H NMR}$   $\delta$  5.86 (m, 1 H), 5.13 (s, 1 H), 5.09 (d, 1 H,  $J = 7.4$ ), 3.29 (s, 3 H), 2.31 (d, 2 H,  $J = 7.2$ ), 1.23 (s, 6 H);  $^{13}\text{C NMR}$   $\delta$  134.1, 117.6, 75.4, 49.1, 44.2, 24.7.

**4-Methylpent-1-enyl 2-methoxyethyl ether (8c)**

$R_f = 0.13$  (2.5% EA–hex);  $^1\text{H NMR}$  (500 MHz)  $\delta$  5.87–5.79 (m, 1 H), 5.05 (s, 1 H), 5.02 (d, 1 H,  $J = 6.4$ ), 3.50 (dt, 4 H,  $J = 2.8$ , 2.4), 3.37 (s, 3 H), 2.25 (d, 2 H,  $J = 7.3$ ), 1.16 (s, 6 H);  $^{13}\text{C NMR}$  (125 MHz)  $\delta$  134.7, 117.1, 74.7, 72.6, 60.8, 59.1, 44.8, 25.3; HRMS Calcd for  $\text{C}_9\text{H}_{18}\text{O}_2$  ( $\text{M}^+$ ): 158.1307. Found: ( $\text{M} - \text{C}_3\text{H}_5^+$ ): 117.09146.

**1-Allyl-1,2,3,4,5,6,7,8-octahydronaphthalene (9)**

$R_f = 0.83$  (1% EA–hex);  $^1\text{H NMR}$   $\delta$  5.80–5.72 (m, 1 H), 4.99 (d, 1 H,  $J = 17.7$ ), 4.97 (d, 1 H,  $J = 8.5$ ), 2.32–2.28 (m, 1 H), 2.11–2.06 (m, 1 H), 2.00–1.72 (m, 7 H), 1.67–1.43 (m, 8 H);  $^{13}\text{C NMR}$   $\delta$  138.2, 130.5, 129.2, 115.4, 38.8, 37.2, 30.9, 30.8, 28.4, 27.6, 23.5, 23.2, 20.0; Anal. Calcd for  $\text{C}_{13}\text{H}_{20}$ : C, 88.57; H, 11.43. Found: C, 88.72; H, 11.50%.

**7-Hydroxybicyclo[5.3]decan-2-one (10a)**

$R_f = 0.26$  (20% EA–hex);  $^1\text{H NMR}$  (500 MHz)  $\delta$  3.13 (t, 1 H,  $J = 8.5$ ), 2.53 (d, 1 H,  $J = 18.9$ ), 2.42–2.36 (m, 2 H), 2.16 (d, 1 H,  $J = 10.1$ ), 1.93–1.60 (m, 10 H), 1.24 (br s, 1 H);  $^{13}\text{C NMR}$  (125 MHz)  $\delta$  211.6, 81.8, 59.5, 44.3, 43.7, 41.7, 24.8, 24.1, 23.4, 21.6; IR (neat)/ $\text{cm}^{-1}$  3463 (br, OH), 1693 (C=O).

**7-Methoxybicyclo[5.3]decan-2-one (10b)**

$R_f = 0.66$  (20% EA–hex);  $^1\text{H NMR}$  (500 MHz)  $\delta$  3.04 (s, 3 H), 2.97 (t, 1 H,  $J = 8.5$ ), 2.49 (d, 1 H,  $J = 17.7$ ), 2.41–2.30 (m, 2 H), 2.08 (ddd, 1 H,  $J = 13.7$ , 6.9, 3.6), 1.87–1.78 (m, 2 H), 1.70–1.58 (m, 3 H), 1.57–1.30 (m, 4 H);  $^{13}\text{C NMR}$  (125 MHz)  $\delta$  211.1, 85.9, 60.8, 48.3, 43.6, 36.1, 35.7, 25.0, 24.9, 23.5, 21.6; IR (neat)/ $\text{cm}^{-1}$  1700 (C=O).

**Bicyclo[5.3]dec-1(7)-en-2-one (11)**

$R_f = 0.28$  (10% EA–hex);  $^1\text{H NMR}$  (500 MHz)  $\delta$  2.65–2.54 (m, 6 H), 2.44 (m, 2 H), 1.82–1.75 (m, 6 H);  $^{13}\text{C NMR}$  (125 MHz)  $\delta$  201.4, 159.0, 138.4, 43.9, 41.6, 33.9, 31.6, 25.8, 22.4, 21.2; IR (neat)/ $\text{cm}^{-1}$  1643 (C=O), 1625 (C=C); Anal. Calcd for  $\text{C}_{10}\text{H}_{14}\text{O}$ : C, 79.96; H, 9.36. Found: C, 79.99; H, 9.31%.

**1-Methoxyspiro[4.5]decan-6-one (12)**

Addition of  $\text{SnCl}_4$  (0.88 mL, 0.88 mmol, 1 M in  $\text{CH}_2\text{Cl}_2$ ) to a 78 °C solution of methyl peroxide **4b** (159.7 mg, 0.88 mmol) and allyltrimethylsilane (0.28 mL, 1.7 mmol) in  $\text{CH}_2\text{Cl}_2$  (4.5 mL), furnished, following a standard work-up and flash chromatography (5% ether–pentane) **9** (27.6 mg, 18%) and **12** (54.0 mg, 34%);  $R_f = 0.23$  (5% EA–hex);  $^1\text{H NMR}$  (500 MHz)  $\delta$  3.84

(dd, 1 H,  $J = 4.4$ , 2.8), 3.21 (s, 3 H), 2.55 (dt, 1 H,  $J = 13.3$ , 7.3), 2.43 (td, 1 H,  $J = 13.7$ , 6.1), 2.39–2.36 (m, 1 H), 2.04 (sextet of d, 1 H,  $J = 9.7$ , 2.8), 1.85–1.53 (m, 8 H), 1.42 (td, 1 H,  $J = 14.1$ , 4.0), 1.07 (ddd, 1 H,  $J = 12.9$ , 7.7, 5.6);  $^{13}\text{C NMR}$  (125 MHz)  $\delta$  211.5, 87.2, 61.2, 56.5, 41.6, 38.5, 32.0, 29.1, 27.6, 22.6, 21.1; IR (neat)/ $\text{cm}^{-1}$  1712 C=O; Anal. Calcd for  $\text{C}_{11}\text{H}_{18}\text{O}_2$ : C, 72.49; H, 9.95. Found: C, 72.50; H, 9.79%.

**1-Allyl-1,6-epoxycyclodecane (13) and 7-allyl-1-oxabicyclo[5.4.0]undecane (14)**

$\text{SnCl}_4$ -mediated allylation (5 h,  $-78$  °C) of a mixture of *cis*- and *trans*-**3b** (369.7 mg, 2.0 mmol) furnished, following chromatography on alumina (1% ether–pentane) and HPLC on silica (hexane), 1-allyl-1,6-epoxycyclodecane **13** (103 mg, 26%, retention time = 21 min), 7-allyl-1-oxabicyclo[5.4.0]undecane **14** (6%, 24 min), and 180 mg of recovered *trans*-**3b**.

**1-Allyl-1,6-epoxycyclodecane (13)**

$R_f = 0.13$  (1% EA–hex);  $^1\text{H NMR}$  (500 MHz)  $\delta$  5.77 (m, 1 H), 5.01 (d, 1 H,  $J = 7.3$ ), 4.99 (d, 1 H,  $J = 16.9$ ), 4.06 (m, 1 H), 2.23 (d, 2 H,  $J = 7.7$ ), 1.78–1.49 (m, 16 H);  $^{13}\text{C NMR}$  (125 MHz)  $\delta$  135.6, 117.2, 80.1, 76.8, 47.6, 37.2, 34.8, 25.3, 24.9; HRMS Calcd for  $\text{C}_{13}\text{H}_{22}\text{O}$  ( $\text{M}^+$ ): 194.1671. Found: 194.1671.

**7-Allyl-1-oxabicyclo[5.4.0]undecane (14)**

$R_f = 0.13$  (1% EA–hex);  $^1\text{H NMR}$  (500 MHz)  $\delta$  5.87 (m, 1 H), 5.05 (d, 1 H,  $J = 10.1$ ), 5.03 (d, 1 H,  $J = 6.9$ ), 3.57 (m, 2 H), 2.47 (dd, 1 H,  $J = 14.5$ , 8.1), 2.25 (dd, 1 H,  $J = 13.3$ , 4.8), 1.92–1.20 (m, 15 H);  $^{13}\text{C NMR}$  (125 MHz)  $\delta$  134.7, 116.7, 61.2, 45.1, 38.9, 32.2, 31.5, 31.3, 30.9, 29.2, 23.2, 21.2; HRMS Calcd for  $\text{C}_{13}\text{H}_{22}\text{O}$  ( $\text{M}^+$ ): 194.1671. Found ( $\text{M} - \text{C}_3\text{H}_5^+$ ): 153.1280.

**Lewis acid-promoted decomposition of ozonides**

A  $-78$  °C solution of ozonide in 1 mL of  $\text{CDCl}_3$  was admixed in an NMR tube with a solution of excess Lewis acid in  $\text{CD}_2\text{Cl}_2$ . The mixture was brought to room temperature and monitored by NMR.

**Reaction of ozonides with allylsilane: general procedure**

To a  $-78$  °C solution of 200 mg (1 equiv., 1.01 mmol) of ozonide **5a** and 0.20 mL (1.3 equiv., 1.3 mmol) and allylsilane in  $\text{CH}_2\text{Cl}_2$  (5 mL) was added  $\text{SnCl}_4$  (1.1 equiv. as a 1 M solution in  $\text{CH}_2\text{Cl}_2$ ). A standard work-up (except  $\text{CH}_2\text{Cl}_2$  extractions) gave 180 mg of a residue, from which was separated 115 mg (0.50 mmol, 50%) of **17**, 13 mg (0.11 mmol, 11%) of **15b** and trace amounts of **16a** and **16b** were isolated. Compounds **15b**, **16a** and **16b** were assigned based upon comparison of  $^1\text{H}$  and  $^{13}\text{C}$  NMR data with authentic samples.

**3-(Trimethylsilylmethyl)-1,2-dioxaspiro[5.4]decane (17)**

$R_f = 0.72$  (10% EA–hex);  $^1\text{H NMR}$   $\delta$  4.40–4.28 (m, 1 H), 2.36 (dd, 1 H,  $J = 11.7$ , 6.9), 1.83 (dd, 1 H,  $J = 11.7$ , 8.1), 1.75–1.50 (m, 6 H), 1.50–1.30 (m, 4 H), 1.14 (dd, 1 H,  $J = 14.1$ , 5.3), 0.83 (dd, 1 H,  $J = 14.1$ , 9.3), 0.05 (s, 9 H);  $^{13}\text{C NMR}$   $\delta$  86.2, 80.3, 53.5, 37.6, 36.0, 25.9, 24.6, 24.1, 21.7,  $-0.4$ . Anal. Calcd for  $\text{C}_{12}\text{H}_{24}\text{O}_2\text{Si}$ : C, 63.10; H, 10.59. Found: C, 62.89; H, 10.40%.  $^1\text{H}$  and  $^{13}\text{C}$  assignments based on DEPT and HETCOR.

**5-(Prop-2-en-1-yl)nonan-5-ol (20e)**

$R_f = 0.66$  (20% EA–hex);  $^1\text{H NMR}$   $\delta$  5.80 (m, 1 H), 5.11 (m, 2 H), 2.20 (d, 2 H,  $J = 7.0$ ), 1.62–1.20 (m, 12 H), 0.87 (t, 6 H,  $J = 7.0$ ).

**Dodec-1-en-4-ol (20f)**

$R_f = 0.50$  (20% EA–hex);  $^1\text{H NMR}$   $\delta$  5.80 (m, 1 H), 5.10 (m, 2 H), 3.62 (m, 1 H), 2.30 (m, 1 H), 2.15 (m, 1 H), 1.60 (s, 1 H, OH), 1.50–1.20 (m, 14 H), 0.84 (t, 3 H,  $J = 7.0$ ).

**1-Phenylbut-3-en-1-ol (20j = 20k)**

$R_f = 0.46$  (20% EA–hex);  $^1\text{H NMR } \delta$  7.40–7.16 (m, 5 H), 5.80 (m, 1 H), 5.15 (m, 2 H), 4.52 (t, 1 H,  $J = 6.1$ ), 2.50 (m, 2 H).

**3-(Trimethylsilylmethyl)-5-tert-butyl-5-methyl-1,2-dioxaspiro[5.4]decane (18)**

$R_f = 0.72$  (10% EA–hex);  $^1\text{H NMR } \delta$  4.22 (m, 1 H), 2.68 (dd, 1 H,  $J = 11.7, 6.9$ ), 1.63 (dd, 1 H,  $J = 11.7, 8.1$ ), 1.35 (s, 3 H), 1.08 (dd, 1 H,  $J = 14.1, 5.3$ ), 0.98 (s, 9 H), 0.82 (dd, 1 H,  $J = 14.1, 9.3$ ), 0.04 (s, 9 H);  $^{13}\text{C NMR } \delta$  90.92, 80.58, 50.40, 36.12, 25.91, 23.06, 20.42, –0.95. Anal. Calcd for  $\text{C}_{12}\text{H}_{26}\text{O}_2\text{Si}$ : C, 62.56; H, 11.37. Found: C, 62.69; H, 11.41%.

**cis-3-Phenyl-5-(trimethylsilylmethyl)-1,2-dioxaspiro[5.4]decane (22j = 22k)**

$R_f = 0.33$  in 5% EA–hexane;  $^1\text{H NMR (500 MHz) } \delta$  7.45–7.25 (m, 5 H), 5.29 (t, 1 H,  $J = 7.5$ ), 4.49 (tt, 1 H,  $J = 9.3, 5.5$ ), 3.14 (ddd, 1 H,  $J = 11.9, 7.9, 6.0$ ), 2.20 (ddd, 1 H,  $J = 11.9, 8.5, 6.9$ ), 1.19 (dd, 1 H,  $J = 14.1, 5.2$ ), 0.90 (dd, 1 H,  $J = 14.1, 9.3$ ), 0.07 (s, 9 H);  $^{13}\text{C NMR } \delta$  141.17, 128.62, 127.88, 126.16, 82.83, 80.11, 51.27, 20.40, –1.01. Anal. Calcd for  $\text{C}_{13}\text{H}_{20}\text{O}_2\text{Si}$ : C, 66.05; H, 8.53. Found: C, 65.94; H, 8.41%.

**trans-3-Phenyl-5-(trimethylsilylmethyl)-1,2-dioxaspiro[5.4]decane (22j = 22k)**

$R_f = 0.33$  (5% EA–hexane);  $^1\text{H NMR (500 MHz) } \delta$  7.45–7.28 (m, 5 H), 5.30 (dd, 1 H,  $J = 8.5, 6.0$ ), 4.62–4.55 (m, 1 H), 2.65 (dt, 1 H,  $J = 11.7, 6.0$ ), 2.55 (dt, 1 H,  $J = 11.7, 8.5$ ), 1.17 (dd, 1 H,  $J = 14.1, 5.4$ ), 0.93 (dd, 1 H,  $J = 14.1, 8.9$ ), 0.08 (s, 9 H);  $^{13}\text{C NMR } \delta$  138.81, 128.66, 128.36, 126.83, 83.17, 80.05, 50.05, 20.90, –0.97. Anal. Calcd for  $\text{C}_{13}\text{H}_{20}\text{O}_2\text{Si}$ : C, 66.05; H, 8.53. Found: C, 65.87; H, 8.34%.

**3-Methyl-3-phenyl-5-(trimethylsilylmethyl)-1,2-dioxaspiro[5.4]decane (22l)**

Compound **22l** was isolated as a 1 : 1 mixture. Pure *cis*-**22l** and a (1 : 1.5) mixture of *cis/trans* **22l** were obtained by HPLC (2% EA–hex).  $R_f = 0.33$  (5% EA–hexane);  $^1\text{H NMR (500 MHz) } \delta$  7.46–7.42 (m, 2 H), 7.32–7.38 (m, 2 H), 7.30–7.24 (m, 1 H), 4.43–4.36 (m, 1 H), 2.93 (dd, 1 H,  $J = 11.7, 6.4$ ), 2.27 (dd, 1 H,  $J = 11.7, 8.1$ ), 1.68 (s, 3 H), 1.17 (dd, 1 H,  $J = 14.1, 5.6$ ), 0.93 (dd, 1 H,  $J = 14.1, 9.3$ ), 0.05 (s, 9 H);  $^{13}\text{C NMR } \delta$  144.80, 128.36, 127.18, 124.98, 86.79, 79.83, 56.00, 27.83, 21.62, –0.99. DEPT 135 NMR  $\delta$  128.36(+), 127.18(+), 124.98(+), 79.83(+), 56.00(–), 27.83(+), 21.62(–), –0.99(+). Anal. Calcd for  $\text{C}_{14}\text{H}_{22}\text{O}_2\text{Si}$ : C, 67.15; H, 8.86. Found: C, 67.22; H, 8.74%.

**cis-3-Propyl-5-(trimethylsilylmethyl)-1,2-dioxaspiro[5.4]decane (23j)**

$R_f = 0.45$  (5% EA–hexane);  $^1\text{H NMR (500 MHz) } \delta$  4.35–4.23 (m, 2 H), 2.77 (dt, 1 H,  $J = 11.7, 6.9$ ), 1.74 (ddd, 1 H,  $J = 11.7, 8.1, 6.5$ ), 1.75–1.30 (m, 4 H), 1.11 (dd, 1 H,  $J = 14.1, 5.4$ ), 0.93 (t, 3 H,  $J = 6.9$ ), 0.85 (dd, 1 H,  $J = 14.1, 9.3$ ), 0.06 (s, 9 H);  $^{13}\text{C NMR } \delta$  81.42, 79.35, 48.33, 37.15, 21.02, 19.52, 13.99, –1.01. Anal. Calcd for  $\text{C}_{10}\text{H}_{22}\text{O}_2\text{Si}$ : C, 59.35; H, 10.96. Found: C, 59.47; H, 11.04%.

**4-Methyl-4-phenylhepta-1,6-diene (24)**

$R_f = 0.75$  in 5% (EA–hex);  $^1\text{H NMR (500 MHz) } \delta$  7.33–7.17 (m, 5 H), 5.63–5.47 (m, 2 H), 5.05–4.92 (m, 4 H), 2.52 (dd, 1 H,  $J = 13.7, 6.6$ ), 2.32 (dd, 2 H,  $J = 11.7, 6.6$ ), 1.29 (s, 3 H);  $^{13}\text{C NMR } \delta$  147.1, 135.1, 128.0, 126.4, 125.6, 117.2, 47.0, 24.2.

**3-Methyl-3-(4-hydroxyhept-6-enyl)-5-(trimethylsilylmethyl)-1,2-dioxolane (25)**

To a –78 °C solution of 145 mg (1.1 mmol) of **14m** in 20 mL of

$\text{CH}_2\text{Cl}_2$  was added 1.1 mL of a 1 M solution of  $\text{SnCl}_4$ . After 2 min, a solution of allylsilane (251 mg, 2 equiv.) in 20 mL of  $\text{CH}_2\text{Cl}_2$  was added. The reaction mixture was stirred for 30 min and subjected to standard work-up. Flash chromatography on silica gel (10% to 20% EA–hex) afforded 151 mg (48%) of **25** as a colorless mixture of four isomers which could be partially separated by HPLC: (10% EA–hex) into two *cis* isomers (35%, 35%) and a mixture of two *trans* isomers (15%, 15%). *cis*-**25a**:  $R_f = 0.30$  (20% EA–hex);  $^1\text{H NMR (500 MHz) } \delta$  5.88–5.78 (m, 1 H), 5.14 (m, 2 H), 4.42–4.34 (m, 1 H), 3.71–3.63 (m, 1 H), 2.364 (dd, 1 H,  $J = 11.7, 6.5$ ), 2.36–2.27 (m, 1 H), 2.20–2.11 (m, 1 H, CH), 1.923 (dd, 1 H,  $J = 11.7, 8.5$ ), 1.80–1.40 (m, 6 H), 1.28 (s, 3 H), 1.079 (dd, 1 H,  $J = 14.1, 5.2$ ), 0.815 (dd, 1 H,  $J = 14.1, 9.3$ ), 0.046 (s, 9 H);  $^{13}\text{C NMR } \delta$  134.8(+), 118.3(–), 85.9, 79.8(+), 70.6(+), 53.56(–), 42.1(–), 40.6(–), 37.2(–), 23.31(+), 21.0(–), 20.8(–), –1.0(+). Anal. Calcd for  $\text{C}_{15}\text{H}_{30}\text{O}_3\text{Si}$ : C, 62.63; H, 10.55. Found: C, 62.51; H, 10.46%. *cis*-**25b**:  $^1\text{H NMR (500 MHz) } \delta$  5.88–5.77 (m, 1 H), 5.17–5.11 (m, 2 H), 4.41–4.33 (m, 1 H), 3.71–3.64 (m, 1 H), 2.357 (dd, 1 H,  $J = 11.67, 6.5$ ), 2.34–2.27 (m, 1 H), 2.19–2.11 (m, 1 H), 1.929 (dd, 1 H,  $J = 11.7, 8.5$ ), 1.75–1.35 (m, 6 H), 1.28 (s, 3 H), 1.08 (dd, 1 H,  $J = 13.7, 5.2$ ), 0.814 (dd, 1 H,  $J = 13.7, 9.3$ ), 0.045 (s, 9 H);  $^{13}\text{C NMR } \delta$  134.8(+), 118.2(–), 85.9, 79.9(+), 70.4(+), 53.5(–), 42.0(–), 40.5(–), 37.0(–), 23.5(+), 20.8(–), 20.6(–), –1.0(+). Anal. Calcd for  $\text{C}_{15}\text{H}_{30}\text{O}_3\text{Si}$ : C, 62.63; H, 10.55. Found: C, 62.44; H, 10.32%. *trans*-**25ab** (mixture of diastereomers):  $R_f = 0.27$  (20% EA–hex);  $^1\text{H NMR (500 MHz) } \delta$  5.87–5.77 (m, 2 H), 5.17–5.11 (m, 4 H), 4.36–4.28 (m, 2 H), 3.70–3.63 (m, 2 H), 2.51–2.45 (m, 2 H), 2.34–2.27 (m, 2 H), 2.19–2.11 (m, 2 H), 1.85–1.80 (m, 2 H), 1.66–1.40 (m, 12 H), 1.34 (s, 3 H), 1.096 (dd, 2 H,  $J = 14.1, 5.2$ ), 0.840 (dd, 2 H,  $J = 13.7, 9.3$ ), 0.047 (s, 18 H);  $^{13}\text{C NMR } \delta$  134.7 (for 2C), 118.4 (for 2C), 86.1 (for 2C), 80.1 (for 2C), 70.5, 70.4, 53.1, 52.9, 42.1, 42.0, 38.8, 38.7, 37.2, 37.1, 26.0, 25.8, 21.1 (for 2C), 21.0, 20.8, –1.0 (for 2C). Anal. Calcd for  $\text{C}_{15}\text{H}_{30}\text{O}_3\text{Si}$ : C, 62.63; H, 10.55. Found: C, 62.67; H, 10.47%.

**3-Methyl-3-(4-acetoxyhept-6-enyl)-5-(trimethylsilylmethyl)-1,2-dioxolane (26)**

Acetylation of **25** (23 mg) with DMAP and acetic anhydride in  $\text{CH}_2\text{Cl}_2$  furnished, after standard work-up and flash chromatography, 25 mg (92%) of **26** as a colorless oil which was resolved by HPLC (3% EA–hex) into two *cis* dioxolanes *cis*-**26a** (35%) and *cis*-**26b** (35%) as well as a mixture of two *trans* dioxolanes *trans*-**26ab** (30%): *cis*-**26a**:  $R_f = 0.58$  (20% EA–hex).  $^1\text{H NMR (500 MHz) } \delta$  5.80–5.70 (m, 1 H), 5.11–5.04 (m, 2 H), 4.97–4.89 (m, 1 H), 4.40–4.32 (m, 1 H), 2.36 (dd, 1 H,  $J = 11.7, 6.8$ ), 2.33–2.28 (m, 2 H), 1.89 (dd, 1 H,  $J = 11.7, 8.5$ ), 2.03 (s, 3 H), 1.75–1.25 (m, 6 H), 1.26 (s, 3 H), 1.08 (dd, 1 H,  $J = 13.9, 5.0$ ), 0.81 (dd, 1 H,  $J = 13.9, 9.5$ ), 0.05 (s, 9 H);  $^{13}\text{C NMR } \delta$  170.8, 133.7(+), 117.7(–), 85.7, 79.8(+), 73.0(+), 53.7(–), 40.3(–), 38.8(–), 33.9(–), 23.2(+), 21.2(+), 20.8(–), 20.3(–), –1.0(+). Calcd. for  $\text{C}_{17}\text{H}_{32}\text{O}_4\text{Si}$ : C, 62.15; H, 9.82. Found: C, 62.19; H, 9.70%. *cis*-**26b**:  $R_f = 0.56$  (20% EA–hex).  $^1\text{H NMR (500 MHz) } \delta$  5.80–5.70 (m, 1 H), 5.11–5.03 (m, 2 H), 4.95–4.88 (m, 1 H), 4.40–4.33 (m, 1 H), 2.35 (dd, 1 H,  $J = 11.7, 6.6$ ), 2.37–2.25 (m, 2 H), 1.90 (dd, 1 H,  $J = 11.7, 8.3$ ), 2.04 (s, 3 H), 1.75–1.67 (m, 1 H), 1.60–1.34 (m, 4 H), 1.43–1.34 (m, 1 H), 1.27 (s, 3 H), 1.07 (dd, 1 H,  $J = 13.7, 5.2$ ), 0.81 (dd, 1 H,  $J = 13.7, 9.5$ ), 0.05 (s, 9 H);  $^{13}\text{C NMR } \delta$  170.8, 133.7(+), 117.7(–), 85.7, 79.8(+), 73.2(+), 53.5(–), 40.3(–), 38.6(–), 33.9(–), 23.4(+), 21.3(+), 20.8(–), 20.4(–), –1.0(+). Anal. Calcd for  $\text{C}_{17}\text{H}_{32}\text{O}_4\text{Si}$ : C, 62.2; H, 9.8. Found: C, 62.6; H, 9.7%. Anal. Calcd for  $\text{C}_{17}\text{H}_{32}\text{O}_4\text{Si}$ : C, 62.15; H, 9.82. Found: C, 62.02; H, 9.67%. Spectral listings for *trans*-**26ab** are included in supplementary material.

**3-Methyl-3-(4-oxohept-6-enyl)-5-(trimethylsilylmethyl)-1,2-dioxolane (27)**

Oxidation of **25** (23 mg) with PDC in  $\text{CH}_2\text{Cl}_2$ , furnished, after



work-up and chromatography (7–16% EA–hex) 19 mg of **27** (83%) and 3.7 mg (16%) of recovered **25**. HPLC (3% EA–hex) resolved the material into pure *cis*-**27** and a mixture of *cis*- and *trans*-**27** (1.9:1.0). *cis*-**27**  $R_f$  = 0.77 (20% EA–hex).  $^1\text{H}$  NMR (500 MHz)  $\delta$  5.97–5.87 (m, 1 H), 5.183 (ddt, 1 H,  $J$  = 10.3, 1.6, 1.2), 5.141 (ddt, 1 H,  $J$  = 17.3, 1.6, 1.2), 4.40–4.33 (m, 1 H), 3.171 (dt, 2 H,  $J$  = 6.9, 1.2), 2.49 (dt, 2 H,  $J$  = 7.3, 6.9), 2.36 (dd, 1 H,  $J$  = 11.7, 6.5), 1.93 (dd, 1 H,  $J$  = 11.7, 8.5), 1.70–1.50 (m, 4 H), 1.28 (s, 3 H), 1.07 (dd, 1 H,  $J$  = 11.7, 5.2), 0.809 (dd, 1 H,  $J$  = 11.7, 9.3), 0.05 (s, 9 H);  $^{13}\text{C}$  NMR  $\delta$  208.5, 130.7(+), 118.8(–), 85.7, 79.9(+), 53.5(–), 47.8(–), 42.3(–), 39.8(–), 23.3(+), 20.8(–), 18.8(–), –1.1(+). Anal. Calcd for  $\text{C}_{15}\text{H}_{28}\text{O}_3\text{Si}$ : C, 63.33; H, 9.92. Found: C, 63.44; H, 9.87%. A spectral listing for the *cis/trans*-**27** mixture is included in supplementary material.

## Acknowledgements

We are grateful for financial support from the NIH and for technical assistance with NMR experiments from Professor Richard Shoemaker.

## References

- H. Kropf, in *Methods in Organic Chemistry (Houben-Weyl) Organische Peroxo-Verbindungen*, ed. H. Kropf, Georg Thieme Verlag, Stuttgart, 1988, p. 1084.
- P. H. Dussault, in *Active Oxygen in Chemistry*, ed. C. S. Foote, J. S. Valentine, A. Greenberg and J. F. Liebman, Blackie A & P, London, 1995, p. 141.
- A. J. Bloodworth, N. Hargreaves and J. C. Mitchell, *J. Chem. Res. (S)*, 1986, 390.
- K. J. McCullough and M. Nojima, in *Organic Peroxides*, ed. W. Ando, John Wiley & Sons, Chichester, 1992, p. 661.
- H. W. Gardner and R. D. Plattner, *Lipids*, 1984, **19**, 294.
- Y.-M. Pu, H. Yeh and H. Ziffer, *Heterocycles*, 1993, **36**, 2099.
- R. M. Goodman and Y. Kishi, *J. Org. Chem.*, 1994, **59**, 5125.
- P. H. Dussault, I. Q. Lee, H.-J. Lee, Q. J. Niu, J. A. Schultz and U. R. Zope, *J. Org. Chem.*, 2000, **65**, submitted.
- R. Lee, Ph.D. Thesis, University of Nebraska–Lincoln, 1996.
- P. H. Dussault and A. Sahli, *J. Org. Chem.*, 1992, **57**, 1009.
- P. H. Dussault, S. Kreifels and I. Q. Lee, *Synth. Commun.*, 1995, **25**, 2613.
- K. Griesbaum, X. Liu and Y. Dong, *Tetrahedron*, 1997, **53**, 5463.
- K. Griesbaum, *Trends Org. Chem.*, 1997, **6**, 145.
- R. C. Corcoran, *Tetrahedron Lett.*, 1990, **31**, 2101.
- H. Nishiyama and K. Itoh, *J. Org. Chem.*, 1982, **47**, 2496.
- W. Adam, A. Griesbeck and E. Staab, *Angew. Chem., Int. Ed. Engl.*, 1986, **25**, 269.
- G. E. Keck, D. E. Abbott, E. P. Boden and E. J. Enholm, *Tetrahedron Lett.*, 1984, **24**, 3927.
- S. C. Vonwiller, J. A. Warren, S. T. Mann and R. K. Haynes, *Tetrahedron Lett.*, 1997, **38**, 2363.
- P. H. Dussault and X. Liu, *Tetrahedron Lett.*, 1999, **40**, 6553.
- M. Miura and M. Nojima, *J. Chem. Soc., Chem. Commun.*, 1979, 467.
- M. Miura, M. Nojima and S. Kusabayashi, *J. Chem. Soc., Perkin Trans. 1*, 1980, 2909.
- M. Miura, S. Nagase, M. Nojima and S. Kusabayashi, *J. Org. Chem.*, 1983, **48**, 2366.
- M. Miura, M. Yoshida, M. Nojima and S. Kusabayashi, *J. Chem. Soc., Chem. Commun.*, 1982, 397.
- P. H. Dussault, C. T. Eary, R. J. Lee and U. R. Zope, *J. Chem. Soc., Perkin Trans. 1*, 1999, 2189.
- E. S. Shanley, in *Organic Peroxides*, ed. D. Swern, Wiley-Interscience, New York, 1970, vol. 3, p. 341.
- P. Patnaik, *A Comprehensive Guide to the Hazardous Properties of Chemical Substances*, Van Nostrand Reinhold, New York, 1992, p. 763.
- L. A. Medard, *Accidental Explosions: Types of Explosive Substances*, Ellis Horwood Limited, Chichester, 1989, vol. 2.
- G. O. Schenck and K.-H. Schulte-Elte, *Justus Liebigs Ann. Chem.*, 1958, **618**, 185.
- H.-S. Dang, A. G. Davies, I. G. E. Davison and C. H. Schiesser, *J. Org. Chem.*, 1990, **55**, 1432.
- N. A. Porter, J. K. Kaplan and P. H. Dussault, *J. Am. Chem. Soc.*, 1990, **112**, 1266.