



# Methyl 3,4,5-trimethoxycinnamate suppresses inflammation in RAW264.7 macrophages and blocks macrophage–adipocyte interaction

Olumayokun A. Olajide<sup>1</sup> · Idowu S. Akande<sup>2</sup> · Carlos da Silva Maia Bezerra Filho<sup>3</sup> · Izabela Lepiarz-Raba<sup>1</sup> · Damião Pergentino de Sousa<sup>3</sup>

Received: 19 March 2020 / Accepted: 2 May 2020  
© The Author(s) 2020

## Abstract

Methyl 3,4,5-trimethoxycinnamate (MTC) is a bioactive natural phenylpropanoid. We evaluated anti-inflammatory effects of synthetic MTC in RAW264.7 macrophages and RAW264.7–3T3-L1 adipocytes co-culture. Levels of cytokines and chemokines, as well as NO and PGE<sub>2</sub> in cell supernatants were analysed using ELISAs, Griess assay and enzyme immunoassays, respectively. In-cell cyto blot was used to assess levels of proteins; while DNA binding and reporter gene assays were used to measure transcription factor DNA binding and transcriptional activities, respectively. Glucose uptake in adipocytes was evaluated with 2-deoxy-2-[(7-nitro-2, 1, 3-benzoxadiazol-4-yl) amino]-D-glucose uptake. MTC (5–20 μM) suppressed LPS + IFNγ-induced release of TNFα, IL-6 and IL-1β, as well as NO/iNOS and PGE<sub>2</sub>/COX-2 levels in RAW264.7 cells. Furthermore, there was a reduction in phospho-IκB and phospho-p65 proteins, accompanied by a reduction in total IκB in RAW264.7 cells. Further studies showed that MTC also produced a reduction in NF-κB DNA binding and luciferase activity. Treatment of RAW264.7 cells with MTC (5–20 μM) resulted in enhanced DNA binding of Nrf2 and an increase in ARE-luciferase activity. In a macrophage–adipocyte co-culture, the compound reduced the release of TNFα, IL-6, IL-1β, MCP-1 and RANTES, while enhancing glucose uptake and activation of AMPKα. Our results suggest that MTC produced anti-inflammatory and antioxidant activities in macrophages. MTC also prevented inflammation in macrophage–adipocyte co-culture. The effect of MTC on glucose uptake in adipocytes is proposed to be linked to activation of AMPK.

**Keywords** Methyl 3,4,5-trimethoxycinnamate · Anti-inflammatory · RAW264.7 macrophages · NF-κB · 3T3-L1 adipocytes · Natural product

## Introduction

Chronic inflammation still remains a fundamental hallmark of a wide variety of diseases, ranging from arthritis, asthma, inflammatory bowel disease, neurodegenerative disorders, cardiovascular diseases, cancer and metabolic disorders. Consequently, chronic inflammation and inflammatory signalling pathways remain prime targets in identifying new treatments for these inflammatory disorders.

Transcription factors such as the nuclear Factor kappa B (NF-κB) and nuclear factor erythroid 2-related factor (Nrf2) play critical roles in inflammation. NF-κB is responsible for the transcriptional control of pro-inflammatory genes, and plays a central role in inflammatory diseases such as rheumatoid arthritis, inflammatory bowel disease, and autoimmunity, as well as diseases comprising a significant inflammatory component such as cancer

**Electronic supplementary material** The online version of this article (<https://doi.org/10.1007/s10787-020-00720-8>) contains supplementary material, which is available to authorized users.

✉ Olumayokun A. Olajide  
o.a.olajide@hud.ac.uk

<sup>1</sup> Department of Pharmacy, School of Applied Sciences, University of Huddersfield, Huddersfield HD1 3DH, UK

<sup>2</sup> Department of Biochemistry, College of Medicine, University of Lagos, Lagos, Nigeria

<sup>3</sup> Laboratory of Pharmaceutical Chemistry, Federal University of Paraíba, João Pessoa 58051-085, Brazil

and atherosclerosis (Mitchell and Carmody 2018). On the other hand, Nrf2 modulates constitutive or inducible molecular systems regulating redox homeostasis, resulting in the activation of antioxidants, anti-inflammatory molecules, as well as phase I and II drug metabolising enzymes (Sivandzade et al. 2019). Nrf2 has been functionally linked to cytoprotection in low-grade stress, chronic inflammation, metabolic alterations, and reactive oxygen species formation (Cuadrado et al. 2018). Interestingly, studies have established that there is a cross-talk between Nrf2 and NF- $\kappa$ B signalling pathways; the absence of Nrf2 enhances NF- $\kappa$ B activity with the resulting increase in pro-inflammatory cytokine production (Wardyn et al. 2015). Furthermore, studies have shown that Nrf2 is able to negate the transcriptional up-regulation of pro-inflammatory genes through mechanisms that are independent of its redox regulatory activity (Kobayashi et al. 2016).

Accumulating evidence suggests that metabolic conditions induce chronic low-grade macrophage-mediated inflammation in colon, liver, muscle and adipose tissue (Li et al. 2018). In this context, pro-inflammatory cytokines are secreted by recruited and resident macrophages in these organs and play a significant role in linking inflammation and insulin signalling in adjacent metabolic cells (Shapiro et al. 2011). Furthermore, it is widely known that obesity, insulin resistance and type two diabetes are closely linked with chronic inflammation, which is accompanied by excessive production of pro-inflammatory cytokines, acute-phase reactants, and activation of inflammatory signalling pathways (Wellen and Hotamisligil 2005; Hotamisligil 2006). Consequently, an important strategy in targeting insulin resistance will involve the use of anti-inflammatory modalities.

Anti-inflammatory natural compounds are now considered to possess significant therapeutic potentials in metabolic syndrome/insulin resistance. Flavonoids are plant-based anti-inflammatory natural products which have been suggested to be suggested to reduce the risk of diabetes by targeting inflammatory signals (Ren et al. 2019).

Methyl 3,4,5-trimethoxycinnamate (Fig. 1) is a natural phenylpropanoid ester found in several plants and has anti-platelet aggregation (Tsai et al. 2005) and antiarrhythmic activities (Zhao et al. 2013). The precursor of this compound 3, 4, 5-trimethoxycinnamic acid, is found in nature and has various pharmacological activities, such as anti-stress (Kawashima et al. 2004) and immunocompetent actions (Yu et al. 2014). Reports have shown that phenylpropanoids isolated from *Polygala tenuifolia* inhibited lipopolysaccharide (LPS)-induced nitric oxide production in BV-2 microglia (Cho et al. 2012). In the present study, methyl 3,4,5-trimethoxycinnamate (MTC) was prepared via fischer esterification using 3,4,5-trimethoxycinnamic acid as the starting material, as previously reported (da

Nóbrega et al. 2018). We then investigated the compound for effects on inflammation and Nrf2 antioxidant mechanisms in RAW264.7 macrophages. We also determined effects of MTC on inflammation in a macrophage–adipocyte co-culture stimulated with LPS and gamma interferon (IFN $\gamma$ ).

## Materials and methods

### Synthesis of methyl 3,4,5-trimethoxycinnamate

MTC was synthesised as earlier described (da Nóbrega et al. 2018). To a solution of 3,4,5-trimethoxycinnamic acid (0.25 g) in 250 mL of MeOH, 0.5 mL 96% (v/v) H<sub>2</sub>SO<sub>4</sub> was added under stirring. The reaction mixture was refluxed for 3 h. Half of MeOH was removed under reduced pressure and the following solution was then diluted with 10-mL water and the product extracted with ethyl acetate. The organic phase of the reaction was successively washed with 5% (w/v) NaHCO<sub>3</sub> (twice) and water, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and filtered. After removal of ethyl acetate under vacuum, the methyl 3,4,5-trimethoxycinnamate was obtained.

### Cell culture

RAW264.7 mouse macrophages (ECACC 91062702) were cultured in Dulbecco's modified eagle's medium (DMEM), supplemented with 2-mM glutamine and 10% foetal bovine serum (FBS). Cell viability experiments were carried out to determine whether incubation of RAW264.7 macrophages with concentrations of MTC used in combination with LPS (1  $\mu$ g/mL) and IFN $\gamma$  (10 ng/mL) would produce cytotoxic effects (Supplementary data 1).

3T3-L1 (ATCC<sup>®</sup> CL-173<sup>™</sup>) mouse pre-adipocytes were cultured in DMEM and supplemented with 2-mM glutamine and 10% FBS. All cells were cultured in an incubator at 37 °C and 5% CO<sub>2</sub>.

### Determination of pro- and anti-inflammatory mediators in culture supernatants of LPS + IFN $\gamma$ -stimulated RAW264.7 cells

Cultured RAW264.7 macrophages were treated with MTC (5–20  $\mu$ M). Thirty minutes later, the cells were stimulated with LPS (1  $\mu$ g/mL) and IFN $\gamma$  (10 ng/mL) for a further 24 h. Culture supernatants were analysed for levels of NO using the Griess assay kit (Promega), according to the manufacturer's instructions. Levels of pro-inflammatory (TNF $\alpha$ , IL-6 and IL-1 $\beta$ ) and anti-inflammatory (IL-10) cytokines in culture supernatants were measured using

mouse ELISA kits (Biolegend), according to the manufacturer's instructions. PGE<sub>2</sub> production was measured using PGE<sub>2</sub> EIA assay kit (Arbor Assays), according to the manufacturer's instructions.

### Luciferase reporter gene assays

RAW264.7 cells were seeded out in a 24-well plate ( $2 \times 10^5$  cells per well) in Opti-MEM (Gibco) and incubated at 37 °C for 4 h. Cells were then transfected with either the pNL3.2.NF- $\kappa$ B-RE [NlucP/NF- $\kappa$ B-RE/Hygro] vector (Promega, UK) or the pGL4.37[luc2P/ARE/Hygro] vector (Promega, UK) using lipofectamine 2000 transfection reagent (Invitrogen) and incubated for a further 4 h. Thereafter, experiments were carried out to evaluate the effects of MTC (5–20  $\mu$ M) on NF- $\kappa$ B and antioxidant response elements (ARE) luciferase activities, respectively. Luciferase activities were measured using Dual-Glo luciferase assay kit (Promega, Southampton, UK) according to the manufacturer's instructions.

### NF- $\kappa$ B and Nrf2 DNA binding assays

DNA binding assays were carried out to evaluate the effects of MTC on DNA binding of the NF- $\kappa$ B transcription factor in LPS + IFN $\gamma$ -stimulated RAW264.7 macrophages. These assays were also used to assess the ability of MTC to induce the binding of Nrf2 transcription factor to ARE consensus sites in the DNA. Cultured RAW264.7 cells were treated with MTC (5–20  $\mu$ M) for 30 min, followed by stimulation with LPS (1  $\mu$ g/mL) and IFN $\gamma$  (10 ng/mL) for a further 24 h in experiments to determine DNA binding of NF- $\kappa$ B. The cells were also treated with MTC (5–20  $\mu$ M) for 24 h to evaluate effects on binding of Nrf2 to ARE consensus sites. Nuclear extracts were collected and analysed for binding of transcription factors to the DNA using the TransAM NF- $\kappa$ B and Nrf2 transcription factor EMSA kits (Activ Motif, Belgium) according to the manufacturer's instructions. The TransAM transcription factor assay kits employed a 96-well plate to which oligonucleotides containing the NF- $\kappa$ B consensus site (5'-GGGACTTTC-3') or the ARE consensus binding site (5'-GTCACAGTGACTCAGCAGAATCTG-3') have been immobilised.

### In-cell western/cytoblot

The in-cell western is now an established method for the rapid quantification of proteins in cells (Velagapudi et al. 2019b), because it combines the specificity of western blotting with the quantification capability of ELISA. In these experiments, RAW264.7 macrophages were seeded into a 96-well plate ( $5 \times 10^4$  cells/mL). At 70% confluence, cells

were treated with MTC (5–20  $\mu$ M) for 30 min, followed by stimulation with LPS (1  $\mu$ g/mL) and IFN $\gamma$  (10 ng/mL) for different incubation periods. Cells were fixed with 8% paraformaldehyde solution (100  $\mu$ L) for 15 min. and then washed with PBS. The cells were then incubated with the primary antibodies overnight at 4 °C. The following antibodies were used: rabbit anti-COX-2 (Abcam), rabbit anti-iNOS (Abcam), rabbit anti-phospho-p65 (Cell Signalling technologies), rabbit anti-phospho-I $\kappa$ B (Santa Cruz Biotechnology), rabbit anti-I $\kappa$ B (Santa Cruz Biotechnology), and rabbit anti-phospho-AMPK $\alpha$  (Santa Cruz Biotechnology). Thereafter, cells were washed with PBS and incubated with anti-rabbit HRP secondary antibody for 2 h at room temperature. Then, 100- $\mu$ L HRP substrate was added to the plate and signal measured at 450 nm with a microplate reader. Readings were normalised with Janus Green stain (Abcam).

### Differentiation of 3T3-L1 pre-adipocytes

Sub-confluent 3T3-L1 pre-adipocytes were differentiated into adipocytes using differentiation medium (DMEM) containing 10% FBS supplemented with 0.5-mM 3-isobutyl-1-methylxanthine (0.5 mM), dexamethasone (1  $\mu$ M) and insulin (200 nM). Adipocyte differentiation was assessed using the oil red O staining quantitative assay kit (Cayman). Culture supernatants were removed from cells, followed by washing with PBS. Thereafter, lipid droplets assay fixative was added to each well and incubated for 15 min at room temperature. Following removal of the fixative, cells were washed and 75- $\mu$ L oil red O solution was added to each well for 20 min at room temperature. This was followed by removal of the oil red O and thorough washing of the wells with 100- $\mu$ L wash solution and the addition of 100- $\mu$ L dye extraction solution and absorbance read on a Tecan F50 microplate at 492 nm. Differentiation was quantitatively confirmed by comparing the absorbance of differentiated and undifferentiated cells (Supplementary data 2).

### RAW264.7-differentiated 3T3-L1 adipocytes co-culture

Following differentiation, 3T3-L1 adipocytes were co-cultured with RAW264.7 macrophages using the Transwell system (0.4  $\mu$ m porous membrane; Corning). In the co-culture, differentiated 3T3-L1 cells ( $1 \times 10^5$ ) were cultured in the lower well; while,  $5 \times 10^4$  RAW264.7 macrophages were cultured in inserts that constituted the upper chamber. Following the establishment of co-culture, RAW264.7 cells in the upper chamber were treated with MTC (5–20  $\mu$ M) for 30 min, followed by stimulation with LPS (1  $\mu$ g/mL) and IFN $\gamma$  (10 ng/mL) for a further 24 h. Culture supernatants were collected and analysed for levels of NO, TNF $\alpha$ , IL-1 $\beta$ , IL-6 using similar protocols described in experiments

on RAW264.7 macrophage mono-culture. Production of monocyte chemoattractant protein-1 (MCP-1/CCL2) was measured using a mouse CCL2 ELISA kit (Invitrogen); while, levels of RANTES in culture supernatants were detected with a mouse CCL5/RANTES ELISA kit (R and D Systems).

Glucose uptake was determined by incubating differentiated 3T3-L1 adipocytes in the lower chamber with 2-deoxy-2-[(7-nitro-2, 1, 3-benzoxadiazol-4-yl) amino]-D-glucose (2-NBDG) (100  $\mu$ M) in glucose-free DMEM for a further 1 h after treatment. Thereafter, cells were washed with PBS and glucose uptake by fluorescence detection at an excitation/emission wavelength of 485 nm/535 nm.


### Statistical analyses

Values of all experiments were represented as a mean  $\pm$  SEM of at least three experiments. Values were compared using one-way ANOVA followed by a post hoc Tukey test. In experiments to confirm differentiation of adipocytes, differences in oil red O staining between undifferentiated and differentiated adipocytes were determined using the Student's *t* test.

## Results

### Structural characterisation of methyl 3,4,5-trimethoxycinnamate

Yield 93%; white solid;  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{H}}$  7.59 (d,  $J = 16.0$  Hz, 1H), 6.73 (s, 2H), 6.33 (d,  $J = 16.0$  Hz, 1H), 3.87 (s, 9H), 3.79 (s, 3H).  $^{13}\text{C}$  NMR (50 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{C}}$  167.5, 153.5, 144.9, 140.1, 129.9, 117.1, 105.2, 61.1, 56.2, 51.8. IR  $\nu_{\text{max}}$  (KBr,  $\text{cm}^{-1}$ ) 3003, 1697, 2943, 2837, 1632, 1040, 1005, 1582, 1506, 851.



**Fig. 1** Chemical structure of Methyl 3,4,5-trimethoxycinnamate

### MTC produced anti-inflammatory activity in LPS + IFN $\gamma$ -stimulated RAW264.7 macrophages


In experiments to determine whether MTC produces anti-inflammatory activity, RAW265.7 cells were stimulated with IFN $\gamma$  (10 ng/mL) and LPS (1  $\mu$ g/mL) for 24 h. This resulted in  $\sim 80$ -fold increase in TNF $\alpha$  secretion into the culture supernatant in comparison with control cells. However, when cells were treated with 5-, 10- and 20- $\mu$ M MTC, there was  $\sim 1.8$ -, 2.1- and 2.5-fold reduction in LPS + IFN $\gamma$ -induced increase in TNF $\alpha$  production, respectively (Fig. 2a). MTC (5–20  $\mu$ M) produced similar effects on the release of IL-6 and IL-1 $\beta$  in LPS + IFN $\gamma$ -stimulated RAW264.7 cells (Fig. 2b, c).

In other experiments, LPS + IFN $\gamma$  stimulation caused a significant ( $p < 0.01$ ) reduction in the levels of IL-10 in RAW264.7 cells (Fig. 2d). On pre-treating the cells with MTC (5  $\mu$ M), there was a statistically insignificant increase in IL-10 release. However, when the concentrations of MTC were increased to 10 and 20  $\mu$ M, significant ( $p < 0.05$ ) increase in IL-10 levels was observed in culture supernatants.

To further confirm the anti-inflammatory activity of MTC, we measured the levels of NO (in form of nitrite) released in culture supernatants obtained from LPS + IFN $\gamma$ -stimulated cells, and observed that pre-treatment with MTC (5–20  $\mu$ M) resulted in significant ( $p < 0.05$ ) reduction in NO production. The results show that NO concentrations in cells stimulated with LPS + IFN $\gamma$  alone was  $\sim 15.4$   $\mu$ M. This was reduced to  $\sim 9.1$ ,  $\sim 9.0$ , and  $\sim 7.1$   $\mu$ M in cells pre-treated with 5, 10 and 20  $\mu$ M of MTC, respectively (Fig. 3a). We also observed that LPS + IFN $\gamma$ -induced increase in iNOS protein levels were markedly reduced in the presence of MTC (5–20  $\mu$ M) (Fig. 3b). It was observed that increases in PGE $_2$  production as well as elevation in COX-2 protein expression caused by stimulating RAW264.7 cells with LPS + IFN $\gamma$  were significantly reduced in the presence of 10 and 20  $\mu$ M of MTC. Reductions observed with 5  $\mu$ M of the compound were not significant (Fig. 4a, b).


### Anti-inflammatory activity of MTC is mediated through inhibition of NF- $\kappa$ B activity

Based on results showing anti-inflammatory activity of MTC, we were next interested to determine whether the compound produced this action by targeting NF- $\kappa$ B signaling in RAW264.7 macrophages. Results of in-cell western blotting in Fig. 5a–c show that stimulating RAW264.7 cells resulted in marked increase in phosphorylation of p65 sub-unit and I $\kappa$ B protein, accompanied by degradation of I $\kappa$ B protein. Pre-treatment of RAW264.7 cells with MTC prior to stimulation with LPS + IFN $\gamma$  resulted in significant ( $p < 0.001$ ) reduction in the expression of phospho-p65



**Fig. 2** Pre-treatment of RAW264.7 macrophages with MTC (5–20  $\mu$ M) prior to stimulation with LPS (1  $\mu$ g/mL) and IFN (10 ng/mL) for 24 h resulted in reduction in the concentrations of TNF $\alpha$  (a), IL-6 (b), and IL-1 $\beta$  (c). Levels of IL-10 were increased, compared

with LPS stimulation (d). Results are mean  $\pm$  SEM of three independent experiments. Data were analysed using one-way ANOVA for multiple comparisons with post-hoc Tukey test. \* $p$  < 0.05, \*\* $p$  < 0.01, \*\*\* $p$  < 0.001, *ns* not significant, compared with LPS + IFN $\gamma$



**Fig. 3** MTC (5–20  $\mu$ M) reduced NO production a and iNOS protein levels b in RAW264.7 macrophages stimulated with LPS (1  $\mu$ g/mL) and IFN (10 ng/mL) for 24 h. Results are mean  $\pm$  SEM of three inde-


pendent experiments. Data were analysed using one-way ANOVA for multiple comparisons with post-hoc Tukey test. \* $p$  < 0.05, \*\* $p$  < 0.01, \*\*\* $p$  < 0.001, compared with LPS + IFN $\gamma$

protein at all the concentrations investigated (Fig. 5a). Interestingly, at 5- $\mu$ M concentration of MTC, reduction in the levels of phospho-I $\kappa$ B was not significant. However, at higher concentrations of 10- and 20- $\mu$ M, phosphorylation of I $\kappa$ B was significantly ( $p$  < 0.05) reduced (Fig. 5b). Furthermore, LPS + IFN $\gamma$ -induced degradation of I $\kappa$ B was significantly ( $p$  < 0.01) reduced by MTC (5–20  $\mu$ M) (Fig. 5c).

In reporter gene assay experiments, LPS + IFN $\gamma$  stimulation of RAW264.7 cells which were transiently

transfected with the luciferase-driven NF- $\kappa$ B reporter resulted in a significant ( $p$  < 0.001) increase in luciferase activity (Fig. 5d). Treatment of the cells with MTC (5–20  $\mu$ M) prior to inflammatory stimulation resulted in significant ( $p$  < 0.01) and concentration-dependent reduction in luciferase activity. Similarly, it was observed that LPS + IFN $\gamma$ -induced DNA binding of NF- $\kappa$ B was significantly ( $p$  < 0.01) reduced in a concentration-dependent fashion (Fig. 5e).





**Fig. 4** MTC (5–20  $\mu$ M) reduced PGE<sub>2</sub> production **a** and COX-2 protein levels **b** in RAW264.7 macrophages stimulated with LPS (1  $\mu$ g/mL) and IFN (10 ng/mL) for 24 h. Results are mean  $\pm$  SEM of three independent experiments. Data were analysed using one-

way ANOVA for multiple comparisons with post-hoc Tukey test. \* $p < 0.05$ , \*\*\* $p < 0.001$ , *ns* not significant, compared with LPS + IFN $\gamma$

### MTC activated Nrf2/HO-1 protective mechanism in RAW264.7 macrophages

We also investigated whether MTC produces antioxidant effects in RAW264.7 cells through activation of Nrf2 and the subsequent increase in the expression of HO-1 protein. We observed that the degree of DNA binding of Nrf2 increased ~threefold from baseline (control) values when the cells were treated with 5- $\mu$ M MTC. However, binding increased ~3.2- and ~5.2-fold with 10- and 20- $\mu$ M concentrations of MTC, respectively (Fig. 6a). To further establish the effect of MTC on Nrf2 activity, we carried out a reporter gene assay to determine whether the compound would have an effect on luciferase-driven ARE activity. Results in Fig. 6b show that treatment with MTC (5–20  $\mu$ M) resulted in significant ( $p < 0.001$ ) increase in ARE-luciferase activity, when compared with untreated control cells. Experiments to evaluate the effect of MTC on HO-1 protein showed that treatment of RAW264.7 cells with MTC (5–20  $\mu$ M) caused a significant ( $p < 0.05$ ) increase in protein levels of HO-1 in the cells (Fig. 6c).


### Treatment with MTC prevents inflammation in RAW264.7–3T3-L1 adipocytes co-culture

Results in Fig. 7a show that addition of LPS (1  $\mu$ g/mL) and IFN $\gamma$  (10 ng/mL) to the RAW264.7 layer of macrophage–adipocyte co-culture resulted in an increase in the concentration of NO in the culture supernatants. However, treatment with MTC (5–20  $\mu$ M) prior to inflammatory stimulation resulted in a significant ( $p < 0.001$ ) and concentration-dependent reduction in NO production. Similar observations were made following analyses of culture supernatants for levels of pro-inflammatory cytokines TNF $\alpha$  (Fig. 7b), IL-6 (Fig. 7c) and IL-1 $\beta$  (Fig. 7d).

To evaluate the effects of MTC on the secretion of chemokines implicated in inflammation-mediated insulin resistance, we analysed culture supernatants and found that following stimulation of the RAW264.7 layer with LPS (1  $\mu$ g/mL) and IFN $\gamma$  (10 ng/mL), there was ~23-fold increase in the levels of MCP-1, in comparison with control cells. Supernatants from cells that were pre-treated with 5, 10 and 20  $\mu$ M of MTC showed ~1.30-, ~1.42- and ~1.71-fold reduction in the concentrations of MCP-1, respectively (Fig. 7e). Similarly, inflammatory stimulation of RAW264.7 layer caused a significant ( $p < 0.001$ ) elevation in the levels of RANTES in the culture supernatants, and these were significantly ( $p < 0.01$ ) reduced following pre-treatment with MTC (5–20  $\mu$ M) (Fig. 7f).

### MTC enhances glucose uptake through activation of AMPK $\alpha$ in 3T3-L1 adipocytes

Based on our observation on the effects of MTC on inflammation-induced MCP-1 and RANTES secretion in a co-culture of RAW264.7 macrophages and 3T3-L1 adipocytes, we became interested in evaluating whether these effects were accompanied by enhancement of glucose uptake in the adipocytes. Expectedly, treatment of adipocytes with only insulin (100 nM) for 24 h caused a marked increase in 2-NBDG uptake by the cells, in comparison with untreated cells (Fig. 8a). However, we also observed that stimulation of the macrophage layer of the co-culture with LPS (1  $\mu$ g/mL) and IFN $\gamma$  (10 ng/mL) for the same period of time resulted in a slight reduction in 2-NBDG uptake when compared with control. Interestingly, treatment with MTC (5–20  $\mu$ M) prior to inflammatory stimulation of RAW264.7 macrophages resulted in significant ( $p < 0.01$ ) increase in 2-NBDG uptake (Fig. 8a).



**Fig. 5** **a** MTC (5–20  $\mu$ M) reduced phosphorylation of p65 in the cytoplasm of LPS (1  $\mu$ g/mL)+IFN $\gamma$  (10 ng/mL)-stimulated RAW264.7 macrophages following 60-h incubation. **b** Reduction in the levels of phospho-I $\kappa$ B $\alpha$  in LPS (1  $\mu$ g/mL)+IFN $\gamma$  (10 ng/mL)-stimulated RAW264.7 macrophages by MTC (5–20  $\mu$ M). **c** MTC (5–20  $\mu$ M) prevented LPS (1  $\mu$ g/mL)+IFN $\gamma$  (10 ng/mL)-induced degradation of I $\kappa$ B $\alpha$  in RAW264.7 macrophages. MTC inhibited


NF- $\kappa$ B-mediated gene transcription (**d**) and DNA binding in LPS (1  $\mu$ g/mL)+IFN $\gamma$  (10 ng/mL)-stimulated RAW264.7 macrophages (**e**). Results are mean  $\pm$  SEM of three independent experiments. Data were analysed using one-way ANOVA for multiple comparisons with post-hoc Tukey test. \* $p$  < 0.05, \*\*\* $p$  < 0.001, *ns* not significant, compared with LPS + IFN $\gamma$

Stimulation of the macrophage layer with LPS (1  $\mu$ g/mL) and IFN $\gamma$  (10 ng/mL) caused a significant ( $p$  < 0.001) reduction in phospho-AMPK $\alpha$  protein in adipocytes after 24 h (Fig. 8b). Pre-treatment with MTC (5  $\mu$ M) did not have significant effect on AMPK $\alpha$  activation. However, when the concentration of the compound was increased to 10 and 20  $\mu$ M, there was a significant inhibition of deactivation of adipocyte AMPK $\alpha$  induced by inflammatory stimulation of macrophages (Fig. 8b).

In Fig. 8c we show that in the presence of the AMPK $\alpha$  inhibitor, compound C (10  $\mu$ M), increase in adipocyte glucose uptake following treatment with MTC (20  $\mu$ M) was insignificantly ( $p$  < 0.05) reversed.

## Discussion

Macrophage-mediated chronic inflammation is known to be a major contributor in many diseases, and several strategies are currently being deployed in preventing or blocking inflammation. Secondary metabolites from natural products have been extensively investigated as a potential source of new chemical scaffolds for new anti-inflammatory drugs. In this study, we have shown that a synthetic form of natural 3, 4, 5-trimethoxycinnamic acid, MTC reduced the release of pro-inflammatory cytokines TNF $\alpha$ , IL-6 and IL-1 $\beta$  in LPS + IFN $\gamma$ -stimulated RAW264.7 cells, while increasing the production of the anti-inflammatory cytokine IL-10. We



**Fig. 6** Activation of Nrf2/HO-1 by MTC in RAW264.7 macrophages. MTC (5–20 μM) treatment increased DNA binding of Nrf2 to immobilised ARE consensus binding site in BV-2 microglia (**a**). MTC activated ARE-luciferase activity in RAW264.7 macrophages transfected with ARE construct (**b**). Increase in HO-1 protein levels by treatment

with MTC (5–20 μM). Results are mean ± SEM of three independent experiments. Data were analysed using one-way ANOVA for multiple comparisons with post-hoc Tukey test. \* $p < 0.05$ , \*\*\* $p < 0.001$ , ns not significant, compared with untreated control

further observed that MTC treatment resulted in the reduction of both NO and PGE<sub>2</sub> release through mechanisms involving inhibition of iNOS and COX-2 proteins, respectively. These effects demonstrate anti-inflammatory activity of MTC.


The release of pro-inflammatory mediators from macrophages has been implicated in many inflammatory and auto-immune diseases. In rheumatoid arthritis, macrophages secrete potent cytokines and chemokines including TNF, IL-1, IL-6, IL-10, IL-12, IL18, IL-15, IL-10, GM-CSF, M-CSF and TGFβ which contribute to the chronic inflammation associated with the disease (Siouti and Andreaskos 2019). Pro-inflammatory macrophages have also been linked to diseases such as atherosclerosis (Bäck et al. 2019), inflammatory bowel diseases (Heinsbroek and Gordon 2009) and the metabolic syndrome (Kumar 2019). The anti-inflammatory activity of MTC, therefore, suggests a potential therapeutic strategy in treating some of these conditions.

Studies have suggested that phenylpropanoids show anti-inflammatory activity in a number of experimental models. Phenylpropanoids isolated from the Chinese olive

(*Canarium album*) and *Polygala tenuifolia* were reported to inhibit neuroinflammation in LPS-activated BV-2 microglia (Cho et al. 2012; Zhang et al. 2019). Similarly, phenylpropanoids isolated from *Ficus hirta* and *Brassica oleracea* inhibited LPS-induced nitric oxide production in RAW 264.7 macrophages (Lee et al. 2014; Cheng et al. 2017). Interestingly, some synthetic derivatives of 3, 4, 5-trimethoxycinnamic acid have been suggested to have anti-inflammatory activity in TNFα-stimulated endothelial cells (Kumar et al. 2011). To our knowledge, this is the first study demonstrating anti-inflammatory activity of MTC in murine macrophages, and suggests that this compound is a potential chemical scaffold for the identification of novel anti-inflammatory agents.

The transcription factor NF-κB is now widely accepted as the master regulator of the genes encoding inflammatory proteins, including iNOS, COX-2, TNFα, IL-6 and IL-1β. NF-κB is also an important target for the discovery of new drugs for treating for various inflammatory disorders. Based on our results showing that MTC produces anti-inflammatory activity in RAW264.7 macrophages, we carried out





**Fig. 7** Pre-treatment of RAW264.7 macrophage–3T3-L1 adipocyte co-culture with MTC (5–20  $\mu$ M) prior to stimulation with LPS (1  $\mu$ g/mL) and IFN (10 ng/mL) for 24 h resulted in reduction in the concentrations of NO (a), TNF $\alpha$  (b), IL-6 (c), IL-1 $\beta$  (d), MCP-1 (e) and


CCL5/RANTES (f). Results are mean  $\pm$  SEM of three independent experiments. Data were analysed using one-way ANOVA for multiple comparisons with post-hoc Tukey test. \* $p$  < 0.05, \*\* $p$  < 0.01, \*\*\* $p$  < 0.001, compared with LPS + IFN $\gamma$

further investigations to determine whether this compound could target the activation of NF- $\kappa$ B. Our experiments revealed that MTC suppressed I $\kappa$ B-mediated activation and nuclear DNA binding of NF- $\kappa$ B, as well as NF- $\kappa$ B-mediated gene transcription in RAW264.7 cells. The effects of the compound on NF- $\kappa$ B further confirms its anti-inflammatory activity in peripheral macrophages.

Investigations by us and others have suggested that the anti-inflammatory action of some compounds could be due, at least in part, to their ability to activate the Nrf2 transcription factor (Onasanwo et al. 2016; Okorji et al. 2016; Velagapudi et al. 2017; 2019a; Zhou et al. 2018). This prompted us to explore and show that MTC increased DNA binding of Nrf2 and enhanced ARE-mediated antioxidant gene transcription in RAW264.7 cells. While this is the

first evidence linking MTC to activation of Nrf2, studies reported by Schädlich et al. (2016) suggest that ginger (*Zingiber officinale*) phenylpropanoids activate Nrf2 in human keratinocytes. Our studies could not establish a link between the anti-inflammatory activity and activation of Nrf2/ARE mechanisms by MTC, thus warranting further investigations.

Studies have shown pro-inflammatory cytokines secreted by recruited and resident macrophages in adipocytes play a significant role in linking inflammation and insulin signalling (Shapiro et al. 2011). Specifically, pro-inflammatory cytokines like TNF $\alpha$ , which are secreted as a result of the interaction between adipocytes and macrophages have been linked to dysfunctions in glucose metabolism and insulin resistance (Guilherme et al. 2008). Consequently, we investigated effects of MTC on



**Fig. 8** Treatment of macrophage–adipocyte co-culture with MTC (5–20 µM) prior to LPS (1 µg/mL) and IFN (10 ng/mL) enhanced 2-NBDG uptake in adipocytes (a), and activated AMPK $\alpha$  (b). The enhancement of 2-NBDG uptake by MTC (20 µM) was partially

prevented in the presence of the AMPK inhibitor, compound C (c). Results are mean  $\pm$  SEM of three independent experiments. Data were analysed using one-way ANOVA for multiple comparisons with post hoc Tukey test. \* $p$  < 0.05, \*\* $p$  < 0.01, \*\*\* $p$  < 0.001, *ns* not significant

inflammatory responses in a co-culture of RAW264.7 macrophages and differentiated 3T3-L1 adipocytes. Our results revealed that stimulating the macrophage layer of the co-culture resulted in significant increases in the concentrations of NO as well as the pro-inflammatory cytokines TNF $\alpha$ , IL-6 and IL-1 $\beta$ . These observations reflect earlier reports linking significant production of pro-inflammatory cytokines following LPS stimulation of macrophage–adipocyte interactions (Yamashita et al. 2007; Fite et al. 2015; Engin 2017; Gothai et al. 2016). Results also showed that MTC prevented macrophage-mediated inflammation in adipocytes, through reduction of elevated levels of adipocyte-specific chemokines MCP-1 and RANTES following inflammatory stimulation of adjacent macrophages. Emerging evidence from several reports have suggested that the use of anti-inflammatory phytochemicals such as flavonoids could be a viable strategy in treating insulin resistance (Miranda et al. 2015; Gun and Lambert 2013; Merone and McDermott 2017). Similar properties have been reported for some novel multi-targeting anti-inflammatory glitazones (Elzahhar et al. 2019). The potential effect of MTC in preventing inflammation-mediated insulin resistance was further confirmed in experiments demonstrating its ability to enhance glucose uptake in adipocytes.

AMPK is an energy sensor that controls glucose metabolism, and has been suggested to suppress the pro-inflammatory environment in adipocytes (Bijland et al. 2013). We showed in this study that MTC suppressed macrophage activation-mediated dephosphorylation of AMPK $\alpha$  in adipocytes, suggesting that this kinase plays a role in the effects of the compound in reducing inflammation and enhancing glucose uptake. This was further confirmed in pharmacological antagonism experiments with the AMPK inhibitor compound C, which show that the glucose uptake promoting effect of MTC in adipocytes was partially reversed.

This study has established that MTC produced anti-inflammatory activity in both RAW264.7 macrophages and in a macrophage–adipocyte co-culture. It is further suggested that the anti-inflammatory activity of this compound possibly contributes to its ability to enhance glucose uptake in adipocytes through mechanisms involving activation of AMPK. It is not clear if this compound is able to affect the levels of other markers of inflammation and insulin resistance in adipocytes; further studies are establishing these possible effects.

**Acknowledgements** This work was supported by the Brazilian agencies: National Council for Scientific and Technological Development

(CNPq), and Coordination for the Improvement of Higher Education Personnel (CAPES). Dr Idowu S Akande received a Travel Grant (Needs Assessment Intervention Fund) from the University of Lagos, Nigeria.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Bäck M, Yurdagül A Jr, Tabas I, Öörni K, Kovanen PT (2019) Inflammation and its resolution in atherosclerosis: mediators and therapeutic opportunities. *Nat Rev Cardiol* 16:389–406
- Bijland S, Mancini SJ, Salt IP (2013) Role of AMP-activated protein kinase in adipose tissue metabolism and inflammation. *Clin Sci (Lond)* 124:491–507
- Cheng J, Yi X, Chen H, Wang Y, He X (2017) Anti-inflammatory phenylpropanoids and phenolics from *Ficus hirta* Vahl. *Fito-terapia* 121:229–234
- Cho N, Huh J, Yang H, Jeong EJ, Kim YC, Kim J, Sung SH (2012) Chemical constituents of *Polygala tenuifolia* roots and their inhibitory activity on lipopolysaccharide-induced nitric oxide production in BV2 microglia. *J Enzyme Inhib Med Chem* 27:1–4
- Cuadrado A, Manda G, Hassan A (2018) Transcription factor NRF2 as a therapeutic target for chronic diseases: a systems medicine approach. *Pharmacol Rev* 70:348–383
- da Nóbrega FR, Ozdemir O, Nascimento Sousa SCS, Barboza JN, Turkez H, de Sousa DP (2018) Piplartine analogues and cytotoxic evaluation against glioblastoma. *Molecules* 23:E1382
- Elzahhar PA, Alaaeddine R, Ibrahim TM et al (2019) Shooting three inflammatory targets with a single bullet: novel multi-targeting anti-inflammatory glitazones. *Eur J Med Chem* 167:562–582
- Engin AB (2017) Adipocyte-macrophage cross-talk in obesity. *Adv Exp Med Biol* 960:327–343
- Fite A, Abou-Samra AB, Seyoum B (2015) Macrophages inhibit insulin signalling in adipocytes: role of inducible nitric oxide synthase and nitric oxide. *Can J Diabetes* 39:36–43
- Gothai S, Ganesan P, Park SY, Fakurazi S, Choi DK, Arulselvan P (2016) Natural phyto-bioactive compounds for the treatment of type 2 diabetes: inflammation as a target. *Nutrients* 8:E461
- Gu Y, Lambert JD (2013) Modulation of metabolic syndrome-related inflammation by cocoa. *Mol Nutr Food Res* 57:948–961
- Guilherme A, Virbasius JV, Puri V, Czech MP (2008) Adipocyte dysfunctions linking obesity to insulin resistance and type 2 diabetes. *Nat Rev Mol Cell Biol* 9:367–377
- Heinsbroek SE, Gordon S (2009) The role of macrophages in inflammatory bowel diseases. *Expert Rev Mol Med* 11:e14
- Hotamisligil GS (2006) Inflammation and metabolic disorders. *Nature* 444:860–867
- Kawashima K, Miyako D, Ishino Y, Makino T, Saito K, Kano Y (2004) Anti-stress effects of 3, 4, 5-trimethoxycinnamic acid, an active constituent of roots of *Polygala tenuifolia* (Onji). *Biol Pharm Bull* 27:1317–1319
- Kobayashi EH, Suzuki T, Funayama R (2016) Nrf2 suppresses macrophage inflammatory response by blocking proinflammatory cytokine transcription. *Nat Commun* 7:11624
- Kumar V (2019) Inflammation research sails through the sea of immunology to reach immunometabolism. *Int Immunopharmacol* 73:128–145
- Kumar S, Singh BK, Arya P et al (2011) Novel natural product-based cinnamates and their thio and thiono analogs as potent inhibitors of cell adhesion molecules on human endothelial cells. *Eur J Med Chem* 46:5498–5511
- Lee JW, Lee DY, Baek DR et al (2014) Phenylpropanoids from red kohlrabi sprouts inhibits nitric oxide production in RAW 264.7 macrophage cells. *Food Sci Biotechnol* 23:965–969
- Li C, Xu MM, Wang K, Adler AJ, Vella AT, Zhou B (2018) Macrophage polarization and meta-inflammation. *Transl Res* 191:29–44
- Merone L, McDermott R (2017) Nutritional anti-inflammatories in the treatment and prevention of type 2 diabetes mellitus and the metabolic syndrome. *Diabetes Res Clin Pract* 127:238–253
- Miranda J, Lasa A, Aguirre L, Fernandez-Quintela A, Milton I, Portillo MP (2015) Potential application of non-flavonoid phenolics in diabetes: anti-inflammatory effects. *Curr Med Chem* 22:112–131
- Mitchell JP, Carmody RJ (2018) NF-κB and the transcriptional control of inflammation. *Int Rev Cell Mol Biol* 335:41–84
- Okorji UP, Velagapudi R, El-Bakoush A, Fiebich BL, Olajide OA (2016) Antimalarial drug artemether inhibits neuroinflammation in BV2 microglia through nrf2-dependent mechanisms. *Mol Neurobiol* 53:6426–6443
- Onasanwo SA, Velagapudi R, El-Bakoush A, Olajide OA (2016) Inhibition of neuroinflammation in BV2 microglia by the biflavonoid kolaviron is dependent on the Nrf2/ARE antioxidant protective mechanism. *Mol Cell Biochem* 414:23–36
- Ren N, Kim E, Li B, Pan H, Tong T, Yang CS, Tu Y (2019) Flavonoids alleviating insulin resistance through inhibition of inflammatory signaling. *J Agric Food Chem* 67:5361–5373
- Schadich E, Hlaváč J, Volná T, Varanasi L, Hajdúch M, Džubák P (2016) Effects of ginger phenylpropanoids and quercetin on Nrf2-ARE pathway in human BJ fibroblasts and HaCaT keratinocytes. *Biomed Res Int* 2016:2173275
- Shapiro H, Lutaty A, Ariel A (2011) Macrophages, meta-inflammation, and immuno-metabolism. *Sci World J* 11:2509–2529
- Siouti E, Andreacos E (2019) The many facets of macrophages in rheumatoid arthritis. *Biochem Pharmacol* 165:152–169
- Sivandzade F, Prasad S, Bhalerao A, Cucullo L (2019) NRF2 and NF-κB interplay in cerebrovascular and neurodegenerative disorders: molecular mechanisms and possible therapeutic approaches. *Redox Biol* 21:101059
- Tsai IL, Lee FP, Wu CC, Duh CY, Ishikawa T, Chen JJ, Chen YC, Seki H, Chen IS (2005) New cytotoxic cyclobutanoid amides, a new furanoid lignan and anti-platelet aggregation constituents from *Piper arborescens*. *Planta Med* 71:535–542
- Velagapudi R, Kumar A, Bhatia HS, El-Bakoush A, Lepiarz I, Fiebich BL, Olajide OA (2017) Inhibition of neuroinflammation by thymoquinone requires activation of Nrf2/ARE signalling. *Int Immunopharmacol* 48:17–29
- Velagapudi R, Jamshaid F, Lepiarz I, Katola FO, Hemming K, Olajide OA (2019a) The tiliroside derivative, 3-O-[(E)-(2-oxo-4-(p-tolyl) but-3-en-1-yl)] kaempferol produced inhibition of neuroinflammation and activation of AMPK and Nrf2/HO-1 pathways in BV-2 microglia. *Int Immunopharmacol* 77:105951
- Velagapudi R, Lepiarz I, El-Bakoush A, Katola FO, Bhatia H, Fiebich BL, Olajide OA (2019b) Induction of autophagy and activation of SIRT-1 deacetylation mechanisms mediate

- neuroprotection by the pomegranate metabolite urolithin A in bv2 microglia and differentiated 3D human neural progenitor cells. *Mol Nutr Food Res* 63:e1801237
- Wardyn JD, Ponsford AH, Sanderson CM (2015) Dissecting molecular cross-talk between Nrf2 and NF- $\kappa$ B response pathways. *Biochem Soc Trans* 43:621–626
- Wellen KE, Hotamisligil GS (2005) Inflammation, stress, and diabetes. *J Clin Invest* 115:1111–1119
- Yamashita A, Soga Y, Iwamoto Y, Yoshizawa S, Iwata H, Koikeguchi S, Takashiba S, Nishimura F (2007) Macrophage–adipocyte interaction: marked interleukin-6 production by lipopolysaccharide. *Obesity (Silver Spring)* 15:2549–2552
- Yu XB, Liu GL, Zhu B, Hao K, Ling F, Wang GX (2014) *In vitro* immunocompetence of two compounds isolated from *Polygala tenuifolia* and development of resistance against grass carp reovirus (GCRV) and *Dactylogyrus intermedius* in respective host. *Fish Shellfish Immunol* 41:541–548
- Zhang S, Huang Y, Li Y, Wang Y, He X (2019) Anti-neuroinflammatory and antioxidant phenylpropanoids from Chinese olive. *Food Chem* 286:421–427
- Zhao Z, Fang M, Xiao D, Liu M, Fefelova N, Huang C, Zang WJ, Xie LH (2013) Potential antiarrhythmic effect of methyl 3,4,5-trimethoxycinnamate, a bioactive substance from roots of *Polygala radix*: suppression of triggered activities in rabbit myocytes. *Biol Pharm Bull* 36:238–244
- Zhou MM, Zhang WY, Li RJ, Guo C, Wei SS, Tian XM, Luo J, Kong LY (2018) Anti-inflammatory activity of Khayandirobilide A from *Khaya senegalensis* via NF- $\kappa$ B, AP-1 and p38 MAPK/Nrf2/HO-1 signaling pathways in lipopolysaccharide-stimulated RAW 264.7 and BV-2 cells. *Phytomedicine* 42:152–163

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.