

Synthesis and Biological Activities of NB-506 Analogues: Effects of the Positions of two Hydroxyl Groups at the Indole Rings

Mitsuru Ohkubo*, Teruyuki Nishimura, Teruki Honma, Ikuko Nishimura, Satoru Ito, Tomoko Yoshinari,
Hiroharu Arakawa Hiroyuki Suda, Hajime Morishima and Susumu Nishimura

*Banyu Tsukuba Research Institute in collaboration with Merck Research Laboratories,
Okubo 3, Tsukuba, Ibaraki, 300-2611, Japan*


Received 25 August 1999; accepted 21 October 1999

Abstract: In the course of a study of 6-*N*-amino-substituted analogues of NB-506 (**1**), a more potent anticancer drug, J-109,404 (**2**), in which the formyl group of NB-506 was replaced with a 1,3-dihydroxypropane group, was reported. A study of further modification in the positions of two hydroxyl groups at the indole rings of **2** resulted in the discovery of a 2,10-dihydroxy analogue, J-107,088 (**3**), which is a promising anticancer agent with a broader therapeutic window than J-109,404. © 1999 Elsevier Science Ltd. All rights reserved.

Keywords: Anticancer agents; Topoisomerase; Indolocarbazole; Substituent effects


DNA topoisomerase I has been reported to be an attractive target for the development of anticancer agents.¹⁾ Recently, NB-506 (**1**)²⁾, a DNA topoisomerase I inhibitor derived from a natural compound, BE-13793C (**4**)³⁾, was reported to be a potent anticancer drug. Previous studies of the 6-*N*-amino analogues of NB-506 to improve the potency as well as aqueous solubility yielded a more potent anticancer drug, J-109,404 (**2**), which has a 1,3-dihydroxypropane group at the 6-*N*-amino position.⁴⁾ This paper reports on the synthesis and biological activities of a new series of analogues of J-109,404 focused on the hydroxyl groups at the indole rings on an indolocarbazole skeleton. The *in vivo* anticancer effects of several potent compounds in mice are also discussed.

Fig. 1




CHEMISTRY

A new series of J-109,404 analogues focused on the position of the hydroxyl groups on the indole rings was synthesized from 6-*N*-methyl compounds **5** by the same method as for J-109,404, shown in **Scheme 1**, and the chemical yield was also summarized. The benzyl groups of the 6-*N*-methyl compounds **5** were removed by hydrogenolysis with palladium hydroxide followed by treatment with 2.0 M aqueous potassium hydroxide to yield anhydride compound **6**. Final compounds, **2**, **3**, **8-24** were obtained by the coupling reaction of **6** with hydrazine **7** in dimethylformamide (DMF) at 80 °C in 42-96% yields.



The most important issues for the effective synthesis of β -glycosides, 6-*N*-methyl compounds **5**, were regioselectivity and stereoselectivity in a glycosylation step. The novel synthetic pathways of an important intermediate, 6-*N*-methyl compounds **5**, are summarized in **Scheme 2**. In the case of symmetric analogues, 6-*N*-methyl compounds **5** could be readily prepared using the same glycosylation reactions as previously reported in the synthesis of NB-506.⁵⁾ In fact, the glycosylation reactions of 2,10- and 3,9-dibenzoyloxy-indolocarbazoles with 1-chloro-2,3,4,6-tetra-*O*-benzylglucose using potassium hydroxide or potassium *tert*-butoxide as a base were each carried out with more than 95% β -selectivity. On the other hand, non-symmetric β -glycosides were effectively obtained by the same method as previously reported,⁶⁾ in which the Mitsunobu reaction was used for the glycosylation reaction of mono-indole compounds or mono-*N*-*tert*-butoxycarbonyl (BOC) bisindole compounds with 2,3,4,6-tetra-*O*-benzylglucose. The stereoselectivity of the Mitsunobu reaction was greater than 90 %.

Scheme 2



Results and Discussion

Several biological activities of J-109,404 analogues are summarized in Table 1. As for topoisomerase-mediated DNA cleavage activity, the high selectivity of these analogues for topoisomerase II and protein kinase C (PKC) was completely maintained, and the hydroxyl group at the C-2 position obviously improved the topoisomerase I-mediated DNA cleavage activity, and in the case of di-hydroxyl analogues, the hydroxyl group at the C-10 position was also effective (**3**, **17** and **21**). The inhibitory activity against topoisomerase I tested by an enzyme assay (Topo-I cleavage, EC₅₀) did not always correlate to that determined by a cellular assay (K⁺/SDS, EC₂₀₀), probably because of their differences in penetration into the cells. However, from the results of the K⁺/SDS assay, two hydroxyl groups seemed to enhance penetration, because neither the mono-hydroxyl analogues nor the tri- or tetra-hydroxyl analogues showed potent EC₂₀₀ values, except for 2-OH compound **9**. As for cytotoxicity (CTX) toward P388 (murine leukemia), MKN-45 (human stomach cancer) and DLD-1

(human colon cancer) cells, the structure-activity relationships (SAR) were nearly the same as that in the K^+ /SDS assay. These results suggested that the number of hydroxyl groups influenced penetration into the cells while the positions of the hydroxyl groups affected the inhibitory activity against topoisomerase I; in particular, hydroxyl group at C-2 position seemed to be most important. Among the compounds, a 2,10-dihydroxy analogue, J-107,088 (**3**), showed not only the greatest activity in stabilizing a DNA-topoisomerase I cleavable complex ($EC_{200} = 0.10 \mu\text{M}$), but also more potent cytotoxicity against human cancer cells than J-109,404.

Table 1 *In vitro* activities of J-109,404 analogues

No.	R ₁	R ₂	Topo-I ^{a)} Cleavage EC ₅₀ (μM)	Topo-II ^{a)} Cleavage EC ₅₀ (μM)	K ⁺ /SDS ^{b)} (P388/S) EC ₂₀₀ (μM)	PKC ^{c)} IC ₅₀ (μM)	CTX ^{d)} P388/S IC ₅₀ (nM)	CTX ^{e)} MKN-45 IC ₅₀ (nM)	CTX ^{e)} DLD-1 IC ₅₀ (nM)
8	1-OH	H	0.65	>50	2.6	>200	22	640	2600
9	2-OH	H	0.23	>50	0.40	>200	3.1	25	1000
10	3-OH	H	1.9	>50	4.50	>200	47	4200	>30000
11	4-OH	H	0.49	NT ^{f)}	>10	>200	910	7100	>30000
12	H	8-OH	0.57	>50	>10	>200	700	2100	>30000
13	H	9-OH	1.1	>50	2.0	>200	130	720	5500
14	H	10-OH	0.65	>50	1.2	>200	13	110	2000
15	H	11-OH	0.21	>50	>10	>200	140	1600	10000
16	1-OH	9-OH	0.51	>50	0.55	>200	32	250	220
17	1-OH	10-OH	0.16	>50	0.40	>200	5.4	87	89
2 (J-109,404)	1-OH	11-OH	0.58	>50	0.45	>200	17	130	520
18	2-OH	9-OH	0.037	>50	0.35	>200	5.2	56	79
3 (J-107,088)	2-OH	10-OH	0.051	>50	0.10	100	1.5	4.8	120
19	2-OH	11-OH	0.055	>50	0.80	>200	10	65	200
20	3-OH	9-OH	0.21	>50	0.60	>200	6.8	250	11000
21	3-OH	10-OH	0.055	>50	0.65	90	3.5	120	340
22	3-OH	11-OH	0.13	>50	3.00	>200	31	300	5100
23	2-OH	9,11-OH	0.090	>50	>10	40	15000	>30000	>30000
24	1,3-OH	9,11-OH	0.35	>50	5.5	>200	>30000	>30000	>30000

a) Topoisomerase-mediated DNA cleavage assay was carried out using supercoiled pBR322 plasmid DNA.^{2b)} b) Effects on the formation of protein-DNA complex in P388 cells were investigated by the K^+ /SDS method.^{2b)} c) The histone II-As was used as a substrate for protein kinase C.^{2b)} d) Cytotoxicity (CTX) against murine leukemia cells (P388) was measured by the colorimetric tetrazolium-formazan method.^{2b)} e) Cytotoxicity (CTX) against human stomach cancer cells (MKN-45) and colon cancer cells (DLD-1) was measured by the colorimetric tetrazolium-formazan method and the sulforhodamine B dye-staining method.^{2b)} f) NT: not tested.

Several analogues were tested for anticancer effects in mice. As shown in Table 2, a good correlation between cytotoxicity and anticancer activity against human stomach cancer cells, MKN-45, was observed for the tested compounds, while toxicity (LD_{10}) did not correlate with cytotoxicity. In conclusion, a 2, 10-dihydroxy analogue, J-107,088 (**3**), was found to have potent anticancer activity and a very wide safety margin. J-107,088 (**3**) is now being tested clinically.⁷⁾

Table 2 Anticancer activity

	R ₁	R ₂	CTX MKN-45 IC ₅₀ (μM)	GID ₇₅ ^{c)} MKN-45 (mg / m ²)	LD ₁₀ ^{d)} (mg / m ²)	Safety Margin ^{e)} LD ₁₀ / GID ₇₅
9^{a)}	2-OH	H	0.025	800	1000	1.3
14^{a)}	H	10-OH	0.11	820	2700	3.3
16^{a)}	1-OH	9-OH	0.25	2500	1900	0.8
17^{a)}	1-OH	10-OH	0.087	220	370	1.7
2 (J-108,404)^{a)}	1-OH	11-OH	0.13	78	390	5.0
18^{b)}	2-OH	9-OH	0.056	290	1000	3.4
3 (J-107,088)^{b)}	2-OH	10-OH	0.0048	45	>1600	>36
19^{a)}	2-OH	11-OH	0.064	320	350	1.1
20^{a)}	3-OH	9-OH	0.25	300	380	1.3
21^{b)}	3-OH	10-OH	0.12	710	1000	1.4
22^{a)}	3-OH	11-OH	0.3	2200	1800	0.8

a) Compounds were injected intravenously five times / week for 2 weeks, and treatment was initiated when tumors grew to 0.2 cm³ or larger. b) Compounds were injected intravenously two times / week for 2 weeks, and treatment was initiated when tumors grew to 0.2 cm³ or larger. c) Anticancer effect on MKN-45 human stomach cancer cells implanted subcutaneously. into flanks of nude mice. GID₇₅; approximate 75% Growth Inhibition Dose. d) LD₁₀; approximate 10% Lethal Dose at the treatment schedule. e) Safety margin: the ratio of LD₁₀ / GID₇₅.

Acknowledgments: We are grateful to Atsushi Hirano, Taro Kakikawa, Hiroshi Kawamoto, Masato Nakano and Masashi Morita of our institute for their collaboration and stimulating discussions. This paper is dedicated to the memory of Dr. Akira Okura, who died during the course of this research. We are also grateful to Ms. D. LeBlanc, Merck & Co., for her critical reading of this manuscript.

References and Notes

1. a) Morham, S. G.; Kluckman, K. D.; Voulomanos, N.; Smithies, O. *Mol. Cell. Biol.* **1996**, *16*, 6804. b) Husain, I.; Mohler, J. L.; Seigler, H. F.; Besterman, J. M. *Cancer Res.* **1994**, *54*, 539.
2. a) Arakawa, H.; Iguchi, T.; Morita, M.; Yoshinari, T.; Kojiri, K.; Suda, H.; Okura, A.; Nishimura, S. *Cancer Res.* **1995**, *55*, 1316. b) Yoshinari, T.; Matsumoto, M.; Arakawa, H.; Okada, H.; Noguchi, K.; Suda, H.; Okura, A.; Nishimura, S. *Cancer Res.* **1995**, *55*, 1310. c) Fukasawa, K.; Komatani, H.; Hara, Y.; Suda, H.; Okura, A.; Nishimura, S.; Yoshinari, T. *Int. J. Cancer* **1998**, *75*, 145.
3. Kojiri, K.; Kondo, H.; Yoshinari, T.; Arakawa, H.; Nakajima, S.; Satoh, F.; Kawamura, K.; Okura, A.; Suda, H.; Okanishi, M. *J. Antibiot.* **1991**, *44*, 723.
4. Ohkubo, M.; Kojiri, K.; Kondo, H.; Tanaka, S.; Kawamoto, H.; Nishimura, T.; Nishimura, I.; Yoshinari, T.; Arakawa, H.; Suda, H.; Morishima, H.; Nishimura, S. *Bioorg. & Med. Chem. Lett.* **1999**, *9*, 1219.
5. a) Ohkubo, M.; Kawamoto, H.; Ohno, T.; Nakano, M.; Morishima, H. *Tetrahedron* **1997**, *53*, 585. b) Ohkubo, M. "Development of a new indolocarbazole anticancer agent (NB-506)". *Yuki Gosei Kagaku Kyokaiishi* **1997**, *55*, 566.
6. Ohkubo, M.; Nishimura, T.; Jona, H.; Honma, T.; Ito, S.; Morishima, H. *Tetrahedron* **1997**, *53*, 5937.
7. a) Yoshinari, T.; Ohkubo, M.; Fukasawa, K.; Egashira, S.; Hara, Y.; Matsumoto, M.; Nakai, K.; Arakawa, H.; Morishima, H.; Nishimura, S. *Cancer Res.* **1999**, *59*, 4271. b) Arakawa, H.; Morita, M.; Kodaera, T.; Okura, A.; Ohkubo, M.; Morishima, H.; Nishimura, S. "In vivo anti-tumor activity of a novel

indolocarbazole compound, J-107088 on murine and human tumors transplanted into mice" *Jap. J. of Cancer Res.* **1999**, in press.

8. Physical data for a representative compound, J-107,088 (**3**): mp >250 °C; $[\alpha]_D^{20} +163^\circ$; $^1\text{H-NMR}$ (300 MHz, DMSO- d_6), δ_{H} (ppm) : 3.2-3.3 (1H, m), 3.4-3.6 (6H, m), 3.78 (1H, m), 3.85-3.95 (2H, m), 4.02 (1H, m), 4.53 (2H, t, J = 5.4 Hz), 4.91 (1H, m), 5.11 (1H, d, J = 5.3 Hz), 5.32 (1H, d, J = 4.6 Hz), 5.55 (1H, d, J = 2.6 Hz), 5.86 (1H, t, J = 3.8 Hz), 5.97 (1H, d, J = 8.3 Hz), 6.80 (1H, dd, J = 2.0, 8.6 Hz), 6.82 (1H, dd, J = 2.0, 8.6 Hz), 6.98 (1H, d, J = 2.0 Hz), 7.18 (1H, d, J = 1.7 Hz), 8.79 (1H, d, J = 8.6 Hz), 8.87 (1H, d, J = 8.6 Hz), 9.75 (1H, s), 9.78 (1H, s), 11.20 (1H, s); HRMS (FAB) calcd for $\text{C}_{29}\text{H}_{29}\text{N}_4\text{O}_{11}$ 609.1833, found 609.1816.