

# RSC Advances

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: J. A. Rombouts, R. Veenboer, C. Vilellas, P. Lu, A. W. Ehlers, K. Andries, A. Koul, H. Lill, E. Ruijter, R. Orru, K. Lammertsma, D. Bald and J. C. Slootweg, *RSC Adv.*, 2016, DOI: 10.1039/C6RA22693K.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



## ARTICLE

# Synthesis, Characterization and Biological Activity of Fluorescently Labeled Bedaquiline Analogues

Jeroen A. Rombouts,<sup>a,b</sup> Richard M.P. Veenboer,<sup>a</sup> Cristina Villellas,<sup>c</sup> Ping Lu,<sup>d</sup> Andreas W. Ehlers,<sup>a,e</sup> Koen Andries,<sup>c</sup> Anil Koul,<sup>c</sup> Holger Lill,<sup>d</sup> Eelco Ruijter,<sup>b</sup> Romano V. A. Orru,<sup>b</sup> Koop Lammertsma,<sup>a,e</sup> Dirk Bald,<sup>\*,d</sup> J. Chris Slootweg<sup>\*,a</sup>

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Diarylquinolines represent a new class of antibiotics with high potency against *Mycobacterium tuberculosis*. As such, they are of urgent importance in the treatment of drug-resistant bacterial pathogens. In this work, we report a strategy for preparing fluorescently labeled derivatives of the FDA-approved diarylquinoline-based tuberculosis drug Bedaquiline. The labeled compounds were capable of blocking bacterial growth and interfered with the function of ATP synthase, the cellular target of diarylquinolines. This indicates that the chosen labeling strategy does not preclude the antibacterial activity of Bedaquiline, and allowed us to investigate the effect of labeling on drug recognition by bacterial efflux pumps in living *M. tuberculosis* strains. These properties, coupled with the efficient fluorescence of the attached BODIPY fluorophore means that these compounds can be used as a research tool to gain deeper understanding into the mechanism of action of this class of drugs.

## Introduction

In 2014, some 9 million individuals were newly infected worldwide with *Mycobacterium tuberculosis*, and ca. 1.5 million patients died from tuberculosis.<sup>[1]</sup> In particular, multi-drug resistant and extremely drug resistant strains of *M. tuberculosis* are unyielding to treatment with current tuberculosis chemotherapy.<sup>[2]</sup> Consequently, new drugs are urgently needed to counteract development of drug resistance. Using small molecules that inhibit energy metabolism is a new approach to kill *M. tuberculosis* and related pathogenic mycobacteria.<sup>[3]</sup> Several new or repurposed drug entities are known to block the oxidative phosphorylation pathway, a key route in energy metabolism. Clofazimine and

related phenothiazines interfere with the type-II NADH dehydrogenase function,<sup>[4]</sup> the imidazopyridines inhibit the cytochrome bc1 complex,<sup>[5]</sup> and the diarylquinolines inhibit ATP synthase in mycobacteria and other Gram-positive pathogenic bacteria.<sup>[6]</sup> Bedaquiline, the lead compound of the diarylquinolines, is a stereo-specific ATP synthase inhibitor<sup>[7]</sup> which acts strongly bactericidal against *M. tuberculosis*.<sup>[8]</sup> Bedaquiline has been approved by the US Food and Drug Administration and by the European Medicines Agency for treatment of multi-drug resistant tuberculosis. Bedaquiline and other drugs targeting energy metabolism may well contribute to shortening and simplifying tuberculosis chemotherapy.<sup>[9]</sup>

However, a detailed understanding of the mechanism of action of Bedaquiline is required in order to fully exploit its potential in chemotherapy, and perhaps develop novel Bedaquiline analogues. Utilization of a fluorescent drug derivative can be a valuable strategy to study antibacterial drugs, as reported earlier for the characterization of the benzothiazidinone class of drugs.<sup>[10]</sup> Two prerequisites must be fulfilled for successful implementation of drug labeling in such investigations, *i.e.*, that the fluorescence of the label is not quenched by interactions with the drug or the environment, and that the antibacterial action and target binding of the drug is not precluded by the presence of the label.

In this study we report on the design and execution of a labeling strategy for Bedaquiline, by covalently attaching a BODIPY fluorophore to the quinoline ring system of Bedaquiline (see Figure 1). The synthesis and photochemical

<sup>a</sup> Department of Chemistry and Pharmaceutical Sciences, Vrije Universiteit Amsterdam, De Boelelaan 1083, 1081 HV Amsterdam, The Netherlands.  
j.c.slootweg@vu.nl

<sup>b</sup> Department of Chemistry & Pharmaceutical Sciences and Amsterdam Institute for Molecules Medicines and Systems (AIMMS), Vrije Universiteit Amsterdam, De Boelelaan 1108, 1081 HZ Amsterdam, The Netherlands

<sup>c</sup> Infectious diseases and vaccines therapeutic area, Janssen Research & Development, Johnson & Johnson Pharmaceuticals, Turnhoutseweg 30, 2340 Beerse, Belgium

<sup>d</sup> Department of Molecular Cell Biology, AIMMS, Faculty of Earth and Life Sciences, Vrije Universiteit Amsterdam, De Boelelaan 1085, 1081 HV Amsterdam, The Netherlands  
d.bald@vu.nl


<sup>e</sup> Department of Chemistry, University of Johannesburg, Auckland Park, Johannesburg, 2006 South Africa

† Electronic Supplementary Information (ESI) available: In the supporting information, a pdf document is available containing all NMR spectroscopic data pertaining to the compounds mentioned, as well photochemical characterization of compounds **C1** and **C2**. See DOI: 10.1039/x0xx00000x

## ARTICLE

## Journal Name

characterization of two such fluorescent derivatives is described, and their antibacterial activity is determined. This specific fluorescent label is chosen by virtue of the compact size, photochemical stability and ease of chemical synthesis and modification of BODIPY-type compounds.<sup>[11]</sup> A linear polar spacer of variable length is used to separate the drug and the label, with the aim of keeping bactericidal activity intact and the fluorescence of the label unperturbed by the coupling to Bedaquiline. Considering the importance of creating labeled compounds that are sufficiently similar to the parent compound under investigation, we also evaluated if the newly prepared compounds share affinity for the same drug efflux pump system in *M. tuberculosis* as the parent compound.



**Figure 1: Structural formula of Bedaquiline and schematic depiction of fluorescent label.** Using a Sonogashira cross-coupling, the alkyne-terminated polar linker attached to the BODIPY fluorophore will be covalently coupled to the aryl bromide position of Bedaquiline.

## Materials and Methods

### Assay of ATP synthesis by mycobacterial membrane vesicles

Inverted membrane vesicles were prepared from *M. smegmatis*<sup>[12]</sup> and ATP synthesis was measured as described earlier.<sup>[13]</sup> Briefly, membrane vesicles (1 mg mL<sup>-1</sup>) from *M. smegmatis* were incubated in 50 mM morpholino-ethanesulfonic acid (MES, pH 6.5, 5 mM MgCl<sub>2</sub>, 2 mM ADP, 20 mM KH<sub>2</sub>PO<sub>4</sub>, 100 μM P<sup>1</sup>,P<sup>5</sup>-di(adenosine-5') pentaphosphate (Ap5A), 25 mM glucose, 11.8 U mL<sup>-1</sup> hexokinase (Sigma), and protease inhibitors (complete, EDTA-free; protease inhibitor cocktail tablets from Roche). Samples (0.25 mL) were incubated at 37 °C with vigorous stirring in 18-mL flasks. The reaction was initiated with 5 mM succinate. After 2 hours, each reaction was stopped with 25 mM EDTA, followed by transfer to ice. Samples were transferred to Eppendorf tubes, boiled for 5 min, and centrifuged (10,000 *g*, 20 min) to remove denatured protein. In supernatants, the synthesized glucose-6-phosphate was quantified by NADP<sup>+</sup> (2.5 mM) reduction in the presence of 3 U mL<sup>-1</sup> of glucose-6-phosphate dehydrogenase (Roche). NADPH formation was monitored with a spectrophotometer at 340 nm.

### MIC determination by Resazurin Microtiter Assay

The MIC of bodipy-labeled Bedaquiline compounds against *M. tuberculosis* in the presence or absence of the efflux inhibitor verapamil (at 40 μg/mL) was measured by the resazurin microtitre assay (REMA).<sup>[14]</sup> Serial 2-fold dilutions of antibiotics were performed in 7H9/0.5% glycerol/10% OADC (7H9 broth), in 96-well microtitre plates, with a final volume of 100 μL per well. A volume of 100 μL of inoculum suspension was added to each well. For

preparation of this inoculum either liquid cultures in logarithmic phase were adjusted to OD<sub>600</sub> ~ 0.3 and diluted 1/100 in 7H9 broth. Plates were incubated 7 days at 37 °C. To each well, 40 μL of a resazurin solution 0.1 mg/mL were added, and plates were incubated for an additional 48 hours at 37 °C. Resazurin is an indicator of bacterial growth; as a consequence of the metabolic activity of bacteria, resazurin (blue color, weakly fluorescent) is reduced to resofurin (pink color, fluorescent). Interpretation of results was based on visual inspection of change of color (blue to pink). MIC was defined as the lowest concentration of antibiotic that prevented growth.

### Synthetic Protocols; General Information

CH<sub>2</sub>Cl<sub>2</sub> (HPLC grade) was obtained from VWR and used as received. Anhydrous N,N-dimethylformamide (DMF) was obtained from Sigma-Aldrich and used as received. THF was distilled from sodium/potassium alloy and stored in a Schlenk flask before use. Solvents used for chromatography were of HPLC or spectrographic grade. All other reagents were obtained from Sigma-Aldrich or VWR and used as received. Column chromatography was performed on Merck "Type E" aluminum oxide (particle size 150 mesh) or SiliCycle silica gel (SiO<sub>2</sub>, 40 - 63 μm particles) using the described eluent systems. Thin Layer Chromatography (TLC) was performed using TLC plates from Merck (Al<sub>2</sub>O<sub>3</sub>: "Neutral F60" with F254 indicator on aluminium backing. SiO<sub>2</sub>: "Kieselgel F60" with F254 indicator on aluminium backing) and compounds were visualized by visual inspection and UV detection at 254/366 nm. Melting points were measured using ΔT = 1 °C min<sup>-1</sup> on a Stuart Scientific SMP3 melting point apparatus on the temperature range 50 °C – 400 °C, and are uncorrected. Boiling points were measured in a similar manner, using an open-ended boiling point tube. Infrared (IR) spectra were recorded neat using a Shimadzu FTIR-8400s spectrophotometer and wavelengths are reported in cm<sup>-1</sup>. <sup>1</sup>H, HSQC, HMBC and <sup>13</sup>C (Attached Proton Test) nuclear magnetic resonance (NMR) spectra were used to characterize prepared compounds, and were recorded on a Bruker Avance 500 (125.78 MHz for <sup>13</sup>C) or Bruker Avance 400 (100.64 MHz for <sup>13</sup>C) using the residual solvent as internal standard. <sup>19</sup>F spectra were recorded on a Bruker 250 spectrometer without any calibration. Electrospray Ionisation (ESI) mass spectrometry was carried out using a Bruker micrOTOF-Q instrument in positive ion mode (capillary potential of 4500 V). Electron Impact (EI) mass spectrometry was carried out using a Shimadzu GCMS-QP2010 Plus. Determination of absorption and emission spectra and the related parameters was performed using a Shimadzu UV-2501PC spectrophotometer (absorption measurements) and a Jobin Yvon Fluorolog spectrophotometer (fluorescence measurements). Bedaquiline fumarate was obtained from Johnson & Johnson; the free base used during synthesis was prepared by extracting a CH<sub>2</sub>Cl<sub>2</sub> solution of the fumarate three times with saturated NaHCO<sub>3</sub> solution. The identity and purity of the free base thus afforded from the material supplied by Johnson & Johnson was verified using HRMS, chiral HPLC and multinuclear NMR spectroscopy.

### Synthesis of 4-(but-3-yn-1-yloxy)benzaldehyde (A1)

To dry THF (200 mL) kept at 0 °C under an N<sub>2</sub> atmosphere was added 3-butyne-1-ol (1.00 g, 8.20 mmol, 1.0 eq.), triphenylphosphine (2.26 g, 8.60 mmol, 1.05 eq.) and 4-hydroxybenzaldehyde (574 mg,


8.20 mmol, 1.0 eq.).<sup>[19]</sup> After five minutes of stirring, diethylazodicarboxylate (DEAD; 2.2 M toluene solution, 3.73 mL, 8.20 mmol) was added via syringe. The solution was warmed up to room temperature overnight; a color change from yellow to colorless was observed. After 48 h, the solution was concentrated *in vacuo*. Column chromatography (SiO<sub>2</sub> stationary phase, gradient elution 5% EtOAc in cyclohexane to 15% EtOAc in cyclohexane) afforded 4-(but-3-yn-1-yloxy)benzaldehyde (354 mg, 2.03 mmol, 25%) as a white crystalline solid. TLC (4:1 cyclohexane:EtOAc): R<sub>f</sub> = 0.50, visualized using 254 nm light. <sup>1</sup>H NMR (500 MHz, DMSO): δ 9.87 (s, 1H, -CHO), 7.86 (d, 2H, <sup>3</sup>J(H,H) = 8.5 Hz, CH), 7.14 (d, 2H, <sup>3</sup>J(H,H) = 8.5 Hz, CH), 4.18 (t, 2H, <sup>3</sup>J(H,H) = 6.4 Hz, -OCH<sub>2</sub>-), 2.92 (t, 1H, <sup>4</sup>J(H,H) = 2.5 Hz, -CCH), 2.68 (dt, 2H, <sup>4</sup>J(H,H) = 2.5 Hz, <sup>3</sup>J(H,H) = 6.4 Hz, -CH<sub>2</sub>CC-) ppm. <sup>13</sup>C NMR (125.8 MHz, DMSO, 293 K): δ 191.4 (-CHO), 163.2, 131.9, 129.8, 115.0, 81.19 (-CCH), 72.7 (-CCH), 66.2 (-OCH<sub>2</sub>-), 18.8 (-CH<sub>2</sub>CC-) ppm. HRMS (ESI): Calc. [M+H]<sup>+</sup>: 175.0754; found 175.0746. MS (EI, + mode): Calc. [M]<sup>+</sup>: 174.1; found 174.1. Calc. [M-CHO]<sup>+</sup>: 145.1; found 145.1. Calc. [M-O(CH<sub>2</sub>)<sub>2</sub>CCH]<sup>+</sup>: 121.0; found 121.0. IR: 3209, 2824, 2806, 2745, 1672, 1601, 1576, 1529, 1508, 1393, 1250, 1215, 1163, 1109, 1038, 893, 827, 729, 652 cm<sup>-1</sup>. MP: 68–70 °C.

### Synthesis of But-3-yn-1-yloxy-BODIPY (B1)

To dry THF (100 mL) under an N<sub>2</sub> atmosphere was added aldehyde **A1** (300 mg, 1.73 mmol, 1.0 eq.), 2,4-dimethylpyrrole (362 mg, 3.81 mmol, 2.2 eq.) and 2 drops of trifluoroacetic acid. After 16 hours of stirring at room temperature, DDQ (398 mg, 1.73 mmol, 1.0 eq.) was added as a solution in 10 mL THF. After 1 hour of stirring, the reaction mixture was cooled to 0 °C using ice, and NEt<sub>3</sub> (2.0 mL, 14.3 mmol, ~ 8 eq.) was added via syringe followed by BF<sub>3</sub>·OEt<sub>2</sub> (2.5 mL, 25.3 mmol, ~ 15 eq.). The reaction mixture was allowed to warm up to room temperature over the course of two hours, after which it is filtered over celite. The celite is washed with 100 mL CH<sub>2</sub>Cl<sub>2</sub>; pooled organic phases are concentrated *in vacuo*. The brown, sticky solid thus obtained is diluted with 100 mL of EtOAc and washed with satd. NaHCO<sub>3</sub>, satd. NH<sub>4</sub>Cl, water and brine. After drying over Na<sub>2</sub>SO<sub>4</sub>, the organic layer is concentrated *in vacuo*. Column chromatography (SiO<sub>2</sub> stationary phase, gradient elution cyclohexane to 20% EtOAc in cyclohexane) afforded but-3-yn-1-yloxy-BODIPY **B1** (335 mg, 0.38 mmol, 22%) as an orange crystalline solid. TLC (2:1 cyclohexane:EtOAc): R<sub>f</sub> = 0.85, red spot with green fluorescence. <sup>1</sup>H NMR (500.2 MHz, CDCl<sub>3</sub>, 293 K): δ 7.17 (d, 2H, <sup>3</sup>J(H,H) = 8.5 Hz, phenyl CH), 7.12 (d, <sup>3</sup>J(H,H) = 8.5 Hz, 2H, phenyl CH), 5.98 (s, 2H, pyrrole CH), 4.16 (t, <sup>3</sup>J(H,H) = 7.0 Hz, 2H, -OCH<sub>2</sub>-), 2.74 (dt, 2H, <sup>4</sup>J(H,H) = 2.0 Hz, <sup>3</sup>J(H,H) = 7.0 Hz, -CH<sub>2</sub>CC-), 2.55 (s, 6H, pyrrole -CH<sub>3</sub>), 2.08 (d, <sup>4</sup>J(H,H) = 2.0 Hz, 2H, -CCH), 1.43 (s, 6H, pyrrole -CH<sub>3</sub>) ppm. <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>, 293 K): δ 159.0, 155.3, 143.1, 141.7, 131.8, 129.3 (phenyl CH), 127.5, 121.1 (pyrrole CH), 115.2 (phenyl CH), 80.2 (-CCH), 70.1 (-CCH), 66.0 (-OCH<sub>2</sub>-), 19.6 (CH<sub>2</sub>CCH), 14.6 (both pyrrole -CH<sub>3</sub> groups, signals overlap) ppm. <sup>19</sup>F NMR (235.4 MHz, CDCl<sub>3</sub>, 293 K): δ -146.3 (q, <sup>1</sup>J(F,B) = 33.2 Hz, BF<sub>2</sub>) ppm. HRMS (ESI): Calc. [M+H]<sup>+</sup>: 393.1872; found 393.1919. MS (EI, + mode): Calc. [M]<sup>+</sup>: 392.2; found 392.3. Calc. [M-HF]<sup>+</sup>: 372.2; found 372.3. IR: 3283, 2957, 2943, 2926, 2910, 2887, 1607, 1541, 1506, 1456, 1433, 1408, 1372, 1364, 1340, 1306, 1283, 1248, 1109, 1022, 1013, 892, 581 cm<sup>-1</sup>. MP: 151–154 °C (decomposition).

### Synthesis of Bedaquiline-butynyloxy-BODIPY (C1)

To a 20 mL glass reactor was added Bedaquiline fumarate (68 mg, 102 μmol, 1.0 eq.) and Butynyloxy-BODIPY **B1** (60 mg, 153 μmol, 1.5 eq.) in NEt<sub>3</sub> (3 mL). The solution was degassed with argon for 5 minutes, after which CuI (2 mg, 10 μmol, 0.1 eq.) and Pd(PPh<sub>3</sub>)<sub>4</sub> (12 mg, 10 μmol, 0.1 eq.) in 3 mL argon-degassed DMF was added via syringe from stock solutions prepared and kept under argon. The reactor was capped with a Teflon-lined plug. After stirring the reaction mixture at 90 °C for 2 hours, the reaction mixture was poured into 1:1 ethyl acetate : satd. NaHCO<sub>3</sub>. The ethyl acetate layer was washed twice with water and once with brine. The organic phase was dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated *in vacuo*. Column chromatography (first column SiO<sub>2</sub> stationary phase using 6:1 pentane:EtOAc to 1:1 pentane:EtOAc; second column SiO<sub>2</sub> stationary phase using 1:1 pentane:EtOAc to 100% EtOAc) afforded Bedaquiline-butynyloxy-BODIPY **C1** (Figure 2, 32.5 mg, 38 μmol, 37%) as a bright orange solid.



**Figure 2: Structural formula of Bedaquiline-butynyloxy-BODIPY.** The numbering scheme used in this work for elucidation of NMR spectroscopic data is shown.

TLC (Pentane:EtOAc 1:1): R<sub>f</sub> = 0.4, red spot with green fluorescence. <sup>1</sup>H NMR (500.2 MHz, CDCl<sub>3</sub>, 293 K): δ 8.94 (s, 1H, CH<sup>7</sup>), 8.61 (d, 1H, <sup>3</sup>J(H,H) = 8.5 Hz, naphthyl CH), 7.94 (s, 1H, CH<sup>9</sup>), 7.89 (t, 2H, <sup>3</sup>J(H,H) = 6.5 Hz, naphthyl CH), 7.77 (d, 1H, <sup>3</sup>J(H,H) = 8.5 Hz, CH<sup>12</sup>), 7.66 (d, 1H, <sup>3</sup>J(H,H) = 8.0 Hz, naphthyl CH), 7.62 (m, 1H, naphthyl CH; signals overlap), 7.61 (m, 1H, CH<sup>11</sup>; signals overlap), 7.49 (t, 1H, <sup>3</sup>J(H,H) = 7.0 Hz, naphthyl CH), 7.31 (t, 1H, <sup>3</sup>J(H,H) = 7.5 Hz, naphthyl CH), 7.20 (d, 2H, <sup>3</sup>J(H,H) = 8.5 Hz, phenol CH), 7.12 (br. s, 2H, phenyl CH<sup>19</sup>), 7.08 (d, 2H, <sup>3</sup>J(H,H) = 8.5 Hz, phenol CH), 6.89 (br. s, 3H, phenyl CH<sup>20</sup>/CH<sup>21</sup>), 5.98 (s, 2H, pyrrole CH), 5.89 (s, 1H, CH<sup>5</sup>), 4.28 (t, 2H, <sup>3</sup>J(H,H) = 7.0 Hz, OCH<sub>2</sub>-), 4.22 (s, 3H, OCH<sub>3</sub>), 3.01 (t, 2H, <sup>3</sup>J(H,H) = 7.0 Hz, -OCH<sub>2</sub>CH<sub>2</sub>-), 2.56 (s, 6H, pyrrole -CH<sub>3</sub>), 2.5 (1H, -CH<sup>3</sup><sub>2</sub>-, signals overlap), 2.13 (t, 1H, <sup>3</sup>J(H,H) = 13 Hz, -CH<sup>3</sup><sub>2</sub>-), 1.98 – 1.92 (m, 2H, -CH<sup>2</sup><sub>2</sub>-, signals overlap), 1.94 (s, 6H, N(CH<sub>3</sub>)<sub>2</sub>), 1.45 (s, 6H, pyrrole -CH<sub>3</sub>) ppm. <sup>13</sup>C NMR (125.8 MHz, CDCl<sub>3</sub>, 293 K): δ 161.6, 159.1, 155.3, 144.6, 143.2, 141.8, 140.6, 139.4, 134.7, 131.8, 131.8, 131.8, 131.3, 129.9, 129.9, 129.3, 128.2, 127.4, 127.3, 127.1, 126.8, 125.8, 125.3, 125.3, 125.2, 125.1, 124.5, 121.2, 118.5, 115.3, 85.3 (-CH<sub>2</sub>CC-), 82.3 (-CH<sub>2</sub>CC-), 66.4 (-OCH<sub>2</sub>-), 56.4 (C<sup>5</sup>), 54.2 (-OCH<sub>3</sub>), 49.5, 44.7 (N(CH<sub>3</sub>)<sub>2</sub>), 33.4, 20.6 (-CH<sub>2</sub>CC-), 14.7 (pyrrole CH<sub>3</sub>), 14.6 (pyrrole CH<sub>3</sub>) ppm. <sup>19</sup>F NMR (235.4 MHz, CDCl<sub>3</sub>, 293 K): δ -146.3 (q, <sup>1</sup>J(F,B) = 32.4 Hz, BF<sub>2</sub>) ppm. HRMS (ESI): Calc. [M+H]<sup>+</sup>: 867.4252; found 867.4201. No Bedaquiline starting material was found. IR: Broad peak at 3000 – 2700 cm<sup>-1</sup>. 1543, 1508, 1454, 1400, 1369, 1350, 1306, 1286, 1242, 1192, 1155, 1119, 1080, 1038, 1014, 926, 833, 806, 737, 704 cm<sup>-1</sup>. MP: 137–145 °C (decomposition).



## ARTICLE

## Journal Name

Synthesis of Pentynyloxy-eg<sub>4</sub>-OH

To dry DMF (12 mL) and tetraethylene glycol mono THP ether (4.90 g, 17.6 mmol, 1.0 eq), prepared according to Baker *et al.*<sup>[16]</sup> and kept in a dry Schlenk flask under nitrogen atmosphere, was added NaOMe (1.19 g, 22.0 mmol, 1.25 eq) at 0 °C. The clear solution was warmed up to room temperature over the course of 30 minutes. After again cooling down the reaction mixture to 0 °C, 5-chloro-1-pentyne (1.81 g, 17.6 mmol, 1.0 eq) and of sodium iodide (0.15 g, 1 mmol, 5 mol%) were added. The reaction mixture was heated to 80 °C for 16 hours. Subsequently, the reaction mixture was quenched with a little aqueous saturated NaHCO<sub>3</sub> solution, diluted with approximately 100 mL CH<sub>2</sub>Cl<sub>2</sub> and extracted three times with water. The organic phase was concentrated *in vacuo* and dissolved in methanol (100 mL) to which was added para-toluenesulfonic acid monohydrate (150 mg). The methanolic solution was stirred at room temperature for 72 hours, after which it was concentrated *in vacuo*, dissolved in 100 mL CH<sub>2</sub>Cl<sub>2</sub> and extracted once with saturated aqueous NaHCO<sub>3</sub> and twice with water. The organic phase was dried over magnesium sulfate, concentrated *in vacuo* and subjected to column chromatography (SiO<sub>2</sub> stationary phase, 100% EtOAc) to afford pentynyloxy-eg<sub>4</sub>-OH (695 mg, 2.67 mmol, 15%) as a colorless oil. TLC (100% EtOAc): R<sub>f</sub> = 0.20, visualized using iodine vapour or CerMop staining. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ 3.68 (2H, br. s.), 3.6–3.5 (10 H, m), 3.56 (4H, m), 3.51 (2H, t, <sup>3</sup>J(H,H) = 6.3 Hz), 2.24 (2H, dt, <sup>3</sup>J(H,H) = 7 Hz, <sup>4</sup>J(H,H) = 2.5 Hz, HCCCH<sub>2</sub>), 1.91 (1H, t, <sup>3</sup>J(H,H) = 2.5 Hz, HCC-), 1.75 (2H, m, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>O-) ppm. <sup>13</sup>C (125 MHz, CDCl<sub>3</sub>): δ 84.0 (-CCH), 72.6 (-CCH), 70.6, 70.6, 70.6, 70.3, 70.2, 69.6, 68.5, 61.7 (-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-), 28.5 (-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-), 15.2 (-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-) ppm. HRMS (ESI): Calc. [M+H]<sup>+</sup>: 261.1690 m/z. Found: 261.1697 m/z. Calc. [M+Na]<sup>+</sup>: 283.1518 m/z. Found: 283.1516 m/z. IR: ν 3485.1, 3427.3, 2908.5, 2358.8, 1249.8, 1120.6, 1110.9, 1072.4, 1033.8, 731.0, 530.4, 505.3, 493.7 m 418.5, 405.0 cm<sup>-1</sup>. Boiling point: 374 °C.

Synthesis of Pentynyloxy-eg<sub>4</sub>-benzaldehyde (A2)

To dry THF (20 mL) kept at 0 °C under an N<sub>2</sub> atmosphere was added pentynyloxy-eg<sub>4</sub>-OH (550 mg, 2.11 mmol, 1.0 eq.), triphenylphosphine (583 mg, 2.11 mmol, 1.0 eq.) and 4-hydroxybenzaldehyde (257 mg, 2.11 mmol, 1.0 eq.). After five minutes of stirring, DEAD (918 mg, 2.11 mmol, 1.0 eq) was added as a 40 wt% solution in toluene via a syringe. The solution was warmed up to room temperature over night; a color change from yellow to colorless was observed in the first 15 minutes. After 48 h, the solution was concentrated *in vacuo*. Column chromatography (SiO<sub>2</sub> stationary phase, gradient elution 100% cyclohexane to 1:1 EtOAc:cyclohexane followed by a second SiO<sub>2</sub> column using 8:1 EtOAc:CH<sub>2</sub>Cl<sub>2</sub>) afforded pentynyloxy-eg<sub>4</sub>-benzaldehyde **A2** (259 mg, 0.71 mmol, 34%) as a colorless oil. TLC (1:1 EtOAc:cyclohexane): R<sub>f</sub> = 0.25, visualized using 254 nm light. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 9.83 (s, 1H, -CHO), 7.78 (d, 1H, <sup>3</sup>J(H,H) = 8.8 Hz, CH), 6.97 (d, 1H, <sup>3</sup>J(H,H) = 8.8 Hz, CH), 4.16 (t, 2H, <sup>3</sup>J(H,H) = 4.8 Hz, benzaldehyde-OCH<sub>2</sub>-), 3.84 (t, 2H, <sup>3</sup>J(H,H) = 4.8 Hz, benzaldehyde-OCH<sub>2</sub>CH<sub>2</sub>-), 3.7–3.6 (m, 12H, -(OCH<sub>2</sub>CH<sub>2</sub>)<sub>3</sub>), 3.57 (t, 2H, <sup>3</sup>J(H,H) = 6.3 Hz, -OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CCH), 2.23 (dt, 2H, <sup>3</sup>J(H,H) = 6.9 Hz, <sup>4</sup>J(H,H) = 2.8 Hz, HCCCH<sub>2</sub>), 1.91 (t, 1H, <sup>4</sup>J(H,H) = 2.8 Hz, HCC-), 1.74 (m, 2H, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>O-) ppm. <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 190.9 (-CHO), 163.9 (C<sup>1</sup>), 132.0 (C<sup>3</sup>), 130.00 (C<sup>4</sup>), 114.9 (C<sup>2</sup>), 84.0 (-CCH), 70.9, 70.6, 70.6,

70.6, 70.6, 70.2, 69.5, 69.4, 67.8, 63.6 (-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-), 28.5 (-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-), 15.2 (-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-) ppm. HRMS (ESI): Calc. [M+H]<sup>+</sup>: 365.1951 m/z. Found: 365.1959 m/z. Calc. [M+Na]<sup>+</sup>: 387.1784 m/z. Found: 387.1778 m/z. IR: ν 2869.9, 1685.7, 1600.8, 1577.7, 1311.5, 1258.5, 1217.0, 1161.1, 1057.0, 833.2, 651.9 cm<sup>-1</sup>. Boiling point: 337 °C (decomposition).

## Sonogashira Test Reaction forming 3-(quinolin-6-yl)prop-2-yn-1-ol

In a 10 mL glass reactor, 6-bromoquinoline (210.5 mg, 1.01 mmol, 1 eq), 3-butyne-1-ol (141.0 mg, 2.01 mmol, 2 eq), Pd(PPh<sub>3</sub>)<sub>4</sub> (61.0 mg, 0.05 mmol, 5 mol%), CuBr (9.1 mg, 0.06 mmol, 6 mol%) and triethylamine (4 mL, 0.25 M) were dissolved, degassed using N<sub>2</sub> and reacted by heating the mixture to 90 °C for 70 minutes.<sup>[20]</sup> The reaction mixture was partitioned between a saturated aqueous solution of (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> and diethyl ether. The water layer was made alkaline by addition of a saturated aqueous solution of NaHCO<sub>3</sub> followed by extraction with diethyl ether. The combined organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>), volatiles were removed *in vacuo* and the crude was subjected to column chromatography (SiO<sub>2</sub>, gradient of pure cyclohexane to ethyl acetate) to yield the product, which was obtained as an oil (151 mg, 0.78 mmol, 77%). TLC (2:1 cyclohexane : EtOAc): R<sub>f</sub> = 0.50. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ 8.89 (m, 1H), 8.08 (d, 1H, <sup>3</sup>J(H,H) = 8.5 Hz), 8.01 (d, 1H, <sup>3</sup>J(H,H) = 8.5 Hz), 7.88 (s, 1H), 7.60 (d, 1H, <sup>3</sup>J(H,H) = 8.5 Hz), 7.40 (q, 1H, <sup>3</sup>J(H,H) = 4 Hz), 3.87 (t, 2H, <sup>3</sup>J(H,H) = 6.0 Hz), 2.75 (t, 2H, <sup>3</sup>J(H,H) = 6.0 Hz), 2.28 (br, 1H, OH). <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>): δ 150.9, 147.6, 135.8, 132.5, 131.2, 129.6, 128.1, 121.9, 121.8, 88.2, 82.1, 61.224.1.


Synthesis of Pentynyloxy-eg<sub>4</sub>-BODIPY (B2)

To dry THF (30 mL) under an N<sub>2</sub> atmosphere was added aldehyde **A2** (200 mg, 0.55 mmol, 1.0 eq.), 2,4-dimethylpyrrole (115 mg, 1.21 mmol, 2.2 eq.) and 1 drop of trifluoroacetic acid. After 16 hours of stirring at room temperature, DDQ (125 mg, 0.55 mmol, 1.0 eq.) was added as a solution in 1 mL THF. After 1 hour of stirring, the reaction mixture was cooled to 0 °C using ice, and NEt<sub>3</sub> (444 mg, 4.4 mmol, 8 eq) of NEt<sub>3</sub> was added via syringe followed by BF<sub>3</sub>·OEt<sub>2</sub> (780 mg, 5.5 mmol, 10 eq.). The reaction mixture was allowed to warm up to room temperature over the course of two hours, after which it was filtered over Celite. The celite was washed with CH<sub>2</sub>Cl<sub>2</sub>; pooled organic phases are concentrated *in vacuo*. The brown, sticky solid thus obtained was diluted with CH<sub>2</sub>Cl<sub>2</sub> and washed with satd. NaHCO<sub>3</sub>, satd. NH<sub>4</sub>Cl, water and brine. The pooled aqueous layers were back-extracted once with CH<sub>2</sub>Cl<sub>2</sub>. After drying over Na<sub>2</sub>SO<sub>4</sub>, the organic layer was concentrated *in vacuo*. Column chromatography (SiO<sub>2</sub> stationary phase, isocratic elution with 1:1 EtOAc:cyclohexane) gave pentynyloxy-eg<sub>4</sub>-BODIPY **B2** (141.7 mg, 0.23 mmol, 54%) as a brown-orange waxy solid. TLC (1:1 EtOAc:cyclohexane): R<sub>f</sub> = 0.20, red spot with green fluorescence. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ 7.14 (d, 2H, <sup>3</sup>J(H,H) = 8.5 Hz, phenol CH), 7.01 (d, <sup>3</sup>J(H,H) = 8.5 Hz, 2H, phenol CH), 5.97 (s, 2H, pyrrole CH), 4.17 (t, 2H, <sup>3</sup>J(H,H) = 5.0 Hz, -CH<sub>2</sub>-), 3.90 (t, 2H, <sup>3</sup>J(H,H) = 5.0 Hz, -CH<sub>2</sub>-), 3.76–3.59 (m, 12 H, -OCH<sub>2</sub>CH<sub>2</sub>O-), 3.55 (t, 2H, <sup>3</sup>J(H,H) = 6.3 Hz, -OCH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>CCH), 2.54 (s, 6H, pyrrole CH<sub>3</sub>), 2.28 (dt, 2H, -CH<sub>2</sub>CCH), 1.93 (t, 1H, -CCH), 1.79 (m, 2H, -OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-), 1.41 (s, 6H, pyrrole) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>): δ 159.4, 155.3, 143.3, 141.9, 132.0, 129.3 (phenol CH), 127.3, 121.2 (pyrrole CH), 115.3 (phenol CH), 84.1 (-CCH), 71.0, 70.8, 70.7, 70.4, 69.8, 69.7, 68.6,

67.6, 28.6 ( $-\text{OCH}_2\text{CH}_2\text{CH}_2-$ ), 15.3 ( $-\text{CH}_2\text{CCH}$ ), 14.8 (both pyrrole  $\text{CH}_3$  groups, signals overlap) ppm.  $^{19}\text{F}$  NMR (235.4 MHz,  $\text{CDCl}_3$ , 293 K):  $\delta$  -146.3 (q,  $^1J(\text{F},\text{B}) = 32.9$  Hz,  $\text{BF}_2$ ) ppm. HRMS (ESI): Calc.  $[\text{M}+\text{Na}]^+$ : 605.2969 m/z. Found: 605.2982 m/z. IR:  $\nu$  2922.0, 2864.1, 2358.8, 2337.6, 1608.5, 1541.0, 1508.2, 1467.7, 1407.9, 1363.6, 1305.7, 1286.4, 1245.9, 1191.9, 1155.3, 1982.0, 1049.2, 1012.6, 974.0, 853.1, 763.8, 704.0, 601.8, 582.5  $\text{cm}^{-1}$ . Melting point: 158–164  $^\circ\text{C}$  (decomposition).

### Synthesis of Bedaquiline-pentynoxy-eg<sub>4</sub>-BODIPY (C2)

To a 10 mL glass reactor was added Bedaquiline fumarate (76.6 mg, 114.5  $\mu\text{mol}$ , 1.0 eq.) and pentynoxy-eg<sub>4</sub>-BODIPY **B2** (99.7 mg, 172  $\mu\text{mol}$ , 1.5 eq.) in 5 mL  $\text{NEt}_3$ . The solution was degassed with argon for 5 minutes, after which  $\text{CuI}$  (1.1 mg, 6  $\mu\text{mol}$ , 5 mol%) and  $\text{Pd}(\text{PPh}_3)_4$  (26.4 mg, 23  $\mu\text{mol}$ , 0.2 eq.) in 5 mL argon-degassed DMF were added via syringe from stock solutions prepared and kept under argon. The reactor was capped with a Teflon-lined plug. After stirring the reaction mixture at 90  $^\circ\text{C}$  for 2 hours, the reaction mixture was poured into 1:1 ethyl acetate : satd.  $\text{NaHCO}_3$ . The ethyl acetate layer was washed twice with water and once with brine. The organic phase was dried over  $\text{Na}_2\text{SO}_4$  and concentrated *in vacuo*. Extensive column chromatography ( $\text{SiO}_2$  stationary phase using 3:1 pentane:EtOAc to 1:1 pentane:EtOAc; followed by  $\text{Al}_2\text{O}_3$  stationary phase using 8:1 hexane:EtOAc to 100% EtOAc) followed by washing of the obtained solid with about 5 mL of isopropanol gave Bedaquiline-pentynoxy-eg<sub>4</sub>-BODIPY **C2** (Figure 3, 17.9 mg, 17.0  $\mu\text{mol}$ , 15%) as a bright orange solid.



**Figure 3:** Structural formula of Bedaquiline-pentynoxy-eg<sub>4</sub>-BODIPY **C2**. The numbering scheme used in this work for elucidation of NMR spectroscopic data is shown.

TLC (3:1 EtOAc:cyclohexane):  $R_f = 0.22$ , red spot, green fluorescence.  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.86 (s, 1H), 8.54 (d, 1H,  $^3J(\text{H},\text{H}) = 9.0$  Hz), 8.31 (br. s, 1H, -OH), 7.81 (m, 3H), 7.67 (d, 1H,  $^3J(\text{H},\text{H}) = 8.5$  Hz), 7.58 (d, 1H,  $^3J(\text{H},\text{H}) = 8.0$  Hz), 7.54 (t, 1H,  $^3J(\text{H},\text{H}) = 7.75$  Hz), 7.50 (d, 1H,  $^3J(\text{H},\text{H}) = 8.5$  Hz), 7.42 (t, 1H,  $^3J(\text{H},\text{H}) = 7.5$  Hz), 7.23 (t, 1H,  $^3J(\text{H},\text{H}) = 7.75$  Hz), 7.08 (d, 2H,  $^3J(\text{H},\text{H}) = 9.0$  Hz, phenol CH), 6.95 (br. s, 2H), 6.94 (d, 2H,  $^3J(\text{H},\text{H}) = 8.5$  Hz, phenol CH), 6.81 (t, 3H,  $^3J(\text{H},\text{H}) = 3$  Hz), 5.98 (s, 2H, pyrrole CH), 5.82 (s, 1H,  $\text{CH}^5$ ), 4.13 (s, 3H, -OMe), 4.10 (t, 2H,  $^3J(\text{H},\text{H}) = 4.6$  Hz), 3.83 (t, 2H,  $^3J(\text{H},\text{H}) = 5$  Hz,  $-\text{CH}_2-$ ), 3.7 – 3.5 (m, 14 H,  $-\text{CH}_2-$ ), 2.49 (s, 6H, pyrrole  $\text{CH}_3$ ), 2.5 (m, 2H,  $-\text{CH}^2-$ ), 2.5 (m, 1H, diastereotopic  $-\text{CH}^3$  protons), 2.05 (t, 1H,  $^3J(\text{H},\text{H}) = 12.5$  Hz, diastereotopic  $-\text{CH}^3$  protons), 1.9 (m, 2H,  $-\text{CH}^2-$ ), 1.90 (s, 6H,  $-\text{NMe}_2$ ), 1.9 – 1.8 (m, 2H,  $-\text{CH}^2-$ ), 1.34 (s, 6H, pyrrole  $\text{CH}_3$ ) ppm.  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ ):  $\delta$  161.6, 159.5, 155.3, 144.5, 143.3, 142.0, 141.9, 140.8, 139.4, 134.8, 132.0, 131.1, 130.1, 130.0, 129.3 (phenol CH), 128.3, 127.5, 127.3, 127.2, 126.9, 125.5,

125.4, 125.4, 124.6, 121.2 (pyrrole CH), 119.3, 115.3 (phenol CH), 89.5, 82.8, 81.1, 71.0, 70.8, 70.5, 70.0, 69.9, 67.6, 56.5 ( $\text{C}^2$ ), 54.3 (OMe), 49.6 ( $\text{C}^5$ ), 44.9 ( $\text{C}^1$ ), 33.5 ( $\text{C}^3$ ), 29.0 ( $\text{C}^6$ ), 16.4 ( $\text{C}^4$ ), 14.8 (pyrrole  $\text{CH}_3$ ), 14.7 (pyrrole  $\text{CH}_3$ ) ppm.  $^{19}\text{F}$  NMR (235.4 MHz,  $\text{CDCl}_3$ , 293 K):  $\delta$  -146.3 (q,  $^1J(\text{F},\text{B}) = 32.9$  Hz,  $\text{BF}_2$ ) ppm. HRMS (ESI): Calc  $[\text{M}+\text{H}]^+$ : 1057.5457,  $[\text{M}+\text{H}+\text{Na}]^{2+}$ : 540.2675,  $[\text{M}+2\text{H}]^{2+}$ : 529.2765,  $[\text{M}+\text{H}-\text{F}]^{2+}$ : 519.2734 m/z. Found resp. 1057.5426, 540.2656, 529.2739, 519.2715 m/z. IR:  $\nu$  2922.0, 2860.2, 2358.8, 2331.8, 1606.6, 1543.0, 1508.2, 1456.2, 1438.8, 1400.2, 1305.7, 1245.9, 1193.9, 1155.3, 1116.7, 1105.1, 1080.1, 1062.7, 1051.1, 1012.6, 975.9, 910.3, 835.1, 729.0, 704.0, 640.3, 603.7, 582.5, 455.2  $\text{cm}^{-1}$  MP: 165–170  $^\circ\text{C}$  (decomposition).

## Results and discussion


### Synthesis of Fluorescent Alkynyl-BODIPY Labels

Considering the presence of an aryl bromide moiety in the molecular structure of Bedaquiline, we arrived at a labeling strategy using Sonogashira cross-coupling chemistry to form a carbon-carbon bond between the bromoquinoline position and an alkyne-functionalized fluorophore.<sup>[15]</sup> In principle, labeling of Bedaquiline on the 6-position of the quinoline ring is expected to keep binding affinities towards the microbial target protein intact, as modification of the 6-position of the quinoline moiety has been previously applied for covalent linkage to a Biacore chip and an affinity resin.<sup>[6c]</sup> Furthermore, the bromoquinoline moiety has been shown to project away from the central binding pocket in the c-subunit of ATP synthase, with the bromo-moiety located at the distal position.<sup>[14]</sup> Notwithstanding the apparent feasibility of this approach, it is obvious that close proximity of the fluorophore will lead to undesired fluorophore-protein interactions. Thus, we chose to investigate the synthesis of two alkyne-functionalized BODIPY<sup>[13]</sup> fluorophores, which were subsequently used to prepare two fluorescently labeled Bedaquiline analogues having a shorter ( $\sim 6$  Å) or longer ( $\sim 22$  Å) linear spacer between drug and fluorophore.

Fluorescent alkynyl-BODIPY derivatives **B1** and **B2** were prepared using a two-step route (see Figure 4) starting from the Mitsunobu aryl ether synthesis of aldehydes **A1** and **A2**, where the phenolic oxygen of 4-hydroxybenzaldehyde is alkylated by a butynyloxy or pentynyloxy-tetraethyleneglycol terminal alkyne, respectively. Subsequent acid-catalyzed condensation of the benzaldehyde with 2,4-dimethylpyrrole followed by oxidation and boron trifluoride chelation allowed us to obtain the target fluorophores, albeit in rather poor overall yields of 5% (**B1**) and 18% (**B2**) due to the typically poor yields of BODIPY preparations,<sup>[13]</sup> and the difficulties encountered in separating the aryl ethers from the byproducts of the Mitsunobu reaction.

ARTICLE


Journal Name



**Figure 4: Synthesis of alkyne-functionalized BODIPY derivatives.** Reaction conditions: i. 4-hydroxybenzaldehyde (1 eq.), alcohol R<sup>1</sup>-OH or R<sup>2</sup>-OH (1 eq.), PPh<sub>3</sub> (1.05 eq.), DEAD (40 wt% in toluene, 1 eq.), THF, 0 °C → rt, overnight; ii. A (1 eq.), 2,4-dimethylpyrrole (2.2 eq.), TFA (cat.), THF, rt, 16h; then DDQ 1 eq.), rt, 1h; then NEt<sub>3</sub> (excess), BF<sub>3</sub>·OEt<sub>2</sub> (excess), 0 °C → rt, 2h.

### Sonogashira-based Labeling of Bedaquiline


Initially, we ascertained the synthetic feasibility of our labeling strategy by reacting 3-butyn-1-ol with 6-bromoquinoline as model for Bedaquiline under typical<sup>[18]</sup> Sonogashira conditions (1.2:1.0 Cu:Pd stoichiometry), affording the desired coupling product in 77% isolated yield. We found that a direct translation of these conditions to the reaction of our BODIPY labels with Bedaquiline led to the formation of the labeled drug, contaminated with a significant amount of the homo-coupled diyne as a side product. We speculate that the presence of the Bedaquiline moiety is to blame for the appearance of this side product through the action of its Lewis basic sites, which can serve as a competitive bidentate ligand to one of the catalytically active metal species.<sup>[19]</sup> However, encouraged by the presence of the target molecule, we then attempted to prepare the labeled Bedaquiline analogues using relatively little Cu<sup>I</sup> catalyst with respect to Pd<sup>0</sup> (1:4 Cu:Pd stoichiometry); considering that the homocoupled bisalkyne is likely formed via the copper-catalysed Glaser coupling mechanism,<sup>[20]</sup> lowering the relative copper concentration should suppress this side reaction.<sup>[21]</sup> Under these conditions, a satisfactory conversion to the target compound was observed. Repeated purification was required to remove any traces of the unreacted parent drug, which accounts for the poor yields of, respectively, 37% and 15% for **C1** and **C2** (see Figure 5).



**Figure 5: Labeling of Bedaquiline using a Sonogashira Coupling.** Reaction conditions: iii. Bedaquiline fumarate 1 eq, BODIPY 1.5 eq, CuI 0.05 eq, Pd(PPh<sub>3</sub>)<sub>4</sub> 0.2 eq in 10 mL 1:1 DMF:NEt<sub>3</sub> added under Argon atmosphere at rt; reaction heated at 90 °C for 2h.

### Photophysical Properties of Fluorescent Bedaquiline Analogues

Using steady-state light absorption and fluorescence measurements, acetonitrile solutions of compounds **C1** and **C2** were characterized photophysically (see Figure 6). In the chosen solvent, varying distance between label and drug appears to have no influence on the properties under study, as virtually identical wavelength maxima  $\lambda_{\text{abs}}$  and  $\lambda_{\text{em}}$  were observed, with a Stokes shift of 10 nm for both compounds. The quantum yield of fluorescence is also independent of linker identity, showing the expected isolation of the fluorophore from linker and drug. The absorption and emission wavelength maxima of **C1** and **C2** are essentially identical in methanol, a protic solvent of similar polarity to acetonitrile (see Supporting Information). We also confirmed that the parent compound Bedaquiline did not show emission peaks in the 400 – 600 nm wavelength range (not shown in Figure 6). As is typical for BODIPY derivatives<sup>[13]</sup> no changes in the absorption spectrum were seen when a cuvette of **C1** was left standing under ambient illumination for several weeks; nor could we over time observe any degradation in the <sup>1</sup>H NMR spectrum of **C1**. The labeled Bedaquiline derivatives described can thus be regarded as stable, suitable fluorescent probes for biological applications.




**Figure 6: Emission and Absorption curves for compounds C1 and C2.** Normalized absorption (solid lines) and emission (dashed lines) plots, recorded in acetonitrile solution. The inset tabulates absorption and emission wavelength maxima  $\lambda_{\text{abs}}$  and  $\lambda_{\text{em}}$  and the corresponding quantum yield of fluorescence  $\Phi$ , calculated using fluorescein in 0.1 M aqueous NaOH as a reference. Peak normalized instrument responses for absorption measurements correspond to an extinction coefficient of respectively  $\epsilon = 7.6 \cdot 10^3$  (**C1**) or  $6.8 \cdot 10^3$  (**C2**) cm<sup>2</sup> mol<sup>-1</sup>.

### BODIPY-Labeled Bedaquiline Compounds inhibit ATP synthesis

After successful preparation of the BODIPY-Bedaquiline compounds, we sought to ascertain their antibacterial action, and to investigate if their mode of action is sufficiently similar to the parent compound; only sufficiently parent-like probe molecules are expected to yield useful results in follow-up studies on the mode of action of diarylquinoline antibiotics. To do so, the ability of **C1** and **C2** to inhibit the function of the biological target of the parent drug, ATP synthase, was evaluated. We measured the impact of the compounds on ATP synthesis activity using inverted membrane vesicles derived from *Mycobacterium smegmatis*. As depicted in

Figure 7, both **C1** and **C2** blocked the ATP synthesis activity in a dose-dependent manner. However, inhibition was clearly more efficient in case of **C2** ( $IC_{50} \sim 2.0 \mu M$ ). The lower activity of **C1** ( $IC_{50} > 10 \mu M$ ) may be due to the shorter linker, interfering with effective binding of the compound to its binding pocket, which is located close to the center of the hydrophobic, membrane-spanning part of ATP synthase.<sup>[16]</sup> These  $IC_{50}$  values are higher than those reported for the parent compound ( $IC_{50}$  between 10 - 100 nM),<sup>[7, 24]</sup> but comparable to and consistent with values reported for fluorescent derivatives of other anti-tuberculosis drugs, e.g. for benzothiadiazinones.<sup>[13]</sup> The results conclusively prove that the presented labeling strategy allows the preparation of Bedaquiline analogues that retain the ability to inhibit ATP synthesis, given the correct spacing distance between label and drug.



**Figure 7: Inhibition of mycobacterial ATP synthesis activity by BODIPY-Bedaquiline (BDQ) compounds.** ATP synthesis activity was measured in inverted membrane vesicles from *M. smegmatis* using the glucose-6-phosphate dehydrogenase method. BODIPY-Bedaquiline compounds (open bars: C1, butynyloxy linker, closed bars: C2, pentynyloxy-eg<sub>4</sub> linker) were added at the indicated concentrations. 100 nM BDQ and the ATP synthase inhibitor DCCD were used as control.

#### BODIPY-Labeled Bedaquiline Compounds Inhibit Bacterial Growth

Next, we evaluated their ability to interfere with growth of living bacteria. As shown in Table 1, both **C1** and **C2** suppressed growth of *M. tuberculosis*, with Minimal Inhibitory Concentrations of 1.8  $\mu M$  and 7.3  $\mu M$ , respectively (Table 1), compared to 0.1  $\mu M$  for the parent compound Bedaquiline.<sup>[6a]</sup> This indicates that labeling of Bedaquiline with the BODIPY moiety and introduction of the linker decreased, but did not abolish, the antibacterial activity of the drug. In *M. tuberculosis*, efflux pumps can be an important determinant for the sensitivity towards antibacterials.<sup>[17]</sup> Recently it was reported that the efflux pump inhibitor verapamil enhanced the susceptibility of *M. tuberculosis* for Bedaquiline<sup>[17d]</sup> whereas upregulation of the mycobacterial efflux pump MmpS5-MmpL5 decreased the sensitivity for this drug.<sup>[14,17e]</sup> Therefore we evaluated the metabolic similarity of the labeled compounds to Bedaquiline by checking if they are also recognized by the efflux system. Addition of the efflux pump inhibitor verapamil increased the sensitivity of *M. tuberculosis* reference strains H37Rv for **C1** and **C2** (respectively twofold and eightfold, Table 1), suggesting that both BODIPY-Bedaquiline analogues are indeed transported by an

efflux pump. We also tested growth inhibition of two *M. tuberculosis* strains with elevated levels of the MmpS5-MmpL5 efflux pump: one strain carrying a mutation in the regulator of MmpS5-MmpL5 (CV37) and one laboratory strain carrying a plasmid for over-expression of MmpS5-MmpL5 (H37Rv pCVGA30). Both efflux pump overexpressing strains displayed considerably lower sensitivity towards **C1** and **C2**, as well as to the parent compound (Table 1), suggesting that the fluorescent analogues are substrates of the same efflux system as Bedaquiline.

**Table 1: Growth inhibition of *M. tuberculosis* by BODIPY-Bedaquelines.** Strains tested were a *M. tuberculosis* reference strain (H37Rv) an *in vitro* generated Rv0678 mutant with increased efflux pump activity against BDQ (CV37), the reference strain with over-expressed efflux pumps (H37Rv pCVGA30) and with empty overexpression plasmid as control (H37Rv pSD5). VER: efflux pump inhibitor verapamil. BDQ = Bedaquiline. Minimal inhibitory concentrations (MICs) were measured using the rezurin method as in reference [14] and are expressed in mol L<sup>-1</sup>.

| Strain        | BDQ                  | BDQ + VER             | C1                    | C1 + VER              | C2                    | C2 + VER              |
|---------------|----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| H37Rv         | 1.1 10 <sup>-7</sup> | 7.0 10 <sup>-9</sup>  | 1.8 10 <sup>-6</sup>  | 9.1 10 <sup>-7</sup>  | 7.3 10 <sup>-6</sup>  | 9.1 10 <sup>-7</sup>  |
| CV37          | 9.1 10 <sup>-7</sup> | 2.3 10 <sup>-7</sup>  | >2.9 10 <sup>-5</sup> | >2.9 10 <sup>-5</sup> | >2.9 10 <sup>-5</sup> | >2.9 10 <sup>-5</sup> |
| H37Rv pCVGA30 | 4.5 10 <sup>-7</sup> | <5.7 10 <sup>-8</sup> | 7.3 10 <sup>-6</sup>  | 9.1 10 <sup>-7</sup>  | >2.9 10 <sup>-5</sup> | >2.9 10 <sup>-5</sup> |
| H37Rv pSD5    | 1.1 10 <sup>-7</sup> | 7.0 10 <sup>-9</sup>  | 1.8 10 <sup>-6</sup>  | 4.5 10 <sup>-7</sup>  | 3.6 10 <sup>-6</sup>  | 9.1 10 <sup>-7</sup>  |

#### Conclusions

The obtained results show that we successfully designed a labeling strategy for Bedaquiline that generates compounds sufficiently similar to the parent drug to be of use in scientific research. The obtained fluorescent analogues **C1** and **C2** can successfully interfere with ATP synthesis, the biological target of Bedaquiline. Growth of living *M. tuberculosis* bacteria is suppressed at approximately 20- or 70-fold higher Minimum Inhibitory Concentrations compared to Bedaquiline, dependent on the spacing (6 Å versus 22 Å, respectively) between fluorescent label and drug. The probe compounds also share the affinity for the MmpS5-MmpL5 drug efflux system with the parent compound. Thus, fluorescent antibiotics prepared using the method described in this work are promising tools for deeper fluorescence-based experimentation on diarylquinoline-based tuberculosis drugs. Applications may include investigation of drug in/efflux at the level of either the bacterial population or the individual mycobacterial cell, determination of sub-cellular drug or target localization in the bacterium, as well as biochemical studies on binding of the drug to its target.

#### Acknowledgements


P.L. is indebted to a fellowship from the Chinese Scholarship council. J.A.R. was supported by the BioSolar Cells Programme of the Dutch Ministry of Economic Affairs. D.B. and J.C.S. thank AIMMS (Vrije Universiteit Amsterdam) for a stimulus grant (2011.A503). The authors wish to thank Peggy Janssens for technical assistance and W. Bas Wittekoek and Yvon van der Hulst for preliminary experiments.



## Notes and references

- World Health Organization, 2015, "Global tuberculosis report 2015", *World Health Organization*.
- A. Koul, E. Arnoult, N. Lounis, J. Guillemont, K. Andries, *Nature*, 2011, **469**, 483-490.
- D. Bald and A. Koul, *FEMS Microbiol. Lett.*, 2010, **308**, 1-7; J.G. Hurdle, A.J. O'Neill, I. Chopra, R.E.E. Lee, *Nat. Rev. Microbiol.*, 2011, **9**, 62-75; P.A. Black, *Antimicrob. Agents Chemother.*, 2014, **58**, 2491-2503; P. Lu, H. Lill and D. Bald, *Biochim. Biophys. Acta*, 2014, **1837**, 1208-1218; G.M. Cook, K. Hards, C. Vilcheze, T. Hartman and M. Berney, *Microbiol. Spectr.*, 2014, **2**.
- H.I. Boshoff, T.G. Myers, B.R. Copp, M.R. McNeil, M.A. Wilson, C.E. Barry, *J. Biol. Chem.*, 2004, **279**, 40174-40184; E.A. Weinstein, T. Yano, L.S. Li, D. Avarbock, A. Avarbock, D. Helm, A.A. McCollm, K. Duncan, J.T. Lonsdale, H. Rubin, *Proc. Natl. Acad. Sci.*, 2005, **102**, 4548-4553; T. Yano, S. Kassovska-Bratinova, J.S. Teh, J. Winkler, K. Sullivan, A. Isaacs, N. M. Schechter, H. Rubin, *Biol. Chem.*, 2011, **286**, 10276-10287.
- K.A. Abrahams, J.A.G. Cox, V.L. Spivey, N.J. Loman, M.J. Pallen, C. Constantinidou, R. Fernández, C. Alemparte, M.J. Remuñán, D. Barros, L. Ballell, G.S. Besra, *PLoS. One.*, 2012, **7**, e52951; K. Pethe, P. Bifani, J. Jang, S. Kang, S. Park, S. Ahn, J. Jiricek, J. Jung, H. Kyoung Jeon, J. Cechetto, T. Christophe, H. Lee, M. Kempf, M. Jackson, A.J. Lenaerts, H. Pham, V. Jones, M.J. Seo, Y.M. Kim, M. Seo, J.J. Seo, D. Park, Y. Ko, I. Choi, R. Kim, S.Y. Kim, S. Lim, S. Yim, J. Nam, H. Kang, H. Kwon, C. Oh, Y. Cho, Y. Jang, J. Kim, A. Chua, B. Tan, M.B. Nanjundappa, S.P.S. Rao, W.S. Barnes, R. Wintjens, J.R. Walker, S. Alonso, S. Lee, J. Kim, Soohyun Oh, T. Oh, U. Nehrbass, S. Han, Z. No, J. Lee, P. Brodin, S. Cho, K. Nam, J. Kim, *Nat. Med.*, 2013, **19**, 1157-1160.
- K. Andries, P. Verhasselt, J. Guillemont, H.W. Göhlmann, J.M. Neefs, H. Winkler, J. Van Gestel, P. Timmerman, M. Zhu, E. Lee, P. Williams, D. de Chaffoy, E. Huitric, S. Hoffner, E. Cambau, C. Truffot-Pernot, N. Lounis, V. Jarlier, *Science*, 2005, **307**, 223-227; A.C. Haagsma, R. Abdillahi-Ibrahim, M.J. Wagner, K. Krab, K. Vergauwen, J. Guillemont, K. Andries, H. Lill, A. Koul, D. Bald, *Antimicrob. Agents Chemother.*, 2009, **53**, 1290-1292; A.C. Haagsma, I. Podasca, A. Koul, K. Andries, J. Guillemont, H. Lill, D. Bald, *PLoS. One.*, 2011, **6**, e23575; W. Balemans, L. Vranckx, N. Lounis, O. Pop, J. Guillemont, K. Vergauwen, S. Mol, R. Gilissen, M. Motte, D. Lançois, M. De Bolle, K. Bonroy, H. Lill, K. Andries, D. Bald and A. Koul, *Antimicrob Agents Chemother.*, 2012, **56**, 4131-4139.
- A. Koul, N. Dendouga, K. Vergauwen, B. Molenberghs, L. Vranckx, R. Willebrords, Z. Ristic, H. Lill, I. Dorange, J. Guillemont, D. Bald, K. Andries, *Nat. Chem. Biol.*, 2007, **3**, 323-324;
- A. Koul, L. Vranckx, N. Dhar, H.W. Göhlmann, E. Özdemir, J.M. Neefs, M. Schulz, P. Lu, E. Mørtz, J.D. McKinney, K. Andries, D. Bald, *Nat Commun.*, 2014, **5**, 3369; J. Dhillion, K. Andries, P.P. Phillips, D.A. Mitchison, *Tuberculosis*, 2010, **5**, 301-305.
- A.H. Diacon, A. Pym, M. Grobusch, *New Engl. J. Med.*, 2009, **360**, 2397-2405; A.H. Diacon, *Am. J. Respir. Crit. Care. Med.*, 2015, **191**, 943-953; S. Tyagi, N.C. Ammerman, S.Y. Li, J. Adamson, P.J. Converse, R.V. Swanson, D.V. Almeida, J.H. Grosset, *Proc. Natl. Acad. Sci.* 2015, **112**, 869-374.
- J. Neres, F. Pojer, E. Molteni, L.R. Chiarelli, N. Dhar, S. Boy-Röttger, S. Buroni, E. Fullam, G. Degiacomi, A.P. Lucarelli, R.J. Read, G. Zanoni, D.E. Edmondson, E. De Rossi, M.R. Pasca, J.D. McKinney, P.J. Dyson, G. Riccardi, A. Mattevi, S.T. Cole and C. Binda, *Sci. Transl. Med.*, 2012, **150**, 150ra121.
- A. Loudet, K. Burgess, *Chem. Rev.*, 2007, **107**, 4891-4932.
- P. Lu, M.H. Heineke, A. Koul, K. Andries, G.M. Cook, H. Lill, R. van Spanning, D. Bald, *Sci. Rep.*, 2015, **5**, 10333.
- P. Lu P, A.C. Haagsma, H. Pham, J.J. Maaskant, S. Mol, H. Lill, D. Bald, *Antimicrob. Agents Chemother.*, 2011, **55**, 5354-5357.
- K. Andries, C. Villellas, N. Coeck, K. Thys, T. Gevers, L. Vranckx, N. Lounis, B.C. de Jong, A. Koul, *PLoS One.*, 2014, **9**, e102135.
- R. Chinchilla, C. Najera, *Chem. Rev.*, 2007, **107**, 874-922.
- L. Preiss et al., *Science advances*, 2015, **1**, e1500106; S. Singh, K.K. Roy, S.R. Khan, V.K. Kashyap, A. Sharma, S. Jaiswal, S.K. Sharma, M.Y. Krishnan, V. Chaturvedi, J. Lal, S. Sinha, A. Dasgupta, R. Srivastava, A.K. Saxena, *Bioorg. Med. Chem.*, 2015, **4**, 742-752.
- M. Viveiros, M. Martins, L. Rodrigues, D. Machado, I. Couto, J. Ainsa, L. Amaral, *Expert Rev Anti Infect Ther.*, 2012, **9**, 983-998; K.N. Adams, J.D. Szumowski, L. Ramakrishnan, *J Infect. Dis.*, 2014, **3**, 456-466; T. Coelho, D. Machado, I. Couto, R. Maschmann, D. Ramos, A. von Groll, M.L. Rossetti, P.A. Silva, M. Viveiros, *Front Microbiol.*, 2015, **6**, 330; S. Gupta, K.A. Cohen, K. Winglee, M. Maiga, B. Diarra, W.R. Bishai, *Antimicrob. Agents Chemother.*, 2014, **58**, 574-576; R.C. Hartkoorn, S. Uplekar, S.T. Cole, *Antimicrob. Agents Chemother.*, 2014, **58**, 2979-2981.
- Q. Zhang, H. Ren, G.L. Baker, *J. Org. Chem.*, 2014, **79**, 9546-9555.
- M. Meyer, S. Kuntz, I. Grillier-Vuissoz, H. Martin, L. Richert, S. Flament, Y. Chapleur, M. Boisbrun, *Letters in Drug Design and Discovery*, 2014, **11**, 256-264.
- L. Guandalini, E. Martini, S. Dei, D. Manetti, S. Scapecchi, E. Teodori, M. Romanelli, K. Varani, G. Greco, L. Spadola, E. Novellino, *Bioorg. Med. Chem.*, 2005, **13**, 799-807.
- G. Chen, W. Gong, Z. Zhuang, M.S. Andra, Y. Chen, X. Hong, Y. Yang, T. Liu, K.N. Houk, J. Yu, *Science*, 2016, **6303**, 1023-1027.
- I.D. Campbell, G. Eglinton, *Org. Synth.*, 1965, **45**, 39.
- M. Karak, L.C.A. Barbosa, G.C. Hargaden, *RSC Adv.*, 2014, **4**, 53442-53466.
- Bedaquiline showed approx. 60% inhibition of ATP synthesis under our conditions at 100 nM, suggesting a 25-fold difference in activity between Bedaquiline and **C2**.

Table of Contents



Labeling the tuberculosis drug Bedaquiline with a fluorescent alkynyl-BODIPY moiety does not disrupt its antibacterial properties