

pubs.acs.org/joc

Bioinspired Total Synthesis and Human Proteasome Inhibitory Activity of (-)-Salinosporamide A, (-)-Homosalinosporamide A, and Derivatives Obtained via Organonucleophile Promoted Bis-cyclizations

Henry Nguyen,[†] Gil Ma,^{†,§} Tatiana Gladysheva,[‡] Trisha Fremgen,[‡] and Daniel Romo^{*,†}

[†]Department of Chemistry, Texas A&M University, P.O. Box 30012, College Station, Texas 77842-3012, United States, and [‡]Genzyme Corporation, Drug and Biomaterial R & D, 153 Second Avenue, Waltham, Massachusetts 02451, United States. [§]Current address: Lundbeck Research USA, Inc., Paramus, NJ 07652.

romo@tamu.edu

Received August 25, 2010



A full account of concise, enantioselective syntheses of the anticancer agent (-)-salinosporamide A and derivatives, including (-)-homosalinosporamide, that was inspired by biosynthetic considerations is described. The brevity of the synthetic strategy stems from a key bis-cyclization of a β -keto tertiary amide, which retains optical purity enabled by A^{1,3}-strain rendering slow epimerization relative to the rate of bis-cyclization. Optimization studies of the key bis-cyclization, enabled through byproduct isolation and characterization, are described that ultimately allowed for a gram scale synthesis of a versatile bicyclic core structure with a high degree of stereoretention. An optimized procedure for zincate generation by the method of Knochel, generally useful for the synthesis of salino A derivatives, led to dramatic improvements in side-chain attachment and a novel diastereomer of salino A. The versatility of the described strategy is demonstrated by the synthesis of designed derivatives including (-)-homosalinosporamide A. Inhibition of the human 20S and 26S proteasome by these derivatives using an enzymatic assay are also reported. The described total synthesis of salino A raises interesting questions regarding how biosynthetic enzymes leading to the salinosporamides proceeding via optically active β -keto secondary amides, are able to maintain the stereochemical integrity at the labile C2 stereocenter or if a dynamic kinetic resolution is operative.

Introduction

The development of practical, scaleable total syntheses of complex, bioactive natural products has recently become of

2 J. Org. Chem. 2011, 76, 2–12

great interest due to a desire to further explore biological activity or to facilitate development of structure-activity relationship profiles. Inevitably, one of the best ways to achieve such a practical synthesis involves consideration of

Published on Web 11/03/2010



FIGURE 1. Structures of naturally occurring bicyclic β -lactone proteasome inhibitors.

potential biosynthetic strategies utilized by the producing organism to assemble such stereochemically complex structures.1

The potential of human 20S proteasome inhibitors continues to be of interest for cancer chemotherapy and the recent FDA approval of bortezomib (Velcade) validates the proteasome as a target for cancer chemotherapy.² Isolated by Fenical and co-workers³ from the marine actinomycete, Salinispora tropica, salinosporamides A and B (1a.b, salino A and B) are unique bicyclo [3.2.0] β -lactone-containing natural products (Figure 1). Phase I human clinical studies for multiple myeloma followed the finding that salino A exhibited great potential in mouse models toward several cancers when administered intravenously despite the potentially labile β -lactone.⁴ The terrestrial metabolites, cinnabaramides

(2) (a) Voorhees, P. M.; Dees, E. C.; O'Neil, B.; Orlowski, R. Z. Clin. Cancer Res. **2003**, *9*, 6316. (b) Rajkumar, S. V.; Richardson, P. G.; Hideshima, T.; Anderson, K. C. J. Clin. Oncol. **2005**, *23*, 630. (c) Joazeiro, C. A. P.; Anderson, K. C.; Hunter, T. Cancer Res. 2006, 66, 7840.

(3) Feling, R. H.; Buchanan, G. O.; Mincer, T. J.; Kauffman, C. A.; Jensen, P. R.; Fenical, W. F. Angew. Chem., Int. Ed. 2003, 42, 355.

(4) For a review describing the development of salinosporamide A as an anticancer agent, see: Fenical, W.; Jensen, P. R.; Palladino, M. A.; Lam, K. S.; Lloyd, G. K.; Potts, B. C. *Bioorg. Med. Chem.* **2009**, *17*, 2175.

(5) Stadler, M.; Bitzer, J.; Mayer-Bartschmid, A.; Muller, H.; Benet-Buchholz, J.; Gantner, F.; Tichy, H.-V.; Reinemer, P.; Bacon, K. B. J. Nat. Prod. 2007, 70, 246.

(6) Omura, S.; Fujimoto, T.; Otoguro, K.; Matsuzaki, K.; Moriguchi, R.;

Tanaka, H.; Sasaki, Y. J. Antibiot. 1991, 44, 117.
 (7) Manam, R. R.; Macherla, V. R.; Tsueng, G.; Dring, C. W.; Weiss, J.;
 Neuteboom, S. T. C.; Lam, K. S.; Potts, B. C. J. Nat. Prod. 2009, 72, 295.

(9) (a) Caubert, V.; Masse, J.; Retailleau, P.; Langlois, N. Tetrahedron *Lett.* 2007, 48, 381. (b) Villanueva, M. I.; Rupnicki, L.; Lam, H. W. *Tetrahedron* 2008, 64, 7896. (c) Momose, T.; Kaiya, Y.; Hasegawa, J.; Sato, T.; Chida, N. Synthesis 2009, 17, 2983. (d) Struble, J. R.; Bode, J. W. Tetrahedron 2009, 65, 4957. (e) Mosey, R. A.; Tepe, J. J. Tetrahedron Lett. 2009, 50, 295. (f) Ling, T.; Potts, B. C.; Macherla, V. R. J. Org. Chem. 2010, 75, 3882.

OCFeatured Article



FIGURE 2. Enantioselective strategy to salino A and derivatives.

(e.g., A, 2),⁵ lactacystin- β -lactone (omuralide, 3a) derived from the prodrug lactacystin (4),⁶ and the recently isolated mixed congener antiprotealide (3b),⁷ bear close structural similarities to salino A and congeners, which have been targets for total⁸ and formal⁹ syntheses,¹⁰ structure–activity studies,¹¹ biosyn-thetic engineering,¹² and crystallographic studies with the 20S proteasome.13

Our synthetic strategy to salino A and derivatives utilizes a nucleophile-catalyzed aldol-lactonization (NCAL) process, a reaction we previously developed for the synthesis of carbocycle-fused β -lactones (Figure 1).¹⁴ Indeed, salino A and related natural products were the initial inspiration for the development of the NCAL methodology in our group (Figure 2). In particular, the study of an intramolecular aldol lactonization was inspired by biosynthetic considerations of

⁽¹⁾ Sorensen, E. J.; Theodorakis, E. Tetrahedron 2006, 62, 5159 and papers following this Preface to the Tetrahedron Symposia-In-Print "Nature-Inspired Approaches to Chemical Synthesis

^{(8) (}a) Reddy, L. R.; Saravanan, P.; Corey, E. J. J. Am. Chem. Soc. 2004, 126, 6230. (b) Reddy, L. R.; Fournier, J.-F.; Reddy, B. V. S.; Corey, E. J. Org. Lett. 2005, 7, 2699. (c) Endo, A.; Danishefsky, S. J. J. Am. Chem. Soc. 2005, 127, 8298. (d) Mulholland, N. P.; Pattenden, G.; Walters, I. A. S. Org. *Biomol. Chem.* **2006**, *4*, 2845. (e) Ma, G.; Nguyen, H.; Romo, D. *Org. Lett.* **2007**, *9*, 2143. (f) Ling, T.; Macheria, V. R.; Manam, R. R.; McArthur, K. A.; Ports, B. C. M. Org. Lett. 2007, 9, 2289. (g) Mulholland, N. P.; Pattenden, G.; Walters, I. A. S. Org. Biomol. Chem. 2008, 6, 2782. (h) Takahashi, K.; Midori, M.; Kawano, K.; Ishihara, J.; Hatakeyama, S. Angew. Chem., Int. Ed. 2008, 47, 6244. (i) Fukuda, T.; Sugiyama, K.; Arima, S.; Harigaya, Y.; Nagamitsu, T.; Omura, S. Org. Lett. 2008, 10, 4239. (j) Nguyen, H.; Ma, G.; Romo, D. Chem. Commun. 2010, 46, 4803.

⁽¹⁰⁾ For a recent review on syntheses of lactacystin and salino A, see: Shibasaki, M.; Kanai, M.; Fukuda, N. Chem. Asian J. 2007, 2, 20.

^{(11) (}a) Macherla, V. R.; Mitchell, S. S.; Rama Rao Manam, R. R.; Reed, K. A.; Chao, T.-H.; Nicholson, B.; Deyanat-Yazdi, G.; Mai, B.; Jensen, P. R.; Fenical, W. F.; Neuteboom, S. T. C.; Lam, K. S.; Palladino, M. A.; Potts, B. M. J. Med. Chem. 2005, 48, 3864. (b) Hogan, P. C.; Corey, E. J. J. Am. Chem. Soc. 2005, 127, 15386.

^{(12) (}a) Beer, L. L.; Moore, B. S. Org. Lett. 2007, 9, 845. (b) Eustaquio, A. S.; Moore, B. S. Angew. Chem., Int. Ed. 2008, 47, 3936. (c) Nett, M.; Gulder; Tobias, A. M.; Kale; Andrew, J.; Hughes, C. C.; Moore, B. S. J. Med. Chem. 2009, 52, 6163. (d) Liu, Y.; Hazzard, C.; Eustaquio, A. S.; Reynolds, K. A.; Moore, B. S. J. Am. Chem. Soc. 2009, 131, 10376. (e) Eusthquio, A. S.; McGlinchey, R. P.; Liu, Y.; Hazzard, C.; Beer, L. L.; Florova, G.; Alhamadsheh, M. M.; Lechner, A.; Kale, A. J.; Kobayashi, Y.; Reynolds, K. A.; Moore, B. S. Proc. Nat. Acad. Sci. U.S.A. 2009, 106, 12295.

^{(13) (}a) Groll, M.; Huber, R.; Potts, B. C. M. J. Am. Chem. Soc. 2006, 128, 5136. (b) Mosey, R. A.; Tepe, J. J. Tetrahedron Lett. 2009, 50, 295. (c) Groll, M.; McArthur, K. A.; Macherla, V. R.; Manam, R. R.; Potts, B. C. J. Med. Chem. 2009, 52, 5420. (d) Manam, R.; McArthur, K. A.; Chao, T.-H.; Weiss, J.; Ali, J. A.; Palombella, V. J.; Groll, M.; Lloyd, G. K.; Palladino, M. A.; Neuteboom, S. T. C.; Macherla, V. R.; Potts, B. C. M. J. Med. Chem. 2008, 51, 6711.

^{(14) (}a) Cortez, G. S.; Tennyson, R.; Romo, D. J. Am. Chem. Soc. 2001, 123, 7945. (b) Oh, S. H.; Cortez, G. S.; Romo, D. J. Org. Chem. 2005, 70, 2835. (c) Henry-Riyad, H.; Lee, C. S.; Purohit, V. C.; Romo, D. Org. Lett. 2006, 8, 4363. (d) Purohit, V. C.; Matla, A. S.; Romo, D. J. Am. Chem. Soc. 2008, 130, 10478.

SCHEME 1. Synthesis of Ketene Homodimers 12/13 and Heterodimers 14/15



this class of proteasome inhibitors, and this general biosynthetic pathway has recently been supported by detailed biosynthetic studies.¹² We previously reported racemic syntheses of cinnabaramide A and salino A^{8e} and more recently an enantioselective synthesis of (–)-salino A and (–)-homosalino A on the basis of these strategies.^{8j} Retention of optical purity during the bis-cyclization process was made possible by the A^{1,3}-strain of β -keto tertiary amides.¹⁵ Herein, we disclose a full account of our bioinspired syntheses of salino A and derivatives including optimization of the key bis-cyclization process that simultaneously forms the γ -lactam and fused β -lactone core. Furthermore, the bioactivity of these derivatives was assayed against the chymotripsin-like, caspase-like, and trypsin-like activities of the human proteasome.

Our primary retrosynthetic disconnections, namely the biscyclization $(7 \rightarrow 6)$ and amide coupling steps $(8 + 9 \rightarrow 7)$, which were inspired by biosynthetic considerations, are key steps in our retrosynthetic analysis. The optical purity of β -lactone **6** would reflect the diastereometric purity of β -ketoamide 7 derived from acylation of serine derivatives with racemic ketene dimer 9. The C4 stereocenter in β -ketoamide 7 is lost during the bis-cyclization process due to formation of an ammonium enolate intermediate, thus the integrity of the C2 stereocenter primarily dictates the optical purity of β -lactone 6 which we proposed would be preserved by A^{1,3} strain.¹⁵ Introduction of the C4-side chain initially relied on the strategy developed by Corey,^{8a} and in this report, we describe the use of Knochel's procedure for zincate formation;¹⁶ however, the success of this process and subsequent manipulations was initially not guaranteed given the potential reactivity of the β -lactone.¹⁷

Results and Discussion

Initial Studies of the Bis-cyclization Leading to C4-Unsubstituted Bicyclic β -Lactones. We first targeted simple C4unsubstituted substrates for the bis-cyclization. Ketene homodimers 12/13 and subsequently employed (vide infra) heterodimers 14/15 were prepared by dimerization of acid chlorides 10/11 by the method of Sauer (Scheme 1).^{8e,j,18} As expected, homodimers 12/13 were obtained in moderate to TABLE 1.

Derivatives 21a-e from Homodimers 12a-e

Nguyen et al.

PMBHN、 CO₂Bn 16 THE, 50 °C 18a-e: R² = Bn Pd/C **19a-e**: R² = H 12а-е (1.5 equiv) Br 20 Ņ́⊕ n-Pr ⊖ OTf 4-PPY (1.5 equiv) R i-Pr2NEt, CH2Cl2 21а-е 21a'-e' 0 °C, 2 h % vield % vield **B**-lactone dr ' entry $(19)^{a,l}$ (**21** $)^{l}$ 1 84 93 2.2:1 21a (19a) (21a)CH₂Cy 2 80 90 21b 2.2:1 (19b) (21b) 3 72 85 2.5:1 21c (**19c**) (21c) (>19:1)^d 4 77 25 (19d) (21d) 5 81 70 4:1 (19e) (21e) CI(CH2)2 CH2)2C

^{*a*}Yield is for two steps. ^{*b*}Yields refer to isolated, purified (SiO₂) product. ^{*c*}Determined by ¹H NMR analysis of crude reaction mixtures. ^{*d*}Observed diastereomeric ratio (dr) if reaction is allowed to proceed at 25 °C for 36 h (54% yield). PMB = *p*-methoxybenzyl, 4-PPY = 4-pyrrolidinopyridine.

good yields (43-65%) following silica gel purification; however, heterodimers **14/15** provided the expected statistical mixtures, and further reductions in yields were also noted during purification leading to low absolute yields (5-13%) yield, 20-52% based on a statistical, theoretical yield of 25\%). Importantly, these dimerizations are readily run on multigram scale from inexpensive, commercially available acid chlorides, and heterodimers **14/15**, despite low absolute yields, can also be obtained in multigram quantities following silica gel purification.

With racemic ketene homodimers 12a-e in hand, coupling with *N*-PMB-glycine benzyl ester (16) by the method of Calter¹⁹ proceeded efficiently to provide keto acid substrates 19a-e following hydrogenolysis (Table 1). Bis-cyclization of these keto acids was achieved using conditions similar to those developed for carbocycles, with 4-pyrrolidinopyridine (4-PPY) as nucleophilic promoter at 0 °C and proceeded efficiently (70–93%) in most cases to give bicyclic lactones 21a-ewith moderate diastereoselectivity (dr, 2.2–4:1). Keto acid

⁽¹⁵⁾ Evans, D. A.; Ennis, M. D.; Le, T.; Mandel, N.; Mandel, G. J. Am. Chem. Soc. 1984, 106, 1154.

⁽¹⁶⁾ Ren, H.; Dunet, G.; Mayer, P.; Knochel, P. J. Am. Chem. Soc. 2007, 129, 5376.

⁽¹⁷⁾ Jacobsen and co-workers had previously demonstrated the stability of a related spiro- β -lactone in their studies toward omuralide; see: Balskus, E. P.; Jacobsen, E. N. J. Am. Chem. Soc. **2006**, 128, 6810.

^{(18) (}a) Sauer, J. C. J. Am. Chem. Soc. **1947**, 69, 2444. (b) Purohit, V. C.; Richardson, R. D.; Smith, J. W.; Romo, D. J. Org. Chem. **2006**, 71, 4549. (c) Duffy, R. J.; Morris, K. A.; Romo, D. J. Am. Chem. Soc. **2005**, 127, 16754.

⁽¹⁹⁾ Calter, M. A.; Orr, R. K.; Song, W. Org. Lett. 2003, 5, 4745.

SCHEME 2. Total Synthesis of *rac*-Cinnabaramide A $((\pm)$ -2)



19c gave high diastereoselectivity (> 19:1) but in reduced yield (54%; Table 1, entry 3); however, monitoring this reaction by ¹H NMR at 25 °C indicated that the high selectivity was due to selective degradation of the minor diastereomer by prolonged reaction times. The C2-unsubstituted keto acid **19d** gave only 25% yield (Table 1, entry 4) which may result from facile enolization due to the absence of the C2-substitutent (R). This likely leads to diminished rates of the initial aldol step. Importantly, keto acid **19e** bearing two primary chlorides thus mimicking the substrate required for the proposed salino A synthesis also proceeded efficiently (Table 1, entry 5).

Total Synthesis of rac-Cinnabaramide A (2). Encouraged by the results with C4-unsubstituted keto acid substrates, we next studied more sterically demanding substrates (i.e., bearing C4-substitution) by targeting the synthesis of cinnabaramide A (2). The synthesis commenced with reductive amination of commercially available (S)-O-benzylserine ((S)-22) with *p*-anisaldehyde (Scheme 2). Subsequent esterification provided the protected serine derivative 23a in 58% overall yield (two steps). The required unsymmetrical ketene dimer 14a, derived from heterodimerization of acetyl and octanoyl chlorides (5% yield),^{18a} was coupled with serine derivative 23a to provide diastereomeric esters 24 (dr 1:1), and a Me₃SnOH-mediated hydrolysis²⁰ provided keto acid 25. The key bis-cyclization provided diastereomeric β -lactones 26/ 27 in 45% yield with moderate diastereoselectivity (dr 3.3:1), and NOE analysis confirmed that the major diastereomer corresponded to that found in the cinnabaramides.²¹ Hydrogenolysis of the benzyl ether facilitated separation of the major alcohol diastereomer 28 (79%, dr > 19:1), and this was followed by Parikh-Doering oxidation²² to give an intermediate aldehyde, which was used directly in the next step. Applying the method developed by Corey with zinc reagent 29 gave alcohol 30 in 57% yield (two steps, dr 4.7:1). Oxidative PMB deprotection with ceric ammonium nitrate gave rac-cinnabaramide A $((\pm)-2)$, which could be isolated diastereometrically pure in 48% yield. Spectral data for the synthetic material correlated with the published data and further verification of the relative

SCHEME 3. Total Synthesis of *rac*-Salino A $((\pm)$ -1a)



stereochemistry was possible by X-ray analysis.²¹ This synthesis demonstrated that several reactions could be performed in the presence of the β -lactone including organozinc additions to aldehydes.

Total Synthesis of *rac*-Salinosporamide A ((\pm)-1a). In a highly analogous manner to that described for cinnabaramide A, the synthesis of racemic salino A was accomplished, and thus only key reactions are highlighted (Scheme 3). The required heteroketene dimer 14b derived from acetyl chloride and commercially available 4-chlorobutanoyl chloride could be obtained in multigram quantities by the method of Sauer.¹⁸ The keto amide substrate 31 was then prepared by coupling heteroketene dimer 14b and serine derivative 23b, bearing an allyl ester which aided subsequent deprotection. In this case, the unoptimized bis-cyclization proceeded to give β -lactone 32 in 25–35% yield (dr 2–3:1); however, the relative stereochemistry of the major diastereomer corresponded to

⁽²⁰⁾ Furlám, R. L. E.; Emesto, G.; Mata, E. G.; Masearetti, O. A.

Tetrahedron **1998**, *54*, 13023. (21) See the Supporting Information for experimental details.

 ⁽²²⁾ Parikh, J. P.; Doering, W. E. J. Am. Chem. Soc. 1967, 89, 5505.

SCHEME 4. Proposed Mechanism for the Bis-cyclization with Two Diastereomeric Keto Acid Substrates (a) (2R,4R)-34 from (*R*)-Serine (b) (2R,4S)-40 from (*S*)-Serine



salino A as confirmed by subsequent conversion to the natural product. Bicyclic β -lactone **32** was processed to racemic salino A (**1a**) using a sequence identical to that described for cinnabaramide A (cf. Scheme 2). Once again, it was gratifying to verify that the potentially labile β -lactone and primary chloride were compatible with several final reaction steps in the synthesis.

Optimization Studies of the Bis-cyclization. Our racemic syntheses of cinnabaramide A and salino A demonstrated the viability of our synthetic approach. However, with an ultimate goal of developing an efficient, enantioselective route to a versatile salino A core (i.e., β -lactone 32) amenable to the synthesis of derivatives, we set out to identify reaction conditions that would minimize formation of observed byproducts and most critically, epimerization of the C2 stereocenter of β -keto acid 34 during the bis-cyclization. Our working mechanism for this bis-cyclization invokes a nucleophile catalyzed aldol-lactonization process and follows from our previous studies with carbocycle-fused β -lactones (Scheme 4).¹⁴ Activation of the carboxylic acid is followed by transacylation with 4-PPY to provide acyl ammonium 36, and deprotonation provides ammonium enolate 37. Equilibration to the reactive conformer 37" enables subsequent aldol reaction leading to aldolate **38**. Lactonization, which may occur via an " S_N 2" process,^{14c,23} then delivers β -lactone **39**. While a [2 + 2] cycloaddition mechanism via an intermediate ketene has not been excluded,²⁴ evidence to date including reduced conversions upon varying nucleophilic promoters (vide infra) is suggestive of nucleophile involvement in the rate-determining or prior step. Indirect evidence comes from previously reported highly enantioselective biscyclization routes to carbocycle-fused β -lactones employing chiral nucleophile.^{14a,c}

At the outset of our studies, we recognized that several energetically favored conformers dictated by $A^{1,3}$ -strain, e.g., conformers **37** and **37'**, while unproductive for aldol lactonization since the ketone is not proximal to the ammonium enolate,²⁵ would preserve the enantiopurity of the single stereocenter (C2). This would be possible since the α -proton is nearly in plane with the amide carbonyl in these conformations due to $A^{1,3}$ -strain (Scheme 4a).¹⁵ Furthermore, an expectedly higher energy conformation such as that represented by **37**'' is required to achieve bis-cyclization, and because of the proximity and favorable formation of a

⁽²³⁾ For lead references to this mechanistic possibility, see: (a) Adler, M.; Adler, S.; Boche, G. J. Phys. Org. Chem. **2005**, 18, 193. (b) Fox, J. M.; Dmitrenko, O.; Liao, L.; Bach, R. D. J. Org. Chem. **2004**, 69, 7317.

⁽²⁴⁾ For previous reports of β -lactones from keto acid derivatives via proposed [2 + 2] mechanisms, see: (a) Boswell, G. A.; Dauben, W. G.; Ourisson, G.; Rull, T. *Bull. Soc. Chim. Fr.* **1958**, 1598. (b) Kagan, H. B.; Jacques, J. *Bull. Soc. Chim. Fr.* **1958**, 1600. (c) Brady, W. T.; Gu, Y. Q. *J. Org. Chem.* **1988**, *53*, 1353. (d) Reddy, L. R.; Corey, E. J. *Org. Lett.* **2006**, *8*, 1717For a previous report of an aldol-lactonzation pathway, see:. (e) Merlic, C. A.; Marlog, B. C J. Org. Chem. **2003**, *68*, 6056. For an anionic version, see: (f) Shindo, M.; Matsumoto, K.; Sato, Y.; Shishido, K. Org. Lett. **2001**, *3*, 2029.

⁽²⁵⁾ The alternate N-C1 amide rotamer, which is also accessible energetically but also unproductive for aldol lactonization, is not considered in this analysis.



^{*a*}All reactions were performed at -5 to -10 °C except entry 1, which was performed at 23 °C. ^{*b*}Carboxylic acid activating agent. ^{*c*}Nucleophilic promoter. ^{*d*}Yields are given for two steps (bis-cyclization/allyl deprotection) and refer to isolated, purified (SiO₂) products. ^{*e*}Byproducts were not easily separated, and thus, approximate yields are provided on the basis of ¹H NMR (500 MHz) integration. ^{*f*}Recovered starting material, keto acid **52**.

5-membered lactam ring, aldol reaction might be expected to be facile. What we did not anticipate was the impact of the C4 absolute stereochemistry on the diastereoselectivity of the bis-cyclization. In our early optimization studies, we determined that diastereoselectivity was highly dependent on whether (S)- or (R)-serine-derived keto acids were employed leading to different diastereoselectivities, 1:2 vs 8:1, respectively. We suggest that the moderate diastereoselectivity observed for the bis-cyclization is governed by developing torsional strain between the methyl group (C3) of the ketone and the C2 substituent (cf. 41, R⇔Me) during the initial aldol reaction, leading to the preferred relative stereochemistry following lactonization (cf. 39). Note that only a moderate preference for this conformation (cf. 37") over the alternative (cf. 41) is expected,²⁶ and this is reflected in the maximum observed diastereoselectivity for the bis-cyclization (dr 4-7:1, vide infra). Regarding the C4 absolute stereochemistry dependence, this is likely a result of required conformations for deprotonation imposed by the N-C4 bond rotamer of the two diastereomeric, acyl ammonium species 36 and 40 leading to ammonium enolates 37 and 41 and suggests the intriguing possibility of a memory of chirality effect which we are currently studying in related systems. With acyl ammonium 36 derived from (R)-serine, deprotonation leads directly to a N-C4 conformer that can access reactive conformer 37" leading to major diastereomer 39. Alternatively, deprotonation of the diastereomeric acyl ammonium 40 derived from (S)-serine leads directly to reactive conformer 41 that delivers the diastereometric β -lactone 42. This analysis assumes a rapid bis-cyclization since enantiopurity could indeed be maintained in a non-Curtin-Hammett situation, if once the higher energy reactive conformation is achieved, biscyclization immediately takes place. For these reasons, we utilized (R)-serine for the enantioselective synthesis (vide infra).

TABLE 2. Optimization of the Bis-cyclization Leading to Bicyclic β -Lactone (\pm)-32

In early optimization studies of the bis-cyclization of keto acid **52** directed toward salino A, we isolated and identified three byproducts: enol lactone **43**, cyclopropyl ketoamide **44**,





and γ -lactam 45 (Table 1). Only the former two byproducts were observed when both 4-PPY and Hünig's base were employed accompanied by low yields of β -lactones 32 and 32' (Table 1, entries 1 and 2). Enol lactone 43 is likely derived from intramolecular acylation of enol 46 of the β -ketoamide with the activated ester (Scheme 5). Cyclopropyl ketoamide 44 may be derived from a condensation reaction of enol 46 onto the activated acid leading to β -diketone 47, followed by a retro-Claisen process induced by 4-PPY via 48, and finally cyclization via enol 49 leading to the cyclopropane (Scheme 5, red mechanism arrows). We next studied the use of 4-PPY (7.0 equiv) as both base and nucleophilic promoter, which reduced the amount of byproduct 43 and 44 with attendant dramatic improvements in yield of β -lactone but formation of a new byproduct, unsaturated γ -lactam 45 (Table 2, entry 3). Consideration of possible mechanisms for formation of this byproduct suggested control experiments to determine if lactam 45 was derived from decomposition of β -lactones 32 and 32' via a decarboxylation pathway. Subjecting β -lactones 32/32' to 4-PPY at 23 °C indeed led to lactam 45 (31%) and interestingly recovered β -lactones 32 and 32' with improved diastereometric ratio $(dr, 5:1 \rightarrow 9:1, Scheme 6)$. Decomposition was much slower at -5 to -10 °C and was not observed if only Hünig's base was added to β -lactones 32 and 32'. The increased diastereomeric

⁽²⁶⁾ The preferred conformation of 2-butanone places the methyl and carbonyl oxygen nearly in plane (eclipsed conformation). However, the energy difference is low (~1 kcal/mol); see: Eliel; E. L.; Wilen, S. H. *Stereochemistry of Oganic Compounds*; Wiley & Sons: New York, 1994; pp 616–617.





ratio suggests a differential rate of decomposition for the two diastereomers. This may be explained by a lower barrier synelimination (endothermic, late transition state) for β -lactone **32**' due to a more accessible proton and release of torsional strain ($\mathbf{R} \Leftrightarrow \mathbf{M}\mathbf{e}$) which is absent in diastereomeric β -lactone **32**. Finally, decarboxylation and protonation leads to the observed unsaturated γ -lactam **45**.

These studies highlighted the importance of minimizing reaction time to avoid decomposition of β -lactone and to minimize C2-deprotonation which avoids these byproducts and maintains optical purity during the bis-cyclization process. Dramatic improvements were therefore realized when alternative activating agents and most importantly a less polar solvent (e.g., toluene), which may slow C2-deprotonation, were employed. Use of MsCl in toluene avoided formation of all byproduct and yields of the β -lactones **32**, **32'** increased to 61% (Table 2, entry 4). Employing the less nucleophilic dimethylaminopyridine (DMAP) also led to β -lactone **32**, **32'** but in slightly reduced yields (54%) under similar conditions suggestive of nucleophile involvement in the rate-determining or prior step (Table 2, entry 5).

The impact of the chlorine atom on the acidity of the C2proton was investigated under the original bis-cyclization conditions with a dual purpose of studying inductive effects on conversion and byproduct formation, and demonstrating the versatility of this synthetic strategy for C2-derivative synthesis. Ketoamide substrate 50a was prepared from a β -keto thioester while **50b** was prepared from heteroketene dimer 14c (not shown).²¹ Bis-cyclization of the keto acid derived from allyl ester 50a bearing a γ -siloxy group (Table 3, entry 2) led to similar results as those from ketoamide 31 bearing a γ -chloro substituent (salino core shown for comparison, Table 3, entry 1); however the δ -chloro keto acid derived from allyl ester 50b led to a significant increase in yield (Table 3, entry 3). The yields obtained appear to correlate with the ¹H NMR chemical shifts of the C2 protons of β -ketoamides 31 and **50a,b** (Table 3, entries 1-3). Comparison of $H_{a,b}$ chemical shifts of the γ -heteroatom substituted ketoamide **31** (Table 3, entry 1) leading to the salino A core with that of the δ -substituted ketoamides **50a-b** (Table 3, entries 2, 3) shows a downfield shift ($\Delta \delta \sim 0.4$) pointing to the inductive effects and attendant increased acidity leading to byproduct formation imposed by the heteroatoms. Inductive effects leading to reduced yields (70-85%) and greater byproduct formation were also observed in simpler C4-unsubstituted substrates bearing α -phenyl and β -chloro C2 substituents (Table 1,

 TABLE 3.
 Variation of the C2-Side Chain: Inductive Effects on Efficiency of Bis-cyclization



entry	R^1	β -lactone	% yield ^a , ^b	δ H _a , H _b ^c
1	$(CH_2)_2Cl$	32/32′	40	3.92, 3.94
2	(CH ₂) ₂ OTBS	51a/51a'	42	3.92, 3.95
3	$(CH_2)_3Cl$	51b/51b'	62	3.52, 3.57

^{*a*}Yield for two steps (deprotection and bis-cyclization). ^{*b*}Yields refer to isolated, purified β -lactones. ^{*c*}Chemical shift of C2 protons (H_{a/b}) of diastereometric ketoamide esters **31**, **50**a,b.

entries 3 and 5) in comparison to other C2-alkyl substrates in that series (90-93% yield).

Total Synthesis of (-)-Salinosporamide A. Having identified bis-cyclization conditions that improved yields and diastereoselectivity, we turned our attention to the asymmetric version of the bis-cyclization. The synthesis of salino A began with reductive amination of commercially available (R)-O-benzyl serine ((R)-22) (99% ee) with p-anisaldehyde and subsequent esterification provided allyl ester (+)-23b (Scheme 7). After extensive optimization, the desired ester (+)-23b could be obtained with minimal racemization (98%) ee) in 79% yield (two steps) by careful control of reaction temperature and duration.²⁷ Acylation of serine derivative (+)-23b with unsymmetrical ketene dimer (\pm)-14b under microwave conditions gave diastereomeric β -ketoamides 31/31' (dr 1:1) in 80% yield. Separation of the diastereomers by MPLC on mulitgram scale (3-4 g/run) provided the required (R,R)- β -ketoamide **31** (dr 30:1, 98% ee, 46%). Under acidic conditions, the undesired diastereomer (2S, 4R)-31' could be transformed to a 1:1 mixture of diastereomers via epimerization of the C2 but not the C4 stereocenter (verified by chiral HPLC), thus achieving an effective resolution of ketene dimer (\pm) -14b²⁸ and greater material throughput.²⁹

⁽²⁷⁾ Optical purities were determined by chiral HPLC analysis (ref 21).

⁽²⁸⁾ Attempts to prepare the required optically active heteroketene dimer 14b by the method of Calter (ref 19) led to very low yields.

⁽²⁹⁾ Chiral HPLC analysis verified that epimerization only occurred at C2 within limits of detection (ref 21).

SCHEME 7. Nine-Step Enantioselective Synthesis of (–)-Salino A (1a) from (*R*)-OBn Serine ((+)-22)



SCHEME 8. Mechanism of Inhibition of the Proteasome (CT Site) with Salino A $(1a^{13a})$ and Proposed for Homosalino A (56)



SCHEME 9. Varying the C2-Side Chain: Synthesis of (-)-Homosalino A (54)



Optimization of the Pd(0)-mediated allyl deprotection of ester (R,R)-31 was also required to minimize epimerization of the C2 stereocenter and ultimately delivered keto acid 52 with neglibile erosion of the diastereomeric purity.

At the outset of the studies toward an asymmetric process, it was unclear whether the A^{1,3}-strain induced conformational bias in β -keto tertiary amides (~4 kcal/mol) would be sufficient to avoid epimerization of this center in the time frame and under the basic conditions of the bis-cyclization. In our initial studies, application of bis-cyclization conditions developed for the racemic series^{8e} to keto acid **52** (dr ~29:1, 98% ee) using

modified Mukaiyama's reagent but without Hünig's base led to similar yields and diastereoselectivity (40%, dr 2.5:1) obtained previously, with an attendant erosion of optical purity of β -lactone **32** (85% ee). As described in our optimization studies above, we ultimately found that MsCl at lower temperature with less polar solvents led to increased diastereoselectivity. When these conditions were utilized, the diasteroeselectivity improved (7:1) and minimal epimerization (~3%) was observed providing bicyclic β -lactone **32** in 35% yield and 92% ee. Further improvements in yield up to 60% were realized with longer reaction times under these same conditions; however, this led to reduced diastereoselectivity and enantiopurity (dr 4:1, 88% ee). Importantly, the bis-cyclization could be performed on

SCHEME 10. Varying the C4-Side Chain of Salino A



TABLE 4. IC₅₀ Values for Inhibition of the Human 20S and 26S Proteasome by Salino A Derivatives in a Luminogenic Enzymatic Assay^a

	compd	1050		
entry		CT-L activity 20S/26S (nM)	C-L activity 20S/26S (nM)	T-L activity 20S/26S (nM)
1	<i>rac</i> -cinnabaramide A, (\pm) -2	$2.8 \pm 0.6/2.2 \pm 0.04$	$125 \pm 12^{b}/230 \pm 24^{b}$	$260 \pm 29^{b}/412 \pm 83$
2	(-)-salino A (1a) (92% ee)	$0.8 \pm 0.08/2.5 \pm 1.3$	$111 \pm 22/156 \pm 18$	$39 \pm 7/37 \pm 16$
3	<i>rac</i> -phenyl deriv (61)	$29 \pm 0.7/57 \pm 20$	$788 \pm 50/1585 \pm 633$	$702 \pm 235/589 \pm 120$
4	rac-diMePh (63)	inactive ^b , ^c	inactive ^b , ^c	inactive ^b , ^c
5	(-)-homosalino (54) (88% ee)	$0.7 \pm 0.04/2.3 \pm 1.1$	$144 \pm 12/188 \pm 3$	$118 \pm 28/125 \pm 30$
6	velcade	$2.6 \pm 0.7/7.4 \pm 1.9$	$25 \pm 7/72 \pm 17$	$254 \pm 24/680 \pm 150$
$^{a}IC_{50}$	values are the mean \pm standard deviat	tion of 4 or more experiments. ^b The	average of two experiments. ^c Esse	ntially inactive at >99,000 nM.
(CT-L:	chymotrypsin-like activity; C-L: caspas	e-like activity; T-L: trypsin-like activ	vity)	•

gram scale with comparable diastereoselectivity and retention of coul enantiopurity (52% over two steps, dr 5:1, 90% ee).

Completion of the salino A synthesis entailed hydrogenolysis to provide alcohol (-)-53, which could be separated at this stage from the minor diastereomer produced during the bis-cyclization process (75% yield). Modified Moffatt oxidation³⁰ and addition of the zinc reagent **29** derived from the reaction of *n*-butyllithium with cyclohexenyltributyltin and ZnCl₂ following the procedure of Corey^{8a} gave a mixture of C5,C6-diastereomeric alcohols (dr 11:3:1:1) in 62% yield (two steps, Scheme 7, procedure A ("proc A")) with the desired diastereomer 33 as the major adduct. Alternately, zinc reagent 29' could be prepared directly from commercially available 3-bromocyclohexene and activated zinc by the method of Knochel,¹⁶ which greatly simplified product purification since tin byproducts were avoided (Scheme 7, proc B). Most importantly, this protocol gave only two diastereomers 33/33' (dr 4:1), improved the yield to 74%, and avoided the use of toxic tin reagents. This procedure also enabled isolation of a novel salino A diastereomer 1a' (C5, C6-bis-epi-salino A, not shown) whose relative stereochemistry was verified by X-ray analysis.²¹ Finally, deprotection of PMB-protected lactam (-)-33 led to pure (-)-salino A (1a) which correlated well with spectroscopic data reported previously (syn $\left[\alpha\right]_{D}^{23}$ -71.3; lit. $[\alpha]^{25}_{D} - 72.9$).³

Total Synthesis of (–)-Homosalinosporamide A (Homosalino A). In our optimization studies described above, we altered the C2-side chain with the dual purpose of studying inductive effects (cf. Table 3) and providing the one carbon homologated side chain that could ultimately lead to (–)-homosalino A (54). Groll and co-workers have proposed that acylation of the proteasome by the β-lactone of salino A at the catalytic N-terminus (Thr 1) is followed by cyclization of the incipient alcohol with the pendant chloroethyl group leading to the formation of a tetrahydrofuran as verified by X-ray analysis (Scheme 8).^{13a} This led us to consider the synthesis of homosalino A (54) to determine if a tetrahydropyran (cf. 57, n = 2)

could be formed at the active site of the proteasome in analogy to the known formation of a tetrahydrofuran.

The synthesis followed an identical route as for salino A but employed heteroketene dimer (\pm)-14c with serine derivative (+)-23b to provide the homologous ketoester (-)-58 (Scheme 9). Bis-cyclization following ester deprotection provided the bicyclic- β -lactone (-)-59 in 60% yield (dr 3.5:1). Using an identical sequence as described for salino A, the β -lactone core was converted to (-)-homosalino A ((-)-54, 88% ee).²¹

Variation of the C4-Side Chain of the Salino Core. Our synthetic strategy also enables facile variation of the C4-side chain by addition of organometallic reagents to the versatile aldehyde obtained by oxidation of alcohol (\pm)-53 (Scheme 10). Addition of Grignard reagents (dr ~3:1) followed by PMB deprotection provided the known phenyl^{12c} and novel dimethylphenyl salino derivatives (\pm)-61 and (\pm)-63, respectively. The relative stereochemistry of the major diastereomer of phenyl derivative (\pm)-61 was verified by single-crystal X-ray analysis.²¹

Biological Studies. The salino A derivatives synthesized were analyzed for their ability to inhibit the chymotrypsin-like (CT-L), caspase-like (C-L), and trypsin-like (T-L) activities of the 20S and 26S human proteasome in a luminogenic enzymatic assay (Table 4). Both velcade (entry 6) and synthetic salino A (92% ee, entry 2) were included as controls and data obtained correlates well with previous analyses.¹³ As expected on the basis of known structure-activity relationships,^{7,13a} changes made to the C5-side chain in general led to lower activity with drastic loss of activity observed for the racdimethylphenyl derivative 63. The potency of the racemic phenyl derivative against the human 20A proteasome was much greater than that reported previously for the optically active phenyl derivative against the yeast 20S proteasome.^{12c} However, as previously noted, changes in the C2-side chain are more tolerable as indicated by (-)-homosalino A (54, Table 4, entry 5), and indeed, this derivative exhibited very similar activity to the natural product with the exception of decreased inhibition of the trypsin-like activity of the 20S proteasome ($\sim 3 \times$ decrease).

^{(30) (}a) Pfitzner, K. E.; Moffatt, J. G. J. Am. Chem. Soc. 1965, 87, 5661.
(b) Coulton, S.; Southgate, R. J. Chem. Soc., Perkin Trans. 1 1992, 961.

In summary, we developed a concise, nine-step enantioselective route to (-)-salino A and derivatives from O-benzyl serine. The key bis-cyclization of a β -ketoamide, amenable to gram scale, constructs both the γ -lactam and the fused- β lactone in one operation leading to the brevity of the synthesis. The flexibility of the described strategy derives from the versatility of the bicyclic core (-)-32 enabling attachment of various C4-side chains, even in the presence of the β -lactone, and the ability to use alternative ketene dimers 14 to vary the C2 side chain. Several derivatives including (-)-homosalino A were synthesized, and their activity toward proteasome inhibition was measured. The ability of the described β -keto tertiary amide substrates to maintain stereochemical integrity by virtue of A^{1,3} strain raises interesting questions regarding how such integrity is maintained with β -keto secondary amides, known salino A precursors,4c or if a dynamic kinetic resolution is operative during related biosynthetic aldol lactonizations.³¹ The synthesis of additional hypothesis-driven salino derivatives by the described strategy, their bioactivity, and crystallographic studies of their interaction with the proteasome will be reported in due course.

Experimental Section

(2*R*,4*R*)-*β*-Keto Acid, 52. To a solution of ketoamide (2*R*,4*R*)-31 (2.30 g, 4.55 mmol, ~32:1 dr) in THF (91.0 mL) at -5 °C (ice and saturated NaCl solution) was added Pd(PPh₃)₄ (526 mg, 0.455 mmol), followed by immediate addition of morpholine (0.475 mL, 5.46 mmol). The reaction mixture was stirred at -5 °C for 70 min and then diluted with ice-cold Et₂O (800 mL). A 0.02 N HCl solution was added until the pH was measured to be ~3. The layers were separated, and the organic layer was washed with brine (400 mL), dried over MgSO₄, and concentrated. The crude keto acid (2*R*,4*R*)-**52** (~ 29:1 dr according to 500 MHz ¹H NMR) was used directly in the subsequent step without further purification. (Note: longer reaction times led to epimerization).

Gram-Scale Bis-cyclization Leading to Benzyloxy β -Lactone (-)-32/32'. To a solution of 4-pyrrolidinopyridine (2.60 g, 18.2) mmol, 4.0 equiv) in toluene (84 mL) at -5 °C (ice and saturated NaCl solution) was added MsCl (0.53 mL, 6.83 mmol, 1.5 equiv). Immediately, a solution of the freshly prepared keto acid (R,R)-5a (total amount from above, ~4.55 mmol) in toluene (25 mL) was added to the resulting suspension via syringe pump over 45 min, and 5 mL of additional toluene was used to ensure complete transfer. After 3 h, the reaction mixture was diluted with ice-cold Et₂O (700 mL) and washed with 20% CuSO₄ solution (500 mL) to remove most of the 4-pyrrolidinopyridine and saturated NH₄Cl (500 mL), and then washed with water (2×500 mL). The organic layer was dried over MgSO₄, filtered, and concentrated. The residue was purified by flash chromatography (1:9 \rightarrow 3:7 EtOAc/ hexanes) to give a mixture of two inseparable β -lactones (-)-32/ 32' (1.05 g, 52%, dr 5:1, 500 MHz ¹H NMR) as a yellow oil and recovered keto acid (10%, 4:1 dr): $R_f = 0.36$ (20% EtOAc/ hexanes). The enantiomeric excess of β -lactone 32 was determined to be 90%. The diasterometric β -lactones were carried directly forward to the deprotection at which point they could be separated. Data for (-)-32: IR (neat) 1830, 1703 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.32-7.36 (m, 3H), 7.13-7.15 (m, 4H), 6.80 (d, J=8.5 Hz, 2H), 4.73 (d, J=15.5 Hz, 1H), 4.31 (d, J=15.5 Hz, 1H), 4.17 (d, J = 12.0 Hz, 1H), 4.13 (d, J = 11.5 Hz, 1H), 4.01 (ddd,J = 5.0, 7.5, 12.5 Hz, 1H), 3.77 - 3.81 (m, 1H), 3.77 (s, 3H), 3.73(d, J = 11.5 Hz, 1H), 3.57 (d, J = 11.5 Hz, 1H), 2.91 (t, J =7.5 Hz, 1H), 2.31-2.38 (m, 1H), 2.10-2.16 (m, 1H), 1.72 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 173.8, 166.1, 159.2, 136.4,

JOC Featured Article

129.2(2C), 128.6, 128.5(2C), 128.2, 128.0(2C), 113.9(2C), 83.4, 79.3, 73.5, 61.6, 55.2, 45.0, 44.3, 42.5, 28.4, 19.2; HRMS (ESI) Calcd. For $C_{24}H_{27}CINO_5[M + H]$ 444.1578, found 444.1590. The enantiomeric excess of (-)-**32** was determined to be 92% by chiral HPLC (CHIRALPAK IA, 250 × 4.6 mm (L × I.D.), solvent (isocratic) 87:13 hexanes/2-propanol, flow rate 1.0 mL/min, $\lambda =$ 230 nm). Retention times: (-)-**32**, 13.68 min; (+)-**32**, 16.12 min.

Hydroxy β -Lactone (-)-53. To a mixture of β -lactones (-)-32 and 32' (260 mg, 0.586 mmol, dr 7:1) in THF was added palladium on carbon (52 mg, 20 wt %). After the flask was evacuated twice by aspirator vacuum and refilled with H_2 , a balloon of H₂ was attached to the flask and the heterogeneous solution was stirred vigorously at 23 °C for 12 h. The reaction mixture was then diluted with Et₂O and dried over MgSO₄. The organics were filtered through a pad of Celite, concentrated, and purified by MPLC (SiO₂, 5:95 EtOAc/CH₂Cl₂) to give the desired alcohol(-)-53 in 75% yield (155 mg, dr > 19:1, 92% ee) as a waxy solid: $R_f = 0.29$ (5:95 EtOAc/CH₂Cl₂); $[\alpha]^{23}_{D} = -67.0$ (c = 0.95, CHCl₃). Data for (-)-53: IR (neat) 3449, 1831, 1687 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.30 (d, J = 8.5 Hz, 2H), 6.89 (d, J =8.5 Hz, 2H, 5.13 (d, J = 15.0 Hz, 1H), 4.06 (d, J = 15.5 Hz, 1H),4.03 (ddd, J = 5.5, 7.5, 12.5 Hz, 1H), 3.92 (dd, J = 9.0, 13.5 Hz,1H), 3.85 (dd, J = 4.5, 13.5 Hz, 1H), 3.80 (s, 3H), 3.78–3.82 (m, 1H), 2.94 (t, J = 7.0 Hz, 1H), 2.32–2.38 (m, 1H), 2.01–2.18 (m, 1H), 1.77 (s, 3H), 0.86 (dd, J = 5.0, 9.5 Hz, 1H); ¹³C NMR (125 MHz, CDCl₃) δ 174.2, 166.7, 159.6, 129.0(2C), 128.7, 114.7(2C), 83.6, 80.2, 55.3, 55.1, 44.9, 44.1, 42.4, 28.4, 19.1; HRMS (ESI) Calcd. For C₁₇H₂₁ClNO₅ [M + H] 354.1108, found 354.1098.

Representative Procedure for Modified Moffatt Oxidation and Zincate Formation/Addition As Described for *N*-PMB-salino A, (–)-33. To a solution of alcohol (–)-53 (42 mg, 0.119 mmol, dr >19:1) in DMSO/toluene (0.9 mL/0.9 mL) were added EDCI (0.114 mg, 0.595 mmol) and dichloroacetic acid (5 μ L, 0.060 mmol) at 23 °C. The reaction mixture was stirred for 6 h and then diluted with EtOAc (100 mL). The reaction mixture was acidified using 0.1 N HCl to pH 3. The organic layer was then washed with brine (60 mL), dried over MgSO₄, filtered, and concentrated. The crude aldehyde was used for the next step without further purification due to some instability of the aldehyde observed upon flash column chromatography.

Preparation of a stock solution of zincate 29': Zinc dust (1.64 g, 25.0 mmol, Baker, 20 mesh) was activated by the addition of an HCl solution (5 mL, 1.0 M) in a 25 mL round-bottomed flask. After the mixture was stirred vigorously for 15 min at 23 °C, the acid solution was decanted and the zinc was repeatedly washed with dry THF $(3 \times 5 \text{ mL})$. The zinc was then dried under high vacuum for 2 h. Addition of dry THF (5 mL) was followed by slow addition of 3-bromocyclohexene (0.806 g, 5.0 mmol) over 15 min. The resulting mixture was stirred at 23 °C for 18 h and then transferred via syringe to a dry centrifuge tube to remove the excess zinc suspension (10 min, 80 rpm). The resulting clear solution was then utilized directly, following titration as described below, being careful not to disturb the pellet at the bottom during transfers. The concentration of the zincate 29' solution was determined as follows: A stock solution of iodine in THF was prepared by addition of iodine (254 mg, 1.0 mmol) in 10 mL of THF leading to a $100 \,\mu$ M solution. A portion (500 μ L) of this iodine solution was placed into a dry 5 mL round-bottomed flask equipped with a magnetic stir bar. The zincate solution, after being centrifuged, was added dropwise into the iodine stock solution until the red color disappeared. Using this procedure, the prepared zincate solution was determined to be $\sim 200 \ \mu M$ and was used (in excess) directly in the following reaction.

To a solution of the crude aldehyde (\sim 0.119 mmol) in THF (1.0 mL) at -78 °C was added \sim 1.2 mL (0.238 mmol, \sim 200 μ M in THF, \sim 2.0 equiv) of zincate **29**' via syringe pump over 30 min. The resulting mixture was slowly warmed to -20 °C for 1 h, quenched with water and diluted with Et₂O (120 mL). Saturated

⁽³¹⁾ The C2-epimer of salino A has been isolated; see ref 11.

NH₄Cl was added until a pH of 7 was achieved. The layers were separated, and then the organic layer was washed with brine (80 mL). The filtrate was dried over MgSO₄, filtered, and concentrated. The residue was purified by flash chromatography (95:5 \rightarrow 85:5 EtOAc/hexanes) to give a mixture of two inseparable diastereomers of *N*-PMB salino A (-)-33 and C5, C6-bis*epi-N*-PMB salino A (-)-33' (38 mg, 74%, dr = 4:1 according to 500 MHz ¹H NMR) as a colorless oil, which was carried directly to the next step without further purification.

Salino A ((-)-1a) and C5,C6-Bis-epi-salino A ((-)-1a'). To a methanolic solution (0.8 mL) containing the mixture of alcohols (-)-33/33' (38 mg, 0.088 mmol) was added an aqueous solution of CAN (0.240 g, 0.44 mmol) in H₂O (0.2 mL) at 0 °C dropwise. After being stirred at 0 °C for 6 h, the reaction mixture was diluted with EtOAc (100 mL) and washed with saturated solution of NaHCO₃ and brine. The organic layer was dried over MgSO₄, filtered, and concentrated. The residue was purified by flash chromatography (5:95 \rightarrow 15:85 EtOAc/CH₂Cl₂) to give salino A (-)-1a (14 mg, 51%, dr > 19:1, 500 MHz ¹H NMR) as a white solid (previously described, see ref 8j) in addition to a novel diastereomeric salino A 1a' (3.3 mg, 12%, dr > 19:1, 500 MHz ¹H NMR). Data for C5,C6-bis-*epi*-salino A (1a'): $R_f =$ 0.56 (33% EtOAc/hexanes); IR (neat) 2931, 1830, 1702 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 6.28 (s, 1H), 6.10 – 6.14 (m, 1H), 5.85 - 5.88 (m, 1H), 3.99 (ddd, J = 5.0, 7.5, 12.5 Hz, 1H), 3.83 (dd, J = 7.5, 9.0 Hz, 1H), 3.78 (ddd, J = 5.0, 7.5, 12.5 Hz, 1H),2.83 (t, J = 7.5 Hz, 1H), 2.48 - 2.50 (m, 1H), 2.27 - 2.34 (m, 1H),2.11-2.18 (m, 1H), 2.06-2.08 (m, 2H), 1.98 (d, J = 9 Hz, 1H), 1.80 – 1.91 (m, 2H), 1.87 (s, 3H), 1.60 – 1.67 (m, 2H); ¹³C NMR (125 MHz, CDCl₃) δ 175.8, 165.0, 135.2, 123.5, 85.4, 78.7, 71.8, 44.9, 42.7, 38.4, 28.5 26.6, 25.2, 21.8, 19.4; HRMS (MALDI) calcd for $C_{15}H_{20}CINO_4Na$ [M + Na] 336.0973, found 336.0957.

Proteasome Inhibition Assay. The Proteasome-Glo Assay (Promega) was used, which is a homogeneous coupled-enzyme system that utilizes luminogenic substrates to measure the three proteolytic activities (chymotrypsin-like(CT-L), trypsin-like (T-L), and caspase-like (C-L)) associated with the proteasome. Purified human 20S and 26S proteasome (BioMol/Enzo Life Sciences, Inc.) at $0.5-2 \ \mu g/mL$ was incubated in 10 μL of assay buffer (10 mM Hepes, pH 7.6) at room temperature for $20-30 \ min$ in the presence of the salino A derivatives at various concentrations or equivalent volume of solvent DMSO as a control along with salino A and Velcade. The reaction was initiated by addition of $10 \ \mu L$ of 2X solution of various substrates. Reactions were performed at room temperature. After 1 h, inhibition of each proteasomal activity was measured on a multilabel microtiter plate reader.

Acknowledgment. We thank the NIH (GM069784) and the Welch Foundation (A-1280) for support. We thank Drs. Joe Reibenspies and Nattamai Bhuvanesh (TAMU) for X-ray analysis. We thank Mr. Nicholas Krudy (TAMU NSF REU, CHE-0755207) for technical assistance. We thank Mr. Ranjith C. Perumalil and Mr. Mikail C. Abbasov for graphics assistance for journal cover art. The U.S. Department of Energy Genomics: GTL Program, http://doegenomestolife. org, is acknowledged for the proteasome graphic used in the Table of Contents entry and Abstract graphic.

Supporting Information Available: General procedures for synthesis and the human proteasome inhibiton assay and characterization data including ¹H and ¹³C NMR spectra (for compounds **1a'**, **21e**, **44**, **45**, **61**, **63**) and X-ray analysis (**61**, **1a'**). This material is available free of charge via the Internet at http:// pubs.acs.org.