LETTERS

C–H Oxidation/Michael Addition/Cyclization Cascade for Enantioselective Synthesis of Functionalized 2-Amino-4*H*-chromenes

Bo Wu,[†] Xiang Gao,[†] Zhong Yan,[†] Mu-Wang Chen,[†] and Yong-Gui Zhou^{*,†,‡}

[†]State Key Laboratory of Catalysis, Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, P. R. China [‡]State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, Shanghai 200032, P. R. China

(5) Supporting Information

ABSTRACT: A streamlined method for the enantioselective synthesis of 2-amino-4*H*-chromenes from readily available 2-alkyl-substituted phenols and active methylene compounds bearing a cyano group with up to 97% ee is presented. This reaction is a cascade procedure including manganese dioxide mediated C–H oxidation for the generation of *o*-quinone methides and bifunctional squaramide-catalyzed Michael addition/cyclization.



• he development of efficient and novel chemical reactions that allow the rapid and direct construction of complex and diversified molecules from available and inexpensive starting materials stands at the forefront of synthetic chemistry, and it also has impressive significance in industrial processes. One of the most powerful methods to realize these goals is the application of cascade reactions which carry out multiple transformations in a single step, accompanied by undeniable benefits including atom economy, economy of labor, and sustainability of resources.² In particular, these cascade strategies are useful and effective in the total synthesis of natural products and bioactive molecules.^{2c-e,3} The significant importance of the 2-amino-4H-chromene scaffold in biologically active molecules, natural products, and synthetic drugs⁴ has motivated chemists to develop various methods for the asymmetric synthesis of 2-amino-4H-chromenes, involving organocatalytic tandem Michael addition/cyclization,⁵ Mannich cyclization/tautomerization cascade sequences,⁶ three-component cascade reaction,⁷ conjugate addition of nitroalkanes to 2iminochromenes,⁸ and metal complex catalyzed cascade reactions.9 Although considerable progress has been made in the formation of chiral 2-amino-4H-chromenes, there are several disadvantages including low catalytic efficacy, poor stereoselectivity, and unsatisfactory substrate scope. Hence, the development of facile and enantioselective approaches to such scaffolds with a broad substrate scope is still highly desirable.

o-Quinone methides (*o*-QMs) are a prominent class of intermediates in a large number of biological processes¹⁰ and are recognized as a highly reactive chemical species.¹¹ In this decade, a series of organocatalytic enantioselective reactions of o-QMs^{12,13} have been reported due to their inherent reactivity. Organocatalytic formal [4 + 2] cycloaddition of *o*-QMs with active methylene compounds bearing a cyano group is also a concise method for the synthesis of chiral 2-amino-4*H*-chromenes. Recently, the Han group reported quinine-catalyzed annulation of the electron-rich and stable *o*-QMs

with malononitrile, giving 2-amino-3-cyano-4*H*-chromenes with excellent enantioselectivity (eq 1, Scheme 1).^{12m}





However, the substrate scope was limited. Fochi's group and our groups independently described a similar bifunctional squaramide-catalyzed reaction of *o*-QMs generated *in situ* from 2-(1-arylsulfonyl-alkyl)phenols with active methylene compounds bearing a cyano group under basic conditions for the preparation of chiral 2-amino-4*H*-chromenes (eq 2, Scheme 1).^{12n,o}

2-Alkyl-substituted phenol derivatives are easily available and can be oxidized to generate *o*-QM intermediates.¹⁴ Additionally, we have also disclosed that *o*-QM intermediates could be

Received: October 30, 2015

generated in situ from 2-alkyl-substituted phenol derivatives by metal oxidant mediated oxidation, which could be rapidly trapped by the sulfur ylides to allow stereoselective synthesis of trans-2,3-dihydrobenzofurans.^{14g} Therefore, we envisioned a metal oxidant mediating the C-H oxidation of 2-alkylsubstituted phenol derivatives to produce the o-QM intermediates that underwent asymmetric Michael addition with active methylene compounds bearing a cyano group via an organocatalytic system followed by cyclization for the construction of optical active 2-amino-4H-chromenes. The combination of a metal oxidant and organocatalyst in this cascade reaction could effectively impede background reactions and ensure high enantioselectivity. Generally, a simple combination of metal oxidant and organocatalyst remains challenging in organic chemistry. Organocatalysts experience difficulty in tolerating highly oxidizing conditions.¹⁵ Moreover, a metal oxidant might bind tightly with the organocatalyst, which causes the deactivation of the organocatalyst.¹⁶ Herein, we reported cascade C-H oxidation/Michael addition/ cyclization reactions with the combination of a metal oxidant and an organocatalyst for the enantioselective synthesis of 2amino-4H-chromenes (eq 3, Scheme 1).

At the outset, 2-alkyl-substituted phenol 1a and malononitrile 2a (1.2 equiv) were chosen as model substrates. The oxidant had a significant influence in the reaction and triggered the generation of o-QM intermediates. In this respect, we initially turned our attention to the evaluation of oxidants. To our delight, the anticipated product 2-amino-4H-chromene 3a was obtained in 81% isolated yield and 78% ee in the presence of silver oxide and a catalytic amount of quinine (Table 1, entry 1). However, other metal oxidants, AgOAc and $K_3Fe(CN)_{6i}$ exhibited lower reactivity (Table 1, entries 2 and 3). Organic oxidants such as DDQ and PhI(OAc)₂ afforded mixed products. Additionally, the yield could be further improved, and the enantioselectivity was maintained by using MnO_2 as the oxidant (Table 1, entry 4). Subsequently, various solvents were extensively examined, and it was found that solvent effect played a crucial role in the reactivity and enantioselectivity. Among them, CHCl₃ proved to be a suitable solvent in view of the excellent yield and good enantioselectivity (Table 1, entries 5-10). Next, a series of organocatalysts were explored. Cinchona alkaloid-based thiourea catalysts 4b and 4c provided disappointing results with moderate enantioselectivity (Table 1, entries 11 and 12). Gratifyingly, quinine-based squaramide catalyst 4d gave the best enantioselectivity (91% ee) (Table 1, entry 13). Satisfactorily, no decrease in enantioselectivity and yield was observed when the catalyst loading was reduced to 5 mol % (Table 1, entry 16).

With the aforementioned optimal reaction conditions, we next explored the substrate scope of the current cascade reaction between 2-alkyl-substituted phenols 1 and active methylene compounds bearing a cyano group 2. As summarized in Scheme 2, in all cases, the reaction performed very well, delivering the corresponding chiral 2-amino-4*H*-chromenes in good yields and high enantioselectivities. For 2-alkyl substituted sesamols 1a-1g, electronic and steric properties had no obvious influence on the enantioinduction. Notably, vinyl substrates 1h and 1j were favorable reaction partners, and the reaction proceeded smoothly with good enantioselectivities and moderate yields. In addition, 2-alkyl-substituted 4,5-dimethoxyphenol substrates 1i-k were employed in the cascade reaction with excellent enantiocontrol. Interestingly, 2-alkyl-substituted 4-methoxynaphthols 11 and





^{*a*}Conditions: **1a** (0.10 mmol), **2a** (0.12 mmol), cat. (0.01 mmol), oxidant (2.0 equiv), while MnO₂ was used in 4.0 equiv, solvent (1.5 mL), 30 °C. ^{*b*}Isolated yields. ^{*c*}Determined by chiral HPLC. ^{*d*}Catalyst loading was reduced to 5 mol %.

Im could also be oxidized to generate *o*-QMs and reacted with malononitrile to afford the corresponding adducts in high yield and enantioselectivity. Furthermore, a train of active methylene compounds bearing a cyano group proceeded successfully, achieving moderate to good yields and excellent enantioselectivities. It is worth noting that when the substituents R were alkyl groups, mixed products were obtained. The absolute configuration of compounds **3a** and **3k** was determined to be *S* by comparison of the specific rotations with the reported literature data (see the Supporting Information).^{12m,o}

Considering that both amino and cyano groups are versatile functional groups, further derivatizations were conducted for the production of biologically active molecules. The reaction of (S)-6-amino-8-(4-methoxyphenyl)-8*H*-[1,3]dioxolo[4,5-*g*]-chromene-7-carbonitrile **3a** with cyclohexanone afforded polycyclic compound **5** in 63% yield with 89% ee in the presence of aluminum chloride according to the known literature method (Scheme 3),¹⁷ which is a tacrine analogue and acts as a potential drug for the treatment of Alzheimer's disease.^{17b}

In conclusion, we have developed a streamlined method for the asymmetric synthesis of chiral 2-amino-4*H*-chromenes from readily available 2-alkyl-substituted phenols and active methylene compounds bearing a cyano group with excellent

в

Scheme 2. Substrate Scope for the Synthesis of 2-Amino-4*H*-chromenes



Scheme 3. Synthesis of Bioactive Compound 5



enantioselectivity in the presence of a metal oxidant and an organocatalyst. This reaction is recognized as a cascade procedure including manganese dioxide mediated C-H oxidation for the generation of *o*-quinone methides and a bifunctional squaramide-catalyzed Michael addition/cyclization process. This protocol not only provides new access to chiral 2-amino-4*H*-chromenes but also suggests a useful strategy for the combination of metal oxidants and organocatalysts.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.or-glett.Sb03148.

Experimental procedures, characterization data, and NMR spectra (PDF)

AUTHOR INFORMATION

Corresponding Author

*E-mail: ygzhou@dicp.ac.cn.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We are grateful for the financial support from National Natural Science Foundation of China (21372220 and 21532006).

REFERENCES

(1) Walji, A. M.; MacMillan, D. W. C. Synlett 2007, 2007, 1477.
(b) Young, I. S.; Baran, P. S. Nat. Chem. 2009, 1, 193.

(2) For reviews, see: (a) Xu, P.-F., Wang, W., Eds. Catalytic Cascade Reactions; Wiley: Hoboken, NJ, 2014. (b) Wang, Y.; Lu, H.; Xu, P.-F. Acc. Chem. Res. 2015, 48, 1832. (c) Nicolaou, K. C.; Montagnon, T.; Snyder, S. A. Chem. Commun. 2003, 551. (d) Nicolaou, K. C.; Edmonds, D. J.; Bulger, P. G. Angew. Chem., Int. Ed. 2006, 45, 7134. (e) Smith, J. M.; Moreno, J.; Boal, B. W.; Garg, N. K. Angew. Chem., Int. Ed. 2015, 54, 400.

(3) For selected examples, see: (a) Beemelmanns, C.; Reissig, H.-U. Angew. Chem., Int. Ed. 2010, 49, 8021. (b) Granger, B. A.; Jewett, I. T.; Butler, J. D.; Hua, B.; Knezevic, C. E.; Parkinson, E. I.; Hergenrother, P. J.; Martin, S. F. J. Am. Chem. Soc. 2013, 135, 12984. (c) Rodríguez-López, J.; Crisóstomo, F. P.; Ortega, N.; López-Rodríguez, M.; Martín, V. S.; Martín, T. Angew. Chem., Int. Ed. 2013, 52, 3659. (d) Yue, G.; Zhang, Y.; Fang, L.; Li, C.; Luo, T.; Yang, Z. Angew. Chem., Int. Ed. 2014, 53, 1837. (e) Liu, J.; Ng, T.; Rui, Z.; Ad, O.; Zhang, W. Angew. Chem., Int. Ed. 2014, 53, 136.

(4) Foe selected examples, see: (a) Patchett, A. A.; Nargund, R. P. Annu. Rep. Med. Chem. 2000, 35, 289. (b) Das, S. G.; Doshi, J. M.; Tian, D.; Addo, S. N.; Srinivasan, B.; Hermanson, D. L.; Xing, C. J. Med. Chem. 2009, 52, 5937. (c) Kemnitzer, W.; Drewe, J.; Jiang, S.; Zhang, H.; Wang, Y.; Zhao, J.; Jia, S.; Herich, J.; Labreque, D.; Storer, R.; Meerovitch, K.; Bouffard, D.; Rej, R.; Denis, R.; Blais, C.; Lamothe, S.; Attardo, G.; Gourdeau, H.; Tseng, B.; Kasibhatla, S.; Cai, S. X. J. Med. Chem. 2004, 47, 6299. (d) Wang, J.-L.; Liu, D.; Zhang, Z.-J.; Shan, S.; Han, X.; Srinivasula, S. M.; Croce, C. M.; Alnemri, E. S.; Huang, Z. Proc. Natl. Acad. Sci. U. S. A. 2000, 97, 7124. (e) Kasibhatla, S.; Gourdeau, H.; Meerovitch, K.; Drewe, J.; Reddy, S.; Qiu, L.; Zhang, H.; Bergeron, F.; Bouffard, D.; Yang, Q.; Herich, J.; Lamothe, S.; Cai, S. X.; Tseng, B. Mol. Cancer Ther. 2004, 3, 1365.

(5) (a) Wang, X.-S.; Yang, G.-S.; Zhao, G. Tetrahedron: Asymmetry 2008, 19, 709. (b) Gogoi, S.; Zhao, C.-G. Tetrahedron Lett. 2009, 50, 2252. (c) Ding, D.; Zhao, C.-G. Tetrahedron Lett. 2010, 51, 1322. (d) Ramireddy, N.; Abbaraju, S.; Zhao, C.-G. Tetrahedron Lett. 2011, 52, 6792. (e) Xie, J.-W.; Huang, X.; Fan, L.-P.; Xu, D.-C.; Li, X.-S.; Su, H.; Wen, Y.-H. Adv. Synth. Catal. 2009, 351, 3077. (f) Hu, Z.-P.; Lou, C.-L.; Wang, J.-J.; Chen, C.-X.; Yan, M. J. Org. Chem. 2011, 76, 3797. (g) Ren, Q.; Gao, Y.; Wang, J. Chem. - Eur. J. 2010, 16, 13594. (h) Du, Z.; Siau, W.-Y.; Wang, J. Tetrahedron Lett. 2011, 52, 6137. (i) Gao, Y.; Yang, W.; Du, D.-M. Tetrahedron: Asymmetry 2012, 23, 339. (j) Hu, K.; Lu, A.; Wang, Y.; Zhou, Z.; Tang, C. Tetrahedron 2014, 70, 181.

(6) Ren, Q.; Siau, W.-Y.; Du, Z.; Zhang, K.; Wang, J. Chem. - Eur. J. 2011, 17, 7781.

(7) (a) Yang, G.; Luo, C.; Mu, X.; Wang, T.; Liu, X.-Y. *Chem. Commun.* **2012**, *48*, 5880. (b) Zhang, G.; Zhang, Y.; Yan, J.; Chen, R.; Wang, S.; Ma, Y.; Wang, R. *J. Org. Chem.* **2012**, *77*, 878.

(8) Li, W.; Liu, H.; Jiang, X.; Wang, J. ACS Catal. 2012, 2, 1535.

(9) (a) Dong, Z.; Liu, X.; Feng, J.; Wang, M.; Lin, L.; Feng, X. *Eur. J. Org. Chem.* **2011**, 2011, 137. (b) Chen, W.; Cai, Y.; Fu, X.; Liu, X.; Lin, L.; Feng, X. *Org. Lett.* **2011**, *13*, 4910.

(10) (a) Rokita, S. E., Ed. Quinone Methides; Wiley: Hoboken, NJ, 2009. (b) Basarić, N.; Mlinarić-Majerski, K.; Kralj, M. Curr. Org. Chem. **2014**, 18, 3. (c) Doria, F.; Nadai, M.; Folini, M.; Scalabrin, M.; Germani, L.; Sattin, G.; Mella, M.; Palumbo, M.; Zaffaroni, N.; Fabris, D.; Freccero, M.; Richter, S. N. Chem. - Eur. J. **2013**, 19, 78. (d) Doria, F.; Nadai, M.; Folini, M.; Di Antonio, M.; Germani, L.; Percivalle, C.; Sissi, C.; Zaffaroni, N.; Alcaro, S.; Artese, A.; Richter, S. N.; Freecero,

M. Org. Biomol. Chem. 2012, 10, 2798. (e) Nadai, M.; Doria, F.; Di Antonio, M.; Sattin, G.; Germani, L.; Percivalle, C.; Palumbo, M.; Richter, S. N.; Freccero, M. Biochimie 2011, 93, 1328.

(11) For reviews, see: (a) Van de Water, R. W.; Pettus, T. R. R. *Tetrahedron* 2002, 58, 5367. (b) Willis, N. J.; Bray, C. D. *Chem. - Eur. J.* 2012, 18, 9160. (c) Amouri, H.; Le Bras, J. *Acc. Chem. Res.* 2002, 35, 501. (d) Ferreira, S. B.; da Silva, F. d. C.; Pinto, A. C.; Gonzaga, D. T. G.; Ferreira, V. F. *J. Heterocycl. Chem.* 2009, 46, 1080. (e) Bai, W.-J.; David, J. G.; Feng, Z.-G.; Weaver, M. G.; Wu, K.-L.; Pettus, T. R. R. *Acc. Chem. Res.* 2014, 47, 3655.

(12) For asymmetric annulations of o-quinone methides, see: (a) Alden-Danforth, E.; Scerba, M. T.; Lectka, T. Org. Lett. 2008, 10, 4951. (b) Lv, H.; You, L.; Ye, S. Adv. Synth. Catal. 2009, 351, 2822. (c) Rueping, M.; Uria, U.; Lin, M.-Y.; Atodiresei, I. J. Am. Chem. Soc. 2011, 133, 3732. (d) Lv, H.; Jia, W.-Q.; Sun, L.-H.; Ye, S. Angew. Chem., Int. Ed. 2013, 52, 8607. (e) Izquierdo, J.; Orue, A.; Scheidt, K. A. J. Am. Chem. Soc. 2013, 135, 10634. (f) El-Sepelgy, O.; Haseloff, S.; Alamsetti, S. K.; Schneider, C. Angew. Chem., Int. Ed. 2014, 53, 7923. (g) Hsiao, C.-C.; Liao, H.-H.; Rueping, M. Angew. Chem., Int. Ed. 2014, 53, 13258. (h) Lee, A.; Scheidt, K. A. Chem. Commun. 2015, 51, 3407. (i) Saha, S.; Schneider, C. Chem. - Eur. J. 2015, 21, 2348. (j) Zhao, J.-J.; Sun, S.-B.; He, S.-H.; Wu, Q.; Shi, F. Angew. Chem., Int. Ed. 2015, 54, 5460. (k) Hsiao, C.-C.; Raja, S.; Liao, H.-H.; Atodiresei, I.; Rueping, M. Angew. Chem., Int. Ed. 2015, 54, 5762. (1) Saha, S.; Schneider, C. Org. Lett. 2015, 17, 648. (m) Adili, A.; Tao, Z.-L.; Chen, D.-F.; Han, Z.-Y. Org. Biomol. Chem. 2015, 13, 2247. (n) Caruana, L.; Mondatori, M.; Corti, V.; Morales, S.; Mazzanti, A.; Fochi, M.; Bernardi, L. Chem. - Eur. J. 2015, 21, 6037. (o) Wu, B.; Gao, X.; Yan, Z.; Huang, W.-X.; Zhou, Y.-G. Tetrahedron Lett. 2015, 56, 4334.

(13) For other asymmetric reactions of *o*-quinone methides, see:
(a) Luan, Y.; Schaus, S. E. J. Am. Chem. Soc. 2012, 134, 19965.
(b) Guo, W.; Wu, B.; Zhou, X.; Chen, P.; Wang, X.; Zhou, Y.-G.; Liu, Y.; Li, C. Angew. Chem., Int. Ed. 2015, 54, 4522. (c) Zhao, W.; Wang, Z.; Chu, B.; Sun, J. Angew. Chem., Int. Ed. 2015, 54, 1910. (d) Wang, Z.; Ai, F.; Wang, Z.; Zhao, W.; Zhu, G.; Lin, Z.; Sun, J. J. Am. Chem. Soc. 2015, 137, 383. (e) Dai, W.; Lu, H.; Jiang, X.-L.; Gao, T.-T.; Shi, F. Tetrahedron: Asymmetry 2015, 26, 109. (f) Li, M.-L.; Chen, D.-F.; Luo, S.-W.; Wu, X. Tetrahedron: Asymmetry 2015, 26, 219. (g) Wilcke, D.; Herdtweck, E.; Bach, T. Synlett 2011, 2011, 1235. (h) Saha, S.; Alamsetti, S. K.; Schneider, C. Chem. Commun. 2015, 51, 1461.

(14) For selected examples, see: (a) Bolon, D. A. J. Org. Chem. 1970, 35, 3666. (b) Jurd, L. Tetrahedron 1977, 33, 163. (c) Angle, S. R.; Yang, W. J. Am. Chem. Soc. 1990, 112, 4524. (d) Angle, S. R.; Rainier, J. D.; Woytowicz, C. J. Org. Chem. 1997, 62, 5884. (e) Bishop, L. M.; Winkler, M.; Houk, K. N.; Bergman, R. G.; Trauner, D. Chem. - Eur. J. 2008, 14, 5405. (f) Liao, D.; Li, H.; Lei, X. Org. Lett. 2012, 14, 18. (g) Wu, B.; Chen, M.-W.; Ye, Z.-S.; Yu, C.-B.; Zhou, Y.-G. Adv. Synth. Catal. 2014, 356, 383.

(15) Guo, C.; Song, J.; Luo, S.-W.; Gong, L.-Z. Angew. Chem., Int. Ed. **2010**, 49, 5558.

(16) Yang, D.; Wang, L.; Zhao, D.; Han, F.; Zhang, B.; Wang, R. Chem. - Eur. J. 2013, 19, 4691.

(17) (a) Kamdar, N. R.; Haveliwala, D. D.; Mistry, P. T.; Patel, S. K. *Med. Chem. Res.* **2011**, *20*, 854. (b) Maalej, E.; Chabchoub, F.; Oset-Gasque, M. J.; Esquivias-Pérez, M.; González, M. P.; Monjas, L.; Pérez, C.; de los Ríos, C.; Rodríguez-Franco, M. I.; Iriepa, I.; Moraleda, I.; Chioua, M.; Romero, A.; Marco-Contelles, J.; Samadi, A. *Eur. J. Med. Chem.* **2012**, *54*, 750.