

Development of a chimeric c-Src kinase and HDAC inhibitor

Kristin S. Ko, Michael E Steffey, Kristoffer R Brandvold, and Matthew B. Soellner

ACS Med. Chem. Lett., **Just Accepted Manuscript** • DOI: 10.1021/ml400175d • Publication Date (Web): 03 Jul 2013

Downloaded from <http://pubs.acs.org> on July 3, 2013

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



Development of a chimeric c-Src kinase and HDAC inhibitor

Kristin S. Ko,² Michael E. Steffey,¹ Kristoffer R. Brandvold,¹ and Matthew B. Soellner^{1,2}

¹Department of Medicinal Chemistry and ²Department of Chemistry, University of Michigan, 930 N. University Avenue, Ann Arbor, Michigan 48109, United States

KEYWORDS: c-Src, kinase, chimera, HDAC, combination therapy

ABSTRACT: On the basis of synergism observed between a selective c-Src kinase inhibitor with an HDAC inhibitor, the development of the first chimeric c-Src kinase and HDAC inhibitor is described. The optimized chimeric inhibitor is shown to be a potent c-Src and HDAC inhibitor. Chimeric inhibitor **4** is further shown to be highly efficacious in cancer cell lines and significantly more efficacious than a dual-targeting strategy using discrete c-Src and HDAC inhibitors.

The non-receptor tyrosine kinase c-Src plays an important role in many aspects of cell physiology, regulating diverse cellular processes including division, motility, adhesion, angiogenesis, and survival.^{1,2} c-Src was the first proto-oncogene identified, is frequently over-expressed in cancer, and the extent of over-expression of c-Src correlates with malignant potential.^{1,2} Furthermore, c-Src expression levels inversely correlate with patient survival.^{1,2} Recently, c-Src activity was shown to be a main mode of resistance to Herceptin, a first line therapy for Her2+ breast cancer.³ Therefore, c-Src kinase is an attractive therapeutic target in cancer.

We recently reported the first highly selective inhibitor of c-Src (Figure 1).⁴ Despite potent biochemical activity against c-Src, our selective c-Src inhibitor (**1**) is only modestly potent in cellular proliferation assays using breast cancer cell lines.⁴ Following the success of combinatorial drug therapies in the treatment of HIV,⁵ tuberculosis,⁶ and other microbial infections,⁷ the use of multiple targeted drugs for cancer chemotherapy is increasingly being pursued.⁸ We reasoned that multi-target inhibition using our selective c-Src inhibitor might lead to improved cellular efficacy.

To identify drug combinations that would be synergistic with c-Src inhibition, we examined a small library of targeted inhibitors in combination with our selective c-Src inhibitor **1**. These studies were performed in SK-BR-3 cells, a Her2+ breast cancer cell line previously shown to be growth dependent upon c-Src kinase activity.^{4,9} From these experiments, we identified that panobinostat, a histone deacetylase (HDAC) inhibitor in clinical trials,¹⁰ was highly synergistic with c-Src inhibitor **1** (Figure 2). HDAC inhibitors have been shown to promote the growth arrest and apoptosis of cancer cells with minimal toxicity.¹¹ We believe that the observed synergy is due to previously reported mechanisms whereby HDAC inhibitors can down-regulate c-Src levels through repression of SRC transcription.¹²




Figure 1. Structures of highly selective c-Src inhibitor **1**, vorinostat, and panobinostat.

To determine whether the synergy observed with c-Src inhibition and panobinostat was general for any HDAC inhibitor, we performed combination experiments with vorinostat,¹³ an FDA approved HDAC inhibitor, and c-Src inhibitor **1** (Table 1). c-Src inhibitor **1** and vorinostat have a GI₅₀ of 4.8 μ M and 1.2 μ M, respectively, for SK-BR-3 proliferation. In combination, c-Src inhibitor **1** + vorinostat (1:1) has a GI₅₀ for SK-BR-3 proliferation of 0.8 μ M, which is an improvement over either inhibitor dosed alone.¹⁴ Next, as a measure of cellular toxicity, we examined each compound's ability to inhibit proliferation of primary human mammary epithelial cells (HMEC). c-Src inhibitor **1** and vorinostat have a GI₅₀ of 4.3 μ M and 5.8 μ M, respectively, for HMEC proliferation. The combination of **1** + vorinostat (1:1) has a GI₅₀ of 5.4 μ M against primary mammary epithelial cells.




Figure 2. Synergy studies of selective c-Src inhibitor **1** (2 μ M), panobinostat (HDACi, 10 nM), and combination (1+HDACi, 2 μ M **1**, 10 nM panobinostat) in SK-BR-3 cell line. Red line denotes predicted additivity [(eA+eB)-(eA*eB)] of **1** + panobinostat. The higher level of inhibition than the predicted additivity indicated synergism between **1** and panobinostat.

Using the SK-BR-3 and HMEC data, we calculated a therapeutic index (GI_{50} HMEC / GI_{50} SK-BR-3) for c-Src inhibitor **1**, vorinostat, and the combination of **1** + vorinostat (Table 1).¹⁵ c-Src inhibitor **1** has a poor therapeutic index of 0.9 while vorinostat's therapeutic index is 4.8. Disappointingly, the combination of **1** + vorinostat has an insignificant improvement in therapeutic index (6.8) relative to vorinostat alone (4.8).

Table 1. Cellular efficacy of selective c-Src inhibitor **1**, vorinostat, **1**:vorinostat (1:1), and chimera **4**


	GI_{50} (μ M), SK-BR-3	GI_{50} (μ M), HMEC	Therapeutic Index
Compound 1	4.8 ± 0.2	4.3 ± 0.4	0.9
Vorinostat	1.2 ± 0.1	5.8 ± 0.2	4.8
1 + Vorinostat	0.80 ± 0.03	5.4 ± 0.2	6.8
Chimera 4	0.20 ± 0.03	4.7 ± 0.3	23.5

We wondered whether there would be any advantage for a chimeric inhibitor, where a single molecule could serve as both a c-Src kinase and HDAC inhibitor, rather than using two separate agents in combination. For example, we thought that we might obtain improved cellular efficacy. In addition, using a single agent to inhibit both c-Src and HDAC does not lead to the additive toxicity that is often observed with combination therapy.¹⁴ Chimeric kinase-HDAC inhibitors have previously been developed, however, no Src-HDAC chimeric compounds have been reported.^{16–18} In addition, previously reported studies of kinase-HDAC chimeras lack a comparison of therapeutic indices between combination therapy and chimeric inhibition.^{16–18}

We previously reported PP2~alkyne (**2**), a modular and selective c-Src inhibitor scaffold.⁴ We envisioned using this kinase inhibitor scaffold to append HDAC pharmacophores. The classic pharmacophore for HDAC inhibitors consists of a zinc-binding motif, a hydrophobic linker, and a recognition cap.¹⁹ Using PP2~alkyne, HDAC elements can readily be appended using “click” chemistry.²⁰ Importantly, the use of a triazole ring as the recognition cap in HDAC inhibitors has previously

been reported and shown to be highly efficacious both *in vitro* and *in cellulo*.²¹ Previous reports with triazole-based HDAC inhibitors have demonstrated that a 6-carbon hydrophobic linker will provide potent HDAC inhibition.²¹ While only 1,4-[1,2,3]-triazoles have been reported as HDAC inhibitors,²¹ we reasoned that because our selective c-Src inhibitor **1** contains a 1,5-[1,2,3]-triazole,⁴ we would synthesize and evaluate both regioisomers.


Chart 1. Structure of PP2~Alkyne (**2**) and chimeric inhibitors **3** and **4**



We synthesized compounds **3** and **4** as chimeric Src/HDAC inhibitors. Compound **3** has a 1,4-triazole and was synthesized using a copper-mediated cycloaddition reaction,²⁰ while compound **4** has a 1,5-triazole synthesized using a ruthenium-mediated cycloaddition reaction (Chart 1, see Supporting Information for synthetic details).²² Using a previously reported fluorescence assay for c-Src kinase activity,²³ we found that **3** and **4** were competent c-Src kinase inhibitors (K_i = 371 and 138 nM, respectively). We next examined the ability of **3** and **4** to inhibit HDAC1 using a Fluor de Lys based-assay²⁴ and found both compounds were potent inhibitors of HDAC1 (K_i = 0.62 and 0.26 nM, respectively). In our assays, compound **4** was a better inhibitor of both c-Src and HDAC1. Thus, the 1,5-triazole regiochemistry was used exclusively for subsequent linker optimization.

In an effort to optimize potency for both c-Src and HDAC1, we synthesized a series of chimeric HDAC-Src inhibitors containing varied hydrophobic linkers (Table 2). This series included alkyl linkers of varied length as well as styrene-containing linkers that are found in panobinostat.¹⁰ The six-carbon alkyl linker (compound **4**) was found to be optimal for inhibition of both c-Src kinase and HDAC1. Of note, we found the styrene linkers (compounds **7** and **8**) were ineffective as c-Src inhibitors and only modest inhibitors of HDAC1 compared to the *n*-alkyl linkers.

Table 2. SAR of Linker




Compound	R =	K_i , c-Src (nM)	K_i , HDAC1 (nM)
4		138 ± 15	0.26 ± 0.02
5		190 ± 23	0.82 ± 0.29
6		407 ± 46	9.8 ± 0.8
7		4300 ± 440	35 ± 0.6
8		2140 ± 390	23 ± 1.5


Chimeric inhibitor **4** is one of the most potent HDAC1 inhibitors reported to date ($K_i = 260$ pM) and is also a potent c-Src inhibitor ($K_i = 138$ nM). To decipher the binding contributions for each half of the chimera, two fragments of inhibitor **4** were synthesized (Chart 2). Compound **10** contains only the HDAC inhibitor pharmacophore, while compound **9** includes the c-Src kinase binding elements. Interestingly, we observe a marked decrease in affinity for both c-Src and HDAC1 when both elements are not present. Specifically, compound **10**, which retains all of the HDAC inhibitor pharmacophore elements, has a K_i for HDAC1 that is >170-times higher than found with chimeric inhibitor **4**. These data imply that the c-Src binding elements enhance HDAC1 inhibition observed with compound **4**. Likewise, the c-Src inhibitor fragment **9** has nearly 10x less affinity for c-Src than chimera **4**, suggesting that the addition of the HDAC fragment is important for c-Src inhibition. Together, these data demonstrate that chimera **4** is not simply two inhibitors linked together, but rather represents a merged inhibitor where both elements are required for affinity against each target.

Our chimeric inhibitor was initially optimized for HDAC inhibition using HDAC1, however, we assumed it could be a promiscuous inhibitor of HDACs. Profiling of compound **4** against a panel of 11 HDACs was performed by Reaction Biology (Malvern, PA). The HDAC profiling revealed that our chimera is a potent and non-selective inhibitor against class I, IIa, and IV HDACs (Table 3). Consistent with vorinostat's selectivity, chimera **4** is not an effective inhibitor of class IIb HDACs (Table 3). Relative to vorinostat, chimera **4** has improved affinity to all HDACs except HDAC8 and HDAC11.

Chart 2. c-Src inhibitor **9** and HDAC inhibitor **10**



9
 K_i , c-Src = 605 ± 90 nM
 K_i , HDAC1 = $>12,500$ μ M



10
 K_i , c-Src = $>12,500$ μ M
 K_i , HDAC1 = 45 ± 2 nM

Table 3. HDAC Profiling of Chimera **4** and Vorinostat

	HDAC Class	IC ₅₀ (nM), Chimera 4	IC ₅₀ (nM), Vorinostat ^a
HDAC1	I	86 ± 11	306
HDAC2	I	231 ± 39	232
HDAC3	I	19 ± 1	132
HDAC4	IIa	$3,982 \pm 920$	76,000
HDAC5	IIa	$3,891 \pm 681$	27,200
HDAC6	IIb	2.7 ± 0.5	20
HDAC7	IIa	$13,220 \pm 860$	105,000
HDAC8	I	$2,311 \pm 251$	306
HDAC9	IIa	$28,020 \pm 1,910$	141,000
HDAC10	IIb	51 ± 3	432
HDAC11	IV	224 ± 26	200

^a Data from Reaction Biology (Malvern, PA).

In previously published work, we found that c-Src inhibitors that are selective for c-Src over c-Abl are more efficacious in cell culture with non-hematopoietic cancers.⁴ Thus, we wanted to determine whether chimera **4** has selectivity for c-Src over c-Abl. Gratifyingly, in our biochemical assay, chimera **4** was selective for c-Src over c-Abl (K_i for c-Src = 138 nM, K_i for c-Abl = 2,350 nM). We next tested the ability of **4** to inhibit Hck, a SRC-family kinase with 85% similarity across the kinase domain to c-Src, and found it has a $K_i = 504$ nM. Together, these data suggest that chimera **4** is selective for c-Src over homologous kinases. Given that our compound shares many features with our highly selective c-Src inhibitor **1**,⁴ it is likely that chimera **4** is also selective for c-Src.

In an effort to compare chimeric inhibition to dual-targeting c-Src and HDAC1, we examined the efficacy of chimera **4** *in cellulo*. Combination dosing of selective c-Src inhibitor **1** + vorinostat (1:1) was found to have a $GI_{50} = 0.78$ μ M for SK-BR-3 cells and a $GI_{50} = 5.4$ μ M for non-cancer HME cells. This resulted in a therapeutic index of 6.8 (*vide supra*). In comparison, chimeric inhibitor **4** was more efficacious at inhibiting the growth of SK-BR-3 cells ($GI_{50} = 0.2$ μ M) and has similar non-cancer cellular toxicity ($GI_{50} = 4.7$ μ M for HME

cells), resulting in a cellular therapeutic index of 23.5 (Table 1). This corresponds to chimeric inhibitor **4** having an improvement in therapeutic index significantly higher than dual targeting c-Src and HDACs with two distinct compounds (23.5 versus 6.8, respectively). These results highlight an important advantage for chimeric inhibition over dual-agent targeting: we observe synergistic activity against cancer cells while not increasing the cellular toxicity relative to the single agent counterparts.

To better characterize the cellular efficacy of our chimeric c-Src/HDAC inhibitor, compound **4** was submitted to the National Cancer Institute for screening in the NCI-60 panel (see Supporting Information for full NCI-60 data).²⁵ From this panel, chimera **4** has an average $GI_{50} = 0.26 \mu\text{M}$. Significantly, the efficacy of chimera **4** across the NCI-60 is better than vorinostat (NCI-60 average $GI_{50} = 0.53 \mu\text{M}$) and a FDA-approved c-Src inhibitor (dasatinib, NCI-60 average $GI_{50} = 5.7 \mu\text{M}$). In addition to the improved efficacy across the NCI-60 panel, chimera **4** does not have increased toxicity relative to primary human mammary cells (chimera **4**, HMEC $GI_{50} = 4.7 \mu\text{M}$; vorinostat, HMEC $GI_{50} = 5.8 \mu\text{M}$; dasatinib, HMEC $GI_{50} = 1.8 \mu\text{M}$).

Table 4. NCI-60 Panel Data for Chimera 4, Vorinostat, and Dasatinib against Select Cell Lines

	Cell line	GI_{50} (μM), chimera 4	GI_{50} (μM), vorinostat	GI_{50} (μM), dasatinib
a)	KM12	0.47	1.88	7.44
	MCF7	0.35	2.19	8.32
	U251	0.28	1.53	2.81
b)	DU-145	0.39	1.36	0.16
	HS 578T	0.17	4.83	0.03
	MDA-MB-231	0.39	2.32	0.02
c)	HCT-116	0.22	0.37	3.70
	MALME-3M	0.07	0.37	6.61
	SW-620	0.25	0.54	8.43

Analysis of the NCI-60 data demonstrates that chimera **4** is a highly efficacious agent in cell lines where vorinostat and dasatinib are ineffective alone (Table 4a). Furthermore, chimera **4** is more efficacious than vorinostat when c-Src inhibition is shown to be efficacious (Table 4b). For example, chimera **4** is an efficacious inhibitor of Hs 578T, a triple negative breast cancer cell line, cell growth ($GI_{50} = 0.17 \mu\text{M}$) while vorinostat is not ($GI_{50} = 4.83 \mu\text{M}$), due to c-Src inhibition having an important role in Hs 578T cell proliferation (dasatinib $GI_{50} = 0.03 \mu\text{M}$). This dramatic increase in efficacy demonstrates that chimera **4** is acting as more than a HDAC inhibitor alone in cellulo. Finally, chimera **4** is observed to be more effective than dasatinib in cell lines where cellular proliferation is dependent upon HDAC1 activity (Table 4c). Together, these data demonstrate that compound **4**'s impressive cellular efficacy in the NCI-60 panel is inherent in its chimeric nature and the

ability to inhibit both c-Src kinase and HDAC1 is required for the cellular potency observed.

In summary, we have reported the first chimeric c-Src kinase and HDAC inhibitor. Furthermore, we have performed detailed studies that demonstrate that chimera **4** is a potent and selective c-Src kinase inhibitor as well as a potent and non-selective HDAC inhibitor. We demonstrated that our chimeric inhibitor has improved efficacy in cellular experiments compared to dosing two individual inhibitors targeting c-Src and HDACs. Chimera **4** has significant efficacy in the NCI-60 panel, while not possessing significant toxicity to primary human cells, and represents a novel small molecule probe that can provide simultaneous inhibition of c-Src and HDACs. Our approach to constructing kinase-HDAC inhibitor hybrids using a triazole linkage should be general and readily adapted to any kinase and/or HDAC pair of interest.

ASSOCIATED CONTENT

Supporting Information. Synthetic procedures and characterization data, assay conditions, and cell data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

*Tel: 734-615-2767 Email: soellner@med.umich.edu

Author Contributions

The manuscript was written through contributions of all authors and all authors have given approval to the final version of the manuscript.

Funding Sources

Funding of this research was provided in part by NIH grant R01GM088546 to M.B.S. and the University of Michigan College of Pharmacy.

Notes

The authors have no competing financial interests.


ACKNOWLEDGMENT

We would like to thank Christel Fox for preparation of c-Src, c-Abl, and Hck kinase domain proteins.


REFERENCES

- (1) Thomas, S. E.; Brugge, J. S. Cellular functions regulated by Src family kinases. *Annu. Rev. Cell Dev. Biol.* **1997**, *13*, 513–609.
- (2) Martin, G. S. The hunting of the Src. *Nat. Rev. Mol. Cell Biol.* **2001**, *2*, 467–475.
- (3) Zhan, S.; Huang, W.-C.; Li, P.; Guo, H.; Poh, S.-B.; Brady, S. W.; Xiong, Y.; Tesng, L.-M.; Li, S.-H.; Ding, Z.; Sahin, A. A.; Esteve, F. J.; Hortobagti, G. N.; Yu, D. Combating trastuzumab resistance by targeting SRC, a common node downstream of multiple resistance pathways. *Nat. Med.* **2011**, *17*, 461–469.
- (4) Brandvold, K. B.; Steffey, M. E.; Fox, C. C.; Soellner, M. B. Development of a highly selective c-Src kinase inhibitor. *ACS Chem. Biol.* **2012**, *7*, 1393–1398.
- (5) Pirrone, V.; Thakkar, N.; Jacobson, J. M. Combinatorial approaches to the prevention and treatment of HIV-1 infection. *Antimicrob. Agents Chemother.* **2011**, *55*, 1831–1842.
- (6) Harries, A. D.; Dye, C. Tuberculosis. *Ann. Rheumatic Diseases* **2006**, *5–6*, 415–431.
- (7) Bouza, E.; Muñoz, P. Monotherapy versus combination therapy for bacterial infections. *Med. Clin. North Am.* **2000**, *84*, 1357–1389.



- (8) Dancey, J. E.; Chen, H. X. Strategies for optimizing combinations of molecularly targeted anticancer agents. *Nat. Rev. Drug Discov.* **2006**, *5*, 649–659.
- (9) Zheng, X.; Resnick, R. J.; Shalloway, D. Apoptosis of estrogen-receptor negative breast cancer and colon cancer cell lines by PTP α and Src RNAi. *Int. J. Cancer* **2008**, *122*, 1999–2007.
- (10) Atadja, P. Development of the pan-DAC inhibitor panobinostat (LBH589): Successes and challenges. *Cancer Lett.* **2009**, *280*, 233–241.
- (11) Bolden, J. E.; Peart, M. J.; Johnstone, R. W. Anticancer activities of histone deacetylase inhibitors. *Nat. Rev. Drug Discov.* **2006**, *5*, 769–784.
- (12) Kostyniuk, C.; Dehm, S.; Batten, D.; Bonham, K. The ubiquitous and tissue specific promoters of the human SRC gene are repressed by inhibitors of tissue deacetylases. *Oncogene* **2002**, *21*, 6340–6347.
- (13) Marks, P. A.; Breslow, R. Dimethyl sulfoxide to vorinostat: development of this histone deacetylase inhibitor as an anticancer drug. *Nat. Biotechnol.* **2007**, *25*, 84–90.
- (14) Zimmermann, G. R.; Lehár, J.; Keith, C. T. Multi-target therapeutics: When the whole is greater than the sum of the parts. *Drug Discov. Today* **2007**, *12*, 34–42.
- (15) Muller, Y.; Milton, M. The determination and interpretation of the therapeutic index in drug development. *Nat. Rev. Drug Discov.* **2012**, *11*, 751–761.
- (16) Cai, X.; Zhai, H.-X.; Wang, J.; Forrester, J.; Qu, H.; Yin, L.; Lai, C.-J.; Bao, R.; Qian, C. Discovery of 7-(4-(3-ethynylphenylamino)-7-methoxyquinazolin-6-yloxy)-N-hydroxyheptanamide (CUDC-101) as a Potent Multi-Acting HDAC, EGFR, and HER2 Inhibitor for the Treatment of Cancer. *J. Med. Chem.* **2010**, *53*, 2000–2009.
- (17) Mahboobi, S.; Dove, S.; Sellmer, A.; Winkler, M.; Eichhorn, E.; Pongratz, H.; Ciossek, T.; Baer, T.; Maier, T.; Beckers, T. Design of chimeric histone deacetylase- and tyrosine kinase-inhibitors: A series of imatinib hybrids as potent inhibitors of wild-type and mutant BCR-ABL, PDGFR β , and histone deacetylases. *J. Med. Chem.* **2009**, *52*, 2265–2344.
- (18) Mahboobi, S.; Sellmer, A.; Winkler, M.; Eichhorn, E.; Pongratz, H.; Ciossek, T.; Baer, T.; Maier, T.; Beckers, T. Novel chimeric histone deacetylase inhibitors: a series of lapatinib hybrids as potent inhibitors of epidermal growth factor receptor (EGFR), human epidermal growth factor receptor 2 (HER2), and histone deacetylase activity. *J. Med. Chem.* **2010**, *53*, 8546–8601.
- (19) Paris, M.; Porcelloni, M.; Binaschi, M.; Fattori, D. Histone deacetylase inhibitors: From bench to clinic. *J. Med. Chem.* **2008**, *51*, 1505–1534.
- (20) Kolb, H. C.; Finn, M. G.; Sharpless, K. B. Click chemistry: Diverse chemical function from a few good reactions. *Angew. Chem., Int. Ed.* **2001**, *40*, 2004–2021.
- (21) Chen, P.; Patil, V.; Guerrant, W.; Green, P.; Oyeler, A. Synthesis and structure-activity relationship of histone deacetylase (HDAC) inhibitors with triazole-linked cap group. *Bioorg. Med. Chem.* **2008**, *16*, 4839–4892.
- (22) Zhang, L.; Chen, X. G.; Xue, P.; Sun, H. Y. Y.; Williams, I. D.; Sharpless, K. B.; Fokin, V. V.; Jia, G. C. Ruthenium-catalyzed cycloaddition of alkynes and organic azides. *J. Am. Chem. Soc.* **2005**, *127*, 15998–15999.
- (23) Wang, Q.; Cahill, S. M.; Blumenstein, M.; Lawrence, D. S. Self-reporting fluorescent substrates of protein tyrosine kinases. *J. Am. Chem. Soc.* **2006**, *128*, 1808–1809.
- (24) Shultz, B. E.; Misialek, S.; Wu, J.; Tang, J.; Conn, M. T., R.; Wong, L. Kinetics and comparative reactivity of human class I and IIb histone deacetylases. *Biochemistry* **2004**, *43*, 11083–11091.
- (25) Shoemaker, R. H. The NCI60 human tumour cell line anti-cancer drug screen. *Nat. Rev. Cancer* **2006**, *6*, 813–823.



**Selective c-Src
kinase inhibitor**



HDAC inhibitor



**Chimeric c-Src and
HDAC inhibitor**