J.C.S. Снем. Сомм., 1980

A Simple, Stereocontrolled Synthesis of a Thromboxane B₂ Synthon

By Alan G Kelly and James S Roberts*

(Department of Chemistry, University of Stirling, Stirling FK9 4LA)

Summary A short stereocontrolled synthesis of a key intermediate in thromboxane B_2 synthesis has been accomplished starting from the carbohydrate laevo-glucosan

sis of starch ⁴ This compound has several advantages including cost, absolute configuration, and chemical reactivity arising from its rigid bicyclic framework and the three axial hydroxy groups In the first step, laevoglucosan was converted into the known epoxy-tosylate (**3**)⁵ which, on treatment with allylmagnesium chloride in the presence of a catalytic amount of cuprous iodide⁶ [inverse addition, 6 equiv CH₂=CHCH₂MgCl, 0 l equiv CuI, dry tetrahydrofuran (1HF), 0 °C, 20 h], gave the allyl derivative (**4**),† m p. 65—67 °C, $[\alpha]_{2^{D}}^{2^{D}} = -58^{\circ}$ (*c*, 0·9, CHCl₃), in 88% yield There is ample literature precedent for the stereo- and regiospecific opening of this epoxide ⁴ Treatment of (**4**) with lithium triethylborohydride⁷ (3 equiv LiEt₃BH, dry THF,

In recent years an impressive number of natural products have been synthesised from carbohydrate precursors ¹ We report a simple, carbohydrate-based synthesis of the key intermediate (11) which has been used previously in thromboxane B_2 syntheses ²,³

The starting material, laevoglucosan (1,6-anhydro- β -D-glucopyranose) (2), is readily available from a wide variety of carbohydrate sources, for example, by controlled pyroly-

[†] A'l new compounds had satisfactory spectral and analytical data

room temperature, 8 h) followed by tosylation of the resultant alcohol (oil, b p ca 85 °C/04 mmHg, $[\alpha]_{\rm p}^{25} =$ -88° (c, 0.7, CHCl₃)} produced (5), † m p 100-100.5 °C, $[\alpha]_{\mathbf{p}}^{25} = -70.0^{\circ}$ (c, 1.0, CH₂Cl₂), in 51% yield



229

Previously we had demonstrated⁸ that lithium triethylborohydride reduction of the di-tosylate (6) gives a high yield of the two alcohols (7) and (8) in the ratio of 5.25:1The mechanism of this reaction does not involve direct displacement of the two tosylate groups but proceeds by way of intermediate formation of (3) followed by hydride attack at C-4 to produce (9) and subsequent hydride attack at C-2 to give predominantly (7) In the case of (4) the 4α -allyl group serves to make the reduction completely regiospecific such that only the 3β -alcohol is obtained with no trace of the isomeric 2β -alcohol

Oxidation of (5) with ruthenium dioxide-excess of sodium metaperiodate in aqueous acetone⁹ (room temperature, 2.5 h) gave directly the highly crystalline tricyclic lactone (10)[†] in 80% yield, m p 151–151.5 °C, $[\alpha]_{D}^{25} = -6.4^{\circ}$ (c, 03, CH₂Cl₂) Amberlyst 15 acid resin in methanol proved to be the method of choice for cleavage of the 1,6-anhydro bridge to produce the two known bicyclic lactones (11) and (12) in 81% yield, in the ratio of 1.55:1 The α -(11) and β -isomers (12) could be readily separated by flash column chromatography on silica gel¹⁰ (Et₂O-MeOH, 9:1) to give pure samples of each. Although pure, the α -isomer (11) was obtained as an oil⁺ $[\alpha]_D^{25} = + 92.7^\circ$ (c, 1.53, CHCl₃) (lit $+86.6^{\circ},^{2b}+100^{\circ},^{2c}+85^{\circ},^{2d}$ and this proved to be identical to an authentic sample, by 1 r and n m r spectroscopy as well as tlc mobility In addition, this isomer was converted quantitatively into the *p*-phenylbenzoate derivative, m p 154—155 °C, $[\alpha]_{\rm D}^{25} = +50.3^{\circ}$ (c, 0.6, CHCl₃) (lit, ²⁰ m p 149—150 °C, $[\alpha]_{\rm D}^{20} = +48.6^{\circ}$ The β -isomer is a crystalline compound, m p 125–126 °C, $[\alpha]_D^{25} = -99^\circ$ (c, 0.42,CHCl₃) and has identical i r and n m r spectra to those reported 3a,b The α -isomer (11)^{3a,b} and its t-butyldiphenvlsilyl derivative^{2a} have been converted by standard 'prostaglandin' methodology into thromboxane B₂ and its C-15 epimer Apart from its simplicity, this synthesis offers the additional potential advantage of permitting the stereospecific introduction of other C-4 side chains by organometallic-induced opening of the epoxide (3) with subsequent inversion of the C-3 configuration [e g], by superoxide displacement of the tosylate group in a derivative of (5)]

We thank the Northern Ireland Ministry of Education (AGK) for a grant, Drs N A Nelson and R C Kelly, The Upjohn Company, Kalamazoo, for an authentic sample of (\pm) -(11), together with 1r and nmr spectra of (11) and (12), and Professor M Černý, Charles University, Prague, for a generous gift, at one stage, of laevoglucosan

(Received, 10th December 1979, Com 1280.)

Two references (ref 2c, 3b) state that this compound is an oil, while two other references (ref 2b, 2d) give a m p of 101-101-5 and 102-103 °C, respectively

¹S Hanessian, Acc Chem Res , 1979, 12, 159

² (a) S Hanessian and P Lavallée, Can J Chem, 1977, 55, 562, (b) E J Corey, M Shibasaki, and J Knolle, Tetrahedron Lett, 1977, 1625, (c) H Ohrui and S Emoto, Agric Biol Chem, 1977, 41, 1773, (d) O Hernandez, Tetrahedron Lett, 1978, 219

³ For non-carbohydrate-based syntheses, see (a) N A Nelson and R W Jackson, *Tetrahedron Lett*, 1976, 3275, (b) R C Kelly, I Schletter, and S J Stein, *ibid*, p 3279, (c) W P Schneider and R A Morge, *ibid*, p 3283, (d) E J Corey, M Shibasaki, J Knolle, and T Sugahara, *ibid*, 1977, 785

For an excellent review of 1,6-anhydro derivatives of aldohexoses, see M Černý and J Staněk, Jr, Adv Carbohydr Chem Biochem, 1977, 34, 23

- ⁵L J Carlson, J Org Chem, 1965, **30**, 3953 ⁶C Huynh, F Derguini-Boumechal, and G Linstrumelle, Tetrahedron Lett, 1979, 1503
- ⁷S Krishnamurthy and H C Brown, J Org Chem, 1976, 41, 3064
- ⁸ A G Kelly and J S Roberts, Carbohydr Res, 1979, **77**, 231 ⁹ Cf G Stork and A R Schoofs, J Am Chem Soc, 1979, **101**, 5081
- ¹⁰ W C Still, M Kahn, and A Mitra, J Org Chem, 1978, 43, 2923