# **Catalytic Asymmetric Synthesis of α-Quaternary Proline Derivatives** by 1,3-Dipolar Cycloaddition of α-Silylimines\*\*

Jorge Hernández-Toribio, Silvia Padilla, Javier Adrio,\* and Juan C. Carretero\*

#### In memory of Christian G. Claessens

Pyrrolidines and derivatives are privileged scaffolds in synthetic and medicinal chemistry, and are present in a myriad of natural products and biologically active compounds.<sup>[1]</sup> In particular, modified proline derivatives have been extensively used as conformationally rigid cores in peptidomimetics.<sup>[2]</sup> In this area, α-quaternary amino acids have received great attention since this type of unit improves the lipophilicity and restricts the conformational flexibility of the peptidic chain, and therefore has a great impact in the biological activity.<sup>[3]</sup> Furthermore,  $\alpha$ -quaternary prolines have found wide application as chiral synthons and organocatalysts in organic synthesis.<sup>[4]</sup> As a consequence, a variety of strategies for their enantioselective preparation, mainly based on the funtionalization of L-proline, have been reported.<sup>[5]</sup> Despite this progress, the development of efficient asymmetric methodologies to access densely substituted aquaternary prolines are still in high demand.

The catalytic asymmetric 1,3-dipolar cycloaddition of azomethine ylides with activated olefins is one of the most reliable and straightforward approaches to the preparation of optically active highly substituted 2-carboxylate pyrrolidine derivatives. Since the seminal contributions of Zhang and co-workers<sup>[6]</sup> and Jørgensen and co-workers<sup>[7]</sup> in the metal-catalyzed enantioselective preparation of 2,5-disubstituted pyrrolidine derivatives using Schiff bases of amino acid esters as azomethine precursors, outstanding progress has been achieved in this research area.<sup>[8,9]</sup> In this context, and because of its high reactivity, glycinate imines are the most frequently employed azomethine ylide precursors in this reaction.<sup>[10,11]</sup>

 $\alpha$ -Silylimines constitute a different and much less studied type of azomethine ylide precursor, which leads to 5-

 [\*] J. Hernández-Toribio, S. Padilla, Dr. J. Adrio, Prof. Dr. J. C. Carretero Departamento de Química Orgánica, Facultad de Ciencias, Universidad Autónoma de Madrid, Cantoblanco 28049 Madrid (Spain)
 E-mail: javier.adrio@uam.es

juancarlos.carretero@uam.es

[\*\*\*] Financial support of this work from the Ministerio de Economía y Competitividad (MINECO, CTQ2009-07791), the Consejería de Educación de la Comunidad de Madrid (project AVANCAT; S2009/ PPQ-1634), and Consejería de Educación de la Comunidad de Madrid, Universidad Autónoma de Madrid (UAM/CAM project CCG-10-UAM/PPQ-5853) is gratefully acknowledged. S.P. thanks the MINECO for a predoctoral contract. We thank the Takasago Company (Dr. Taichiro Touge) for a generous loan of Segphos chiral ligands.

Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.201203828.

unsubstituted pyrrolidines,<sup>[12]</sup> a substitution pattern not accessible by the typical process from Schiff bases of amino acid esters (Scheme 1 a). However, despite this structural interest, as far as we are aware, their use in catalytic asymmetric processes remains undocumented.



**Scheme 1.** 1,3-Dipolar cycloaddition of  $\alpha$ -silylimines.

The great efficiency of Schiff bases of amino acid esters as azomethine ylide precursors relies on the formation of a rigid five-membered N,O-bidentate metalated azomethine (Scheme 1 b). Thus, we reasoned that the presence of a coordinating group in the  $\alpha$ -silylimine, such as an ester moiety, could enhance the reactivity and stereoselectivity of the process by means of the formation of a presumed five-membered metalated intermediate, which would approach the chiral ligand to the reactive center (Scheme 1 c). With the aim of expanding the scope of the pyrrolidine adducts accessible by catalytic asymmetric 1,3-dipolar cycloaddition, we report herein the first enantioselective procedure involving  $\alpha$ silylimines as azomethine precursors to provide 5-unsubstituted quaternary proline derivatives.

To evaluate the viability of the process, we chose as a model reaction the cycloaddition of ethyl-(trimethylsilyl)methyliminopropanoate (1a) with *N*-phenylmaleimide (2;Table 1). We began our investigation by examining the silvercatalyzed reaction in the presence of a variety of chiral ligands in toluene as the solvent. Although good yields and high diastereoselectivities were obtained with the chiral ligands **4**–



🕏 WILEY 🛍

Angewandte ommunications





<sup>[</sup>a] Yield of isolated product. [b] Determined by <sup>1</sup>H NMR spectroscopy. [c] Determined by HPLC. [d]  $CuPF_6 = [Cu(CH_3CN)_4]PF_6$ . [e] Reaction carried out with 1.5 equiv of 1a. [f] 5 mol % of catalysts. TMS = trimethylsilyl.

7, disappointingly, the pyrrolidine adduct **3a** was obtained in racemic form (entries 1-4). Interestingly, a complete inversion in the cis/trans diastereoselectivity and a remarkable 81% ee was observed in the presence of the bulkier and more electron-rich DTBM-Segphos ligand 8 (entry 5). Further optimization of the reaction conditions with this ligand was then undertaken. We observed a substantial enhancement in the enantioselectivity upon employing Cu<sup>I</sup> salts as catalysts (entries 7-9), and in particular [Cu(CH<sub>3</sub>CN)<sub>4</sub>]PF<sub>6</sub> provided the best result (entry 9). The use of other solvents<sup>[13]</sup> or the presence of an external base was less effective. We also confirmed that the presence of 3 equivalents of the silylimine was necessary to obtain good yields (a moderate 40% yield was obtained using 1.5 equivalents of the silvlimine; entry 10). The cycloaddition can be also performed with a lower catalyst loading (5 mol%), albeit with significant erosion in the reactivity and the enantioselectivity (entry 11).<sup>[14]</sup>

With these optimized reaction conditions, we next studied this cycloaddition with a variety of substituted silvlimines. As summarized in Table 2, the reaction worked well for alkylsubstituted  $\alpha$ -silylimines, thus affording the *trans* isomers with nearly complete diastereoselectivity (only the trans isomer was detected by <sup>1</sup>H NMR spectroscopy) and excellent enantioselectivity (entries 1-3). Similarly, the catalytic system was effective with a variety of para-substituted aromatic silvlimines having both electron-donating and electron-withdrawing substituents (entries 4-9), as well as meta-substituted aromatic substrates (entries 10 and 11). The stereochemical **Table 2:** Scope of the 1.3-dipolar cycloaddition of  $\alpha$ -silvlimines with Nphenylmaleimide (2).



[a] Yield of isolated product. [b] Only the trans isomer was detected by <sup>1</sup>H NMR spectroscopy. [c] Determined by HPLC. [d] No cycloaddition product was detected. RT = room temperature.

and configurational assignment of 3g was unequivocally established by X-ray diffraction analysis.<sup>[15]</sup> In contrast, no cycloaddition product was isolated from the sterically more demanding ortho-substituted aromatic silvlimines 1m and 1n (entries 12 and 13). Finally, the unsubstituted silylimine **1p**, derived from ethyl glyoxalate, was also a suitable substrate, albeit proceeding with a lower enantioselectivity (entry 14).

Interestingly, the cycloaddition also proceeded with the nonsymmetrical and more sterically demanding N-phenyl-3methylmaleimide (9). The reaction with silvlimine 1c afforded a 2:1 mixture of regioisomers of the corresponding pyrrolidines bearing two quaternary stererocenters [Scheme 2, Eq. (1)]. The major regioisomer, 10 a, was isolated in 42% yield and excellent enantioselectivity (95% ee), whereas the isomer 10b having two adjacent quaternary





© 2012 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

R R These are not the final page numbers!

www.angewandte.org

2

stereocenters was obtained in 21 % yield and 45 % *ee.* In view of the good reactivity profile of this catalyst system, we next turned our attention to the challenging cyclic silylimines. Pleasingly, the reaction with the  $\alpha$ -silylimine **1q** proceeded smoothly, thus leading to the pyrrolidine spirolactone<sup>[16]</sup> **11** with excellent yield and complete diastereo- and enantiose-lectivity [Scheme 2, Eq. (2)].

Finally, to further explore the scope of this asymmetric [3+2] cycloaddition of  $\alpha$ -silylimines we extended the study to acyclic dipolarophiles. Given the excellent properties of the sulfonyl group both as an electron-withdrawing group and as an easily removed substituent,<sup>[17]</sup> we tested a variety of  $\alpha$ , $\beta$ -unsaturated sulfones as olefin partners (Table 3). The reaction of the silylimines **1a** and **1c** with phenylvinylsulfone (**12**)

**Table 3:** 1,3-Dipolar cycloaddition of  $\alpha$ -silylimines with vinylsulfones.

			[Cu(CH <sub>3</sub> CN) <sub>4</sub> ]PF <sub>6</sub> /				
TMS		2R <sup>2</sup> +	PhO <sub>2</sub> S	_∕ <sup>R³(</sup> 	igand <b>8</b> 10 mol%) IF, RT, 12h	PhO <sub>2</sub> S R <sup>4</sup>	R <sup>1</sup>
R <sup>1</sup> =Me, R <sup>2</sup> =Et (1a)			R <sup>3</sup> =R <sup>4</sup> =H ( <b>12</b> )			H	00 <sub>2</sub> R-
R <sup>1</sup> =Ph(CH <sub>2</sub> ) <sub>2</sub> , R <sup>2</sup> =Et ( <b>1c</b> )			R <sup>3</sup> =H, R <sup>4</sup> =SO <sub>2</sub> Ph ( <b>13</b> )			trans-15–17	
R <sup>1</sup> =Me, R <sup>2</sup> =Me ( <b>1r</b> )			R <sup>3</sup> =CO <sub>2</sub> Me, R <sup>4</sup> =H ( <b>14</b> )				
Entry	R <sup>1</sup>	$\mathbb{R}^2$	R <sup>3</sup>	$R^4$	Product	Yield [%] <sup>[a]</sup>	ee [%] <sup>[b]</sup>
1 <sup>[c]</sup>	Me	Et	Н	н	15 a	48	55
2	Me	Et	н	Н	15 a	75	83
3	$Ph(CH_2)_2$	Et	н	Н	15 c	72	94
4	Me	Et	Н	$SO_2Ph$	16a	70	73
5 <sup>[d]</sup>	Me	Et	Н	$SO_2Ph$	16a	78	79
6	Me	Et	$CO_2Me$	Н	17 a	80	97
7	Ph(CH <sub>2</sub> ) <sub>2</sub>	Et	$CO_2Me$	Н	17 c	78	97
8	Me	Me	$\rm CO_2Me$	Н	17 r	89	99

[a] Yield of isolated product. [b] Determined by HPLC. [c] Reaction performed in toluene. [d] (*R*)-MeO-biphep was used as the ligand; see the Supporting Information for the ligand structure.

under the optimized reaction conditions (but using THF as the solvent), afforded the corresponding sulfonyl pyrrolidines 15a and 15c with excellent diastereoselectivity (only the trans diastereomers were detected by <sup>1</sup>H NMR spectroscopy), and 83% and 94% ee, respectively (entries 2 and 3). The reaction of 1a with 1,1-(bissulfonyl)ethylene (13) also proceeded efficiently, thus leading to the adduct 16a in 73% ee (entry 4). In this case, the asymmetric induction was enhanced to 79% ee when the MeO-biphep ligand was used (entry 5). Encouraged by these results we also tested the unsymmetrically disubstituted Z-sulfonyl acrylate 14.<sup>[18]</sup> The [3+2] cycloadditions with the silvlimines 1a, 1c, and 1r, catalyzed by Cu<sup>I</sup>/ DTBM-segphos, took place with nearly complete regioselectivity, diastereoselectivity, and enantioselectivity (entries 6-8). According to the previous results from our research group in other 1,3-dipolar cycloadditions of Z-sulfonyl acrylates, the regioselectivity was controlled by the sulfonyl group.<sup>[19]</sup> We also tested the cycloaddition of the cyclic  $\alpha$ -silylimine **1** q with dipolarophile 14, thereby providing the pyrrolidine spirolactone **17q** as a single isomer in 81% *ee* (Scheme 3).

The usefulness of sulfonylpyrrolidines to provide  $\alpha$ quaternary proline derivatives was next demonstrated by the desulfonylation reactions shown in Scheme 4. The pyrro-



Scheme 3.



**Scheme 4.** Desulfonylation reactions. DMAP = 4-dimethylaminopyridine

lidines **19** and **20** were obtained by reductive cleavage of the sulfonyl group without any detectable epimerization by treatment with Na(Hg). It is interesting to note that the 2,3-dicarboxylate ester pyrrolidine **20** displays the opposite regioselectivity to that obtained using typical acrylate dipolar-ophiles.<sup>[20]</sup> In contrast, the basic elimination of the sulfonyl group of **17r** provided the pyrroline **21** selectively.

To gain some insight into the reaction mechanism of this cycloaddition, some experiments were conducted. First, the reaction of the silylimine **1a** with *N*-phenylmaleimide (**2**) catalyzed by  $[Cu(CH_3CN)_4]PF_6/DTBM-Segphos was monitored by ESI-MS (Scheme 5; R = Me). After 10 minutes we were able to detect the presumed ions corresponding to the Cu/DTBM-Segphos complex$ **22**(*m*/*z*1241.6), Cu/DTBM-Segphos/desilylated azomethine complex**23**(*m*/*z*1370.7), and



Scheme 5. Proposed mechanism.

© 2012 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

einheim www.angewandte.org These are not the final page numbers!

## Angewandte Communications

Cu/DTBM-Segphos/cycloaddition product 24 (m/z 1543.7; see the Supporting Information for details). These data suggest that the desilylation step, presumably promoted by the excess of imine substrate, takes place before the cycloaddition reaction. Accordingly, when silyl scavengers such as pyridine, 2,6-lutidine or CsF were used as additives the amount of the starting sylilimine 1 could be reduced from 3 to 1.5 equivalents to achieve full conversion.<sup>[21]</sup> On the basis of these data, a simplified plausible mechanism is shown in Scheme 5. Coordination of the silvlimine 1 to the chiral DTBM-Segphos/Cu<sup>I</sup> catalyst 22 and desilylation (promoted by the excess of starting imine 1 or an external base such as pyridine, 2,6-lutidine, or CsF) would lead to the key metalbound azomethine ylide dipole 23, whose subsequent cycloaddition with 2 would provide the pyrrolidine product and the recovery of the catalyst 22.

In summary, an efficient catalytic asymmetric protocol for the Cu<sup>I</sup>-catalyzed 1,3-dipolar cycloaddition of  $\alpha$ -silylimines and activated olefins has been described. A variety of  $\alpha$ quaternary proline derivatives were obtained with good yields and excellent levels of diastereoselectivity and enantiocontrol with a variety of dipolarophiles, such as maleimides (up to 99% *ee*) and  $\alpha$ , $\beta$ -unsaturated sulfones (up to 99% *ee*). The use of the bulky DTBM-Segphos ligand proved to be crucial for attaining this high enantioselectivity.

Received: May 17, 2012 Revised: June 28, 2012 Published online:

**Keywords:** asymmetric catalysis · azomethine ylides · copper · cycloaddition · silanes

- For reviews, see: a) J. P. Michael, *Nat. Prod. Rep.* 2008, 25, 139;
  b) S. G. Pyne, A. S. Davis, N. J. Gates, K. B. Lindsay, T. Machan, M. Tang, *Synlett* 2004, 2670; c) Y. Cheng, Z.-T. Huang, M.-X. Wang, *Curr. Org. Chem.* 2004, 8, 325.
- [2] For recent examples, see: a) B. Song, M. G. Bomar, P. Kibler, K. Kodukula, A. K. Galande, *Org. Lett.* 2012, *14*, 732; b) L. R. Whitby, Y. Ando, V. Setola, P. K. Vogt, B. L. Roth, D. L. Boger, *J. Am. Chem. Soc.* 2011, *133*, 10184.
- [3] a) P. Maity, B. König, *Biopolymers* 2008, 90, 8; b) T. Kan, Y. Kawamoto, T. Asakawa, T. Furuta, T. Fukuyama, *Org. Lett.* 2008, 10, 169; c) S. Pizzarello, *Acc. Chem. Res.* 2006, 39, 231.
- [4] For reviews, see: a) C. Grondal, M. Jeanty, D. Enders, *Nat. Chem.* 2010, 2, 167; b) *Proc. Natl. Acad. Sci. USA* 2010, 107, special feature issue on organocatalysis.
- [5] For a review, see: a) M. I. Calaza, C. C. Cativiela, *Eur. J. Org. Chem.* 2008, 3427. For recent references, see: b) P. Tuzina, P. Somfai, *Org. Lett.* 2009, 11, 919; c) C. Caupène, G. Chaume, L. Ricard, T. Brigaud, *Org. Lett.* 2009, 11, 209.
- [6] J. M. Longmire, B. Wang, X. Zhang, J. Am. Chem. Soc. 2002, 124, 13400.
- [7] A. S. Gothelf, K. V. Gothelf, R. J. Hazell, K. A. Jørgensen, Angew. Chem. 2002, 114, 4410; Angew. Chem. Int. Ed. 2002, 41, 4236.
- [8] For recent reviews, see: a) J. Adrio, J. C. Carretero, *Chem. Commun.* 2011, 47, 6784; b) C. Nájera, J. M. Sansano, *Top. Heterocycl. Chem.* 2008, 12, 117; c) L. M. Stanley, M. P. Sibi, *Chem. Rev.* 2008, 108, 2887; d) H. Pellissier, *Tetrahedron* 2007, 63, 3235; e) G. Pandey, P. Banerjee, S. R. Gadre, *Chem. Rev.* 2006, 106, 4484; f) T. M. V. D. Pinho e Melo, *Eur. J. Org. Chem.*

**2006**, 2873; g) M. Bonin, A. Chauveau, L. Micouin, *Synlett* **2006**, 2349; h) S. Husinec, V. Savic, *Tetrahedron: Asymmetry* **2005**, *16*, 2047; i) I. Coldham, R. Hufton, *Chem. Rev.* **2005**, *105*, 2765; j) C. Nájera, J. M. Sansano, *Angew. Chem.* **2005**, *117*, 6428; *Angew. Chem. Int. Ed.* **2005**, *44*, 6272.

- [9] For selected recent references, see: copper catalysts: a) M. Potowski, M. Schürmann, H. Preut, A.P. Antonchick, H. Waldmann, Nat. Chem. Biol. 2012, 8, 428; b) E. Conde, D. Bello, A. de Cózar, M. Sánchez, M. A. Vázquez, F. P. Cossío, Chem. Sci. 2012, 3, 1486; c) M. González-Esguevillas, J. Adrio, J. C. Carretero, Chem. Commun. 2012, 48, 2149; silver catalysts: d) M.-C. Tong, J. Li, H.-Y. Tao, Y.-X. Li, C.-J. Wang, Chem. Eur. J. 2011, 17, 12922; e) T.-L. Liu, Z.-L. He, Q.-H. Li, H.-Y. Tao, C.-J. Wang, Adv. Synth. Catal. 2011, 353, 1713; f) Y. Yamashita, T. Imaizumi, S. Kobayashi, Angew. Chem. 2011, 123, 4995; Angew. Chem. Int. Ed. 2011, 50, 4893; g) Z.-Y. Xue, X. Fanga, C.-J. Wang, Org. Biomol. Chem. 2011, 9, 3622; nickel catalysts: h) T. Arai, N. Yokoyama, A. Mishiro, H. Sato, Angew. Chem. 2010, 122, 8067; Angew. Chem. Int. Ed. 2010, 49, 7895; calcium catalysts: i) T. Tsubogo, S. Saito, K. Seki, Y. Yamashita, S. Kobayashi, J. Am. Chem. Soc. 2008, 130, 13321; iron catalysts: i) H. Wu, B. Wang, H. Liu, L. Wang, Tetrahedron 2011, 67, 1210; gold catalysts: k) M. Martín-Rodríguez, C. Nájera, J. M. Sansano, A. de Cózar, F. P. Cossío, Chem. Eur. J. 2011, 17, 14224.
- [10] For a systematic study on other types of Schiff bases of amino acid esters, other than glycinates, see C. Nájera, M. de Gracia Retamosa, M. J. Sansano, *Angew. Chem.* 2008, *120*, 6144; *Angew. Chem. Int. Ed.* 2008, *47*, 6055. See also Ref. [8].
- [11] Other types of azomethine ylide precursors have been recently applied in catalytic asymmetric processes. Azlactones: a) A. D. Melhado, M. Luparia, F. D. Toste, J. Am. Chem. Soc. 2007, 129, 12638. α-iminophosphonates: b) Y. Yamashita, X.-X. Guo, R. Takashita, S. Kobayashi, J. Am. Chem. Soc. 2010, 132, 3262. α-iminonitriles: c) R. Robles-Machín, I. Alonso, J. Adrio, J. C. Carretero, Chem. Eur. J. 2010, 16, 5286. N-(2-pyridylmethyl)imines: d) S. Padilla, R. Tejero, J. Adrio, J. C. Carretero, Org. Lett. 2010, 12, 5608.
- [12] a) K. Achiwa, M. Sekiya, *Tetrahedron Lett.* 1982, 23, 2589; b) N. Imai, Y. Terao, K. Achiwa, *Chem. Pharm. Bull.* 1987, 35, 2085; c) O. Tsuge, S. Kanemasa, A. Hatada, K. Matsuda, *Bull. Chem. Soc. Jpn.* 1986, 59, 2537; d) M. Komatsu, H. Okada, T. Akaki, Y. Oderaotoshi, S. Minakata, *Org. Lett.* 2002, 4, 3505.
- [13] Poorer results were obtained with other solvents such as THF (65% conv., 12 h, 91% ee), α,α,α-trifluorotoluene (50% conv., 12 h, 90% ee), and CH<sub>2</sub>Cl<sub>2</sub> (85% conv., 12 h, 88% ee).
- [14] No cycloaddition was observed under the optimal reaction conditions with the phenyl-substituted  $\alpha$ -silylmine **1s**, thus showing the key activating role of the ester moiety.



- [15] CCDC 846895 (3g) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www. ccdc.cam.ac.uk/data\_request/cif.
- [16] Pyrrolidine spirolactones have been used as intermediates in natural product synthesis. For instance, see: a) M. Sun, H. Lu, Y.-Z. Wang, H. Yang, H.-M. Liu, J. Org. Chem. 2009, 74, 2213; b) M. H. Becker, P. Chua, R. Downham, C. J. Douglas, N. K. Garg, S. Hiebert, S. Jaroch, R. T. Matsuoka, J. A. Middleton, F. W. Ng, L. E. Overman, J. Am. Chem. Soc. 2007, 129, 11987.

www.angewandte.org

© 2012 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

These are not the final page numbers!

- [17] a) N. S. Simpkins, *Sulfones in Organic Synthesis*, Pergamon Press, Oxford, **1993**. For a review on desulfonylation reactions, see: b) C. Nájera, M. Yus, *Tetrahedron* **1999**, *55*, 10547.
- [18] In contrast, no cycloaddition was observed when typical *trans*substituted dipolarophiles such as dimethyl fumarate or βnitrostyrene were used as dipolarophiles.
- [19] A. López-Pérez, J. Adrio, J. C. Carretero, Angew. Chem. 2009, 121, 346; Angew. Chem. Int. Ed. 2009, 48, 340.
- [20] This pyrrolidine substitution pattern is found in natural and biologically active compounds. For selected references, see: a) K. Shimamoto, Y. Shigeri, *Cent. Nerv. Syst. Agents Med.*

*Chem.* **2006**, *6*, 59; b) H. Bräuner-Osborne, L. Bunch, N. Chopin, F. Couty, G. Evano, A. A. Jensen, M. Kusk, B. Nielsen, N. Rabaso, *Org. Biomol. Chem.* **2005**, *3*, 3926; c) J. M. Ndungu, J. P. Cain, P. Davis, S.-W. Ma, T. W. Vanderah, J. Lai, F. Porreca, V. J. Hruby, *Tetrahedron Lett.* **2006**, *47*, 2233.

[21] Compared with the standard reaction conditions involving 3 equivalents of silyl imine, in the presence of pyridine, 2,6-lutidine, or CsF desilylating agents we observed an erosion in the enantioselectivity of the process (91%, 90%, and 51% *ee*, respectively).



## **Communications**



### Asymmetric Catalysis

J. Hernández-Toribio, S. Padilla, J. Adrio,\* J. C. Carretero\* \_\_\_\_\_\_ IIII--IIII

Catalytic Asymmetric Synthesis of  $\alpha$ -Quaternary Proline Derivatives by 1,3-Dipolar Cycloaddition of  $\alpha$ -Silylimines



**Going pro**: The title reaction between  $\alpha$ silylimines and activated olefins proceeds in the presence of a Cu<sup>1</sup>/DTBM-Segphos catalyst system with excellent diastereoselectivity and enantioselectivity. This process provides straightforward access



to highly enantioenriched 5-unsubstituted  $\alpha$ -quaternary proline derivatives. TMS = trimethylsilyl, DTBM-Segphos = 5,5'-bis[di(3,5-di-*tert*-butyl-4-methoxyphenyl)phosphino]-4,4'-bi-1,3-benzodioxole.

These are not the final page numbers!