

CHEMISTRY A European Journal



WILEY-VCH

Accepted Article Title: Boron-nitrogen-doped nanographenes: a synthetic tale from borazine precursors. Authors: Jacopo Dosso, Tommaso Battisti, Benjamin D Ward, Nicola Demitri, Colan Hughes, Andrew P. Williams, Kenneth D.M Harris, and Davide Bonifazi This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article. To be cited as: Chem. Eur. J. 10.1002/chem.201905794 Link to VoR: http://dx.doi.org/10.1002/chem.201905794 **Supported by** ACES

- Chemistry - A European Journal -

- Full Paper -

Boron-nitrogen-doped nanographenes: a synthetic tale from borazine precursors.

Jacopo Dosso, Tommaso Battisti, Benjamin D. Ward, Nicola Demitri, Colan E. Hughes, P. Andrew Williams, Kenneth D. M. Harris, and Davide Bonifazi*

- [a] J. Dosso, T. Battisti, Dr. B. D. Ward, Dr. C. E. Hughes, Dr. P.
 A. Williams, Prof. Dr. K. D. M. Harris, Prof. Dr. D. Bonifazi;
 School of Chemistry, Cardiff University, Park Place, Cardiff
 CF10 3AT (UK). E-mail: bonifazid@cardiff.ac.uk.
- [b] Dr. N. Demitri; Elettra Sincrotrone Trieste, S.S. 14 Km 163.5 in Area Science Park, 34149 Basovizza - Trieste, Italy.
- [**] D.B. gratefully acknowledges Cardiff University and the EU through the MSCA-RISE funding scheme (project INFUSION and PHOTOTRAIN) for the financial support. TB thanks the project ITN-ETN PHOTOTRAIN project for his PhD fellowship. The authors also acknowledge the use of the Advanced Computing (ARCCA) at Cardiff University, and associated support services.

Supporting information for this article is available on the <u>https://onlinelibrary.wiley.com</u> or from the author. Crystallographic data for the structures reported in this paper have been deposited at the Cambridge Crystallographic Data Centre as supplementary publication no. **1963744**, **1963743**, **1963745** and **1973297** for compounds **18**, **7**, **11** and **1** respectively.

Keywords: polycyclic aromatic hydrocarbons, heteroatom doping, BNdoping, borazine, coronenes, gulf-type peripheries, optoelectronics, powder X-ray diffraction.

10.1002/chem.201905794

Abstract. In this paper we report a comprehensive account of our synthetic efforts to prepare borazino-doped hexabenzocoronenes using the Friedel-Crafts-type electrophilic aromatic substitution. We showed that hexafluoro-functionalized aryl-borazines, bearing an ortho fluoride leaving group on each N- and B-aryl rings, could lead to cascade-type electrophilic aromatic substitution events in the yields stepwise C-C bond formation giving higher of borazinocoronenes than those obtained with borazine precursors bearing fluoride leaving groups at the ortho positions of the B-aryl substituents. It is with this pathway that an unprecedented boroxadizine-doped PAH featuring a gulf-type periphery could be isolated, and its structure proved through single crystal X-ray diffraction analysis. Mechanistic studies on the stepwise Friedelcyclisation suggest mechanism Crafts-type that the of the planarization reaction proceeds through extension of the π system. To appraise the doping effect of the boroxadizine unit on the optoelectronic properties of topology-equivalent molecular graphenes, the all-carbon and pyrylium PAH analogues, all featuring a gulf-type periphery, were also prepared. As already shown for the borazino-doped hexabenzocoronene, the replacement of the central benzene ring by its B_3N_2O congener widens the HOMO-LUMO gap and dramatically enhances the fluorescence quantum yield.

Figure to table of contents



Introduction.

The doping of polycyclic aromatic hydrocarbons (PAHs) is currently emerging as one of the most promising approaches to tailor the organic semiconductors.^[1-4] optoelectronic properties of The substitution of C=C bonds with isoelectronic and isostructural dipolar boron-nitrogen (BN)^[5-7] and boron-oxygen (BO)^[8-11] couples is an efficient method to gain control on the HOMO-LUMO band gap without significant structural modifications of the molecular scaffold.^[12,13] The use of polar BN and BO bonds typically widen the HOMO-LUMO gap^[14,15] and imparts dipole-driven self-assembly behaviors in the solid state^[3,16] and on surfaces.^[17-19] Noticeable examples pyrenes,^[16] phenanthrenes,^[20] include BN-doped anthracenes^[21], perylenes^[22,23] and nanographenes.^[9,24-26] In these PAHs, the doping units are isolated BN couples selectively placed either at the peripheries or in the core of aromatic precursors bearing amino groups at specific positions. Another strategy consists in the use of preorganized doping units, in which the BN and BO couples are clustered in six-membered rings and used to replace benzenoid rings. Examples include the long-known inorganic $B_3N_nO_0$ -based six-membered rings displayed in Figure 1: borazine (B_3N_3) , ^[31] boroxadizine (B_3N_2O) , ^[28-30] bordioxazine (B_3NO_2) , ^[28,29] and boroxine (B_3O_3) . ^[27]



Figure 1. Inorganic $B_3N_nO_0$ -based six-membered rings that can be used for doping graphenoid structures.

Amongst all inorganic rings, only the B_3N_3 , B_3N_2O and B_3NO_2 units can be fully integrated into fully planarized nanographene systems. One can hardly fail to notice that the B_3N_3 ring can replace any benzenoic units in a molecular graphene scaffold, whereas the B_3N_2O and B_3NO_2 rings can only be placed at the peripheries. For instance, a bordioxazine-fused PAH displaying catalytic activity in the dehydrative amidation of carboxylic acids and amines has been very recently reported by Shibasaki and Kumagai.^[32] Borazine derivatives have been mainly used as precursors to build BN-doped graphitic-like carbon materials. Important examples include graphenes^[33-35] and polymers.^[36-40] On the other hand, the use of the boroxadizine ring to dope PAHs remains elusive to the best of our knowledge.

Certainly, the most intriguing example in the field is that of hexabenzoborazinocoronene (1), the B_3N_3 -doped congener of hexabenzocoronene (HBC), that can be considered as the smallest borazino-doped nanographene structure (Scheme 1). Bettinger and coworkers firstly reported the thermolytic synthesis of 1, in which the central benzene ring has been replaced by a borazine core (Scheme 1).^[34] In a parallel avenue, our group contributed to the field with the engineering of borazine-doped polyphenylenes^[37,40] and reported the rational synthesis of a soluble derivative of 1, **2**,^[33] molecule through six intramolecular Friedel-Crafts-type reactions^[33,41] using a hexafluoro-functionalized borazine precursor (Schemes 1-2). In our synthetic approach, the fluoride leaving groups are placed at the ortho positions of the B-aryl substituents borazine precursors and the electrophilic aromatic in the substitution reaction led to molecule $\mathbf{2}$ in 5% yield along with a non-cyclic helicoidal side-product 7 (Scheme 2). Capitalizing on the Friedel-Crafts-type electrophilic aromatic substitution, [41] herein we report a comprehensive account of our synthetic efforts to prepare borazino-doped hexabenzocoronenes. Aiming at enhancing the yield of soluble borazinocoronene 2, we reasoned that the hexafluorofunctionalized isomer of molecule 5, bearing an ortho fluoride leaving group on each N- and B-aryl rings, could lead to cascadetype electrophilic aromatic substitution events in the stepwise C-C bond formation. It is during these synthetic investigations that we could isolate the first boroxadizine-doped PAH featuring a gulf-type periphery and prove its structure through single crystal X-ray diffraction analysis. Propelled by these findings, we decided to investigate further the doping effect of the boroxadizine unit and prepared all-carbon and pyrylium^[42-45] PAH analogues, all featuring a gulf-type^[46] periphery. As observed for the case of molecule 2, the replacement of the central benzene ring by its B₃N₂O congener widens the HOMO-LUMO gap and it enhances the fluorescence quantum yield. Xylyl moieties were introduced on all PAHs to favor solubility.

Current synthetic approaches towards borazino-doped Nanographenes:



Smallest borazino-doped Nanographenes:



Y = H or reactive group

Scheme 1. Synthetic planarization protocols for preparing borazino-doped
hexabenzocoronenes 1 and 2.

Results and discussion.

Synthesis and structure determination of BN-doped nanographenes. In

our early report,^[33] we prepared B_3N_3 -doped hexabenzocoronene **2** capitalizing on the Friedel-Crafts reaction using difluoroarenes (route A, Scheme 2).^[41] In this strategy, hexafluoro borazine **5** was obtained after reaction of 4-xylyl aniline **3** with BCl₃ followed by addition of 2,6 difluorophenyl lithium **4** (Scheme 2).^[33] Subsequent planarization with [iPr₃Si-CB₁₁H₆Cl₆] and Me₂SiMes₂ at 110 °C in PhCl gave molecule **2** (5% yield) and partially fused BN-derivative **7** as the major product (17% yield). Further purification of the reaction mixture allowed us to isolate partially fused derivative **A1**. Also,

mass analysis investigations showed that molecular ions corresponding to ${\bf A2}$ were present in the reaction crude (see also below for the hypothesized stepwise cyclization mechanism).



Scheme 2. Routes A and B for preparing BN-doped nanographenes.



Figure 2. a) Front view of 7 with BN distances. b), c) front view and side view with axial enantiomers (M and P highlighted) of 7. Atom colors: pink B, blue N, gray C; space group: $P 2_1/n$.

10.1002/chem.201905794

The structure of partially planarized derivative 7 was further confirmed by single-crystal X-ray analysis (Figure 2), which displayed an helicene-like arrangement of the aromatic rings, with a 37° tilt around a central cove-type periphery. The BN bond lengths for 7 are: 1.414(4)-1.422(3) Å for the BNH couple and 1.448(3)-1.454(4) Å for the NBN fragment, slightly different from those reported for 2 (1.433(3)-1.442(2) Å).

Aiming at improving the reaction yield, we redesigned the borazine precursor and prepared a BN-derivative bearing one fluorine atom on the ortho-position of each ring (route B, scheme 2). In this synthetic route, aniline $\mathbf{8}$ is reacted with BCl₃ in refluxing toluene, followed by the addition of 2-fluorophenyl lithium 9 to give hexafluoro borazine derivative 10 in a 36% yield. Subsequent ringclosure reaction in the presence of Me_2SiMes_2 and $[iPr_3Si-CB_{11}Cl_6H_6]$ at 110°C in degassed PhCl afforded borazino-coronene 2 in 15% yield. While purifying the reaction mixture, an additional product was isolated in a 4% yield. To our surprise the product proved to be B₃N₂O-doped PAH **11** featuring a gulf-type periphery. To the best of our knowledge, this molecule represents the first example of a PAH embedding a B_3N_2O cycle. To validate its chemical structure, crystals suitable for X-ray diffraction were grown, and the resulting structure is displayed in Figure 3. Focusing on the bond lengths in the B_3N_2O ring, the BN distances varies between 1.437(4) Å and 1.441(4) Å, similarly to the values reported for molecule 2.^[33] The BO bond lengths are 1.386(4) and 1.375(4), very close to the distances measured for boric acid and phenyl boroxine.^[47]



Figure 3. Front view of molecule **11** with BN and BO bond lengths. Atom colors: pink B, blue N, red O, gray C; Space group: $P 2_1/c$.

To probe the versatility of route B, we undertook the synthesis of unsubstituted borazinocoronene 1, and prepared hexafluoro borazine

10.1002/chem.201905794

12. As for intermediate 10, each B- and N-aryl rings in precursor 12 bear an *ortho*-fluoride atom (Scheme 2). Friedel-Crafts reaction led to 1 as an insoluble material, whose structure was confirmed by mass-spectrometry in line with the seminal report by Bettinger and co-workers.^[34] Given the insolubility of molecule 1, we have carried out structure determination from powder X-ray diffraction data.

Crystal Structure Determination of 1 from Powder XRD Data. A previous powder XRD study has been reported^[34] for **1**, involving profile-fitting (using the Le Bail technique) of the powder XRD pattern recorded for a two-phase sample comprising 1 together with another crystalline phase arising from one of the reagents involved in the synthesis. The Le Bail fitting showed that the unit cell of 1 closely matches the unit cell in the reported crystal structure^[48,49] of hexa-peri-hexabenzocoronene (which contains a central C₆ ring instead of a B_3N_3 ring). However, the previous powder XRD study^[34] only considered the preliminary profile-fitting stage of the structure determination process, and did not proceed to carry out structure solution and refinement. The powder XRD pattern of 1 prepared in the present work closely matches the one reported previously, $^{\scriptscriptstyle [34]}$ but in our case it is clear from the powder XRD pattern that our sample is a pure phase of $\mathbf{1}$ (as discussed below). This powder XRD pattern was used for structure determination of 1.

Profile fitting and unit cell refinement from the powder XRD data were carried out using the Le Bail technique^[50] implemented in the program GSAS,^[51] starting from a unit cell (a = 12.929 Å, b = 5.119Å, c = 17.9983 Å, $\beta = 108.984^{\circ}$) and space group (P2₁/n) that were derived from the reported^[49] unit cell of the carbon analogue of **1** by transformation to the conventional setting (which has the shortest lengths of the a- and c-axes and the value of β closest to 90°). A good-quality Le Bail fit was achieved (Figure 4a; $R_{wp} = 0.57$ %, $R_p =$ 0.44%), with the following refined unit cell parameters: a =12.921(5) Å, b = 5.0996(7) Å, c = 18.002(5) Å, $\beta = 109.151(16)^{\circ}$. The fact that this unit cell accounts for all peaks observed in the experimental powder XRD data confirms the assignment of the sample as a pure crystalline phase of **1**. From the unit cell volume and density considerations, it is deduced that there are two molecules of **1** in the unit cell, and therefore the asymmetric unit for space group $P2_1/n$ comprises one half molecule of **1**.

Structure solution of **1** was carried out using the direct-space strategy,^[52,53] implemented using a genetic algorithm (GA) in the EAGER.^[54-61] In the direct-space structure solution program calculations (see SI for more details), the molecular model was derived from the molecule in the crystal structure^[49] of the carbon analogue of $\boldsymbol{1},$ but with the central C_6 ring replaced by a central B_3N_3 ring. The asymmetric unit comprised the whole molecule of ${\bf 1}$ with half occupancy of all atoms, and with the centre of the molecule fixed at a crystallographic inversion centre. Clearly, the inversion symmetry operation creates disorder in the positions of the B and N atoms in the central ring, resulting in an occupancy of $B_{0.5}N_{0.5}$ for each atomic site in the central ring. The trial structure giving the best fit to the experimental powder XRD data (i.e., lowest $R_{\rm WP}$) in the structure solution calculations was used as the initial structural model for Rietveld refinement (see SI for more details), which was carried out using the program GSAS.^[51] The final Rietveld refinement produced a good-quality fit (Figure 4b; $R_{\rm WD} = 0.60$ %, $R_{\rm D} =$ 0.45%), comparable to the quality of fit obtained in the Le Bail fitting procedure (Figure 4a), with the following refined unit cell parameters: a = 12.934(5) Å, b = 5.1022(8) Å, c = 18.009(5) Å, $\beta =$ $109.188(17)^{\circ}$, V = 1122.4(7) Å³.



Figure 4. a) Le Bail fitting of the powder XRD pattern of 1, and b) final Rietveld refinement for 1. Red + marks, experimental data (background

subtracted); green line, calculated data; magenta line, difference plot; black tick marks, predicted peak positions.

The crystal structure of 1 (Figure 5a,b), is essentially isostructural to the crystal structure of the carbon analogue, [49] as demonstrated by the overlay of the two structures in Figure 5c. According to the classification of aromatic hydrocarbon crystal structures,^[62] the structure is assigned to the γ packing type. The molecules are arranged in "slabs" parallel to the *ab*-plane; all molecules in a given slab are related by translation and hence have the same orientation. Adjacent slabs are related by the 21 screw axis (parallel to the *b*-axis), and the molecules in adjacent slabs have different orientations. As discussed above, the crystal structure of 1 is disordered, with each atomic site in the central ring represented as $B_{0.5}N_{0.5}$. This representation of the average structure may be readily interpreted in terms of the molecule of 1 adopting two orientations with equal probability (the two orientations differ by 60° rotation about an axis perpendicular to the plane of the molecule). In the average structure, the positions of the C and H atoms for the two orientations occupy the same set of sites.

To assess whether there is any significant energy difference between the two molecular orientations in the disordered (average) structure, periodic DFT-D calculations were carried out for two ordered structures, in each case containing only one of the two molecular orientations. To generate these ordered structures, the crystallographic inversion centre was removed to give space group P21 (with one molecule in the asymmetric unit), and the central ring was an ordered B_3N_3 ring (i.e., with each atomic site occupied by only B or only N, and with the B and N atoms alternating around the ring). The two ordered structures differ by 60° rotation about an axis perpendicular to the plane of the molecule. Each structure was subjected to DFT-D geometry optimization (with fixed unit cell). The resulting geometry optimized structures are very similar (except in regard to the assignment of B and N atoms to the atomic sites of the central ring); the largest displacement between atomic sites in the two structures is 0.111 Å, with an RMSD of 0.072 Å. The energy difference between the two optimized structures is only 4 \times 10⁻⁴ kJ mol⁻¹. This very small energy difference is fully compatible with the existence of disorder involving equal populations of the two molecular orientations in the average crystal structure determined from powder XRD data.



Figure 5. Crystal structure of 1 viewed a) along the *b*-axis and b) along the *a*-axis. In this disordered structure, the occupancy of each atomic site in the central ring is $B_{0.5}N_{0.5}$. The slabs of molecules discussed in the text are parallel to the *ab*-plane, which is horizontal in both (a) and (b). c) Overlay of the crystal structures of 1 (magenta) and the carbon analogue of 1 (cyan). The overlay was constructed by matching the unit cell axes (not shown) of the two structures.

Computational investigations of the stepwise Friedel-Crafts planarization reaction. To rationalize the synthetic outcomes of the Friedel-Crafts reaction and set future bases to improve the reaction protocols for preparing borazine-doped nanographenes, we decided to study the mechanistic details of the planarization reaction. When looking at the structures of side-products 7 and 11, one hardly fails to see that these compounds can only derive from partially cyclized intermediates, in which a substituent is cleaved possibly following a hydrolysis. Similarly, the presence of side-products A1 and A2 suggests that hydrolysable intermediates with a lower degree of annulation are also formed in the reaction mixture. Taken all together, these observations prompted us to hypothesize that the ring-closing Friedel-Crafts reactions occur stepwise (i.e., through six cyclization reactions, labelled here as c_1-c_6). Considering that the transition state for each endothermic cyclization resembles the intermediate more than the precursor, [63] we studied the planarization reactions through computational simulations of the intermediates that are possibly formed in each annulation step. The structures of all postulated intermediates in the Friedel-Crafts reactions were

10.1002/chem.201905794

calculated using density functional theory calculations (with B3LYP hybrid functional,^[64] with a 6-31(d,p) basis set^[65] on all centres) with Gaussian 09.^[66] Solvent THF was incorporated using the polarized continuum model, in which the molecular cavity is defined by the united atom model that incorporates hydrogen into the parent heavy atom. $^{[67]}$ The relative energies (AAG) are provided in Schemes 3 and 4 and are reported relative to the lowest energy isomer in each cyclization subset. To ease the nomenclature of the intermediates, the structures are labelled as 5^n_{ci} and 10^n_{ci} for synthetic routes A and B, respectively, with n standing for the intermediate within the cyclization subset, and ci the cyclization step. For pathway A, it is clear that only an intermediate can be envisaged for cyclization reactions c1 and two for c5, whereas up to four isomers could be formed within cyclization subsets c2-c4 (Scheme 3). Since we used an isomeric mixture of the starting hexafluoro borazine for pathway B, one can envisage a complicated pattern of intermediates, including two for cyclization subsets c1 and c5, eight for c2, fourteen for c3, and seven for c4 (Scheme 4). The fact that the formation of the B₃N₂O-doped product should derive from the hydrolysis of an intermediate formed within the cyclization subset c4, led us to restrict our calculations to the intermediates formed in steps c4and c5. When looking at the relative energies of the intermediates, it is evident that the lowest energy intermediate for each cyclization subset $(5^{a}_{c2}, 5^{a}_{c3} \text{ and } 5^{a}_{c4} \text{ for pathway A; } 10^{a}_{c4} \text{ and } 10^{a}_{ci} \text{ for }$ pathway B) is that displaying the greatest degree of planarity in the central B_3N_3 ring. A convenient method to quantify the planarity of the B_3N_3 ring is to calculate the RMS deviation of the B atoms from a plane defined by the three N atoms. RMSD analysis show that the smallest values (0.083, 0.242 and 0.130) are observed for the lowest energy intermediates 5^{a}_{c2} , 5^{a}_{c3} , and 5^{a}_{c4} (Table S2). Interestingly, when looking at the isolated side-products in route A (Scheme 3), it is clear that products A1, A2, and 7 derive from the hydrolysis of intermediates 5^{a}_{c2} , 5^{a}_{c3} , and 5^{a}_{c4} , respectively, which lie on the lowest energy pathway. These observations suggest the occurrence of a planarity-driven cyclization mechanism. After the first electrophilic substitution leading to a flat phenanthrene-like substructure $(\mathbf{5}_{c1})$, the second favored reaction occurs at one of the phenanthrene periphery, further extending the conjugation of the π -surface $(\mathbf{5^{a}}_{c2})$. The subsequent cyclization reactions (c3-c5) occur stepwise progressively extending the outer π -conjugated rim until the last annulation c6 lead to final hexabenzoborazinocoronene derivative (Scheme 3).



Scheme 3. Step wise Friedel-Crafts-based planarization mechanism according to route A. Calculated Gibbs free energies ($\Delta\Delta G$) for the possible intermediates formed at each cyclization step [B3LYP/6-31G(d,p)].



Scheme 4. Friedel-Crafts-based stepwise planarization mechanism according to route B. In this case, the calculated Gibbs free energies ($\Delta\Delta G$) of the intermediates intervening in the fourth and fifth cyclization steps are considered. The presence of different isomers in steps 1-3 deterred us from tackling the calculation of all steps [B3LYP/6-31G(d,p)].

A similar planarity-driven cyclization mechanism to that proposed for route A, can be envisaged for route B (Scheme 4), in which the lowest energy intermediates for c4 and c5 are intermediates $10^{a}{}_{c4}$ and $10^{a}{}_{c5}.$ One can hardly fail to notice that the $\text{B}_{3}\text{N}_{2}\text{O}\text{-doped}$ **11** possibly derives from the derivative hydrolysis of intermediate 10^{a}_{c4} after replacement of a fluorine with a hydrogen atom. At the present time, further investigations are needed to unravel the possible source of hydride yielding the fluoridehydrogen exchange reaction. Complementary mass-spectrometric analysis of the reaction mixtures depicted the presence of the molecular ion associated to one of the intermediates from c5, further supporting our theoretical investigations and mechanistic hypothesis.

The greater stability of a planar borazine ring is somewhat intuitive, when considering that a planar system has a greater degree of delocalization in the π-system. However, this electronic explanation can be given increased rigor using Natural Bonding Orbital (NBO) analyses of the calculated structures. NBO analysis of the parent borazine 5 indicates two 3-centre, 4electron hyperbonds in the B_3N_3 core, each centered about a NBN unit and described as a combination of a N-centered p orbital and a BN π bond as NB:N \leftrightarrow N:BN (Figure 6, left), thus describing a strongly delocalized structure. The NBO analyses of those species deviate from planarity indicate substantially lessthat delocalized structures. Using 5^{c}_{c2} as a representative example in which the boron atoms deviate most from planarity within the series, the NBO analysis does not find the 3-centre 4-electron hyperbond (a significant indicator of extensive delocalization), but second-order perturbation analysis indicates a number of donor-acceptor interactions that involve the N-based occupied porbitals and the B-based vacant p orbitals, thereby indicating a lesser degree of delocalization (Figure 6, right). In intermediate 5^{c}_{c2} one of the boron atoms lies on the plane defined by the three N atoms (B_{plane} , deviation from plane = 0.001 Å),

whereas the other two lie above and below the plane in equal and opposite directions ($B_{out-of-plane}$, deviation from plane = 0.441 Å); the NB_{plane} p-p interactions are calculated to have a greater strength (84 kcal) than the equivalent NB_{out-of-plane} interactions (73 and 28 kcal), thus giving a rational explanation for the calculated stabilities of the two intermediates.



Figure 6. Left: A 3-centre, 4-electron bond described by NBO analysis of precursor 5; right: NBO-derived donor-acceptor interaction between a fluorine lone pair and an ipso carbon of the aryl group in 5^{c}_{c2} [B3LYP/6-31G(d,p)].

At last, it is noteworthy that the mean deviation of the borazine ring from planarity does not directly correlate with their relative free energies; for example, 5^{c}_{c2} has the greatest deviation from planarity (RMSD = 0.294 Å; $\Delta\Delta G$ = 44.6 kJ mol⁻¹) from the $5^a{}_{c2} - 5^d{}_{c2}$ series but $5^d{}_{c2}$ has the highest relative free energy (RMSD = 0.250 Å; $\Delta\Delta G$ = 48.4 kJ mol⁻¹). The relative free energy of 5^{c}_{c2} is lower than expected owing to an extraneous stabilizing effect relating to the unlinked non-fluorinated aryl group via two donor-acceptor interactions from the proximal fluorine lone pairs to a lone valence (vacant p orbital) on the ipso-carbon of the aryl moiety (Figure 6). These analyses give an overall picture of the BN delocalization being the dominant factor in determining stability, but where comparatively minor stereoelectronic factors can have a significant contribution to the overall stability of intermediate species in the reaction process.

Synthesis of pyrylium and all-carbon gulf-type congeners of the B_3N_2O -doped PAH. Following the unexpected isolation of B_3N_2O -doped PAH 11, we decided to prepare its all-carbon (18) and pyrylium (26) gulf-type analogues and evaluate the effect of doping of the ring. Our syntheses central benzenoid started with the preparation of 18 (Scheme 5). The synthetic path started with a trimerization reaction of a 1:2 mixture of acetophenone and 2'bromoacetophenone in the presence of a catalytic amount of TfOH at 130 °C. The reaction produced an inseparable mixture of bromoterphenyl derivatives (13), that was directly used in a Suzuki cross coupling with 14. Subsequently, the isomeric mixture was demethylated using BBr_3 in CH_2Cl_2 , yielding a combination of hydroxyl derivatives that could be separated using silica gel column chromatography to give targeted molecule 15 in 23% yield (over 2 steps). Reaction of 15 with $(Tf)_2O$ in pyridine afforded triflate 16, which was cross-coupled with (2,6-dimethylphenyl)boronic acid under typical Suzuki conditions to give final polyphenylene derivative 17. Finally, oxidative ring closure reaction was achieved using DDQ and TfOH to give planarized derivative 18 with a 14% yield.



Next, we proceeded with the preparation of the pyrylium gulf-type 6).^[45] 26 (Scheme Suzuki analogue, molecule cross-coupling reaction between (2,6-dimethylphenyl)-boronic acid and 4-iodo-2bromo-toluene gave bromo-aryl derivative 19 in the presence of Ba(OH)₂ as base.^[68] Molecule **19** was then converted to boronic ester 20 (63%) under Miyaura borylation reaction conditions. Successive Pd-catalyzed cross-coupling reaction led to the formation of amide 22 in a 73% yield (Scheme 6). Cyclization reaction triggered by an amide-directed ortho-lithiation and subsequent addition-elimination reaction on the amide group, gave phenanthrol 23 in almost quantitative yield. Aldolic condensation between phenanthrol 23 and benzaldehyde in the presence of HCl yielded xanthene derivative 24 that, by treatment with PbO₂, gave desired xanthenol 25 with a 41% yield (Scheme 6). At last, xanthenol 25 was treated with $HBF_4 \cdot Et_2O$ and the resulting orange precipitate irradiated at 365 nm in a CH₂Cl₂/AcOH solution to give desired pyrylium 26 in high yield (80%).



Scheme 5. Synthetic pathway for preparing full-carbon gulf congener 18.

Scheme 6. Synthetic pathways for preparing gulf pyrylium congener 26.

The structures of all intermediates and targeted compounds were characterized by spectroscopic and spectrometric techniques. Suitable crystals for X-ray diffraction analysis for the all-carbon structure 18 were obtained (Figure 7d). No crystals for pyrylium 26 could be formed despite numerous attempts. To better understand the differences in the charge distribution between 11 and 18, ESP surfaces were plotted (Figure 7a-7b). Molecule 18 shows an extended π -surface with low electrostatic potential, instead the presence of the inorganic ring in congener 11 induces a relatively high potential region with a strong polarization, with positively charged B atoms and negatively charged N and O atoms (Figures 7a-b). Interestingly, the aromatic sextets in 11 depict considerably negative electrostatic potentials compared to those in 18.



Figure 7. ESP surfaces mapped on the vdW surface up to an electron density of 0.05 electron/bohr⁻³ for a) **11** and b) **18**. Side (up) and top (bottom) view of the packing of c) **11** and d) **18**. Atom colors: pink B, blue N, red O, gray C. Space groups: $P 2_1/c$ and P-1.

The X-ray crystal structures of **11** (Figure 7c) and **18** (Figure 7d) provide further confirmation of the flat nature of both PAHs (average RMSD of the atoms in the aromatic scaffold from their mean plane is 0.13(9) Å for **11** and 0.15(12) Å for **18**). The molecules undergo $\pi \cdots \pi$ stacking interactions, with the molecular units arranged antiparallelly into columns with average distances of 3.52(20) Å and 3.43(16) Å for **18** and **11**, respectively.

NMR properties of the heteroatom-doped nanographenes. To study the effect of doping units on the ring current properties, we compared the ¹H-NMR spectra of PAHs **11**, **18** and **26** (Figure $8b_1-b_3$).

ccepted Manuscri

shift (δ_{Ha}) of the H_a proton, which is 8.82 ppm for **11**, 9.30 ppm and 9.77 ppm for 18 and 26, respectively. To shed further light on the effects of different doping units on the magnetic properties, GIAO-NICS(1) calculations were also performed (Figure 8a). The replacement of the central benzenoid ring with the B_3N_2O cluster causes a strong change in the NICS value from -12 ppm (aromatic) to -0.95 ppm (non-aromatic) in the central region of the molecule. A similar change in the NICS value has also been observed for the non-sextet rings of the Clar resonance structure surrounding the B_3N_2O system, where a shift from an average of -2.68 in 18 to -0.79 ppm in 11 was observed.^[69] This effect seems to confirm the enhancement of the localization of the aromatic sextets induced by the presence of the central B_3N_2O ring.^[12] Molecule 26 depicts intermediate magnetic properties between those observed for 11 and 18, with a -6.5 ppm shift for the central ring (Figure 8a, molecule 26), whereas the other allcarbon rings feature an overall magnetic anisotropy similar to that calculated for 18 (Figure 8a). ACID calculations were also performed for the derivatives (Figure S74), revealing to be in full agreement with the NICS(1) predictions. The experimental data for the $\delta_{\scriptscriptstyle {\rm Ha}}$ of molecules 11 and 18 are fully consistent with both ACID and NICS calculations, while further discussion is required for the $\delta_{ ext{Ha}}$ value observed for the pyrylium congener. If one solely considers the NICS calculations, the $\delta_{\scriptscriptstyle
m Ha}$ value for molecule 26 should fall in 8.87-9.30 ppm range. However, the electron deficient character of the carbon atom attached to H_a proton further deshields the latter, leading to the highest experimental $\delta_{\scriptscriptstyle {
m Ha}}$ value in the series.

As the diagnostic proton resonance, we selected the chemical



Figure 8. a) NICS(1)-GIAO surfaces for 11, 18, and 26 calculated from the B3LYP/6-31+G** optimized geometry in vacuum; b) Detail of the ¹H-NMR spectra (400 MHz, CDCl₃) in the aromatic region for 11 (b₁), 18 (b₂) and 26 (b₃).

As previously observed for molecules **2** and **HBC**,^[33] these data strengthen our hypothesis for which the replacement of an allcarbon fully benzenoid six-membered ring with an inorganic congener, e.g. B_3N_2O and B_3N_3 , reduces the aromatic character of the π -conjugated scaffold. Optoelectronic properties of the heteroatom-doped nanographenes. The steady-state UV-Vis absorption and emission properties of compounds 7, 11, 18 and 26 in CH_2Cl_2 are displayed in Figure 9 and summarized in Table 1. The data for reference compounds HBC and 2 are also reported for comparison purposes.^[33] The absorption envelop for gulf-type molecule 18 appears between 280 and 440 nm, with a very weak lowest-energy α -band centered at 430 nm ($\epsilon < 1000 \text{ cm}^{-1}\text{M}^{-1}$).^[70,71]

Molecule	Absorption	Emission						
	λ [nm] ($arepsilon$ [L mol ⁻¹ cm ⁻¹]) ^{a)}	$\lambda_{ m max, fl} \ [nm]^{ m a)}$	E _{opt} [eV]	$ au_{ m fl}$ [ns]	$arPsi_{fl}$	λ _{max,ph} [nm] ^{e)}	$ au_{ph}$ [s] ^{e)}	$\Delta E_{(H-L)}$ [eV]
11	367 (16250) 362 (11860) 348 (13600) 292 (18500)	371 389 409 430	3.36 ^{b)}	3.37	0.90	_	-	4.23
18	430 (800) 377 (10900) 366 (31300) 332 (70400) 319 (45000) 306 (41500)	408 433 449 458 476	3.11 ^{°)}	<1.0	0.11 ^{d)}	540 584 634	2.7	3.71
26	575 (25027) 539 (9635) 480 (4117) 364 (18116) 343 (23378)	595	2.12 ^{b)}	8.1	0.48	_	_	4.67
7	393 (27730) 373 (17800) 356 (15300) 340 (15700)	403 423 443	3.12 ^{b)}	3.5	0.83	486 522 559	1.45	3.73
HBC	446 (1000) 388 (53800) 358 (159100)	485	2.76	16.4	0.03	570	0.8	3.99
2	375 (24000) 314 (31200)	404	3.29	8.2	0.43	492	4.0	3.51

Table 1. Photophysical data in aerated CH_2Cl_2 and in glassy matrix for molecules 11, 18, 26, 7, HBC and 2.

Accepted Manuscrii

^{a)} Recorded in an aerated CH_2Cl_2 solution at r.t. ^{b)}Calculated from the lowest energy intersection between absorption and emission spectra normalized on the lowest energy peak (E_{opt} [eV] = 1240.5/ λ_{cross} [nm]). ^{c)}Calculated from the average between the energies of the highest emission and lowest absorption peak. ^{d)}9,10-Diphenylanthracene was used as reference standard (0.97 in aerated Cyclohexane).^[1] ^{e)}Recorded at 77K in a 1:1 CH₂Cl₂:CHCl₃ glassy matrix.

Upon replacement of the central benzene ring with a B_3N_2O congener, a significant blue-shift of the absorption bands is observed, and the lowest-energy structured electronic $\pi\pi^*$ transition appears with a maximum at 367 nm ($\varepsilon \approx 16000 \text{ cm}^{-1}\text{M}^{-1}$). As expected, the spectral envelops of both gulf-type molecules **11** and **18** are blue shifted when compared to those of reference benzocoronenes **2** and **HBC** (Figure 9).



Figure 9. Absorption (-), fluorescence (-) in aerated CH_2Cl_2 solutions at r.t. and phosphorescence (•••) at 77 K in a 1:1 $CH_2Cl_2:CHCl_3$ glassy matrix of 2 (λ_{exc} = 315 nm), HBC (λ_{exc} = 355 nm), 11 (λ_{exc} = 265 nm), 7 (λ_{exc} = 350 nm), 18 (λ_{exc} = 265 nm) and 26 (λ_{exc} = 267 nm).

Compound 26 shows a broad unstructured absorption band centered at 575 nm. The substantial redshift of the first electronic

transition of **26** compared to that of the other derivatives is probably ascribable to the presence of the oxonium specie that, being electron deficient, could induce a lowering of the HOMO-LUMO transition energy. Remarkably, the absorption spectrum of helicene 7 closely resembles that of molecule 2. Regarding the emission spectra, boroxadiazine derivative 11 displays the most blue-shifted emission band (λ_{max} = 371 nm, τ_{f1} =3.37 ns, Φ_{f1} = 0.9), followed by 7 (λ_{max} = 403 nm, τ_{f1} = 3.5 ns, Φ_{f1} = 0.83), 18 (λ_{max} = 408 nm, $\tau_{\rm fl}$ < 1 ns, $\phi_{\rm fl}$ = 0.11) and **26** ($\lambda_{\rm max}$ = 595 nm, $\tau_{\rm fl}$ = 8.1 ns, $\Phi_{\rm fl}$ = 0.48). As observed for molecule 2, derivatives 11, 7 and 26 display narrow Stokes shifts (27 meV, 70 meV and 69 meV, respectively), probably due to the rigid molecular scaffold that prevents most of the geometry-changing vibrational relaxations at the excited state. The all-carbon gulf-type derivative 18 displays a low emission quantum yield ($\phi_{f1} = 0.11$), which is, however, three times larger than its parent benzocoronene HBC ($\phi_{\rm fl}$ = 0.03). Interestingly, molecules 11 and 26 show strong emissive properties, with Φ_{f1} values equal to 0.9 and 0.5, respectively. In particular, B_3N_2O -doped nanographene **11** shows a two-fold increase of the \varPhi_{fl} value compared to $\mathbf{2}$. Notably, no phosphorescence was detected for molecules 11 and 26 at 77K, while 18 ($\lambda_{\rm max}$ = 540 nm, $\tau_{\rm ph}$ = 2.7s) and helicene 7 ($\lambda_{\rm max}$ = 486 nm, $\tau_{\rm ph}$ = 1.45 s) depict longlasting phosphorescence emission signals.

Electrochemical studies were carried out by means of cyclic voltammetry (CV). Surprisingly, only irreversible or quasireversible oxidative events were detected, with cathodic peaks ranging between 0.8 and 1.2 V vs Fc⁺/Fc (1.09 V, 1.01 V, 0.757 V, and 0.5 V for **11**, **18**, **7**, and **26**, respectively) at the scan rate of 0.2 V s⁻¹ (Figure S71). HOMO-LUMO profiles and energy gap were calculated in vacuum at the B3LYP/6-31+G** level of theory (Figure 10 and S73). As expected, the HOMO-LUMO profiles suggest that the replacement of the central benzene ring with a B_3N_2O cluster causes a widening of the bandgap, resulting in a strong blue-shift of the absorption and emission properties. Surprisingly, helicene 7 shows a similar HOMO-LUMO gap to that of full-carbon gulf-type nanographene 18. Finally, pyrylium 26 presents the narrowest gap in the series, probably due to the presence of a charge separation state.



Figure 10. Frontier orbital energies estimated from the CV and photophysical data for 11, 7, 18 and 26. HOMO and LUMO orbitals have been plotted over optimized structures calculated in vacuum at the $B3LYP/6-31+G^{**}$ level of theory using Gaussian09/D01.

Conclusions

In conclusion, it was possible to develop a new strategy towards molecule 2 using a borazine bearing one fluorine atom on the *ortho*-position of each ring. This approach leads to higher yields of 2 and to the unexpected formation of unprecedented B_3N_2O doped PAH 11. Mechanistic studies on the stepwise cyclisation for both pathways led to a clearer understanding of the reaction intermediates. A stepwise mechanism proceeding through extension of the π system is highlighted, providing an effective

explanation for the presence of byproducts 7, 11, A1 and A2 in the reaction mixtures. Derivative 11 represents the first example of a B_3N_2O doped PAH and to shed light on the properties arising from this kind of doping, NMR, photophysical and computational studies were performed on 11 and its congeners 18 and 26. These studies underlined for the first time the widening of the HOMO-LUMO gap provided by the substitution of a full-aromatic sextet of the Clar resonance structure with a $B_3N_nO_0$ based ring. NICS(1) studies allowed us to evaluate the effect of doping on aromaticity of this systems, and along with solution NMR, made possible to appreciate the decrease in the magnetic anisotropy arising from borazine based doping of PAHs. Finally, the presence of the gulf region led to an increase in the quantum yield of derivatives 11 and 18 compared to parent compounds 2 and HBC. Given the growing importance of doped PAHs and nanographene systems in science, this study represents an important step towards the synthesis of tailored borazino-doped nanographenes. Moreover, the extensive studies performed on all the obtained derivatives provide an important guideline for the design of precisely doped PAHs presenting tailored properties. This result is, therefore, of fundamental importance in the growing field of organic optoelectronic devices.

Additional Information

The authors declare no conflict of interest.

References

- [1] A. Narita, X. Wang, X. Feng, K. Müllen, Chem. Soc. Rev. 2015, 44, 6616-6643.
- [2] J. Wang, J. Pei, Chin. Chem. Lett. 2016, 27, 1139-1146.

10.1002/chem.201905794

- [3] D. Bonifazi, F. Fasano, M. M. Lorenzo-Garcia, D. Marinelli,
 H. Oubaha, J. Tasseroul, Chem. Commun. 2015, 51, 15222-15236.
- [4] M. Stępień, E. Gońka, M. Żyła, N. Sprutta, Chem. Rev. 2017, 117, 3479-3716.
- [5] M. J. D. Bosdet, W. E. Piers, Can. J. Chem. 2009, 87, 8-29.
- [6] X. Y. Wang, J. Y. Wang, J. Pei, Chem. Eur. J. 2015, 21, 3528-3539.
- [7] Z. Liu, T. B. Marder, Angew. Chem. Int. Ed. 2008, 47, 242-244.
- [8] X. Wang, A. Narita, W. Zhang, X. Feng, J. Am. Chem. Soc. 2016, 138, 9021-9024.
- [9] X. Wang, J. I. Urgel, G. B. Barin, K. Eimre, M. Di Giovannantonio, A. Milani, M. Tommasini, C. A. Pignedoli, P. Ru, X. Feng, R. Fasel, K. Müllen, A. Narita, J. Am. Chem. Soc. 2018, 140, 9104-9107.
- [10] X. Wang, T. Dienel, M. Di Giovannantonio, G. B. Barin, N. Kharche, O. Deniz, J. I. Urgel, R. Widmer, S. Stolz, L. H. De Lima, M. Muntwiler, M. Tommasini, V. Meunier, P. Ruffieux, X. Feng, R. Fasel, K. Müllen, A. Narita, J. Am. Chem. Soc. 2017, 139, 4671-4674.
- [11] M. Numano, N. Nagami, S. Nakatsuka, T. Katayama, Chem. Eur. J. 2016, 22, 11574-11577.
- [12] N. Otero, P. Karamanis, K. E. El-Kelany, M. Rérat, L. Maschio, B. Civalleri, B. Kirtman, J. Phys. Chem. C 2017, 121, 709-722.
- [13] P. Karamanis, N. D. Charistos, M. P. Sigalas, M. Re, J. Phys. Chem. C 2019, 123, 21135-21149.
- [14] H. Helten, Chem. Eur. J. 2016, 22, 12972-12982.

10.1002/chem.201905794

- [15] M. M. Lorenzo-García, D. Bonifazi, Chim. Int. J. Chem. 2017, 71, 550-557.
- [16] M. J. D. Bosdet, W. E. Piers, T. S. Sorensen, M. Parvez, Angew. Chem. Int. Ed. 2007, 46, 4940-4943.
- [17] (a) S. Kervyn, N. Kalashnyk, M. Riello, B. Moreton, J. Tasseroul, J. Wouters, T. S. Jones, A. De Vita, G. Costantini, D. Bonifazi, Angew. Chem. Int. Ed., 2013, 52, 7410-7412; (b) N. Kalashnyk, P. Ganesh Nagaswaran, S. Kervyn, M. Riello, B. Moreton, T. S. Jones, A. De Vita, D. Bonifazi, G. Costantini, Chem. Eur. J. 2014, 20, 11856-11862.
- [18] W. Auwärter, F. Fasano, N. Demitri, D. Bonifazi, M. Garnica, M. Schwarz, Chem. Eur. J. 2018, 24, 9565-9571.
- [19] C. Sánchez-Sánchez, S. Brüller, H. Sachdev, K. Müllen, M. Krieg, H. F. Bettinger, A. Nicolaï, V. Meunier, L. Talirz, R. Fasel, P. Ruffieux, ACS Nano 2015, 9, 9228-9235.
- [20] C. Zhang, L. Zhang, C. Sun, W. Sun, X. Liu, Org. Lett. 2019, 21, 3476-3480.
- [21] J. S. A. Ishibashi, J. L. Marshall, A. Mazie, G. J. Lovinger, B. Li, L. N. Zakharov, A. Dargelos, A. Graciaa, A. Chrostowska, S. Liu, J. Am. Chem. Soc. 2014, 136, 15414-15421.
- [22] a)M. Hirai, N. Tanaka, M. Sakai, S. Yamaguchi, Chem. Rev. 2019, 119, 8291-8331. b) T. Kaehler, M. Bolte, H. Lerner, M. Wagner, Angew. Chem. Int. Ed 2019, 131, 11501-11506.
- [23] M. Müller, S. Behnle, C. Maichle-Mössmer, H. F. Bettinger, Chem. Commun. 2014, 50, 7821-7823.
- [24] D. L. Crossley, R. J. Kahan, S. Endres, A. J. Warner, R. A. Smith, J. Cid, J. J. Dunsford, J. E. Jones, I. Vitorica-Yrezabal, M. J. Ingleson, Chem. Sci. 2017, 8, 7969-7977.

30

- [25] D. T. Yang, T. Nakamura, Z. He, X. Wang, A. Wakamiya, T. Peng, S. Wang, Org. Lett. 2018, 20, 6741-6745.
- [26] (a) X. Wang, F. Zhang, K. S. Schellhammer, P. Machata, F. Ortmann, G. Cuniberti, Y. Fu, J. Hunger, R. Tang, A. A. Popov, R. Berger, K. Müllen, X. Feng, J. Am. Chem. Soc. 2016, 11606-11615; (b) T. Hatakeyama, S. Hashimoto, S. Seki, M. Nakamura, J. Am. Chem. Soc. 2011, 133, 18614-18617; (c) T. Hatakeyama, K. Shiren, K. Nakajima, S. Nomura, S. Nakatsuka, K. Kinoshita, J. Ni, Y. Ono, T. Ikuta, Adv. Mater. 2016, 28, 2777-2781; (d) M. Numano, N. Nagami, S. Nakatsuka, T. Katayama, K. Nakajim, S. Tatsumi, N. Yasuda, T. Hatakeyama Chem. Eur. J. 2016, 22, 11574-11577; (e) K. Matsui, S. Oda, K. Yoshiura, K. Nakajima, N. Yasuda, T. Hatakeyama, J. Am. Chem. Soc. 2018, 140, 1195-1198
- [27] R. Kinney, D. F. Pontz, J. Am. Chem. Soc. 1936, 278, 197.
- [28] R. Oesterle, W. Maringgele, A. Meller, J. Organomet. Chem. 1985, 284, 281-289.
- [29] Y. Kawashima, H. Takeo, C. Matsumura, Inorg. Chem. 1989, 28, 666-669.
- [30] M. Komorowska, K. Niedenzu, W. Weber, Inorg. Chem. 1990, 48, 289-294.
- [31] A. Stock, E. Pohland, Berichte der Dtsch. Chem. Gesellschaft 1926, 59, 2215-2223.
- [32] H. Noda, M. Furutachi, Y. Asada, M. Shibasaki, N. Kumagai, Nat. Chem. 2017, 9, 571-577.
- [33] J. Dosso, J. Tasseroul, F. Fasano, D. Marinelli, N. Biot, A. Fermi, D. Bonifazi, Angew. Chem. Int. Ed. 2017, 56, 4483-4487.
- [34] M. Krieg, F. Reicherter, P. Haiss, M. Ströbele, K. Eichele,M. J. Treanor, R. Schaub, H. F. Bettinger, Angew. Chem. Int.

31

Ed. 2015, 54, 8284-8286.

- [35] C. Chen, K. Guo, Y. Zhu, F. Wang, W. Zhang, H. Qi, Appl. Mater. Interfaces 2019, 11, 33245-33253.
- [36] N. Riensch, A. Deniz, S. Kühl, L. Müller, A. Adams, A. Pich,
 H. Helten, *Polym. Chem.* 2017, *8*, 5255-5446.
- [37] J. Dosso, D. Marinelli, N. Demitri, D. Bonifazi, ACS Omega 2019, 4, 9343-9351.
- [38] S. Behera, K. T. Jackson, H. M. El-Kaderi, T. E. Reich, P. Jena, J. Mater. Chem. 2012, 22, 13524-13528.
- [39] T. E. Reich, K. T. Jackson, S. Li, P. Jena, H. M. El-Kaderi, J. Mater. Chem. 2011, 21, 10629-10632.
- [40] D. Marinelli, F. Fasano, B. Najjari, N. Demitri, D. Bonifazi, J. Am. Chem. Soc. 2017, 139, 5503-5519.
- [41] O. Allemann, S. Duttwyler, P. Romanato, K. K. Baldridge, J. S. Siegel, Science 2011, 332, 574-577.
- [42] O. Anamimoghadam, M. D. Symes, D. Long, S. Sproules, L. Cronin, J. Am. Chem. Soc. 2015, 14944-14951.
- [43] O. Anamimoghadam, M. D. Symes, C. Busche, D. Long, S. T. Caldwell, C. Flors, S. Nonell, L. Cronin, Org. Lett. 2013, 15, 2970-2973.
- [44] F. Pina, M. J. Melo, C. A. T. Laia, A. J. Parola, J. C. Lima, Chem. Soc. Rev. 2012, 869-908.
- [45] D. Wu, W. Pisula, M. C. Haberecht, X. Feng, K. Müllen, Org. Lett. 2009, 11, 5686-5689.
- [46] C. L. Hilton, J. M. Crowfoot, P. Rempala, B. T. King, J. Am. Chem. Soc. 2008, 130, 13392-13399.
- [47] D. K. Straub, J. Chem. Educ. 1995, 72, 494-497.

[48] J. M. Robertson, J. Trotter, J. Chem. Soc. 1961, 1280-1284.

- [49] R. Goddard, M. W. Haenel, W. C. Herndon, C. Kriiger, M. Zander, J. Am. Chem. Soc. 1995, 117, 30-41.
- [50] A. Le Bail, H. Duroy, J. L. Fourquet, Mat. Res. Bull. 1988, 23, 447-452.
- [51] A. C. Larson, R. B. Von Dreele, Los Alamos National Laboratory Report, 2004, LAUR, 86-748.
- [52] K. D. M. Harris, Top Curr. Chem. 2012, 315, 133-178.
- [53] K. D. M. Harris, M. Tremayne, P. Lightfoot, P. G. Bruce, J. Am. Chem. Soc. 1994, 116, 3543-3547.
- [54] K. D. M. Harris, S. Habershon, E. Y. Cheung, R. L. Johnston, Z. Krist. 2004, 219, 838-846.
- [55] B. M. Kariuki, P. Calcagno, D. M. Kenneth, D. Philp, R. L. Johnston, Angew. Chem. Int. Ed 1999, 38, 831-835.
- [56] B. M. Kariuki, K. Psallidas, K. D. M. Harris, R. L. Johnston, R. W. Lancaster, S. E. Staniforth, S. M. Cooper, *Chem. Commun.* 1999, 1677–1678.
- [57] D. Albesa-jove, B. M. Kariuki, S. J. Kitchin, L. Grice, E.Y. Cheung, K. D. M. Harris, ChemPhysChem 2004, 5, 414-418.
- [58] F. Guo, J. Marti, Z. Pan, C. E. Hughes, K. D. M. Harris, J. Phys. Chem. C 2008, 112, 19793-19796.
- [59] P. A. Williams, C. E. Hughes, G. K. Lim, B. M. Kariuki, K.D. M. Harris, Cryst. Growth Des. 2012, 12, 3104-3113.
- [60] C. E. Hughes, G. M. . Reddy, S. Masiero, S. P. Brown, A. P. Williams, K. D. M. Harris, Chem. Sci. 2017, 8, 3971-3979.
- [61] O. Al Rahal, C. E. Hughes, P. A. Williams, A. J. Logsdail,
 Y. Diskin-posner, K. D. M. Harris, Angew. Chem. Int. Ed
 2019, 131, 18964-18968.

33

- [62] R. Desiraju, Gautam, A. Gavezzotti, J. Am. Chem. Soc. 1989, 26, 621-623.
- [63] G. S. Hammond, J. Am. Chem. Soc. 1955, 77, 334-338.
- [64] T. Yanai, D. P. Tew, N. C. Handy, Chem. Phys. Lett. 2004, 393, 51-57.
- [65] W. J. Hehre, R. Ditchfield, J. A. Pople, **1972**, 52, 2257-2561.
- [66] W. C. M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Had, Gaussian 09, Revis. D.01, Gaussian Inc., Wallingford CT 2010.
- [67] J. Tomasi, B. Mennucci, R. Cammi, Chem. Rev. 2005, 105, 2999-3093.
- [68] A. Suzuki, Pure Appl. Chem. 1994, 66, 213-222.
- [69] M. Randic, Chem. Rev. 2003, 3449-3605.
- [70] M. Daigle, A. Picard-Lafond, E. Soligo, J. F. Morin, Angew. Chem. Int. Ed. 2016, 55, 2042-2047.
- [71] C. Kübel, K. Eckhardt, V. Enkelmann, G. Wegner, K. Müllen, J. Mater. Chem. 2000, 10, 879-886.