

# Chemistry A European Journal



Chemistry

## **Accepted Article**

Title: A Hexabenzocoronene-Based Helical Nanographene

Authors: Norbert Jux, Max M. Martin, and Frank Hampel

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: Chem. Eur. J. 10.1002/chem.202001471

Link to VoR: https://doi.org/10.1002/chem.202001471

WILEY-VCH

## COMMUNICATION

## A Hexabenzocoronene-Based Helical Nanographene

Max M. Martin,<sup>[a]</sup> Frank Hampel<sup>[a]</sup> and Norbert Jux\*<sup>[a]</sup>

**Abstract:** A synthetic route towards a novel hexabenzocoronenebased helical nanographene motif was developed. A hexaphenylbenzene precursor was therefore designed, which cannot undergo, due to steric restrictions, a complete planarization reaction. This precursor was transformed under oxidative cyclodehydrogenation conditions to a  $\pi$ -extended [5]helicene, which was fully characterized including X-ray diffraction analysis.

The first preparation of buckminsterfullerenes that was reported 35 years ago (1985)<sup>[1]</sup> marked the beginning of the era of synthetic carbon allotropes.<sup>[2]</sup> In the following years, intense carbon-based material research was conducted that resulted in the discovery of two more members of the carbon family: carbon nanotubes<sup>[3]</sup> in 1991 and graphene<sup>[4]</sup> in 2004. The aim to explore and discover new compositions of pure carbon is still unabated. Only recently the cyclic sp-hybridized cyclo[18]carbon molecule<sup>[5]</sup> was visualized on a surface and confirmed as a member of the carbon allotrope family.<sup>[6]</sup> Additionally, hypothetical carbon-only architectures, such as helical or twisted derivatives of graphene, toroidal carbon nanotubes or Mackay crystals, are subject of ongoing research.<sup>[7]</sup> Important steps towards elusive novel carbon-only materials were already achieved by the synthesis of model compounds that possess key characteristics of new carbon allotropes. Especially, polycyclic aromatic hydrocarbons (PAHs) and nanographenes with a non-planar, twisted<sup>[7-13]</sup> or helical<sup>[14-23]</sup> topology are becoming increasingly popular in the field of carbonbased materials.<sup>[24]</sup> Due to the out-of-plane distortion, a third dimension is introduced to the nanographenes that results in unique properties different to the ones of the planar analogues. Herein, we like to extend the synthetic toolbox of non-planar nanographenes by a simple and efficient route that is based on well-established reactions used for hexa-peri-hexabenzocoronene (HBC) type chemistry. Precise control over the formation of the helical unit should be gained by the help of steric repulsive effects that prevent complete planarization during the final oxidative cyclodehydrogenation reaction step. This strategy and all synthetic details are presented in the following paper.

First, a suitable tolane, which is capable of forming a helical HBCbased product in the upcoming reaction steps, had to be designed. For that, tetra-*tert*-butyl-tolane **4** was selected. Although a synthesis of **4** is already literature known,<sup>[25]</sup> we employed a

 [a] Dr. M. M. Martin, Dr. F. Hampel, Prof. Dr. N. Jux Department of Chemistry and Pharmacy & Interdisciplinary Center for Molecular Materials (ICMM), Organic Chemistry II Friedrich-Alexander-University Erlangen-Nuernberg Nikolaus-Fiebiger-Str. 10, 91058 Erlangen, Germany e-mail: max.martin@fau.de, norbert.jux@fau.de

Supporting information for this article is given via a link at the end of the document.

different approach, which easily allows to prepare gram quantities of tolane 4. With this procedure, no precious reagents or expensive purification techniques, e.g., column chromatography, were necessary. At first, 3,5-di-tert-butylbenzaldehyde was reacted under McMurry conditions<sup>[26]</sup> to stilbene 2 (Scheme 1) using zinc powder as the reducing agent and TiCl<sub>4</sub> as the titanium species. After aqueous workup and recrystallization from CH<sub>2</sub>Cl<sub>2</sub>/MeOH, product 2 was obtained in suitable yields of 69 %. Stilbene 2 was subsequently brominated<sup>[27]</sup> using Br<sub>2</sub>, which yielded the double brominated compound 3 after aqueous workup in almost quantitative yields (95 %). Next, a double HBr elimination<sup>[27]</sup> under strong basic conditions (KOtBu in THF) generated the desired tolane 4, which was isolated after aqueous workup and recrystallization from CH<sub>2</sub>Cl<sub>2</sub>/MeOH in 78 % yield. With tolane 4 and tetracyclone 5 a standard [4+2] Diels-Alder reaction was performed. The reaction proceeded smoothly at 260 °C in the microwave reactor, which yielded hexaphenylbenzene (HPB) 6 after recrystallization in 81 % yield. Finally, an oxidative cyclodehydrogenation reaction was applied to form an extended aromatic  $\pi$ -system. However, due to the steric demand of the tert-butyl groups attached to the phenyl rings drawn in red (Scheme 1), it seemed impossible to form all six C-C bonds as it is usually the case for HBCs and related materials. As a result, the twisted nanographene 7, in which five C-C bonds were formed, was generated under standard reaction conditions (FeCl<sub>3</sub>, CH<sub>3</sub>NO<sub>2</sub>/CH<sub>2</sub>Cl<sub>2</sub>) and isolated in 80 % yield. The reaction proceeded smoothly without any side product formation (followed by TLC).



 $\begin{array}{l} \textbf{Scheme 1. Synthesis of HBC-based [5]helicene 7. a) Zn, TiCl_4, THF, 70 \ ^{\circ}C, \\ 22 \ h; b) Br_2, CHCl_3, rt, 30 \ min; c) KOtBu, THF, 0 \ ^{\circ}C, 20 \ min; d) Ph_2O, 260 \ ^{\circ}C \\ in microwave reactor, 12 \ h; e) FeCl_3, CH_3NO_2, CH_2Cl_2, 0 \ ^{\circ}C, 100 \ min. \end{array}$ 

Product 7 as well as all of the precursor compounds were carefully characterized by NMR spectroscopic and mass spectrometric techniques (for details see SI). X-ray crystallography of 2, 4, 6 and 7 confirmed furthermore the successful synthesis of these compounds and allowed to get an insight into the solid-state behavior. Generally, molecules within this project showed a high tendency to form crystalline structures, as some of the single

### WILEY-VCH

## COMMUNICATION

crystals, e.g., the ones of 7, grew within 24 h. The structures of 2, 4 and 6 are depicted in the supporting information (Figure S1 -S4) and a detailed structural analysis of the target compound 7 is shown in Figure 1. HBC-based [5]helicene 7 crystallizes in a monoclinic crystal system with the space group P2<sub>1</sub>/n. The helical character of 7, which is defined, *i.e.*, by the sum of torsion angles (81.5°) and the interplanar angle (42.2°),<sup>[28]</sup> was unambiguously confirmed by the crystal structure. The helicity values of 7 significantly differ to the ones of previously reported [5]helicenes, e.g., to a carbon[5]helicene (sum of torsion angles: 65.9 - 67.8°; interplanar angle: 47.3 – 51.3°).<sup>[29-30][31]</sup> For more details see Table S6 - S7 in the supporting information. This deviation can be explained by two major structural variations that affect the helical properties of the HBC-based [5]helicene. On the one hand, the rigid aromatic backbone of 7 (Figure 1a, drawn in grey) reduces the flexibility, especially of the inner rings B, C and D, and therefore limits the overall twisting capability of the helix. However, on the other hand, the bulky tert-butyl groups counteract, due to steric repulsion, to the effects of the stiff backbone and are leading to an increased distortion of the helical unit, particularly of the outer rings A and E. The sum of these two effects lead to a helical unit with an unevenly distributed helical character with areas of inand decreased degree of distortion. This is also supported by the large bond length variations of 1.41 - 1.48 Å between the inner helical carbon atoms (Figure 1b). The crystal packing (Figure 1d, 1e) reveals that both enantiomers (M) and (P) are present, therefore forming a racemic crystal. The molecules are, compared to the crystal packing of, for example, hexa-tert-butyl-HBC<sup>[32][33]</sup> relatively loosely packed with distances of  $\geq 5$  Å between the aromatic planes of the molecules. Hence, no  $\pi$ - $\pi$  interactions and only CH $-\pi$  as well as London dispersion interactions were found, which seem to be the driving force for the preferential formation of all shown structures in the crystal. Interestingly, layers of a single chirality are found in the crystal of 7, generating an alternating pattern of (M) and (P) enantiomer layers (compare Figure S5).



**Figure 1.** a) Crystal structure of HBC-based [5]helicene 7. Structure is depicted as ORTEP model with thermal ellipsoids drawn at a 50 % probability level; b) torsion angles and bond lengths of the inner helix; c) determination of the interplanar angle; d), e) packing motif; hydrogen atoms, *tert*-butyl groups and solvent molecules are omitted for clarity. Helicenes with the same chirality are drawn in the same colour. CCDC: 1990061.

The differences in the spectroscopic properties between the HBCbased [5]helicene **7** and a planar reference compound, hexa-*tert*butyl-HBC,<sup>[34]</sup> were studied by UV-vis absorption and emission spectroscopy. The UV-vis absorption spectrum of **7** (Figure 2a) features a red-shifted maximum at 367 nm (p-band) with a significantly decreased molar extinction coefficient compared to the reference HBC. Furthermore, a reduced resolution of the fine structure, due to a general broadening of all absorption bands, was observed. This is also true for the fluorescence spectrum (Figure 2b), which shows, aside from broadened signals, hypsochromically shifted emission maxima at 475 and 506 nm for **7**. Therefore, the implementation of a helical unit into a nanographene has clear effects on the photophysical properties, such as on the position but also on the shape of the respective absorption and emission bands.



Figure 2. a) UV-Vis absorption spectra of HBC-based [5]helicene 7 and reference compound hexa-*tert*-butyl-HBC<sup>[34]</sup> in CH<sub>2</sub>Cl<sub>2</sub>; b) normalized steady state fluorescence emission spectra upon irradiation of the absorption maximum.

To conclude, we've presented a new procedure for the synthesis of a HBC-based twisted nanographene that contains a [5]helical unit. The helicity was introduced with the help of sterically demanding groups that prevent a complete closure to planar PAHs during the oxidative cyclodehydrogenation step. These results complement our previous findings, in which related nitrogen-doped HBC-like [5]helicenes were formed, however, controlled by electronic effects.<sup>[35]</sup> The chiral separation and characterization as well as the usage of the twisted nanographene motif of **7** as a platform for further functionalized hybrid materials, *e.g.*, analogue architectures to our previously reported porphyrin-HBCs,<sup>[36-44]</sup> are currently under investigation.

#### Acknowledgements

Funded by the Deutsche Forschungsgemeinschaft (DFG) – Projektnummer 182849149 – SFB 953. M.M.M. thanks the Fonds der Chemischen Industrie (FCI) and the Graduate School Molecular Science (GSMS) for financial support. We thank David Reger for helpful discussions and for providing us with the reference compound hexa-*tert*-butyl-HBC.

**Keywords**: hexabenzocoronene • nanographene • helicene • polycyclic aromatic hydrocarbons • carbon allotrope

- H. W. Kroto, J. R. Heath, S. C. O'Brien, R. F. Curl, R. E. Smalley, *Nature* 1985, *318*, 162-163.
- [2] A. Hirsch, Nat. Mater. 2010, 9, 868-871.
- [3] S. lijima, Nature 1991, 354, 56-58.
- [4] K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, Y. Zhang, S. V. Dubonos, I. V. Grigorieva, A. A. Firsov, *Science* **2004**, *306*, 666-669.

## WILEY-VCH

## COMMUNICATION

- [5] F. Diederich, Y. Rubin, C. B. Knobler, R. L. Whetten, K. E. Schriver, K. N. Houk, Y. Li, *Science* **1989**, *245*, 1088-1090.
- [6] K. Kaiser, L. M. Scriven, F. Schulz, P. Gawel, L. Gross, H. L. Anderson, *Science* 2019, 365, 1299-1301.
- [7] S. H. Pun, Q. Miao, Acc. Chem. Res. 2018, 51, 1630-1642.
- [8] I. R. Márquez, S. Castro-Fernández, A. Millán, A. G. Campaña, Chem. Commun. 2018, 54, 6705-6718.
- [9] K. Y. Cheung, C. K. Chan, Z. Liu, Q. Miao, Angew. Chem. Int. Ed. 2017, 56, 9003-9007.
- [10] S. H. Pun, C. K. Chan, J. Luo, Z. Liu, Q. Miao, Angew. Chem. Int. Ed. 2018, 57, 1581-1586.
- [11] O. Papaianina, V. A. Akhmetov, A. A. Goryunkov, F. Hampel, F. W. Heinemann, K. Y. Amsharov, *Angew. Chem. Int. Ed.* 2017, *56*, 4834-4838.
- [12] V. Akhmetov, M. Feofanov, S. Troyanov, K. Amsharov, *Chem. Eur. J.* 2019, *25*, 7607-7612.
- [13] J. M. Fernandez-Garcia, P. J. Evans, S. Medina Rivero, I. Fernandez, D. Garcia-Fresnadillo, J. Perles, J. Casado, N. Martin, J. Am. Chem. Soc. 2018, 140, 17188-17196.
- [14] T. Fujikawa, Y. Segawa, K. Itami, J. Am. Chem. Soc. 2015, 137, 7763-7768.
- [15] Y. Nakakuki, T. Hirose, H. Sotome, H. Miyasaka, K. Matsuda, J. Am. Chem. Soc. 2018, 140, 4317-4326.
- [16] D. Reger, P. Haines, F. W. Heinemann, D. M. Guldi, N. Jux, Angew. Chem. Int. Ed. 2018, 57, 5938-5942.
- [17] S. H. Pun, Y. Wang, M. Chu, C. K. Chan, Y. Li, Z. Liu, Q. Miao, J. Am. Chem. Soc. 2019, 141, 9680-9686.
- [18] C. M. Cruz, S. Castro-Fernández, E. Maçôas, J. M. Cuerva, A. Campaña, Angew. Chem. Int. Ed. 2018, 57, 14782-14786.
- [19] A. Campaña, C. Cruz, S. Castro-Fernández, E. Maçôas, A. Millán, Synlett 2019.
- [20] J. Ma, K. Zhang, K. S. Schellhammer, Y. Fu, H. Komber, C. Xu, A. A. Popov, F. Hennersdorf, J. J. Weigand, S. Zhou, W. Pisula, F. Ortmann, R. Berger, J. Liu, X. Feng, *Chem Sci* **2019**, *10*, 4025-4031.
- [21] P. J. Evans, J. Ouyang, L. Favereau, J. Crassous, I. Fernandez, J. Perles, N. Martin, Angew. Chem. Int. Ed. 2018, 57, 6774-6779.
- [22] Y. Wang, Z. Yin, Y. Zhu, J. Gu, Y. Li, J. Wang, Angew. Chem. Int. Ed. 2019, 58, 587-591.
- [23] Y. Zhu, X. Guo, Y. Li, J. Wang, J. Am. Chem. Soc. 2019.

- [24] M. A. Majewski, M. Stepien, Angew. Chem. Int. Ed. 2019, 58, 86-116.
- [25] H.-P. Jacquot de Rouville, R. Garbage, R. E. Cook, A. R. Pujol, A. M. Sirven, G. Rapenne, *Chem. Eur. J.* **2012**, *18*, 3023-3031.
- [26] K. S. Jeong, Y. S. Kim, Y. J. Kim, E. Lee, J. H. Yoon, W. H. Park, Y. W. Park, S. J. Jeon, Z. H. Kim, J. Kim, N. Jeong, *Angew. Chem. Int. Ed.* 2006, *45*, 8134-8138.
- [27] S. Ito, M. Wehmeier, J. D. Brand, C. Kübel, R. Epsch, J. P. Rabe, K. Müllen, *Chem. Eur. J.* **2000**, *6*, 4327-4342.
- [28] Y. Shen, C. F. Chen, Chem. Rev. 2012, 112, 1463-1535.
- [29] R. Kuroda, J. Chem. Soc., Perkin Trans. 2 1982, 789-794.
- [30] A.-C. Bédard, A. Vlassova, A. C. Hernandez-Perez, A. Bessette, G. S. Hanan, M. A. Heuft, S. K. Collins, *Chem. Eur. J.* 2013, *19*, 16295-16302.
  [31] CCDC 933556.
- [32] P. T. Herwig, V. Enkelmann, O. Schmelz, K. Müllen, Chem. Eur. J. 2000, 6. 1834-1839.
- [33] CCDC 142008.
- [34] J. Holzwarth, K. Y. Amsharov, D. I. Sharapa, D. Reger, K. Roshchyna, D. Lungerich, N. Jux, F. Hauke, T. Clark, A. Hirsch, *Angew. Chem. Int. Ed. Engl.* **2017**, *56*, 12184-12190.
- [35] F. Ammon, S. T. Sauer, R. Lippert, D. Lungerich, D. Reger, F. Hampel, N. Jux, Org. Chem. Front. 2017, 4, 861-870.
- [36] J. M. Englert, J. Malig, V. A. Zamolo, A. Hirsch, N. Jux, Chem. Commun. 2013, 49, 4827-4829.
- [37] D. Lungerich, J. F. Hitzenberger, M. Marcia, F. Hampel, T. Drewello, N. Jux, Angew. Chem. Int. Ed. 2014, 53, 12231-12235.
- [38] D. Lungerich, J. F. Hitzenberger, W. Donaubauer, T. Drewello, N. Jux, *Chem. Eur. J.* 2016, 22, 16755-16759.
- [39] D. Lungerich, J. F. Hitzenberger, F. Hampel, T. Drewello, N. Jux, Chem. Eur. J. 2018, 24, 15818-15824.
- [40] M. M. Martin, N. Jux, J. Porphyrins Phthalocyanines 2018, 22, 454-460.
- [41] M. M. Martin, M. Dill, J. Langer, N. Jux, J. Org. Chem. 2019, 84, 1489-1499.
- [42] M. M. Martin, D. Lungerich, P. Haines, F. Hampel, N. Jux, Angew. Chem. Int. Ed. 2019, 58, 8932-8937.
- [43] M. M. Martin, D. Lungerich, F. Hampel, J. Langer, T. K. Ronson, N. Jux, *Chem. Eur. J.* 2019, 25, 15083-15090.
- [44] M. M. Martin, C. Dusold, A. Hirsch, N. Jux, J. Porphyrins Phthalocyanines 2020, 24, 268-277.

## WILEY-VCH

## COMMUNICATION

### **Entry for the Table of Contents**

## COMMUNICATION



A twisted nanographene that is based on the well-established hexabenzocoronene motif is presented. A [5]helical unit was implemented into the nanographene structure with the help of steric repulsive effects. The  $\pi$ -extended helicene was characterized, *i.a.*, by UV-vis absorption and emission spectroscopy as well as X-ray crystallography.

M. M. Martin, F. Hampel, N. Jux\*

Page No. – Page No. A Hexabenzocoronene-Based Helical Nanographene