

# Chemical Science

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: H. Park and D. Lee, *Chem. Sci.*, 2020, DOI: 10.1039/D0SC05067A.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

## ARTICLE

Biomimetic hydrogen-bonding cascade for chemical activation:  
telling a nucleophile from a base

Hyunchang Park and Dongwhan Lee\*

Received 00th January 20xx,  
Accepted 00th January 20xx


DOI: 10.1039/x0xx00000x

Hydrogen bonding-assisted polarization is an effective strategy to promote bond-making and bond-breaking chemical reactions. Taking inspirations from the catalytic triad of serine protease active sites, we have devised a conformationally well-defined benzimidazole platform that can be systematically functionalized to install multiple hydrogen bonding donor (HBD) and acceptor (HBA) pairs in a serial fashion. We found that an increasing number of interdependent and mutually reinforcing HBD–HBA contacts facilitate the bond-forming reaction of fluorescence-quenching aldehyde group with cyanide ion, while suppressing the undesired Brønsted acid–base reaction. The most advanced system, evolved through iterative rule-finding studies, reacts rapidly and selectively with  $\text{CN}^-$  to produce a large ( $> 180$ -fold) enhancement in the fluorescence intensity at  $\lambda_{\text{max}} = 450 \text{ nm}$ .


## Introduction

Hydrogen bonding is a versatile functional motif for chemical structure design. An elaborate arrangement of hydrogen bonding donor (HBD) and acceptor (HBA) units have been exploited for molecular recognition,<sup>1–5</sup> signaling,<sup>6–8</sup> self-assembly,<sup>9–12</sup> and chemical activation.<sup>13–15</sup> Beyond the paradigm of simple HBD–HBA pair (Fig. 1a), a serial array of multiple HBD–HBA pairs can also be constructed in a cascade fashion (Fig. 1b).<sup>16–18</sup> With the amphoteric proton mediator in the middle, electronic polarization at one end automatically increase the donor/acceptor ability of the other end of the extended network.<sup>19–23</sup>

In such cooperative arrangement of paired HBD–HBA units, the induced polarization at the mediator site renders the hydrogen bonds shorter and stronger, when a comparison is made to a single HBD–HBA contact.<sup>19</sup> A prominent example of the cascade hydrogen bonding in action is a class of enzymes having catalytic triad, such as serine protease, canonical esterase, and lactonase.<sup>23–27</sup> At the active site of these enzymes, the carboxylate group of aspartate, imidazole group of histidine, and hydroxyl group of serine constitute an extended hydrogen bonding array (Fig. 2a).



**Fig. 1** Schematic representation of (a) simple HBD–HBA pair, and (b) cascade hydrogen bonding with an amphoteric proton mediator, such as imidazole, between widely separated HBD and HBA to create a larger additive dipole to promote the addition/substitution reaction shown in (c).



**Fig. 2** (a) Chemical structure and X-ray structure of the active site of serine protease (PDB: 1HXE). (b) Chemical structure and DFT (B3LYP-D3/6-31G(d,p)) energy-minimized structure of a synthetic "triad" 1.

Department of Chemistry, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul 08826, Korea. E-mail: [dongwhan@snu.ac.kr](mailto:dongwhan@snu.ac.kr)


† Electronic Supplementary Information (ESI) available: Experimental procedures, characterization data and supplementary figures, including  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra, crystallographic data, FT-IR and HR-MS analysis. See DOI: 10.1039/x0xx00000x



In the biological system, the amphoteric imidazole group functions as a mediator of the cascade hydrogen bonding (Fig. 1b). The hydrogen bonding between aspartate and histidine polarizes the amine N–H group to enhance the basicity of the imine nitrogen, which translates to a stronger N...H–O hydrogen bond between histidine and serine. An unusually downfield-shifted proton resonance of the imidazole N–H proton at ca. 15 ppm reflects tight hydrogen bonding, which polarizes the serine O–H group to enhance its nucleophilicity toward electrophilic substrates.<sup>23</sup> For the enzymes, it would be challenging to control the strength of hydrogen bonds by using limited types of functional groups offered by amino acid residues. As such, the construction of hydrogen bonding array, in which the whole is more than the sum of its parts, is an effective strategy to make the best out of what is naturally available.

## Design Principles

Taking inspirations from the chemical activation strategy of such catalytic triad, we have designed a benzimidazole-based synthetic surrogate **1** (Fig. 2b). In the case of enzymes, the three-dimensional polypeptide scaffold enables the precise positioning and alignment of HBD–HBA units. Instead of using peptides, we employed the rigid  $\pi$ -conjugated backbone of benzimidazole as a minimalist artificial platform to build biomimetic hydrogen bonding network. Through facile functionalization at the 2-, 4-, and 7-position of benzimidazole ring (Fig. 3), various HBD and HBA units can be installed around the imidazole core. Additionally, the rich photophysics of 2-arylbenzimidazole motif offers excellent opportunities to develop these biomimetic small molecules into fluorescent probes.<sup>28,29</sup>



**Fig. 3** (a) Cascade hydrogen bonds built on the 4,7-disubstituted benzimidazole platform. (b) Structural “evolution” from the simplest **2** to the most elaborate **1** by installing an increasing number of HBD and HBA units. (c) Mechanism of covalent capture and fluorescence turn-on response of **1** toward CN<sup>-</sup>.

In this paper, we report the chemistry of a latent fluorophore **1** (Fig. 2 and 3) and its application for the detection of toxic cyanide ion. From the simplest molecular prototype **2**, a systematic increase in the number of HBD–HBA units helps polarize the aldehyde carbonyl group (Fig. 3b). The enhanced electrophilicity of the aldehyde moiety of **1** effectively promotes selective covalent capture of cyanide ion (Fig. 3c), and allows rapid fluorescence turn-on detection. More importantly, comparative studies on **1** and its simpler “site models” **2–4** established that multiple interconnected HBD–HBA pairs help distinguish nucleophile from Brønsted base.

## Results and Discussion


### Synthesis of a Minimalist Prototype

A single HBD–HBA pair was first installed onto the benzimidazole platform to prepare **2** (Scheme 1). To maximize the effects of chemical transformation on the emissive properties, it would be ideal if the functional group that is activated by hydrogen bonding (Fig. 1c) functions also as a fluorescence quencher. For such purpose, an aldehyde group was employed as an HBA unit. Through fast intersystem crossing, an aldehyde group facilitates the quenching of  $\pi$ -conjugated fluorophore to which it is directly attached.<sup>30–33</sup>

The chemical structure of **2** (Fig. 3b) satisfies this minimum requirement of this design concept. As outlined in Scheme 1, the synthesis of **2** involved oxidative condensation of diamine **5** and aldehyde, and subsequent oxidation of the primary alcohol group. Except for the reduction of the thiadiazole precursor, all reactions proceeded in moderate to good yield (> 75%). The poor isolation yield of the intermediate **5** might be due to the instability of the electron-rich diamine functionality. The synthetic modularity in the imidazole ring construction aided structural diversification (vide infra).


### Spectroscopic Studies and Response to Cyanide Ion

Compound **2** responds to cyanide anion by large changes in both UV–vis absorption and fluorescence emission spectra. As shown in Fig. 4a, the addition of cyanide to a solution of **2** in DMSO led to a decrease in the absorption at  $\lambda_{\max} = 345$  nm with the development of a new band at  $\lambda_{\max} = 390$  nm. While **2** is only weakly emissive due to the efficient fluorescence quenching by the aldehyde group, the intrinsic blue emission of benzimidazole ( $\lambda_{\max,em} = 465$  nm) was restored upon addition of cyanide (Fig. 4b).



**Scheme 1** Synthetic route to a simple HBD–HBA prototype.






**Fig. 4** (a) Absorption and (b) emission spectra ( $\lambda_{\text{exc}} = 400$  nm) of **2** (0.100 mM) prior to (black dashed lines), and after treatment with NaCN (25 equiv, blue lines) or  $\text{Et}_3\text{N}$  (25 equiv, gray lines) in DMSO at  $T = 298$  K.

In the control experiments with other chemical species, however, we realized that a large enhancement in the emission intensity also occurs with the addition of Brønsted base, such as  $\text{Et}_3\text{N}$  (Fig. 4b). The excitation spectrum of **2**, obtained in the presence of  $\text{Et}_3\text{N}$ , revealed that absorption at  $\lambda \approx 400$  nm is responsible for the emission at  $\lambda_{\text{max,em}} = 465$  nm (Fig. S1a). This finding implicates that the loss of hydrogen bonding by deprotonation of the acidic benzimidazole proton could also contribute to the fluorescence enhancement, presumably by suppressing the intersystem crossing of aldehyde as the main quenching pathway. In other words, both addition and deprotonation reactions could take place in the reaction of **2** with cyanide, but the fluorescence response is essentially indistinguishable.

To delineate the nature of the chemical reaction that is responsible for the fluorescence enhancement (Fig. 4b), we carried out detailed solution  $^1\text{H}$  NMR spectroscopic studies. At room temperature ( $T = 298$  K), the  $^1\text{H}$  NMR spectrum of **2** in DMSO- $d_6$  (Fig. 5a, top) shows a pair of broadened resonances associated with benzimidazole N–H and aldehyde C–H protons.



**Fig. 5** (a)  $^1\text{H}$  NMR spectra of **2** (4.0 mM) prior to (top) and after (bottom) addition of NaCN (10 equiv) in DMSO- $d_6$ ;  $T = 298$  K. (b) Variable-temperature (VT)  $^1\text{H}$  NMR spectra of **2** (4.0 mM) in DMSO- $d_6$  measured at  $T = 293$ – $363$  K.

In stark contrast, the  $^1\text{H}$  NMR spectrum of a solution of **2** in  $\text{CDCl}_3$  (Fig. S2) shows a well-resolved spectral pattern that is consistent with the chemical structure of **2**, thus ruling out the involvement of impurities in the complicated  $^1\text{H}$  NMR spectrum obtained in DMSO- $d_6$ .

We suspect that the intramolecular N–H $\cdots$ O bond of **2** does not provide sufficient thermodynamic bias toward one tautomer over the other in DMSO- $d_6$  (Fig. 5b and Scheme 2). Indeed, the two distinct resonances of the benzimidazole N–H protons at  $\delta = 13.31$  and  $13.04$  ppm converge into a single averaged feature at  $\delta = 12.93$  ppm as the temperature is increased (Fig. 5b, coalescence at  $T_c = 343$  K). From eq (1), the corresponding rate constant ( $k_c$ ) can be determined by the  $\Delta\nu$  value ( $= 108$  Hz) at the slow exchange limit.<sup>34</sup> By applying the Eyring equation in eq (2), the activation energy of  $\Delta G^\ddagger = 16.4$  kcal mol $^{-1}$  was determined, which is typical for the energy barrier for the tautomerization.<sup>35,36</sup> In this process, the resonances of the aldehyde protons at  $\delta = 10.81$  and  $10.34$  ppm, each associated with the two tautomers, also converge into a single resonance at  $\delta = 10.54$  ppm.


$$k_c = \frac{\pi}{\sqrt{2}} |\Delta\nu| \quad (1)$$

$$\Delta G^\ddagger = RT_c \ln \left( \frac{k_B T_c}{k_c h} \right) \quad (2)$$

The addition of cyanide to **2** in DMSO- $d_6$  produced multiple products, presumably from the reactions involving nucleophilic attack as well as simple deprotonation, as deduced from the  $^1\text{H}$  NMR spectrum (Fig. 5a, bottom). We tentatively concluded that the weak and solvent-exposed hydrogen bond of **2** is prone to tautomerization, and subjected to multiple reaction pathways with cyanide functioning as either nucleophilic Lewis base or simple Brønsted base.

### Structural Evolution of Hydrogen Bonding Network


To suppress the undesired acid–base chemistry observed for **2** (Fig. 4 and 5), additional functional groups were introduced to construct tighter hydrogen bonding. We postulated that a stronger bond polarization (Fig. 1) of cascade hydrogen bonding should shift the tautomer equilibrium to make the hydrogen-bonded form dominant even in polar solvent environment (Scheme 3, bottom), thereby enhancing the electrophilicity of the aldehyde group toward covalent capture of cyanide anion. As summarized in Scheme 3, probes **3** and **4** were readily prepared from the common diamine intermediate **5** (Scheme 1).



**Scheme 2** Tautomerization of a simple HBD–HBA pair.








**Scheme 3** Construction of hydrogen bonding network to shift the tautomer equilibrium. (a) aldehyde,  $\text{Na}_2\text{S}_2\text{O}_5$ , EtOH,  $\text{H}_2\text{O}$ ,  $\Delta$ . (b)  $\text{MnO}_2$ , MeCN/ $\text{CH}_2\text{Cl}_2$ , r.t.

For **3**, a hydroxyl group is installed as HBD at the upper ring of the benzimidazole core. Similarly to the catalytic triad (Fig. 2a), the  $\text{O}\cdots\text{H}\cdots\text{N}$  hydrogen bonding between the hydroxyl group and the imine nitrogen atom of benzimidazole is expected to push the tautomer equilibrium to reinforce the  $\text{N}\cdots\text{H}\cdots\text{O}$  hydrogen bonding between the aldehyde and amine moieties of the benzimidazole core. Furthermore, the acidic phenolic  $\text{O}\cdots\text{H}$  ( $\text{pK}_a \sim 10$ )<sup>37</sup> could also function as a sacrificial proton donor in the acid-base reaction to keep the  $\text{N}\cdots\text{H}$  group intact. Even when the acid-base reaction takes place, the deprotonation reaction should occur preferentially at the hydroxyl group, so that the benzimidazole fluorophore would be less perturbed.

As a further structural elaboration, probe **4** has an additional hydrogen bonding acceptor,  $-\text{OMOM}$  group. Here, the hydrogen bonding between the ether oxygen atom of  $-\text{OMOM}$  and the imidazole  $\text{N}\cdots\text{H}$  group is anticipated to help planarize the  $\pi$ -system and strengthen the bifurcated hydrogen bonding. It makes intuitive chemical sense that the multiple HBD–HBA pairs within **4** should work cooperatively in the same direction (Fig. 1b) to preferentially stabilize the desired tautomer which benefits from stronger hydrogen bonding (Scheme 3, bottom).

Indeed, single-crystal X-ray crystallography confirmed the presence of an extensive hydrogen bonding network within **4**. As shown in Fig. 6, the X-ray structure of **4** revealed three hydrogen bonds (dashed lines) around the imidazole core with short  $\text{N}_{\text{imidazole}}\cdots\text{O}_{\text{phenol}}$  (2.568(3) Å) and  $\text{N}_{\text{imidazole}}\cdots\text{O}_{\text{ether}}$  (2.634(4) Å) distances and essentially coplanar arrangement of the extended  $\pi$ -system (torsional angle for  $\text{N1}\cdots\text{C1}\cdots\text{C9}\cdots\text{C10} = 0.65^\circ$ ;  $\text{C2}\cdots\text{C3}\cdots\text{C8}\cdots\text{O4} = 1.92^\circ$ ).




**Fig. 6** X-ray structure of **4** as ORTEP diagrams with thermal ellipsoids at the 50% probability level: (a) face-on and (b) edge-on views. Selected interatomic distances (Å) of the hydrogen bonds (dashed lines):  $\text{N1}\cdots\text{O3}$ , 2.568(3);  $\text{N2}\cdots\text{O1}$ , 2.634(4);  $\text{N2}\cdots\text{O4}$ , 2.901(3).

## Cascade Hydrogen Bonds: Effects on the Reactivity and Solution Structure


DOI: 10.1039/D0SC05067A

Both **3** and **4** exhibited dramatic fluorescence enhancement upon treatment with cyanide anion (Fig. 7). Unlike **2**, however, they remain silent toward  $\text{Et}_3\text{N}$ , thus differentiating Brønsted base from Lewis base. When comparison is made with **2**, slightly blue-shifted but almost superimposable emission spectra ( $\lambda_{\text{max,em}} = 445 \text{ nm}$ ) were observed for the cyanide reaction products of **3** and **4**.

We anticipated that multiple HBD–HBA pairs within **3** and **4** would shift the tautomer equilibrium to benefit from the additive and reinforcing dipole alignment (Scheme 3, bottom). The results from  $^1\text{H}$  NMR spectroscopic studies, however, were rather inconclusive. As shown in Fig. 8, the coexistence of two tautomers was still observed at r.t. for both **3** and **4**. While the broadened proton resonances of **3** resemble those of **2** (Fig. 5a, top), the two tautomers of **4** appear as sharp and well-resolved spectral patterns in ca. 1:1 ratio (Fig. 8a and Fig. 8b, top).



**Fig. 7** Emission spectra ( $\lambda_{\text{exc}} = 343 \text{ nm}$ ) of (a) **3** (0.100 mM) prior to (black dashed lines), and after treatment with NaCN (25 equiv, blue line) or  $\text{Et}_3\text{N}$  (25 equiv, gray line), and (b) **4** (0.100 mM) prior to (black dashed lines), and after treatment with NaCN (25 equiv, blue line) or  $\text{Et}_3\text{N}$  (25 equiv, gray line) in DMSO at  $T = 298 \text{ K}$ .



**Fig. 8** (a)  $^1\text{H}$  NMR spectra of **3** (4.0 mM) in  $\text{DMSO}-d_6$  prior to (top) and after (bottom) addition of NaCN (10 equiv). (b)  $^1\text{H}$  NMR spectra of **4** (4.0 mM) in  $\text{DMSO}-d_6$  prior to (top) and after (bottom) addition of NaCN (10 equiv).  $T = 298 \text{ K}$ .



Similar to the case of **2**, mixtures of products were obtained when cyanide was added to **3** (Fig. 8a, bottom), presumably reflecting the tautomer equilibrium. The sharp and well-resolved  $^1\text{H}$  NMR spectrum of **4** suggests much slower interconversion between the two tautomers (Scheme 3), which is consistent with the more extensive hydrogen bonding array in **4** than its simpler analogues **2** and **3**. Nevertheless, the coexistence of the two tautomers for **4** still resulted in the formation of multiple products upon reaction with  $\text{CN}^-$  (Fig. 8b, bottom). Apparently, the presence of three hydrogen bonds (Fig. 6) is still insufficient for a complete conversion of **4** to a single product.

### Biomimetic Hydrogen Bonding Network

As described above, studies on **2–4** suggested the need for stronger and more tightly regulated hydrogen bonding network to control both the tautomer equilibrium and the reactivity toward nucleophile. We thus decided to install an additional HBD–HBA motif onto **4** to prepare **1** (Fig. 3b). Within a six-membered ring setting, the highly polarized N–H bond of an amide moiety is ideally suited to make a good HBD–HBA pair with the imidazole imine-N atom, as predicted by the energy-minimized DFT (B3LYP-D3/6-31G(d,p)) model shown in Fig. 2b. In line with the schematic diagram shown in Fig. 1, comparative DFT studies on **1** and the simple HBD–HBA pair **2** predicted a large difference in the molecular dipole moment of 6.4413 D (for **1**) vs 0.5614 D (for **2**). The molecular electrostatic potential (MEP) maps of **1** and **2** (Fig. S4) also predict a larger polarization of electron density across cascading dipoles.


To functionalize the 4-position of the benzimidazole core, the overall synthetic scheme needed to be modified to introduce a bromo substituent at the early stage (Scheme 4). From the bromo-functionalized diamine, sequential oxidative condensation and oxidation reactions afforded **6** in high yield (72% for two steps). By cross-coupling reaction onto this bromo position, various functional groups could be installed. An aliphatic amide group was chosen in our molecular design to suppress direct electronic conjugation with the benzimidazole core, thereby minimizing perturbation of the photophysical properties. A palladium-catalyzed Suzuki–Miyaura cross-coupling of **6** with  $\alpha$ -(acetylamino)benzylboronic ester

furnished the target compound **1**. The low isolation yield (ca. 17%) in this final step is due to the purification procedure involving repetitive recrystallization.


The single-crystal X-ray structure of **1** shown in Fig. 9 confirmed the presence of multiple hydrogen bonds. The interatomic distances of hydrogen bonds, 2.5431(1)–2.9007(1) Å, are similar to those of **4** (Fig. 6). The essentially co-planar arrangement of the  $\pi$ -conjugated backbone of **1** (torsional angle for N1–C1–C9–C10 = 4.17°; C2–C3–C8–O4 = 2.03°) further validates the functional role of hydrogen bonding network as conformational lock. Unlike the DFT computational model (Fig. 2b), the amide N–H group of **1** is twisted away from the benzimidazole-N atom. A close inspection of the crystal packing diagram revealed extensive intermolecular  $\text{N}_{\text{amide}}\text{--H}\cdots\text{O}_{\text{carbonyl}}$  hydrogen bonding network between adjacent molecules, which is reinforced further by  $\pi$ – $\pi$  stacking and C–H $\cdots\pi$  contacts (Fig. S5). Apparently, such intermolecular interactions in the condensed phase prevail over the inherent propensity of **1** to make intramolecular  $\text{N}_{\text{amide}}\text{--H}\cdots\text{N}_{\text{imidazole}}$  hydrogen bond as discrete molecular species (Fig. 2b).

### Hydrogen Bonding Network: Effects on the Reactivity and Solution Structure


As shown in Fig. 10a, the addition of cyanide anion to a solution of **1** elicited a rapid and dramatic (> 180-fold) enhancement in the emission intensity at  $\lambda_{\text{max,em}} = 450$  nm, whereas no spectral change was observed with the Brønsted base  $\text{Et}_3\text{N}$ . A large spectral change was also observed in the electronic excitation



**Fig. 9** X-ray structure of **1** as ORTEP diagrams with thermal ellipsoids at the 50% probability level: (a) face-on and (b) edge-on views. Selected interatomic distances (Å) of the hydrogen bonds (dashed lines): N1 $\cdots$ O3, 2.5431(1); N2 $\cdots$ O1, 2.62995(9); N2 $\cdots$ O4, 2.9007(1).



**Scheme 4** Synthetic route to **1**. (a) aldehyde,  $\text{Na}_2\text{S}_2\text{O}_5$ ,  $\text{EtOH}/\text{H}_2\text{O}$ ,  $\Delta$ . (b)  $\text{MnO}_2$ ,  $\text{MeCN}/\text{CH}_2\text{Cl}_2$ , r.t. (c)  $\text{Pd}_2(\text{dba})_3$ ,  $\text{P}(\text{tBu})_3\cdot\text{HBF}_4$ , KF, dioxane/ $\text{H}_2\text{O}$ ,  $\Delta$ .




**Fig. 10** (a) Emission ( $\lambda_{\text{exc}} = 343$  nm) and (b) absorption spectra of **1** (0.100 mM, black dashed lines), and spectral changes induced by treatment with NaCN (25 equiv, blue lines), and with  $\text{Et}_3\text{N}$  (25 equiv, gray lines) in DMSO at  $T = 298$  K. Inset: photographic images of (a) fluorescence response (under 365 nm UV lamp), and (b) color change.



upon the addition of cyanide (Fig. 10b) with color change to yellow. In contrast, only a slight increase in the absorption at  $\lambda \approx 400$  nm region was observed with  $\text{Et}_3\text{N}$ , with the rest of the spectrum remaining essentially superimposable (Fig. 10b). This visually discernible colorimetric change and fluorescence turn-on allow naked-eye detection of cyanide anion (Fig. 10, inset pictures)

To investigate the solution structure of **1**,  $^1\text{H}$  NMR spectroscopic studies were carried out. The  $^1\text{H}$  NMR spectrum of **1** (4.0 mM) in  $\text{DMSO}-d_6$  measured at  $T = 293$  K indicates the dominance of one prevailing tautomer ( $> 90\%$ , based on the peak integration values; Fig. 11a, top), which remain essentially invariant even with increasing temperature up to  $T = 363$  K (Fig. 11b). The 2D ROESY NMR spectrum of **1** (4.0 mM) obtained in  $\text{DMSO}-d_6$  at r.t. revealed prominent ROE signals between (i) benzimidazole N–H and aldehyde C–H, and (ii) benzimidazole N–H and methylene protons of the –OMOM group (Fig. S6). These ROE correlations provide compelling evidence for the dominant tautomeric form of **1**, as predicted by DFT computational studies (Fig. 2b). The gradual up-field shifts of the phenolic O–H ( $\delta = 13.46$  to  $13.30$  ppm) and amide N–H protons ( $\delta = 8.99$  to  $8.68$  ppm) with increasing temperature (Fig. 11b) also suggest their involvement in intramolecular hydrogen bonding.<sup>38,39</sup>




**Fig. 11** (a)  $^1\text{H}$  NMR spectra of **1** (4.0 mM) prior to (top) and after (bottom) addition of NaCN (10 equiv) in  $\text{DMSO}-d_6$  at  $T = 298$  K. Asterisks indicate the resonances of the minor tautomer which is less than 10% based on the peak integration values. (b) Variable-temperature  $^1\text{H}$  NMR spectra of **1** (4.0 mM) in  $\text{DMSO}-d_6$  at  $T = 293$ – $363$  K.

The high conformational stability of **1** was established further by concentration-dependent  $^1\text{H}$  NMR studies. Within the concentration range of  $1.5$ – $4.0$  mM, no noticeable change was observed in the  $^1\text{H}$  NMR spectrum of **1** in  $\text{DMSO}-d_6$ , implying that **1** remains folded in solution against intermolecular hydrogen bonding (Fig. S7). In stark contrast, compound **3** having relatively weak hydrogen bonds undergoes significant broadening of benzimidazole aromatic proton resonances with increasing sample concentration (Fig. S8). Furthermore, the addition of a small amount of  $\text{H}_2\text{O}$  ( $2\ \mu\text{L}$ ) to a solution of **3** in  $\text{DMSO}-d_6$  ( $2.0$  mM,  $500\ \mu\text{L}$ ) sharpened the resonances of these aromatic protons by rapid proton exchange with the N–H group (Fig. S9). No spectral change was observed for **1** under the same condition (Fig. S10).

With the hydrogen-bonded tautomeric form prevailing for **1** in solution, the addition of cyanide anion resulted in clean and complete conversion to the cyanohydrin adduct (Fig. 11a, bottom). To better interpret the  $^1\text{H}$  NMR spectrum of the reaction product, we carried out 2D-COSY NMR studies. As shown in Fig. 12, the cross-peaks at  $\delta = 9.7$  ppm and  $\delta = 6.6$  ppm arise from the  $J$ -coupling of O–H and C–H protons, respectively, of the cyanohydrin.<sup>32,33,40–42</sup> In addition, HPLC-MS analysis (Fig. S11) confirmed the formation of the cyanohydrin adduct of **1** at  $m/z = 473.15$  corresponding to  $[\text{M} + \text{H}]^+$ .

#### Telling a Nucleophile from a Base


To compare the ability of **1**–**4** in distinguishing nucleophile (i.e.  $\text{CN}^-$ ) from Brønsted base (i.e.  $\text{Et}_3\text{N}$ ), each of the probe molecules was treated with either cyanide anion (25 equiv) or  $\text{Et}_3\text{N}$  (25 equiv). The emission intensity was measured at  $\lambda_{\text{em}} = 450$  nm, and the ratio  $I_{\text{CN}}/I_{\text{Et}_3\text{N}}$  was calculated for each molecule. With an increasing number of HBD–HBA units installed around the same fluorogenic benzimidazole core (Fig. 3b), a systematic increase in the  $I_{\text{CN}}/I_{\text{Et}_3\text{N}}$  ratio was observed along the series **2**→**3**→**4**→**1** (Fig. 13). Stronger and networked hydrogen bonds seem to promote bond-forming reaction while effectively suppressing the undesired acid–base reaction. Comparative  $^1\text{H}$  NMR studies (Fig. 5, 8, and 11) establish that such hydrogen bonds can also shift the solution equilibrium toward the more reactive tautomer toward cyanide anion.



**Fig. 12** Partial 2D-COSY contour plot of **1** (4.0 mM) treated with NaCN (10 equiv) in  $\text{DMSO}-d_6$  at  $T = 298$  K. The corresponding 1D NMR spectrum is shown along the ordinate. Inset: DFT (B3LYP-D3/6-31G(d,p)) energy-minimized structure of the cyanohydrin adduct of **1**.








**Fig. 13** Effects of hydrogen bonding on distinguishing Brønsted base from nucleophilic Lewis base. Each bar denotes the ratio of fluorescence intensity at  $\lambda_{\text{em}} = 450$  nm determined for each probes (**1–4**, 0.100 mM) after treating with NaCN (25 equiv,  $I_{\text{NaCN}}$ ) or with Et<sub>3</sub>N (25 equiv,  $I_{\text{Et3N}}$ ) in DMSO at  $T = 298$  K.  $\lambda_{\text{exc}} = 343$  nm for **1**, **3**, and **4**; 400 nm for **2**.

### Selectivity toward Cyanide and Reaction Stoichiometry

To test the selectivity of the probe **1**, aqueous solution samples of 12 different anions, including CN<sup>−</sup>, F<sup>−</sup>, Cl<sup>−</sup>, Br<sup>−</sup>, I<sup>−</sup>, N<sub>3</sub><sup>−</sup>, SCN<sup>−</sup>, OAc<sup>−</sup>, NO<sub>3</sub><sup>−</sup>, ClO<sub>4</sub><sup>−</sup>, PF<sub>6</sub><sup>−</sup>, and OH<sup>−</sup> (25 equiv, delivered as sodium salts except for KPF<sub>6</sub> and KOH) were added to **1** (0.100 mM) in DMSO–H<sub>2</sub>O (99:1, v/v) mixed-solvent system, and the emission spectra recorded under identical conditions. As summarized in Fig. 14a and 14b, the fluorescence turn-on response was observed exclusively for cyanide anion.



**Fig. 14** (a) Emission spectra of **1** after treatment with CN<sup>−</sup> (25 equiv, blue line) or other 11 anions (25 equiv, gray lines; see the list in (b) and (c)). (b) Normalized emission intensity ( $= I/I_0$ ) of **1** in the presence of various anions: 1, CN<sup>−</sup>; 2, F<sup>−</sup>; 3, Cl<sup>−</sup>; 4, Br<sup>−</sup>; 5, I<sup>−</sup>; 6, N<sub>3</sub><sup>−</sup>; 7, SCN<sup>−</sup>; 8, OAc<sup>−</sup>; 9, NO<sub>3</sub><sup>−</sup>; 10, ClO<sub>4</sub><sup>−</sup>; 11, PF<sub>6</sub><sup>−</sup>; 12, OH<sup>−</sup> (delivered as sodium salts except for KPF<sub>6</sub> and KOH). The intensity at  $\lambda = 450$  nm ( $= I$ ) after the treatment of **1** with each anion (25 equiv) is normalized with that of **1** ( $= I_0$ ). (c) Enhancement of the emission intensity after addition of NaCN (25 equiv) to **1** in the presence of each anion (25 equiv). Conditions:  $\lambda_{\text{exc}} = 343$  nm;  $T = 298$  K; [**1**] = 0.100 mM in DMSO–H<sub>2</sub>O (99:1, v/v). (d) Job plot analysis of the reaction of **1** with CN<sup>−</sup> in DMSO at  $T = 298$  K. The absorbance at  $\lambda = 410$  nm was recorded by varying the mole fraction of CN<sup>−</sup> while keeping the total concentration constant ( $= 67$   $\mu$ M).

A large enhancement in the emission intensity was also observed at  $\lambda_{\text{em}} = 450$  nm when cyanide anion was subsequently added to the mixture of **1** and other anions except OH<sup>−</sup> (Fig. 14c). We suspect that deprotonation by strong base could disrupt the cascade hydrogen bonding, thus diminishing the response of **1** toward subsequently added CN<sup>−</sup>. The reaction stoichiometry between **1** and cyanide was determined by Job plot analysis using UV–vis spectroscopy. A sharp maximum at the mole fraction of 0.5 provides compelling evidence for the formation of 1:1 adduct (Fig. 14d), which is also consistent with the results from <sup>1</sup>H NMR (Fig. 11a and 12) and HPLC–MS studies (Fig. S11).

### Kinetic Studies: Rapid Detection of Cyanide under Ambient Conditions

The rate constant for the bimolecular reaction between **1** and cyanide was determined by taking time-dependent UV–vis absorption spectra (Fig. S12). A large enhancement in the absorption at  $\lambda = 400$  nm brought by the addition of cyanide anion helped track the progress of the reaction over time (Fig. 10b). The activation of electrophilic aldehyde group by the hydrogen bonding array led to a rapid chemical transformation. Even at low temperature ( $T = 15$  °C), the reaction of **1** with cyanide in DMSO–MeCN (1:1, v/v) was completed within < 2 seconds (Fig. S12a). Under *pseudo*-first-order kinetic reaction conditions, the kinetic trace was fitted to obtain *pseudo*-first-order rate constants,  $k'$  ( $= k_2[\text{CN}^-]_0$ ); the second-order rate constant  $k_2$  was estimated from the linear relationship between  $k'$  and  $[\text{CN}^-]_0$  (Fig. S12b). While the precise determination of rate constants was hampered by the fast reaction rate, the calculated  $k_2$  value ( $1.3 \times 10^3 \text{ M}^{-1}\text{s}^{-1}$ ) is one of the fastest-responding cyanide probes that operate by covalent capture strategy.<sup>32,43–50</sup> With 20 equiv of cyanide anion, the reaction half-life  $t_{1/2}$  is as short as 0.53 sec.

### Conclusions

As a synthetic mimic of biological hydrogen bonding triad, a T-shaped  $\pi$ -conjugated platform was structurally elaborated. In our molecular design, the amphoteric benzimidazole core reinforces bond-polarizing HBD–HBA networks to activate an electrophilic aldehyde group for covalent capture of toxic CN<sup>−</sup> anion. We found that a systematic increase in the number of hydrogen bonds allows the molecules to distinguish bond-making nucleophile from proton-abstracting Brønsted base. The most advanced molecular probe **1** has four hydrogen bonds around the fluorogenic benzimidazole core, and detects cyanide ion by a rapid and selective turn-on response. Efforts are currently underway in our laboratory to expand the scope of this non-covalent design strategy to other types of chemical transformations of relevance to target-specific signal transduction.

### Conflicts of interest

There are no conflicts to declare.





## Acknowledgements

This work was supported by the National Research Foundation (NRF) of Korea (2020R1A2C2006381 and 2017M3D1A1039558) funded by the Ministry of Science and ICT.


## Notes and references

- 1 A. P. Davis, *Chem. Soc. Rev.*, 2020, **49**, 2531–2545.
- 2 P. Ríos, T. J. Mooibroek, T. S. Carter, C. Williams, M. R. Wilson, M. P. Crump and A. P. Davis, *Chem. Sci.*, 2017, **8**, 4056–4061.
- 3 H. Yao, H. Ke, X. Zhang, S.-J. Pan, M.-S. Li, L.-P. Yang, G. Schreckenbach and W. Jiang, *J. Am. Chem. Soc.*, 2018, **140**, 13466–13477.
- 4 J. F. Neal, W. Zhao, A. J. Grooms, M. A. Smeltzer, B. M. Shook, A. H. Flood and H. C. Allen, *J. Am. Chem. Soc.*, 2019, **141**, 7876–7886.
- 5 C. Guo, H. Wang, V. M. Lynch, X. Ji, Z. A. Page and J. L. Sessler, *Chem. Sci.*, 2020, **11**, 5650–5657.
- 6 D. Basudhar, Y. Madrona, E. T. Yukl, S. Sivaramakrishnan, C. R. Nishida, P. Moëne-Loccoz and P. R. Ortiz de Montellano, *J. Biol. Chem.*, 2016, **291**, 16100–16111.
- 7 E. Odella, S. J. Mora, B. L. Wadsworth, M. T. Huynh, J. J. Goings, P. A. Liddell, T. L. Groy, M. Gervald, L. E. Sereno, D. Gust, T. A. Moore, G. F. Moore, S. Hammes-Schiffer and A. L. Moore, *J. Am. Chem. Soc.*, 2018, **140**, 15450–15460.
- 8 E. Odella, S. J. Mora, B. L. Wadsworth, J. J. Goings, M. A. Gervald, L. E. Sereno, T. L. Groy, D. Gust, T. A. Moore, G. F. Moore, S. Hammes-Schiffer and A. L. Moore, *Chem. Sci.*, 2020, **11**, 3820–3828.
- 9 J. Wang, K. Liu, R. Xing and X. Yan, *Chem. Soc. Rev.*, 2016, **45**, 5589–5604.
- 10 F. Zapata, L. González, A. Caballero, A. Bastida, D. Bautista and P. Molina, *J. Am. Chem. Soc.*, 2018, **140**, 2041–2045.
- 11 L. Gabrielli, D. Núñez-Villanueva and C. A. Hunter, *Chem. Sci.*, 2020, **11**, 561–566.
- 12 D. Núñez-Villanueva, G. Iadevaia, A. E. Stross, M. A. Jinks, J. A. Swain and C. A. Hunter, *J. Am. Chem. Soc.*, 2017, **139**, 6654–6662.
- 13 R. R. Knowles and E. N. Jacobsen, *Proc. Natl. Acad. Sci. U. S. A.*, 2010, **107**, 20678–20685.
- 14 A. G. Doyle and E. N. Jacobsen, *Chem. Rev.*, 2007, **107**, 5713–5743.
- 15 S. Bhunia, A. Rana, P. Roy, D. J. Martin, M. L. Pegis, B. Roy and A. Dey, *J. Am. Chem. Soc.*, 2018, **140**, 9444–9457.
- 16 C. Nagamani, U. Viswanathan, C. Versek, M. T. Tuominen, S. M. Auerbach and S. Thayumanavan, *Chem. Commun.*, 2011, **47**, 6638–6640.
- 17 Y. Zhou, G. Deng, Y.-Z. Zheng, J. Xu, H. Ashraf and Z.-W. Yu, *Sci. Rep.*, 2016, **6**, 36932.
- 18 H. Guo and M. Karplus, *J. Phys. Chem.*, 1994, **98**, 7104–7105.
- 19 A. Shokri, J. Schmidt, X.-B. Wang and S. R. Kass, *J. Am. Chem. Soc.*, 2012, **134**, 2094–2099.
- 20 F. Bartha, O. Kapuy, C. Kozmutza and C. Van Alsenoy, *J. Mol. Struct.: THEOCHEM*, 2003, **666–667**, 117–122.
- 21 A. Bhattacharjee and S. Wategaonkar, *Phys. Chem. Chem. Phys.*, 2016, **18**, 27745–27749.
- 22 R. Ludwig, *J. Mol. Liq.*, 2000, **84**, 65–75.
- 23 P. A. Frey, *J. Phys. Org. Chem.*, 2004, **17**, 511–520.
- 24 A. Rauwerdink and R. J. Kazlauskas, *ACS Catal.*, 2015, **5**, 6153–6176.
- 25 L. Polgár, *Cell. Mol. Life Sci.*, 2005, **62**, 2161–2172.
- 26 C. L. Perrin and J. B. Nielson, *Annu. Rev. Phys. Chem.*, 1997, **48**, 511–544.
- 27 P. Kumar, P. K. Agarwal, M. B. Waddell, T. Mittag, E. H. Serpersu and M. J. Cuneo, *Angew. Chem., Int. Ed.*, 2019, **58**, 16260–16266.
- 28 H. J. Kim, C. H. Heo and H. M. Kim, *J. Am. Chem. Soc.*, 2013, **135**, 17969–17977. DOI: 10.1039/D0SC05067A
- 29 C.-H. Chen, W.-S. Huang, M.-Y. Lai, W.-C. Tsao, J. T. Lin, Y.-H. Wu, T.-H. Ke, L.-Y. Chen and C.-C. Wu, *Adv. Funct. Mater.*, 2009, **19**, 2661–2670.
- 30 S. K. Lower and M. A. El-Sayed, *Chem. Rev.*, 1966, **66**, 199–241.
- 31 Kenry, C. Chen and B. Liu, *Nat. Commun.*, 2019, **10**, 2111.
- 32 J. Jo, A. Olasz, C.-H. Chen and D. Lee, *J. Am. Chem. Soc.*, 2013, **135**, 3620–3632.
- 33 J. Jo and D. Lee, *J. Am. Chem. Soc.*, 2009, **131**, 16283–16291.
- 34 H. S. Gutowsky and C. H. Holm, *J. Chem. Phys.*, 1956, **25**, 1228–1234.
- 35 E. D. Raczynska and W. Kosińska, *Chem. Rev.*, 2005, **105**, 3561–3612.
- 36 C. Zucco, E. L. Dall'Oglio, G. V. Salmória, H. Gallardo, A. Neves and M. C. Rezende, *J. Phys. Org. Chem.*, 1998, **11**, 411–418.
- 37 M. D. Liptak, K. C. Gross, P. G. Seybold, S. Feldgus and G. C. Shields, *J. Am. Chem. Soc.*, 2002, **124**, 6421–6427.
- 38 S. H. Gellman, G. P. Dado, G.-B. Liang and B. R. Adams, *J. Am. Chem. Soc.*, 1991, **113**, 1164–1173.
- 39 B. W. Gung, Z. Zhu and B. Everingham, *J. Org. Chem.*, 1997, **62**, 3436–3437.
- 40 K.-S. Lee, H.-J. Kim, G.-H. Kim, I. Shin and J.-I. Hong, *Org. Lett.*, 2008, **10**, 49–51.
- 41 S. K. Kwon, S. Kou, H. N. Kim, X. Chen, H. Hwang, S.-W. Nam, S. H. Kim, K. M. K. Swamy, S. Park and J. Yoon, *Tetrahedron Lett.*, 2008, **49**, 4102–4105.
- 42 Y. Jhong, W. H. Hsieh, J.-L. Chir and A.-T. Wu, *J. Fluoresc.*, 2014, **24**, 1723–1726.
- 43 Z. Xu, X. Chen, H. N. Kim and J. Yoon, *Chem. Soc. Rev.*, 2010, **39**, 127–137.
- 44 Y.-M. Dong, Y. Peng, M. Dong and Y.-W. Wang, *J. Org. Chem.*, 2011, **76**, 6962–6966.
- 45 L. Hou, F. Li, J. Guo, X. Zhang, X. Kong, X. T. Cui, C. Dong, Y. Wang and S. Shuang, *J. Mater. Chem. B*, 2019, **7**, 4620–4629.
- 46 Y.-D. Lin, Y.-S. Pen, W. Su, K.-L. Liao, Y.-S. Wen, C.-H. Tu, C.-H. Sun and T. J. Chow, *Chem. Asian. J.*, 2012, **7**, 2864–2871.
- 47 S.-J. Hong, J. Yoo, S.-H. Kim, J. S. Kim, J. Yoon and C.-H. Lee, *Chem. Commun.*, 2009, 189–191.
- 48 A. Tigreros, J.-C. Castillo and J. Portilla, *Talanta*, 2020, **215**, 120905.
- 49 S. M. Kim, M. Kang, I. Choi, J. J. Lee and C. Kim, *New J. Chem.*, 2016, **40**, 7768–7778.
- 50 S. Zhu, M. Li, L. Sheng, P. Chen, Y. Zhang and S. X.-A. Zhang, *Analyst*, 2012, **137**, 5581–5585.



## ARTICLE

## Table of Contents



Biomimetic cascade hydrogen bonds promote covalent capture of the nucleophile by polarizing electrophilic reaction site, while suppressing non-productive acid–base chemistry as the competing reaction pathway.

