Asymmetric Catalysis

Organocatalyzed Enantioselective Fluorocyclizations**

Oscar Lozano, George Blessley, Teresa Martinez del Campo, Amber L. Thompson, Guy T. Giuffredi, Michela Bettati, Matthew Walker, Richard Borman, and Véronique Gouverneur*

Asymmetric halogen-promoted cyclizations have been the object of extensive investigations, the ultimate aim being the emulation of nature's remarkable ability to construct architecturally complex enantiopure halogenated compounds from prochiral precursors. Progress has been slow, but significant advances have recently been made in iodo-, bromo-, and chlorocyclizations.^[1] A catalytic enantioselective fluorocyclization induced by an electrophilic fluorinating reagent has not been reported to date. This fact represents a significant gap since such a transformation can streamline synthetic access to enantiopure fluorinated heterocycles, a class of compounds that plays an important role in medicinal chemistry.^[2] We became interested in this problem because of the information available on organocatalytic enantioselective fluorination^[3] and our recent work in cascade fluorination-heterocyclizations.^[4] Herein, we disclose the first successful catalytic process that delivers enantioenriched fluorinated heterocyclic products upon fluorocyclization of prochiral precursors.

For this study, we selected the prochiral indoles 1 and 2 with a pendant heteronucleophile tethered at C3 or at the nitrogen atom, respectively, based on the typical position of this heteroaromatic motif in nitrogen-containing natural products (Figure 1).^[5] The established reactivity profile of indoles towards achiral electrophilic fluorinating reagents offered a starting point to identify suitable reaction conditions to allow for organocatalyzed enantioselective fluorocyclizations.^[6] Two recent discoveries support the use of cinchona alkaloids to induce enantiocontrol. Shibata and co-workers reported that N-F reagents in the presence of catalytic amounts of cinchona alkaloids induce asymmetric fluorination of activated substrates.^[7] In addition, we demonstrated that asymmetric fluoroetherification of a silyl-activated homoallylic alcohol can be performed with stoichiometric amounts of chiral N-F reagents prepared in situ from Selectfluor and (DHQ)₂PHAL. Although the enantioselectivity of

[*] Dr. O. Lozano, G. Blessley, Dr. T. Martinez del Campo, Dr. A. L. Thompson, G. T. Giuffredi, Prof. V. Gouverneur Chemistry Research Laboratory
University of Oxford
12 Mansfield Road OX1 3TA Oxford (UK)
E-mail: veronique.gouverneur@chem.ox.ac.uk
Dr. M. Bettati, Dr. M. Walker, Dr. R. Borman GlaxoSmithKline R&D, Stevenage Herts SG1 2NY (UK)

- [**] We thank the European Union (PIEF-GA-2009-235510 to T.M., PIEF-GA-2008-220034 to O.L.), GSK (G.B.) and the Berrow Foundation for a scholarship to G.T.G. We also thank Dr. S. Fletcher for very helpful discussions.
 - Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.201103151.



Figure 1. Enantioselective fluorocyclization of prochiral indoles. PG = protecting group.

this fluorocyclization was modest, the lead result was encouraging.^[4a]

From the onset, we appreciated the conceptual difficulties in developing such a catalytic asymmetric fluorocyclization. The uncatalyzed reaction must be significantly slower than the catalyzed process, a requirement difficult to meet as any in situ generated transient chiral N–F cinchona species will be of similar reactivity or less reactive with respect to the parent achiral fluorinating agent. In addition, the paucity of information on the factors that control the catalytic asymmetric delivery of halogens onto alkenes in the presence of cinchona alkaloids is limiting.^[1,7] Our studies commenced with the fluorocyclization of the prototypical indole **1a**, a substrate that requires the creation of a stereogenic fluorinated quaternary benzylic carbon center upon fluorocyclization (Table 1).

Both Selectfluor and NFSI induced fluorocyclization at room temperature to afford tetrahydrofuroindole (\pm) -3a as a single *cis* diastereomer (d.r. > 20:1) with average yields of 70% (Table 1, entries 1-3). The process was similarly effective at -78 °C with NFSI in acetone (Table 1, entry 4). Premixing various cinchona alkaloids with Selectfluor at room temperature followed by the addition of 1a at ambient temperature or at -78 °C led to the formation of **3a** with the ee value reaching 74% (Table 1, entries 5-10). The best enantiocontrol was observed when the fluorocyclization was performed with 1.2 equivalents of (DHQ)₂PHAL in acetone at -78°C (Table 1, entry 9), all other alkaloids that were screened were found to be less efficient.^[8] To our great delight, when using a catalytic amount of (DHQ)₂PHAL (20 mol%), **3a** was formed with 66% *ee* (Table 1, entry 12). This reaction was best performed at -78°C in acetone with NFSI and an excess of K₂CO₃. Decreasing the catalyst load, changing the base, or using Selectfluor instead of NFSI proved detrimental (Table 1, entries 11-16). The beneficial effect of the use of an inorganic carbonate base for the



Table 1: Validation and optimization of the fluorocyclization of 1 a.^[8]

			d 0	0.1			
Ľ	Ń	cincho	na alkaloids		N H	u.r. > 2	0.1
	1a Me			3a	м̀е		
Entry	F^+	Alkaloid ^[b]	Base	Solvent	T ^[e]	Yield	ee
	source ^[a]	(mol %)				[%] ^[f]	[%] ^[g]
1	А	-	NaHCO ₃ ^[c]	MeCN	RT	73	-
2	Α	-	NaHCO3 ^[c]	acetone	RT	60	_
3	В	-	NaHCO3 ^[c]	MeCN	RT	76	_
4	В	-	NaHCO ₃ ^[c]	acetone	−78°C	73	-
5	Α	C (100)	NaHCO ₃ ^[c]	MeCN	RT	50	20
6	Α	D (100)	NaHCO ₃ ^[c]	MeCN	RT	53	5
7	Α	E (100)	NaHCO ₃ ^[c]	acetone	−78°C	49	0
8	Α	F (100)	NaHCO ₃ ^[c]	MeCN	RT	72	55
9	Α	F (100)	NaHCO ₃ ^[c]	acetone	−78°C	56	74
10	Α	F (100)	NaHCO ₃ ^[c]	THF	−78°C	64	64
11	В	F (10)	$K_2CO_3^{[d]}$	acetone	−78°C	50	50
12	В	F (20)	$K_2CO_3^{[d]}$	acetone	−78°C	72	66
13	В	F (20)	Cs ₂ CO ₃ ^[d]	acetone	_78°C	58	66
14	В	F (20)	$K_2CO_3^{[d]}$	acetone	RT	50	58
15	В	F (20)	-	acetone	−78°C	36	34
16	А	F (20)	$K_2CO_3^{[d]}$	acetone	−78 °C	47	32

[a] A = 1-chloromethyl-4-fluoro-1,4-diazoniabicyclo[2.2.2]octane bis-(tetrafluoroborate) (Selectfluor), B = N-fluorobenzenesulfonimide (NFSI). [b] C = 2,5-diphenyl-4,6-bis(dihydroquininyl)pyrimidine ((DHQ)₂PYR), D = hydroquinidine-9-phenanthryl ether, E = 1,4bis(dihydroquinidinyl)anthraquinone ((DHQD)₂AQN), F = 1,4-bis(dihydroquininyl)phthalazine ((DHQ)₂PHAL). [c] 1 equiv. [d] 6 equiv. [e] Reaction time up to 5 h at RT and up to 15 h at -78 °C. [f] Yield of isolated product. [g] *ee* value determined by HPLC on a chiral stationary phase.

catalytic asymmetric fluorination with cinchona alkaloids is not unprecedented. $^{\left[7\right] }$

The scope and limitation of this organocatalyzed process was investigated with respect to both the substitution pattern of the indolic structural core and the attached nucleophile. To evaluate the efficiency of the catalytic protocol, the yields and enantiomeric excesses of the products of reactions carried out with either an equimolar (conditions A) or a catalytic amount of the cinchona alkaloid (conditions B) were compared (Table 2).^[8]

The reaction tolerates various substituents on the indolic nitrogen atom, including hydrogen, but the highest enantiomeric excess was observed for the fluorocyclized product 3a derived from N-methyl substituted precursors (Table 2, entries 1-7). The presence of a substituent at position 5 $(\mathbf{R}^1 \neq \mathbf{H})$ led to a markedly improved control over the enantioselectivity (Table 2, entries 8-19). When an equimolar amount of alkaloid was used (conditions A), enantiomeric excesses ranging from 72% to 90% were observed; under catalytic conditions only a slight decrease of enantiomeric excesses (up to 84 % ee) was observed with no compromise on yields (conditions B). Indolic precursors with pendant unprotected primary amines did not afford any fluorinated ringclosed products; this behavior is likely due to a competing Nfluorination reaction. Various N-protected amines were found to be efficient nucleophiles (Table 2, entries 20-37). N-Tosylated indoles underwent fluorocyclization to afford the desired tetrahydropyrroloindoles 31-30 in good yields and *Table 2:* Fluorocyclization of indoles **1 a**–**1 t**.

Tuble .	2. 1100100	~			u 11.	F	. ~	
	R ¹	\checkmark	XH _{con}	ditions	A or B		X	
	\checkmark	-N		d.r. > 2	0:1	\checkmark	-N H	
	10.11	R²				- 01	R²	
	ia-it				3	a-3t		
Entry	Cond. ^[a]	1	R ¹	R ²	ХН	3	Yield [%] ^[b]	ее [%] ^[с]
1	А	la	Н	Me	ОН	3 a	56	74
2	В	la	Н	Me	ОН	3 a	72	66
3	А	1 b	Н	Et	ОН	3 b	51	56
4	В	1 b	Н	Et	ОН	3 b	68	52
5	Α	1c	н	allyl	ОН	3 c	62	60
6	В	1c	Н	allyl	ОН	3 c	76	60
7	A	1 d	Н	Н	ОН	3 d	33	40
8	А	1e	OMe	Me	ОН	3 e	90	86
9	В	le	OMe	Me	ОН	3 e	65	74
10	A	1 f	OBn	Me	ОН	3 f	69	84
11	В	1 f	OBn	Me	ОН	3 f	78	74
12	A	٦g	OEt	Me	ОН	3 g	60	84
13	В	٦g	OEt	Me	ОН	3 g	78	72
14	A	1h	O(allyl)	Me	ОН	3 h	53	86
15	В	1h	O(allyl)	Me	ОН	3 h	65	78
16	Α	1i	Ph	Me	ОН	3 i	55	72
17	В	1i	Ph	Me	ОН	3 i	61	62
18	Α	1 k	Mes	Me	ОН	3 k	57	90
19	В	1 k	Mes	Me	ОН	3 k	55	84
20	Α	11	Н	Me	NHTs	31	54	76
21	В	11	Н	Me	NHTs	31	59	64
22	Α	1 m	OMe	Me	NHTs	3 m	55	78
23	В	1 m	OMe	Me	NHTs	3 m	51	70
24	Α	1n	Ph	Me	NHTs	3 n	50	82
25	В	٦n	Ph	Me	NHTs	3 n	70	70
26	А	10	Mes	Me	NHTs	30	60	92
27	В	10	Mes	Me	NHTs	30	80	84
28	Α	۱p	Н	Me	NHCOMe	3 p	45 ^[d]	66
29	В	۱p	Н	Me	NHCOMe	3 p	95	80
30	А	1 q	Mes	Me	NHCOMe	3 q	38 ^[d]	92
31	В	1 q	Mes	Me	NHCOMe	3 q	65	92
32	Α	1r	Н	Me	NHCO ₂ Me	3 r	56	78
33	В	1r	Н	Me	NHCO ₂ Me	3 r	76	74
34	A	1 s	Н	Me	NHCO ₂ Bn	3 s	40	78
35	В	1 s	Н	Me	NHCO ₂ Bn	3 s	47	77
36	A	1t	Н	Me	NHBoc	3t	67	86
37	В	1t	Н	Me	NHBoc	3 t	70	78

[a] Conditions A: Selectfluor (1.2 equiv), $(DHQ)_2PHAL$ (1.2 equiv), NaHCO₃ (1.2 equiv), acetone, -78 °C; conditions B: NFSI (1.2 equiv), $(DHQ)_2PHAL$ (0.2 equiv), K₂CO₃ (6 equiv), acetone, -78 °C. [b] Yields of isolated products. [c] *ee* value determined by HPLC on a chiral stationary phase. [d] Conversion determined by ¹H NMR spectroscopy of the crude product.



with enantiomeric excesses reaching 84% under conditions B (Table 2, entry 27). Indoles **1p–1q** with a pendant N-acetamido nucleophile engaged very effectively in organocatalytic asymmetric fluorocyclization (Table 2, entries 28–31). This Nprotected nucleophile led to a very efficient control over enantiofacial selectivity (ee = 92%; Table 2, entries 30 and 31). The methyl-, benzyl-, and *tert*-butylcarbamates of *N*-methyl tryptamine (**1r–1t**) were also found to be suitable substrates for organocatalysis, and delivered the products in up to 78% *ee* (Table 2, entries 32-37).

The cyclization products **3a–3t** were formed as single *cis* diastereomers, as judged by ¹H NMR analysis of the crude products of the reactions.^[8] X-ray diffraction studies of the sulfonamide-containing product (3aR,8aS)-**3o** confirmed its structure (Figure 2).^[8,9]



Figure 2. Crystal structure of one molecule of (3a*R*,8a*S*)-**3** o from diffraction data with displacement ellipsoids shown at the 50% probability level.

The fluorocyclization of 6-substituted indoles as well as an indole bearing a phenolic nucleophile were equally successful (Figure 3). The yields and enantiomeric excesses are given for the organocatalytic reactions only. Tetrahydrofuroindoles 3u and 3v were formed in moderate yields and with *ee* values in the range of 60%. The fluorinated tetracyclic dihydrobenzofuro[2,3-*b*]indole 3w was obtained in 80% *ee*.



Figure 3. Organocatalytic asymmetric synthesis of 3 u–3 w.

From a mechanistic point of view, catalytic enantioselective fluorocyclizations benefit from the fluorination being the stereochemistry-determining step. This fact contrasts advantageously with the complications intrinsic to iodo- and bromocyclizations, which arise from the propensity of iodonium and bromonium to undergo degenerate halogen exchange.^[10] Products **3a-3w** are therefore likely to result from an irreversible fluoroquaternization at C3 followed by the intramolecular capture of the transient iminium intermediate by the pendant nucleophile. A series of observations led us to speculate on the origin of enantiocontrol. Our data indicate that similar enantiomeric excesses are observed whether the cinchona alkaloid is used in equimolar or catalytic amounts, although the catalyzed reactions were overall slightly less enantioselective. The same enantiomer is formed preferentially under both conditions (A and B), thus suggesting that the same N-fluoroammonium salt of the cinchona alkaloid is responsible for the enantiocontrol.^[7] ¹⁹F NMR studies confirmed that NFSI (¹⁹F NMR: $\delta =$ -38 ppm) is able to fully transfer fluorine to (DHQ)₂PHAL $(F(DHQ)_2PHAL^{+19}FNMR: \delta = +45 \text{ ppm})$ at room temperature within 30 min.^[11] Under conditions A (stoichiometric amount of alkaloid), fluorination is therefore likely induced by an N-fluoroammonium salt of the cinchona alkaloid as both NFSI and the alkaloid are premixed prior to the addition of the substrate. Fluorine transfer from NFSI to (DHQ)₂PHAL was however found to be ineffective at low temperature (-78°C), with only a trace amount of F-(DHQ)₂PHAL⁺ (less than 2 %) being detected by $^{19}\!F\,NMR$ spectroscopy in the presence or absence of K₂CO₃. Experimentally, we observed that when all components of the organocatalytic reaction are mixed at -78°C, the reaction proceeds as expected with the fluorocyclized product 3a obtained in 50% ee. When the alkaloid (20 mol%) is mixed with NFSI at room temperature prior to reducing the temperature to -78 °C and addition of the prochiral indole 1a (conditions B), 3a is isolated with a slightly improved ee value of 66%. Taken together, these experimental observations suggest that a reaction pathway that does not involve F(DHQ)₂PHAL⁺ may operate, and thus account for the observed enantioselectivity. Associative complexation of the alkaloid with the substrate, for example through hydrogen bonding with the pendant nucleophile, cannot be ruled out.^[12] This hypothesis would be consistent with the level of selectivity found to be dependent on the nature of the nucleophile. Indeed, while 3a and 3l (X = O, NTs) were generated with 66% and 64% ee, respectively, the selectivity was markedly increased for 3p (X = NCOMe, 80% ee), 3r $(X = NCO_2Me, 74\% ee), 3s (X = NCO_2Bn, 77\% ee) and 3t$ (X = NBoc, 78% ee).

Prochiral indole **2**, which bears an O nucleophile at the nitrogen atom, was also subjected to fluorocyclization (Scheme 1). Extensive screening of various alkaloids showed that asymmetric fluorocyclization of **2** using equimolar amounts of Selectfluor and $(DHQ)_2PHAL$ delivered the difluorinated tricyclic tetrahydrooxazolo[3,2-*a*]indole **4** in



Scheme 1. Asymmetric difluorocyclization of **2**, and crystal structure of **4**. Displacement ellipsoids shown at the 50% probability level.

Communications

50% yield and with 68% *ee*. The identity of **4** was unambiguously confirmed by single crystal X-ray diffraction studies.^[8] The stereochemical stability of **4** was assessed by measuring the enantiomeric excess at various times. No variation or decrease was observed, thus testifying to the remarkable stereochemical integrity of these compounds under the reaction conditions. This result is likely due to the presence of the proximal gem-difluoro motif, which prevents postreaction racemization. A plausible reaction pathway accounting for these difluorocyclizations involves the formation of 3-fluoroindole, which subsequently undergoes enantioselective fluorocyclization. Our attempt to prepare **4** under organocatalytic conditions B was equally successful (60% *ee*). This reaction represents a unique example of organocatalytic asymmetric dihalocyclization.

In conclusion, we have developed an organocatalytic route to enantioenriched fluorinated heterocycles. These are the first examples of organocatalyzed asymmetric fluorocyclizations. The process installs the fluorine substituent on a quaternary benzylic stereogenic carbon center and leads to new fluorinated analogues of natural products featuring the hexahydropyrrolo[2,3-b]indole or the tetrahydro-2*H*-furo-[2,3-b]indole skeleton. We have also demonstrated that asymmetric dihalocyclization is a feasible process by taking advantage of the newly formed gem-difluoro motif to prevent postreaction racemization. A catalytic asymmetric variant is also presented. Detailed kinetic studies are under way to elucidate the mechanism and origin of enantioinduction in these processes.

Experimental Section

General procedure: $(DHQ)_2PHAL$ (20 mol%) and NFSI (1.2 equiv) in acetone (1.5 mL) were stirred under argon at RT for 30 min. K₂CO₃ (6.0 equiv) was then added to the solution, and the reaction mixture was stirred for 30 min at -78 °C. A precooled solution (-78 °C) of the indole (1 equiv) in acetone (0.5 mL per 20 mg of indole) was added dropwise to the catalyst solution and the reaction was stirred at the same temperature overnight. The reaction mixture was evaporated and the residue was purified on neutral alumina (hexane/EtOAc = 6:4) to give the fluorocyclized product.

Received: May 8, 2011 Published online: July 12, 2011

Keywords: alkaloids · asymmetric catalysis · fluorocyclizations · indoles · organocatalysis

[1] For recent reviews, see: a) A. Castellanos, S. P. Fletcher, Chem. Eur. J. 2011, 17, 5766-5776; b) G. Chen, S. Ma, Angew. Chem. 2010, 122, 8484-8486; Angew. Chem. Int. Ed. 2010, 49, 8306-8308; for selected iodocyclizations, see: c) G. E. Veitch, E. N. Jacobsen, Angew. Chem. 2010, 122, 7490-7493; Angew. Chem. Int. Ed. 2010, 49, 7332-7335; d) Z. Ning, R. Jin, J. Ding, L. Gao, Synlett 2009, 2291-2294; e) H. Y. Kwon, C. M. Park, S. B. Lee, J.-H. Youn, S. H. Kang, Chem. Eur. J. 2008, 14, 1023-1028; f) A. Sakakura, A. Ukai, K. Ishihara, Nature 2007, 445, 900-903; g) J. M. Garnier, S. Robin, G. Rousseau, Eur. J. Org. Chem. 2007, 3281-3291; h) S. H. Kang S. Y. Kang, C. M. Park, H. Y. Kwon, M. Kim, Pure Appl. Chem. 2005, 77, 1269-1276; i) J. Haas, S. Bissmire, T. Wirth, Chem. Eur. J. 2005, 11, 5777-5785; j) M. Wang, L. X. Gao, W. Yue, W. P. Mai, Synth. Commun. 2004, 34, 1023-1032; k) M. Wang, L. X. Gao, W. P. Mai, A. X. Xia, F. Wang, S. B. Zhang, J. Org. Chem. 2004, 69, 2874-2876; 1) S. H. Kang, C. M. Park, S. B. Lee, M. Kim, Synlett 2004, 1279-1281; m) S. H. Kang, S. B. Lee, C. M. Park, J. Am. Chem. Soc. 2003, 125, 15748-15749; n) J. Haas, S. Piguel, T. Wirth, Org. Lett. 2002, 4, 297-300; o) X.-L. Cui, R. S. Brown, J. Org. Chem. 2000, 65, 5653-5658; p) O. Kitagawa, T. Taguchi, Synlett 1999, 1191-1199; q) R. B. Grossman, R. J. Trupp, Can. J. Chem. 1998, 76, 1233-1237; r) T. Inoue, O. Kitagawa, A. Saito, T. Taguchi, J. Org. Chem. 1997, 62, 7384-7389; s) T. Inoue, O. Kitagawa, S. Kurumizawa, O. Ochiai, T. Taguchi, Tetrahedron Lett. 1995, 36, 1479-1482; t) T. Inoue, O. Kitagawa, O. Ochiai, M. Shiro, T. Taguchi, Tetrahedron Lett. 1995, 36, 9333-9336; u) O. Kitagawa, T. Hanano, K. Tanabe, M. Shiro, T. Taguchi, J. Chem. Soc. Chem. Commun. 1992, 1005-1007; for chlorolactonizations, see: v) A. Jaganathan, A. Garzan, D. C. Whitehead, R. J. Staples, B. Borhan, Angew. Chem. 2011, 123, 2641-2644; Angew. Chem. Int. Ed. 2011, 50, 2593-2596; w) D. C. Whitehead, R. Yousefi, A. Jaganathan, B. Borhan, J. Am. Chem. Soc. 2010, 132, 3298-3300; for bromocyclizations, see: x) W. Zhang, S. Zheng, N. Liu, J. B. Werness, I. A. Guzei, W. Tang, J. Am. Chem. Soc. 2010, 132, 3664-3665; y) K. Murai, T. Matsushita, A. Nakamura, S. Fukushima, M. Shimura, H. Fujioka, Angew. Chem. 2010, 122, 9360-9363; Angew. Chem. Int. Ed. 2010, 49, 9174-9177; z) L. Zhou, C. K. Tan, X. Jiang, F. Chen, Y-Y. Yeung, J. Am. Chem. Soc. 2010, 132, 15474-15476.

- [2] S. Purser, P. R. Moore, S. Swallow, V. Gouverneur, *Chem. Soc. Rev.* 2008, *37*, 320–330.
- [3] For selected reviews, see: a) S. Lectard, Y. Hamashima, M. Sodeoka, Adv. Synth. Catal. 2010, 352, 2708-2732; b) V. A. Brunet, D. O'Hagan, Angew. Chem. 2008, 120, 1198-1201; Angew. Chem. Int. Ed. 2008, 47, 1179-1182; c) J.-A. Ma, D. Cahard, Chem. Rev. 2008, 108, PR1; d) C. Bobbio, V. Gouverneur, Org. Biomol. Chem. 2006, 4, 2065-2075; for a recent example, see: e) P. Kwiatkowski, T. D. Beeson, J. C. Conrad, D. W. C. MacMillan, J. Am. Chem. Soc. 2011, 133, 1738-1741.
- [4] a) S. C. Wilkinson, O. Lozano, M. Schuler, M. Pacheco, R. Salmon, V. Gouverneur, Angew. Chem. 2009, 121, 7217–7220; Angew. Chem. Int. Ed. 2009, 48, 7083–7086; for a recent review on fluorocyclization, see: b) S. C. Wilkinson, R. Salmon, V. Gouverneur, Future Med. Chem. 2009, 1, 847–863.
- [5] For numerous examples, see: K. C. Nicolaou, S. A. Snyder *Classics in Total Synthesis II*, 1st ed., Wiley-VCH, Weinheim, 2003; K. C. Nicolaou, E. J. Sorensen *Classics in Total Synthesis*, 1st ed., Wiley-VCH, New York, 1996.
- [6] a) N. Shibata, T. Tarui, Y. Doi, K. L. Kirk, Angew. Chem. 2001, 113, 4593–4595; Angew. Chem. Int. Ed. 2001, 40, 4461–4463;
 b) Y. Takeuchi, T. Tarui, N. Shibata, Org. Lett. 2000, 2, 639–642.
- [7] T. Ishimaru, N. Shibata, T. Horikawa, N. Yasuda, S. Nakamura, T. Toru, M. Shiro, *Angew. Chem.* 2008, 120, 4225–4229; *Angew. Chem. Int. Ed.* 2008, 47, 4157–4161.
- [8] For full details, see the Supporting Information. Compounds 3 are unstable under acidic conditions or when kept neat over an extended period of time, as they slowly convert into defluorinated oxindoles. These compounds are stable in solution (benzene, acetonitrile, acetone).
- [9] Low-temperature single-crystal diffraction data were collected by using an Oxford Diffraction (Agilent) SuperNova with $Cu_{K\alpha}$ radiation (**30**) or a Nonius KCCD diffractometer with $Mo_{K\alpha}$ radiation (**4** & **5**). Data were reduced by using CrysAlisPro or DENZO/SCALEPACK,^[13] solved by using SIR92^[14] and refined within the CRYSTALS suite.^[15] Data were also collected for **31**, but indicated the structure was modulated at 150 K and room temperature, and thus it could not be solved. Compound **30** was found to be a non-merohedral twin, thus the Rogers, Flack, and Hooft parameters conventionally used as measures of correct-

8108 www.angewandte.org



ness of the absolute structure,^[16] were inappropriate in this context and the crystal was treated as a four-component twin as described in the Supporting Information (CIF). The four twin scale factors refined to c. 0.6 0.4 0 0 (for the identity; rotation about [-1 1 0]; reflection about (-1 1 0) and inversion, respectively). Recrystallization was performed with enantioenriched 30 (e.r. 93:7), it was therefore concluded that the crystal was twinned purely by rotation (i.e., each component is a single enantiomer) and the absolute configuration was tentatively assigned on the basis that a crystal of the major enantiomer was selected. Full refinement details are given in the Supporting Information. CCDC 827531 (30), 827532 (4), 827533 (oxindole of 3m) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac. uk/data_request/cif.; The absolute configuration for 30 corroborates with the sense of enantiocontrol observed upon catalytic asymmetric electrophilic fluorination of structurally related oxindoles with (DHQ)₂PHAL.^[7]

[10] S. E. Denmark, M. T. Burk, Proc. Natl. Acad. Sci. USA 2010, 107, 20655–20660.

- [11] C. Baudequin, J.-F. Loubassou, J.-C. Plaquevent, D. Cahard, J. Fluorine Chem. 2003, 122, 189–193.
- [12] For cinchona alkaloid catalyzed alcoholysis of *meso* anhydrides that proceeds through a general base catalysis mechanism, see: H. Li, X. Liu, F. Wu, L. Tang, L. Deng, *Proc. Natl. Acad. Sci. USA* 2010, *107*, 20625–20629.
- [13] "Processing of X-ray Diffraction Data Collected in Oscillation Mode": Z. Otwinowski, W. Minor in *Methods Enzymology*, *Vol. 276* (Eds.: C. W. Carter, R. M. Sweet), Academic Press, New York, **1997**.
- [14] A. Altomare, G. Cascarano, C. Giacovazzo, A. Guagliardi, M. C. Burla, G. Polidori, M. Camalli, J. Appl. Crystallogr 1994, 27, 435.
- [15] a) P. W. Betteridge, J. R. Carruthers, R. I. Cooper, K. Prout, D. J. Watkin, J. Appl. Crystallogr. 2003, 36, 1487; b) R. I. Cooper, R. O. Gould, S. Parsons, D. J. Watkin, J. Appl. Crystallogr. 2002, 35, 168–174; c) R. I. Cooper, A. L. Thompson, D. J. Watkin, J. Appl. Crystallogr. 2010, 43, 1100–1107.
- [16] A. L. Thompson, D. J. Watkin, *Tetrahedron: Asymmetry* **2009**, 20, 712–717.