

Nonpeptide Angiotensin II Receptor Antagonists. Synthesis and Biological Activity of Potential Prodrugs of Benzimidazole-7-carboxylic Acids¹

Keiji Kubo,*† Yasuhisa Kohara,† Yoshinobu Yoshimura,† Yoshiyuki Inada,† Yumiko Shibouta,† Yoshiyasu Furukawa,† Takeshi Kato,‡ Kohei Nishikawa,† and Takehiko Naka§

Pharmaceutical Research Division, Pharmaceutical Development Division, and Discovery Research Division, Pharmaceutical Group, Takeda Chemical Industries, Ltd., 17-85 Jusohonmachi 2-chome, Yodogawaku, Osaka 532, Japan

Received March 15, 1993

In order to improve the oral bioavailability (BA) of 2-butyl-1-[[2'-(1*H*-tetrazol-5-yl)biphenyl-4-yl]methyl]-1*H*-benzimidazole-7-carboxylic acid (**3**: CV-11194) and 2-ethoxy-1-[[2'-(1*H*-tetrazol-5-yl)biphenyl-4-yl]methyl]-1*H*-benzimidazole-7-carboxylic acid (**4**: CV-11974), novel angiotensin II (AII) receptor antagonists, chemical modification to yield prodrugs has been examined. After selective tritylation of the tetrazole rings in **3** and **4**, treatment of *N*-tritylated benzimidazole-7-carboxylic acids (**6**, **7**) with a variety of alkyl halides, followed by deprotection with hydrochloric acid, afforded esters of **3** and **4**. Mainly 1-(acyloxy)alkyl esters and 1-[(alkoxycarbonyl)oxy]alkyl esters, double ester derivatives, were synthesized. Their inhibitory effect on AII-induced pressor response in rats and oral BA were investigated. (Pivaloyloxy)methyl and (±)-1-[(cyclohexyloxy)carbonyl]oxyethyl esters of **3** and **4** showed marked increases in oral bioavailability which significantly potentiated the inhibitory effect of the parent compounds on AII-induced pressor response. Among them, (±)-1-[(cyclohexyloxy)carbonyl]oxyethyl 2-ethoxy-1-[[2'-(1*H*-tetrazol-5-yl)biphenyl-4-yl]methyl]-1*H*-benzimidazole-7-carboxylate (**10s**, TCV-116) was selected as a candidate for clinical evaluation.

The renin-angiotensin system (RAS) plays an essential role in the regulation of blood pressure and seems to be critically involved in the development and maintenance of hypertension as well as congestive heart failure.² Angiotensin II (AII), an octapeptide produced from angiotensin I by the action of angiotensin converting enzyme localized on the endothelium of blood vessels in the lungs, kidneys, and many other organs, is the primary effector component of the RAS. AII is a powerful vasoconstrictor that exerts its action by interacting with specific receptors which are present on cell membranes. One of the possible modes of interrupting the RAS is AII receptor antagonism, and the discovery of nonpeptide AII antagonists, **1a** (CV-2973), **1b** (CV-2961),³ and **2** (DuP 753: losartan)⁴ (Chart I), has stimulated the recent development of new AII receptor antagonists.

In previous reports,⁵ we have described the discovery of novel nonpeptide AII receptor antagonists 2-butyl-1-[[2'-(1*H*-tetrazol-5-yl)biphenyl-4-yl]methyl]-1*H*-benzimidazole-7-carboxylic acid (**3**, CV-11194)^{5a} and 2-ethoxy-1-[[2'-(1*H*-tetrazol-5-yl)biphenyl-4-yl]methyl]-1*H*-benzimidazole-7-carboxylic acid (**4**, CV-11974)^{5b} (Chart II) which are more potent and have a longer duration of action than **2**.

Although **3** and **4** are very potent AII antagonists, they were found to be absorbed rather inefficiently upon oral administration as indicated by the fact that much lower plasma concentrations were attained after oral administration as compared to parenteral administration. It is reasonable to assume that this is due to the highly polar character of these AII antagonists possessing two acidic groups, a carboxyl group and a tetrazole ring, and that a

Chart I





Chart II



transient masking of these groups could improve oral absorption. Therefore, we investigated the possibility of improving the oral bioavailability (BA) of **3** and **4** by the prodrug approach, and here we describe the results.

Transient masking could be achieved by esterification of the carboxyl group, but preliminary experiments revealed that simple alkyl esters of **3** were hydrolyzed slowly in rat small intestine homogenate, rat plasma, and


* Address for correspondence: Keiji Kubo, Pharmaceutical Research Division, Takeda Chemical Industries, Ltd., 17-85 Jusohonmachi 2-chome, Yodogawaku, Osaka 532, Japan.

† Pharmaceutical Research Division.

‡ Pharmaceutical Development Division.

§ Discovery Research Division.

Scheme I



rat liver homogenate, indicating that a special type of ester would be needed to fulfill the requirement of rapid hydrolysis during passage into the blood stream. Well-known ester prodrugs formed by esterification of a carboxyl group are (acyloxy)methyl esters and [(alkoxycarbonyl)-oxy]methyl esters, double ester types, frequently utilized in the field of β -lactam antibiotics to improve oral absorption.⁶ Double esters are hydrolyzed rapidly in the blood and tissues of several animal species and humans. Hydrolysis of a double ester is thought to proceed *via* a chemically unstable 1-hydroxyalkyl ester which rapidly collapses to the parent compound and the aldehyde. Since an *N*-(1-hydroxyalkyl)tetrazole also seems to be chemically unstable enough to regenerate the parent tetrazole, the tetrazole rings of 3 and 4 can also be used in prodrug synthesis.


As an alternative to finding suitable enzymatically labile esters for acidic drugs, chemically labile esters have also been examined. In this respect, the principle of "neighboring group participation" or intramolecularly assisted cleavage has been an excellent source of ideas for prodrug design.⁷ We have studied the use of disubstituted aminoalkyl esters and a hydroxyalkyl ester of 3 in an effort to improve oral absorption.

Chemistry

The compounds prepared for this study are shown in Tables I and III, and synthetic methods are outlined in Schemes I-III.

Direct alkylation of 3 or 4 with alkyl halides seemed to be the most convenient route to obtain a variety of esters. Since these compounds have three possible positions for alkylation, the carboxyl group and positions N-1 and N-2 in the tetrazole ring, we first examined regioselective methylation of 3 to obtain its methyl ester (8a). Methylation with equimolar amounts of methyl iodide in dimethylformamide (DMF) under ice-cooling and using potassium carbonate as a base gave *N*-methyltetrazoles (5a, 5b), which were separated by flash column chromatography to afford the 1-methyl isomer (5a) in 52% yield and the 2-methyl isomer (5b) in 17% yield (Scheme I). Their structures were assigned based on the fact that in ¹³C NMR spectra the carbons at the 5-position of the tetrazole rings in 5a and 5b appeared at δ 154.32 and δ 164.24, respectively, which are in accord with data reported

Scheme II




for 1- and 2-methyl-5-phenyltetrazole (δ 154.2 and δ 164.25, respectively).⁸ Furthermore, this assignment was supported by the fact that the carbon of the *N*-methyl group in 5a appeared upfield (δ 33.37) from that in 5b (δ 39.20). Similar tendency was reported for 1- and 2-methyl-5-phenyltetrazole (δ 34.95 and δ 39.7, respectively).⁹

Although selective alkylation of the carboxyl group failed, these results suggested the possibility of selective protection of the tetrazole ring. We chose a trityl group as the protecting group for the tetrazole ring, since this protecting group can be removed at the appropriate time by mild acid treatment. Tritylation of 3 with trityl chloride under ice-cooling in the presence of triethylamine gave an *N*-trityltetrazole (6), whose alkylated position was not determined, in high yield. Methylation of 6 with methyl iodide followed by deprotection with 1 N HCl furnished a monomethylated product which was identified as the methyl ester (8a) produced by acid-catalyzed esterification of 3.^{5a} Using this method, a variety of esters of 3 and 4 were synthesized from *N*-trityltetrazoles (6, 7) and alkyl halides¹⁰ as shown in Scheme II and Table I.

Monoalkylation of 3 with iodomethyl pivalate^{10f} afforded *N*-alkylated products (11a, 11b) as an inseparable regioisomer mixture whose ratio was estimated from the integration of the methylene groups (δ 5.39 and δ 6.36) of the (pivaloyloxy)methyl groups in the ¹H NMR spectrum of the mixture as being about 3:1. The assignment of methylene groups was deduced from chemical shifts of the *N*-methyl groups in 5a and 5b (δ 3.15 and δ 4.21, respectively). The compounds whose carboxyl groups and tetrazole rings were masked (12a, 12b) were prepared similarly by reacting 3 with 3.6 equiv of 4-(bromomethyl)-5-methyl-2-oxo-1,3-dioxolene^{10g} as an inseparable regioisomer mixture (Scheme III). The ratio was estimated from the integration of the methylene groups (δ 4.54 and δ 5.41) of the (5-methyl-2-oxo-1,3-dioxol-4-yl)methyl groups in the ¹H NMR spectrum of the mixture as being about 3:2.


Results and Discussion

Initially, analogues of 3 were evaluated *in vivo* for inhibitory effect on the pressor response induced by AII (100 ng/kg, iv) in conscious rats, and the results are listed in Table I.¹¹ Enhancement of the inhibitory activity was observed for double esters (9a-d), whereas simple alkyl (methyl, ethyl, butyl) esters (8a-c),^{5a} 2-aminoethyl esters (8e-g), hydroxyethyl esters (8h), and *N*-alkylated tetrazole analogues (5a, 5b, 11a,b) showed activity comparable to or less than 3. Masking of the tetrazole ring slightly improved the activity (11a, 11b vs 3), but additional masking of the tetrazole ring in 9d failed to increase the activity (9d vs 12a, 12b). These results suggest that masking the carboxyl group is preferable to masking the

Table I. Inhibitory Effects of Analogues of **3** on AII-Induced Pressor Response and Bioavailabilities (BA) in Rats


compd	R ¹	dose, mg/kg po	% inhibition ^a at time, h							BA, ^b %
			0.5	1	2	3	5	7	24	
3 (CD-11194)	H	1	8	33	45	49	55	53	42	5.7
8a	Me	1	11	52	50	47	38	27	11	10.9
8b	Et	3	7	25	46	51	68	63	47	2.4
8c	Bu	30	16	40	80	78	88	90	76	0.4
8d	CH ₂ (4-pyridyl)	3	4	16	18	16	18	8	NT ^c	0.6
8e	CH ₂ CH ₂ NMe ₂	1	0	19	27	32	26	14	9	NT
8f	CH ₂ CH ₂ (morpholino)	10	67	94	96	96	96	96	66	NT
8g	CH ₂ CH ₂ (piperidino)	1	20	47	46	69	62	49	28	NT
8h	CH ₂ CH ₂ OH	1	7	7	10	11	21	25	20	0.5
9a	CH ₂ OCO- <i>t</i> -Bu	1	58	90	96	95	92	86	50	52.8
9b	CH(Me)OCOOEt	1	64	89	84	89	81	75	61	NT
9c	CH(Me)OCOO- <i>c</i> -hexyl	1	58	85	84	81	78	79	47	60.8
9d	CH ₂ (MD) ^d	1	32	63	81	88	75	77	55	NT
5a	H (1-Me) ^e	1	12	29	29	56	59	53	16	7.3
5b	H (2-Me) ^e	1	22	26	6	9	13	26	23	NT
11a, 11b^f	H (1-, 2-CH ₂ OCO- <i>t</i> -Bu) ^e	1	7	4	33	63	67	69	32	NT
12a, 12b^g	CH ₂ (MD) ^d (CH ₂ (MD) ^d) ^e	1	7	9	33	60	59	46	17	NT
2 (DuP 753)		1	5	1	12	21	35	34	17	NT

^a Percent inhibition of the AII (100 ng/kg iv)-induced pressor response at each time after administration of the test compounds in conscious male Sprague-Dawley rats ($n = 2$). The inhibition of the pressor response to AII was calculated from duplicate experiments except **2** (DuP 753) ($n = 4$). The inhibitory effect (percent inhibition) may vary less than 30%. The data in Figure 1 are indicative of the variation measured throughout this study. ^b Bioavailability was calculated from the ratio of the area under the plasma concentration-time curve (AUC) for 0 to infinity after oral dosing (10 mg/kg equivalent to **3**) of the test compounds to that after intravenous administration of **3** at a dose of 1 mg/kg in rats ($n = 3$). The following equation was used for the calculation: $BA = (AUC_{po}/AUC_{iv}) \times (dose_{iv}/dose_{po}) \times 100$. ^c NT means "not tested". ^d MD means "5-Methyl-2-oxo-1,3-dioxolen-4-yl". ^e Substituents on the tetrazole ring. ^f A mixture of the regioisomers (**11a**/**11b** = 3:1) was used. ^g A mixture of the regioisomers (**12a**/**12b** = 3:2) was used.

Scheme III

tetrazole ring. We examined some of these compounds in BA experiments in rats, assessing as **3** (Table I), and determined the rate of hydrolysis to **3** *in vitro* using 1% rat small intestine homogenate, 10% rat plasma, and 2% rat liver homogenate at 37 °C (Table II). The BA as **3** was increased by about 10-fold for double esters (**9a**, **9c**), which had a rapid onset of action. The BA as **3** after administration of the 1-methyltetrazole analogue (**5a**) was slightly improved, but that of the simple alkyl esters (**8b**, **8c**) was lower than that of **3**. The BA as **3** of the simple alkyl esters was decreased by increasing alkyl chain length in R¹. Although the methyl ester (**8a**) has a BA of ca. 30%, its BA as **3** was only 10.9%. These results indicate that the esterification of the carboxyl group in **3** results in an

improvement of the oral BA by increasing the lipophilicity and the absence of a negative charge due to the carboxyl group during transport across the gut lumen. The low BA as **3** seen with **8a** and **8c** could be explained by their high resistance to enzymatic hydrolysis. No amount of **3** was detected after incubating **8a** and **8c** in the tissue homogenate for 60 min. The stability seemed to be ascribed to steric hindrance around the ester groups in these molecules. The compounds bearing a (pivaloyloxy)methyl group at the carboxyl group (**9a**) or the tetrazole ring (mixture of **11a** and **11b**) were hydrolyzed to **3** with a half-life ($t_{1/2}$) of 1.08 and 22.7 min, respectively, in 1% small intestine homogenate. In 10% plasma and 2% liver homogenate, **9a** and **11a,b** were also hydrolyzed rapidly to **3**, but the

Table II. Half-Lives ($t_{1/2}$) of Analogues of **3** and **4** in 1% Rat Small Intestine Homogenate, 2% Rat Liver Homogenate, or 10% Rat Plasma at 37 °C^a


compd	tissue	$t_{1/2}$, min mean \pm SD, $n = 3$
8a	1% SI ^b	stable ^c
	10% plasma	stable
	2% liver	stable
8c	1% SI	stable
8d	1% SI	stable
9a	1% SI	1.08 \pm 0.06
	10% plasma	0.69 \pm 0.01
	2% liver	3.61 \pm 0.23
11a, 11b^d	1% SI	22.7 \pm 1.0
	10% plasma	3.94 \pm 0.32
	2% liver	9.80 \pm 0.19
10e	1% SI	0.30 \pm 0.01
	10% plasma	0.15 \pm 0.00
	2% liver	0.48 \pm 0.01
10s	1% SI	4.92 \pm 0.41
	10% plasma	2.73 \pm 0.15
	2% liver	2.29 \pm 0.06

^a Final concentration was 10 μ g/mL. Half-lives ($t_{1/2}$) were calculated from the concentration change of the analogues. ^b Rat small intestine homogenate. ^c **3** was not detected at 60 min after incubation. ^d A mixture of the regioisomers (**11a/11b** = 3:1) was used.

hydrolysis rate of **9a** was faster than that of **11a,b**. The difference in hydrolysis rate was reflected in their BA and their inhibitory effect on the pressor response induced by AII. These results proved double esters to be efficient prodrugs for **3**.

The goal of this research project was to find an optimal promoiety for **4** which is a more potent AII receptor antagonist than **3**. We applied the "double ester" prodrug technique to **4** as in the case of **3**. The inhibitory effect of **4** and its analogues on AII-induced pressor response in conscious rats and the BA as **4** after administration of these compounds were determined, and the results are shown in Table III. Although faster onset of the inhibitory effect after administration of the (acetyloxy)methyl ester (**10a**), (propionyloxy)methyl ester (**10b**), (butyryloxy)methyl ester (**10c**), and (isovaleryloxy)methyl ester (**10d**) was observed, BA as **4** after administration of these esters was slightly lower than after administration of **4**. The [(cyclohexylcarbonyl)oxy]methyl ester (**10g**) and (benzoyloxy)methyl ester (**10h**) had an inhibitory effect, but the onset of action was slow. The inhibitory effect was significantly greater with 1-[[[(alkyloxy)carbonyl]oxy]ethyl ester (**10o-s**), whereas it was only slightly greater with [[[(alkyloxy)carbonyl]oxy]methyl esters (**10k-n**). Among the double esters, the (pivaloyloxy)methyl ester (**10e**) and (\pm)-1-[[[(cyclohexyloxy)carbonyl]oxy]ethyl ester (**10s**) showed a pronounced inhibitory effect profile, fast onset and prolonged duration of action, and higher BA as **4** which was 7 times that of **4**. These esters (**10e**, **10s**) were hydrolyzed to **4** with a $t_{1/2}$ of 0.30 and 4.92 min, respectively, in 1% small intestine homogenate (Table II).

The (\pm)-1-[[[(cyclohexyloxy)carbonyl]oxy]ethyl ester (**10s**) was selected as a candidate for clinical evaluation under the code name TCV-116, since adverse effects of the promoiety (pivalic acid) of **10e** have been reported.¹² As shown in Figure 1, the inhibitory effects of **10s** at 0.3 and 0.03 mg/kg po were more potent and long lasting than those of the parent compound (**4**). After oral administration of **10s** to rats at a dose of 1 mg/kg, **10s** was not detected in the plasma. In isolated rabbit aortic strips, **10s** and **4** inhibited AII (10^{-8} M)-induced contraction with IC_{50} values of 2.0×10^{-8} M and 2.0×10^{-10} M,^{5b}

**Figure 1.** Inhibitory effects of **10s** (TCV-116) and **4** (0.03 and 0.3 mg/kg po) on AII (100 ng/kg iv)-induced pressor response in conscious normotensive rats ($n = 4-5$).

respectively.¹¹ These results indicate that **10s** is rapidly transformed into **4** after the oral administration and then acts as a potent AII antagonist in the body.

Conclusion


We successfully performed chemical modification of novel AII receptor antagonists, **3** and **4**, to improve oral absorption. Although simple alkyl esters (**8a-c**) and β -substituted ethyl esters (**8e-h**) did not have improved BA as **3** or increased AII antagonistic potency, double esters were found to be effective prodrugs of **3** and **4**. Among them, the (pivaloyloxy)methyl ester (**10e**) and (\pm)-1-[[[(cyclohexyloxy)carbonyl]oxy]ethyl ester (**10s**) of **4** showed significantly enhanced BA as **4** and AII antagonistic potency. On the basis of its profile, **10s** has been selected for clinical evaluation as an antihypertensive agent and is the first example of applying the double ester prodrug technique to an orally effective AII antagonistic agent. We think that this double ester prodrug technique is synthetically easy for good overall yield from diacidic parent compounds and can be used with other diacidic nonpeptide AII antagonists¹³ to increase their oral absorption. Further detailed studies on **10s** are in progress, and the results will be reported elsewhere.¹⁴

Experimental Section

All melting points were determined on a Yanagimoto micro melting point apparatus and are uncorrected. The infrared (IR) spectra were recorded on a Hitachi 215 grating infrared spectrophotometer. The proton nuclear magnetic resonance (¹H NMR) spectra were recorded on a Varian Gemini-200 (200 MHz) spectrometer. The carbon nuclear magnetic resonance (¹³C NMR) spectra were recorded on a JEOL JNM-GX270 (67.8 MHz). Chemical shifts are given in δ values (ppm) using tetramethylsilane as the internal standard, and coupling constants (J) are given in hertz. Column chromatography was performed using silica gel (Wakogel C-300 or Merck Art 9385). The biological assay was performed as previously described.⁵

2-Butyl-1-[[2'-(1-(and 2-methyltetrazol-5-yl)biphenyl-4-yl)methyl]-1H-benzimidazole-7-carboxylic Acid (5a**, **5b**).** To an ice-cooled mixture of **3** (0.94 g, 2.0 mmol) and potassium carbonate (0.28 g, 2.0 mmol) in DMF (6 mL) was added dropwise a solution of methyl iodide (0.28 g, 2.0 mmol) in DMF (2 mL), and the resulting mixture was stirred at that temperature for 20 min. The reaction mixture was diluted with water and extracted with EtOAc. The extract was washed with water and dried (MgSO₄). The solvent was evaporated in vacuo, and the residue was purified by flash column chromatography (CHCl₃/MeOH = 10:1). The first eluent was concentrated in vacuo, and the product was recrystallized from EtOAc-CHCl₃ to give **5b** (0.16 g, 17%)

Table III. Inhibitory Effects of Analogues of 4 on AII-Induced Pressor Response and Bioavailabilities (BA) in Rats



compd	R¹	% inhibition ^a at time, h							BA, ^b %
		0.5	1	2	3	5	7	24	
4 (CV-11974)	H	7	24	51	67	85	91	43	5.0
10a	CH ₂ OCOMe	66	84	79	78	72	59	23	4.1
10b	CH ₂ OCOEt	76	76	60	60	38	31	—	4.3
10c	CH ₂ OCOPr	100	84	78	67	54	48	6	3.5
10d	CH ₂ OCO- <i>i</i> -Pr	80	40	56	52	44	20	6	2.7
10e	CH ₂ OCO- <i>t</i> -Bu	42	77	88	86	87	86	43	34.9
10f	CH ₂ OCO- <i>c</i> -Pen	7	31	61	65	62	54	9	NT ^c
10g	CH ₂ OCO- <i>c</i> -hexyl	0	5	76	85	89	80	14	NT
10h	CH ₂ OCOPh	—	—	23	56	67	76	6	NT
10i	CH ₂ OCOCH=CHPh ^d	—	36	80	80	61	64	3	NT
10j	CH(Me)OCOMe	86	90	92	86	84	75	28	8.8
10k	CH ₂ OCOO- <i>i</i> -Pr	23	55	79	72	68	57	28	NT
10l	CH ₂ OCOOCH ₂ - <i>c</i> -Pr	10	49	57	58	44	30	22	NT
10m	CH ₂ OCOO- <i>c</i> -Pen	5	14	55	68	77	64	14	NT
10n	CH ₂ OCOO- <i>c</i> -hexyl	1	—	4	15	53	54	11	NT
10o	CH(Me)OCOEt	74	87	95	89	81	71	7	4.8
10p	CH(Me)OCOO- <i>i</i> -Pr	70	91	84	79	80	67	12	NT
10q	CH(Me)OCOCH ₂ Et ₂	16	72	86	84	63	53	13	8.0
10r	CH(Me)OCOO- <i>c</i> -Bu	35	77	86	83	79	60	25	10.8
10s	CH(Me)OCOO- <i>c</i> -hexyl	38	76	80	87	82	75	36	33.4
10t	CH ₂ (MD) ^e	49	83	87	85	70	44	6	9.1

^a Percent inhibition of the AII (100 ng/kg iv)-induced pressor response at each time after administration of the test compounds at dose of 0.1 mg/kg po in conscious male Sprague-Dawley rats ($n = 2$). The inhibition of the pressor response to AII was calculated from duplicate experiments except 4 and 10s ($n = 3$). The inhibitory effect (percent inhibition) may vary less than 30%. The data in Figure 1 are indicative of the variation measured throughout this study. ^b Bioavailability was calculated from the ratio of the area under plasma concentration-time curve (AUC) for infinity after oral dosing (1 mg/kg equivalent to 4) of the test compounds to that after intravenous administration of 4 at the same dose in rats ($n = 3$). ^c NT means "not tested". ^d Geometry of the double bond is trans. ^e MD means "5-methyl-2-oxo-1,3-dioxolen-4-yl".

as colorless needles: mp 226–228 °C dec; ¹H NMR (CDCl₃) δ 0.94 (3H, t, $J = 7.2$), 1.38–1.56 (2H, m), 1.79–1.94 (2H, m), 3.07 (2H, t, $J = 7.5$), 4.22 (3H, s), 5.84 (2H, s), 6.81 (2H, d, $J = 8.1$), 7.06 (2H, d, $J = 8.1$), 7.25–7.55 (4H, m), 7.74 (1H, d, $J = 7.6$), 7.81 (1H, dd, $J = 2.2$ and 6.8), 8.00 (1H, d, $J = 7.6$); ¹³C NMR (DMSO-*d*₆) δ 13.52, 21.68, 26.29, 28.59, 39.20, 47.88, 118.36, 121.74 (two peaks were overlapped), 125.26, 125.59, 125.75, 127.65, 129.10, 130.15, 130.18, 130.59, 131.52, 135.76, 139.17, 140.75, 141.37, 157.22, 164.24, 167.31; IR (KBr) 1700 cm⁻¹. Anal. (C₂₇H₂₆N₆O₂·0.7H₂O) C, H, N. The second eluent was concentrated in vacuo, and the product was recrystallized from EtOAc to give 5a (0.49 g, 52%) as colorless prisms: mp 213–214 °C; ¹H NMR (CDCl₃) δ 0.95 (3H, t, $J = 7.4$), 1.38–1.57 (2H, m), 1.80–1.95 (2H, m), 2.99 (2H, t, $J = 8.0$), 3.18 (3H, s), 5.82 (2H, s), 6.80 (2H, d, $J = 8.2$), 6.97 (2H, d, $J = 8.2$), 7.27 (1H, t, $J = 7.7$), 7.48–7.68 (4H, m), 7.80 (1H, d, $J = 7.6$), 7.98 (1H, d, $J = 7.6$); ¹³C NMR (DMSO-*d*₆) δ 13.55, 21.69, 26.49, 28.71, 33.37, 47.57, 117.82, 120.96, 122.07, 122.55, 124.74, 126.10, 127.87, 128.48, 130.16, 131.12, 131.60, 132.18, 137.30, 137.51, 140.83, 143.41, 154.32, 157.36, 167.50; IR (KBr) 1700 cm⁻¹. Anal. (C₂₇H₂₆N₆O₂·0.5H₂O) C, H, N.

2-Butyl-1-[[2'-[1-(and 2-)](pivaloyloxy)methyl]tetrazol-5-yl]biphenyl-4-yl]methyl]-1H-benzimidazole-7-carboxylic Acid (11a, 11b). The mixture of 11a and 11b (3:1) was obtained by a procedure similar to that described above in 23% yield as colorless powder (from acetone-hexane): mp 188–191 °C dec; ¹H NMR (CDCl₃) δ 0.94 (3H, t, $J = 7.4$), 0.96 and 1.18 (9H, 2s), 1.36–1.55 (2H, m), 1.77–1.93 (2H, m), 2.90–2.97 (2H, m), 5.39 and 6.36 (2H, 2 s, 3:1 ratio), 5.81 (2H, s), 6.76–6.83 (2H, m), 6.96–7.03 (2H, m), 7.20–7.29 (1H, m), 7.38–7.97 (6H, m); IR (KBr) 1760, 1600 cm⁻¹. Anal. (C₃₂H₃₄N₆O₄·0.6H₂O) C, H, N: calcd, 14.55; found, 14.13.

(5-Methyl-2-oxo-1,3-dioxolen-4-yl)methyl 2-Butyl-1-[[2'-[1-(and 2-)](5-methyl-2-oxo-1,3-dioxolen-4-yl)methyl]tetrazol-5-yl]biphenyl]methyl]-1H-benzimidazole-7-carboxy-

late (12a, 12b). The mixture of 12a and 12b (4:3) was obtained by a procedure similar to that described for 5a and 5b, except using 3.6 mol equiv of 4-(bromomethyl)-5-methyl-2-oxo-1,3-dioxolene,^{10e} in 32% yield as pale brown amorphous powder: mp 68–71 °C; ¹H NMR (CDCl₃) δ 0.96 (3H, t, $J = 7.3$), 1.38–1.57 (2H, m), 1.80–2.20 (8H, m), 2.87–2.98 (2H, m), 4.54 (1.2H, s), 4.86 and 4.89 (2H, 2s), 5.41 (0.8H, s), 5.71 and 5.76 (2H, 2s), 6.72–6.77 (2H, m), 7.03 (2H, d, $J = 8.2$), 7.21–7.84 (6H, m), 7.97 (1H, d, $J = 8.0$); IR (KBr) 1820, 1735, 1720 cm⁻¹. Anal. (C₃₆H₃₂N₆O₅·0.5H₂O) C, H, N.

2-Ethoxy-1-[[2'-[N-(triphenylmethyl)-1H-tetrazol-5-yl]biphenyl-4-yl]methyl]-1H-benzimidazole-7-carboxylic Acid (7). To an ice-cooled solution of 4 (29.0 g, 66.0 mmol) and triethylamine (7.33 g, 72.4 mmol) in CH₂Cl₂ (300 mL) was added dropwise a solution of trityl chloride (19.9 g, 71.4 mmol) in CH₂Cl₂ (80 mL), and the resulting mixture was stirred at that temperature for 1 h. The reaction mixture was washed with water and dried (MgSO₄). After evaporation of the solvent, the residue was purified by flash column chromatography (CHCl₃/MeOH = 40:1). The product was recrystallized from EtOAc-hexane to give 7 (76%) as colorless needles: mp 168–170 °C; ¹H NMR (CDCl₃) δ 1.40 (3H, t, $J = 7.1$), 4.61 (2H, q, $J = 7.1$), 5.58 (2H, s), 6.76 (2H, d, $J = 8.0$), 6.91–6.96 (8H, m), 7.12 (1H, t, $J = 7.9$), 7.17–7.41 (12H, m), 7.60 (1H, dd, $J = 1.2$ and 7.8), 7.73–7.82 (2H, m); IR (KBr) 1700 cm⁻¹. Anal. (C₄₃H₃₄N₆O₃) C, H, N.

2-Butyl-1-[[2'-[N-(triphenylmethyl)-1H-tetrazol-5-yl]biphenyl-4-yl]methyl]-1H-benzimidazole-7-carboxylic Acid (6). Compound 6 was prepared, by a similar procedure to that described above, in 76% yield as colorless needles (from EtOAc-CHCl₃): mp 185–187 °C dec; ¹H NMR (CDCl₃) δ 0.91 (3H, t, $J = 7.2$), 1.30–1.48 (2H, m), 1.72–1.90 (2H, m), 2.84 (2H, t, $J = 7.9$), 5.72 (2H, s), 6.64 (2H, d, $J = 8.0$), 6.95–7.00 (8H, m), 7.11–7.46 (13H, m), 7.65 (1H, d, $J = 1.2$ and 7.6), 7.79–7.83 (1H,

Table IV. Physicochemical Data of Analogues

compd	X ^a	yield, %	recryst solvent ^b	mp, °C	formula ^c
8d	Cl	85	A	179–180	C ₃₂ H ₂₉ N ₇ O ₂
8e	Cl	60	B	188–192 dec	C ₃₀ H ₃₃ N ₇ O ₂ ·1.5H ₂ O
8f	Cl	83	C	187–188	C ₃₂ H ₃₅ N ₇ O ₃ ·0.2H ₂ O
8g	Cl	67	B	212–214	C ₃₃ H ₃₇ N ₇ O ₂ ·0.2H ₂ O
8h	Cl	25	D	145–147	C ₂₈ H ₂₈ N ₆ O ₃ ·0.5acetone
9a	I	74	amorphous	102–105	C ₃₂ H ₃₄ N ₆ O ₄ ·0.5H ₂ O
9b	I	75	amorphous	92–95	C ₃₁ H ₃₂ N ₆ O ₅ ·0.5H ₂ O
9c	I	74	amorphous	102–105	C ₃₅ H ₃₈ N ₆ O ₅ ·0.5CHCl ₃
9d	Br	71	E	123–125	C ₃₁ H ₂₈ N ₆ O ₅ ·0.5H ₂ O
10a	Cl	38	F	152–154 dec	C ₂₇ H ₂₄ N ₆ O ₅
10b	Cl	60	F	145–150	C ₂₈ H ₂₆ N ₆ O ₅ ·0.2toluene
10c	Cl	36	G	96–100	C ₂₉ H ₂₈ N ₆ O ₅ ·0.4toluene
10d	Cl	53	F	143–145	C ₂₉ H ₂₈ N ₆ O ₅ ·0.1toluene
10e	I	69	A	144–147	C ₃₀ H ₃₀ N ₆ O ₅
10f	Cl	54	A	136–138	C ₃₁ H ₃₀ N ₆ O ₅
10g	Cl	54	A	140–142	C ₃₂ H ₃₂ N ₆ O ₅
10h	Cl	46	A	138–142	C ₃₂ H ₃₂ N ₆ O ₅ ·0.1EtOAc·0.5H ₂ O
10i	Cl	55	A	146–147	C ₃₄ H ₂₈ N ₆ O ₅ ·0.4EtOAc
10j	Cl	31	H	105–107 dec	C ₂₈ H ₂₆ N ₆ O ₅ ·0.5H ₂ O
10k	Cl	67	I	142–144 dec	C ₂₉ H ₂₈ N ₆ O ₆ ·0.5H ₂ O
10l	I	69	I	147–148 dec	C ₃₀ H ₂₈ N ₆ O ₆ ·0.1H ₂ O
10m	I	63	I	142–144 dec	C ₃₁ H ₃₀ N ₆ O ₆
10n	I	64	I	176–177 dec	C ₃₂ H ₃₂ N ₆ O ₆
10o	I	44	amorphous	85–87	C ₂₉ H ₂₈ N ₆ O ₆ ·0.3H ₂ O
10p	Cl	33	amorphous	74–76	C ₃₀ H ₃₀ N ₆ O ₆ ·1.5H ₂ O
10q	I	45	amorphous	86–88 dec	C ₃₂ H ₃₄ N ₆ O ₆ ·0.3H ₂ O
10r	I	36	amorphous	95–97 dec	C ₃₁ H ₃₀ N ₆ O ₆ ·0.5H ₂ O
10s	I	65	I	163 dec	C ₃₃ H ₃₄ N ₆ O ₆
10t	Cl	55	I	122–125 dec	C ₂₉ H ₂₄ N ₆ O ₆ ·CHCl ₃

^a X indicates halides in R¹X used for alkylation of 3 and 4. ^b A = EtOAc–hexane; B = CHCl₃–MeOH; C = EtOAc–CHCl₃; D = acetone; E = benzene–CH₂Cl₂; F = toluene; G = toluene–hexane; H = acetone–ether–hexane; I = EtOH–H₂O. ^c All compounds gave satisfactory analyses C, H, N.

m), 7.93 (1H, dd, *J* = 1.2 and 8.0); IR (KBr) 1690 cm⁻¹. Anal. (C₄₅H₃₈N₆O₂) C, H, N.

Methyl 2-Butyl-1-[[2'-(1*H*-tetrazol-5-yl)biphenyl-4-yl]methyl]-1*H*-benzimidazole-7-carboxylate (8a). To an ice-cooled mixture of 6 (0.21 g, 0.30 mmol) and potassium carbonate (70 mg, 0.51 mmol) in DMF (2 mL) was added dropwise methyl iodide (68 mg, 0.48 mmol), and the resulting mixture was stirred at room temperature for 10 min. The reaction mixture was diluted with water and extracted with EtOAc. The extract was washed with water. After evaporation of the solvent, the residue was treated with 1 N HCl (0.4 mL) at room temperature for 1.5 h in MeOH (2 mL). The reaction mixture was concentrated in vacuo and extracted with CHCl₃. The extract was washed with water and dried (MgSO₄). The solvent was evaporated in vacuo, and the residue was purified by flash column chromatography (CHCl₃/MeOH = 20:1). The product was recrystallized from EtOAc–hexane to give 8a (0.11 g, 79%) as colorless needles which was identified with authentic sample.^{5a}

(±)-1-[[[(Cyclohexyloxy)carbonyl]oxy]ethyl 2-Ethoxy-1-[[2'-(1*H*-tetrazol-5-yl)biphenyl-4-yl]methyl]-1*H*-benzimidazole-7-carboxylate (10s). A mixture of 7 (18.0 g, 26.4 mmol), cyclohexyl 1-chloroethyl carbonate (6.5 g, 31.5 mmol), potassium iodide (2.19 g, 13.2 mmol), and potassium carbonate (4.37 g, 31.6 mmol) in DMF (100 mL) was stirred at 60 °C for 2 h. The reaction mixture was diluted with water and extracted with EtOAc. The extract was washed with water. After evaporation of the solvent, the residue was treated with 1 N HCl (40 mL) at room temperature for 1.5 h in MeOH (150 mL)–CHCl₃ (30 mL). The reaction mixture was concentrated in vacuo and extracted with EtOAc. The extract was washed with water and dried (MgSO₄). The solvent was evaporated in vacuo, and the residue was purified by flash column chromatography (CHCl₃/MeOH = 20:1). The product was recrystallized from EtOH–H₂O to give 10s (10.5 g, 65.3%) as colorless crystals: mp 163 °C dec; ¹H NMR (CDCl₃) δ 1.13–1.84 (16H, m), 4.28–4.55 (3H, m), 5.65 (2H, d, *J* = 3.8), 6.72 (1H, q, *J* = 5.4), 6.81 (2H, d, *J* = 8.4), 6.93 (2H, d, *J* = 8.4), 7.03 (1H, t, *J* = 8.4), 7.22–7.23 (1H, m), 7.31–7.36 (1H, m), 7.52–7.60 (3H, m), 8.02–8.07 (1H, m); IR (KBr) 1750, 1720 cm⁻¹. Anal. (C₃₃H₃₄N₆O₆) C, H, N.

The compounds (8d–g, 9a–d, 10a–r, 10t) were prepared by a procedure similar to that described above and the results were shown in Table IV.

Hydrolysis of Prodrugs to 3 or 4 in Vitro. Each prodrug was incubated in 1% small intestine homogenate, 2% liver homogenate, or 10% plasma prepared from rats at 37 °C. An aliquot sample was withdrawn at 10, 40, 60, 90 sec, 2, 5, 10, and 30 min after incubation and deproteinized with methanol. After centrifugation, the supernatant was evaporated in vacuo under centrifugation. The residue was dissolved in 1/15 M phosphate buffer, and injected into a high-performance liquid chromatograph (HPLC) to determine the concentration of 3 or 4 produced. The HPLC system consisted of a Shimadzu LC-9A pump, ODS column (150 × 4.6 mm i.d., Yamamura Chemical, Kyoto, Japan) and a Shimadzu SPD-6AV UV detector (258 nm). The mobile phase was 0.01 M KH₂PO₄/MeCN (55/45, v/v) with a flow rate of 1 mL/min.

Administration Studies. Male Sprague-Dawley rats (Clea Japan Inc., Tokyo, Japan), weighing about 300 g, were starved but allowed free access to water for 16–18 h before experiments.

Each parent drug was administered intravenously at a dose of 1 mg/kg. The blood was taken from ophthalmic venous plexus at 0.0083, 0.00167, 0.25, 0.5, 1, 2, 3, 5, 7, and 24 h after injection and centrifuged to obtain plasma.

The parent drug (3 or 4) or each ester was given orally to rats at a dose of 10 mg/kg or 1 mg/kg as equivalent to 3 or 4, respectively. The blood was taken to obtain plasma in a similar manner. The concentration of 3 or 4 in the plasma was determined by HPLC as described above.

Acknowledgment. We wish to thank Drs. S. Terao, A. Imada, K. Meguro, and A. Nagaoka for their encouragement and helpful discussions throughout this work, and acknowledge the able technical assistance of Dr. T. Wada, Ms. M. Ojima, Mr. T. Sanada, Mr. N. Tada, Mr. Y. Sugiura, and Mr. E. Imamiya.

References

- (1) Naka, T.; Nishikawa, K.; Kato, T. *Benzimidazole Derivatives, Their Production and Use*. European Patent Application 459136, 1991.
- (2) Ferrario, C. M. The Renin-Angiotensin System: Importance in Physiology and Pathology. *J. Cardiovasc. Pharmacol.* 1990, 15 (Suppl. 3), S1–S5.

- (3) (a) Furukawa, Y.; Kishimoto, S.; Nishikawa, K. Hypotensive Imidazole-5-acetic Acid Derivatives. Jpn. Patent No. 56-71073, 71074, 1981; U.S. Patent 4340598, 4355040, 1982. (b) Furukawa, Y.; Nishikawa, K. 4-Chloro-2-phenyl-imidazole-5-acetic Acid Derivatives. Jpn. Patent No. 58-157768, 1983; U.S. Patent 504049, 1983. (c) Furukawa, Y.; Naka, T.; Kishimoto, S.; Tomimoto, M.; Matsushita, Y.; Miyake, A.; Itoh, K.; Nonpeptide Angiotensin II Receptor Antagonists: Synthetic Studies on Imidazoleacetic Acid Derivatives. *J. Takeda Res. Labs.* 1991, 50, 56-74; *Chem. Abstr.* 1991, 115, 256061t. (d) Nishikawa, K.; Shibuta, Y.; Inada, Y.; Terashita, Z.; Kawazoe, K.; Furukawa, Y.; Nonpeptide Angiotensin II Receptor Antagonists: Pharmacological Studies on Imidazoleacetic Acid Derivatives. *J. Takeda Res. Labs.* 1991, 50, 75-98; *Chem. Abstr.* 1991, 115, 270283y. (e) Chiu, A. T.; Carini, D. J.; Johnson, A. L.; McCall, D. E.; Price, W. A.; Thoolen, M. J. M. C.; Wong, P. C.; Taber, R. L.; Timmermans, P. B. M. W. M. Nonpeptide Angiotensin II Receptor Antagonists. II. Pharmacology of S-8308. *Eur. J. Pharmacol.* 1988, 157, 13-21.
- (4) For reviews, see: (a) Duncia, J. V.; Carini, D. J.; Chiu, A. T.; Johnson, A. L.; Price, W. A.; Wong, P. C.; Wexler, R. R.; Timmermans, P. B. M. W. M. The Discovery of DuP 753, a Potent, Orally Active Nonpeptide Angiotensin II Receptor Antagonist. *Med. Res. Rev.* 1992, 12, 149-191. (b) Wong, P. C.; Barnes, T. B.; Chiu, A. T.; Christ, D. D.; Duncia, J. V.; Herblin, W. F.; Timmermans, P. B. M. W. M. Losartan (DuP 753), An Orally Active Nonpeptide Angiotensin II Receptor Antagonist. *Cardiovasc. Drug. Rev.* 1991, 9, 317-339.
- (5) (a) Kubo, K.; Inada, Y.; Kohara, Y.; Sugiura, Y.; Ojima, M.; Itoh, K.; Furukawa, Y.; Nishikawa, K.; Naka, T. Nonpeptide Angiotensin II Receptor Antagonists. Synthesis and Biological Activity of Benzimidazoles. *J. Med. Chem.*, in press. (b) Kubo, K.; Kohara, Y.; Imamiya, E.; Sugiura, Y.; Inada, Y.; Furukawa, Y.; Nishikawa, K.; Naka, T. Nonpeptide Angiotensin II Receptor Antagonists. Synthesis and Biological Activity of Benzimidazolecarboxylic Acids. *J. Med. Chem.*, in press.
- (6) For a review see: Ferres, H. Pro-drugs of β -Lactam Antibiotics. *Chem. Ind. (London)* 1980, 435-440.
- (7) Bruice, T. C.; Benkovic, S. J. *Bioorganic Mechanisms*; Benjamin, W. A., Ed.; New York, 1966; Vol. 1, pp 134-145.
- (8) Butler, R. N.; McEvoy, T. M. A Carbon-13 and Proton Nuclear Magnetic Resonance Study of Annular Tautomerism and Interannular Conjugation in Some Substituted-5-aryl-tetrazoles. *J. Chem. Soc., Perkin Trans. 2* 1978, 1087-1090.
- (9) Butler, R. N.; Garvin, V. C. A Study of Annular Tautomerism, Interannular Conjugation, and Methylation Reaction of ortho-Substituted-5-aryltetrazoles Using Carbon-13 and Hydrogen-1 N.M.R. Spectroscopy. *J. Chem. Soc., Perkin Trans. 1* 1981, 390-393.
- (10) (a) Ulich, L. H.; Adams, R. The Reaction Between Acid Halides and Aldehydes. *J. Am. Chem. Soc.* 1921, 43, 660-667. (b) Kyburz, R.; Schaltegger, H.; Neuenschwander, M. Eine Neue Fulvensynthese. *Helv. Chem. Acta* 1971, 54, 1037-1046. (c) Nishimura, T.; Yoshimura, Y.; Miyake, A.; Yamaoka, M.; Takanohashi, K.; Hamaguchi, N.; Hirai, S.; Yashiki, T.; Numata, M. Orally Active 1-(Cyclohexyloxycarbonyloxy)alkyl Ester Prodrugs of Cefotiam. *J. Antibiot.* 1987, 40, 81-90. (d) Fujimoto, K.; Ishihara, S.; Yanagisawa, H.; Ide, J.; Nakayama, E.; Nakao, H.; Sugawara, S.; Iwata, M. Studies on Orally Active Cephalosporin Esters. *J. Antibiot.* 1987, 40, 370-384. (e) Sakamoto, F.; Ikeda, S.; Tsukamoto, G. Studies on Prodrugs. II. Preparation and Characterization of (5-Substituted 2-Oxo-1,3-dioxolen-4-yl)methyl Esters of Ampicillin. *Chem. Pharm. Bull.* 1984, 33, 2241-2248. (f) Yoshimura, Y.; Hamaguchi, N.; Yashiki, T. Synthesis and Oral Absorption of Acyloxyethyl Esters of 7 β -(2-(2-Aminothiazol-4-yl)acetamido)-3-((1-(2-dimethylaminoethyl)-1H-tetrazol-5-yl)thio)methyl]ceph-3-em-4-carboxylic Acid (Cefotiam). *Int. J. Pharm.* 1987, 38, 179-190. (g) Yoshimura, Y.; Hamaguchi, N.; Yashiki, T. Preparation of 1-Acyloxyethyl Esters of 7-[[2-(2-Aminothiazol-4-yl)acetamido]-3-[[1-(2-dimethylaminoethyl)-1H-tetrazol-5-yl]thio]methyl]ceph-3-em-4-carboxylic Acid (Cefotiam). *J. Antibiot.* 1986, 39, 1329-1342.
- (11) The biological assay was performed as described previously, see ref 5.
- (12) Editorials: *Lancet* 1990, 335, 631-633.
- (13) (a) Wong, P. C.; Hart, S. D.; Chiu, A. T.; Herblin, W. F.; Carini, D. J.; Smith, R. D.; Wexler, R. R.; Timmermans, P. B. M. W. M. Pharmacology of DuP 532, a Selective and Noncompetitive AT₁ Receptor Antagonist. *J. Pharmacol. Exp. Ther.* 1991, 259, 861-869. (b) Weinstock, J.; Keenan, R. M.; Samanen, J.; Hempel, J.; Finkelstein, J. A.; Franz, R. G.; Gaitanopoulos, D. E.; Girard, G. R.; Gleason, J. G.; Hill, D. T.; Morgan, T. M.; Peishoff, C. E.; Aiyar, N.; Brooks, D. P.; Fredrickson, T. A.; Ohlstein, E. H.; Ruffolo, R. R., Jr.; Stack, E. J.; Sulpizio, A. C.; Weidley, E. F.; Edwards, R. M. 1-(Carboxybenzyl)imidazole-5-acrylic Acids: Potent and Selective Angiotensin II Receptor Antagonists. *J. Med. Chem.* 1991, 34, 1514-1517. (c) Middlemiss, D.; Drew, G. M.; Ross, B. C.; Robertson, M. J.; Scopes, D. I. C.; Dowle, M. D.; Akers, J.; Cardwell, K.; Clark, K. L.; Coote, S.; Eldred, C. D.; Hamblett, J.; Hilditch, A.; Hirst, G. C.; Jack, T.; Montana, J.; Panchal, T. A.; Paton, J. M. S.; Shah, P.; Stuart, G.; Travers, A. Bromobenzofurans: A New Class of Potent, Non-peptide Antagonists of Angiotensin II. *Bioorg. Med. Chem. Lett.* 1991, 1, 711-716. (d) De, B.; Winn, M.; Zydowsky, T. M.; Kerkman, D. J.; DeBernardis, J. F.; Lee, J.; Buckner, S.; Warner, R.; Brune, M.; Hancock, A.; Oppenorth, T.; Marsh, K. Discovery of a Novel Class of Orally Active, Non-peptide Angiotensin II Antagonists. *J. Med. Chem.* 1992, 35, 3714-3717. (e) Middlemiss, D.; Ross, B. C.; Eldred, C.; Montana, J. G.; Shah, P.; Hirst, G. C.; Watson, S. P.; Panchal, T. A.; Paton, J. M. S.; Hubbard, T.; Drew, G. M.; Robertson, M. J.; Hilditch, A.; Clark, K. L. C-Linked Pyrazole Biaryl Tetrazoles as Antagonists of Angiotensin II. *Bioorg. Med. Chem. Lett.* 1992, 2, 1243-1246.
- (14) Shibuta, Y.; Inada, Y.; Ojima, M.; Wada, T.; Noda, M.; Sanada, T.; Kubo, K.; Kohara, Y.; Naka, T.; Nishikawa, K. Pharmacological Profile of a Highly Potent and Long Acting Angiotensin II Receptor Antagonist, CV-11974 and its Prodrug, TCV-116. *J. Pharmacol. Exp. Ther.*, in print.